



**HAL**  
open science

# Connes-Moscovici characteristic map is a Lie algebra morphism

Luc Menichi

► **To cite this version:**

| Luc Menichi. Connes-Moscovici characteristic map is a Lie algebra morphism. 2010. hal-00454042v2

**HAL Id: hal-00454042**

**<https://hal.science/hal-00454042v2>**

Preprint submitted on 10 Feb 2010 (v2), last revised 17 Jun 2010 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

LUC MENICHI

ABSTRACT. Let  $H$  be a Hopf algebra with a modular pair in involution  $(\delta, 1)$ . Let  $A$  be a (module) algebra over  $H$  equipped with a non-degenerated  $\delta$ -invariant 1-trace  $\tau$ . We show that Connes-Moscovici characteristic map  $\varphi_\tau : HC_{(\delta,1)}^*(H) \rightarrow HC_\lambda^*(A)$  is a morphism of graded Lie algebras. We also have a morphism  $\Phi$  of Batalin-Vilkovisky algebras from the cotorsion product of  $H$ ,  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$ , to the Hochschild cohomology of  $A$ ,  $HH^*(A, A)$ . Let  $K$  be both a Hopf algebra and a symmetric Frobenius algebra. Suppose that the square of its antipode is an inner automorphism by a group-like element. Then this morphism of Batalin-Vilkovisky algebras  $\Phi : \text{Cotor}_{K^\vee}^*(\mathbb{F}, \mathbb{F}) \cong \text{Ext}_K(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(K, K)$  is injective.

## 1. INTRODUCTION

Let  $\mathbb{k}$  be any commutative ring and  $\mathbb{F}$  be any field. It is well known that the Hochschild cohomology of an algebra  $A$ ,  $HH^*(A, A)$ , is a Gerstenhaber algebra. It is also well known that the homology of a double pointed loop space,  $H_*(\Omega^2 X)$ , is also a Gerstenhaber algebra [4]. Let  $H$  be a bialgebra. It is not well known (See [18] for a recent paper rediscovering it) that the Cotorsion product of  $H$ ,  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$  has a Gerstenhaber algebra structure (this results from [12, p. 65]). But it should. Indeed, by Adams cobar equivalence, there is an isomorphism  $\text{Cotor}_{S_*(\Omega X)}^*(\mathbb{k}, \mathbb{k}) \cong H_*(\Omega^2 X)$  between the two Gerstenhaber algebras (See the proof of Corollary 26 for details).

The first goal of this paper is to study (Section 4) this Gerstenhaber algebra  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$ . In particular, generalizing a result of Farinati and Solotar [9], we show (Theorem 16) that the exterior product  $\text{Ext}_H^*(\mathbb{k}, \mathbb{k})$  is a sub Gerstenhaber algebra of the Hochschild cohomology of  $H$ ,  $HH^*(H, H)$ .

---

*Key words and phrases.* Batalin-Vilkovisky algebra, Hochschild cohomology, cyclic cohomology, Hopf algebra, Frobenius algebra.

In Section 5, we turn our attention to a particular case of Gerstenhaber algebras: the Batalin-Vilkovisky algebras. In [31], we introduced the notion of cyclic operad with multiplication (Definition 34) and we show (Theorem 35) that every cyclic operad with multiplication  $\mathcal{O}$  gives a cocyclic module such that

- the homology of the associated cochain complex  $H(\mathcal{C}^*(\mathcal{O}))$  is a Batalin-Vilkovisky algebra and

- the negative cyclic cohomology of  $\mathcal{C}^*(\mathcal{O})$ ,  $HC_-^*(\mathcal{O})$ , has a Lie bracket of degree  $-2$ .

Let  $M$  be a simply-connected closed manifold. In [2], Chas and Sullivan showed that  $\mathbb{H}_*(LM)$ , the free loop space homology of  $M$ , is a Batalin-Vilkovisky algebra and that the  $S^1$ -equivariant homology  $H_*^{S^1}(LM)$  has a Lie bracket. The singular cochains of  $M$ ,  $S^*(M)$  is a (derived) symmetric Frobenius algebra. Motivated by Chas-Sullivan string topology, in [31], as first application of Theorem 35, we obtained that the Hochschild cohomology of a symmetric Frobenius algebra  $A$ ,  $HH^*(A, A)$ , is a Batalin-Vilkovisky algebra and that the negative cyclic cohomology of  $A$ ,  $HC_-^*(A)$  has a Lie bracket of degree  $-2$ . It is expected that there is an isomorphism of Batalin-Vilkovisky algebras  $HH^*(S^*(M), S^*(M)) \cong \mathbb{H}_*(LM)$  and an isomorphism of Lie algebras  $HC_-^*(S^*(M)) \cong H_*^{S^1}(LM)$ .

In [14], Getzler showed that the Gerstenhaber algebra  $H_*(\Omega^2 X)$  is in fact a Batalin-Vilkovisky algebra. Therefore as second application of Theorem 35, in [31], we showed that the Cotorsion product of a Hopf algebra  $H$  with an involutive antipode or more generally with a modular pair in involution  $(\delta, 1)$ ,  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$ , is a Batalin-Vilkovisky algebra. In this paper, we give the dual result (Theorem 49) which we believe is far more clear: Let  $K$  be a Hopf algebra such that the square of its antipode is an inner automorphism by a group-like element. Then  $\text{Ext}_K^*(\mathbb{k}, \mathbb{k})$  is a Batalin-Vilkovisky algebra.

In [31], we also had that the negative cyclic cohomology of  $H$ ,  $HC_{-(\delta, 1)}^*(H)$  has a Lie bracket of degree  $-2$ . But Connes and Moscovici never use negative cyclic cohomology: they use the (ordinary) cyclic cohomology. Therefore, in this paper, we show (Corollary 44) that Connes-Moscovici (ordinary) cyclic cohomology of  $H$ ,  $HC_{(\delta, 1)}^*(H)$ , has also a Lie bracket (of degree  $-1$  this time) and we show (Theorem 53 and its variant Theorem 52) that Connes-Moscovici characteristic map  $\chi_\tau : HC_{(\delta, 1)}^*(H) \rightarrow HC_\lambda^*(A)$  is compatible with the Lie brackets of degree  $-1$ . Here  $A$  is a symmetric Frobenius algebra equipped an action of the Hopf algebra  $H$  compatible with the product and the trace.

In fact, we show that Connes-Moscovici characteristic map is induced by a morphism of cyclic operads with multiplication from the cobar construction of  $H, \Omega H$ , to the Hochschild cochain complex of  $A, \mathcal{C}^*(A, A)$ . And we show that the (ordinary) cyclic cohomology of every cyclic operad with multiplication has naturally a Lie bracket of degree  $-1$  (Theorem 36). As a consequence of Theorem 35, we also obtain a morphism of Batalin-Vilkovisky algebras  $H^*(\Phi) : \text{Cotor}_H^*(\mathbb{k}, \mathbb{k}) \rightarrow HH^*(A, A)$  (Theorem 53).

Note that this morphism  $H^*(\Phi)$  is the algebraic counterpart of our very recent morphism of Batalin-Vilkovisky algebras [32, Theorem 3]

$$\text{Cotor}_{S_*(G)}(\mathbb{k}, \mathbb{k}) \cong H_*(\Omega^2 BG) \rightarrow \mathbb{H}_*(LM) \cong HH^*(S^*(M), S^*(M))$$

between the Batalin-Vilkovisky algebra on the homology of double loop space given by Getzler [14], and the Batalin-Vilkovisky algebra on the free loop space homology of a manifold given by Chas and Sullivan. Here  $G$  is a topological group acting on  $M$ .

In the last section, we specialize to the case where the symmetric Frobenius algebra  $A$  is the Hopf algebra  $H$  itself. And we show that the inclusion of Gerstenhaber algebras  $\text{Ext}_H^*(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(H, H)$ , given by Theorem 16, is often an inclusion of Batalin-Vilkovisky algebras (Theorem 62).

## 2. HOCHSCHILD COMPLEX AND (CO)BAR CONSTRUCTION

We work over an arbitrary commutative ring  $\mathbb{k}$ , except for Conjectures 23 -25 in Section 4, for Proposition 45 to Corollary 48 (almost all Section 6) and for all Section 8, where we use an arbitrary field  $\mathbb{F}$  as coefficient.

Let  $A$  be an algebra and  $M$  be a  $A$ -bimodule. The *Hochschild chain complex*  $\mathcal{C}_*(A, M)$  is the chain complex  $\mathcal{C}_n(A, M) = M \otimes A^n$  with differential  $d : \mathcal{C}_n(A, M) \rightarrow \mathcal{C}_{n-1}(A, M)$  given by

$$\begin{aligned} d(m \otimes a_1 \otimes \cdots \otimes a_n) &= ma_1 \otimes a_2 \otimes \cdots \otimes a_n \\ &+ \sum_{i=1}^{n-1} (-1)^i m \otimes a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n + (-1)^n a_n m \otimes a_1 \otimes \cdots \otimes a_{n-1}. \end{aligned}$$

By definition, the *Hochschild homology of  $A$  with coefficients in  $M$* ,  $HH_*(A, M)$  is the homology of  $\mathcal{C}_*(A, M)$ . The *Hochschild cochain complex*  $\mathcal{C}^*(A, M)$  is the cochain complex  $\mathcal{C}^n(A, M) = \text{Hom}(A^n, M)$  with

differential  $d : \mathcal{C}^n(A, M) \rightarrow \mathcal{C}^{n+1}(A, M)$  given by

$$d(f)(a_0 \otimes \cdots \otimes a_n) = a_0 f(a_1 \otimes \cdots \otimes a_n) + \sum_{i=1}^n (-1)^i f(a_0 \otimes \cdots \otimes a_{i-1} a_i \otimes \cdots \otimes a_n) + (-1)^{n+1} f(a_0 \otimes \cdots \otimes a_{n-1}) a_n.$$

By definition, the *Hochschild cohomology of  $A$  with coefficients in  $M$* ,  $HH^*(A, M)$  is the homology of  $\mathcal{C}^*(A, M)$ . Suppose that  $A$  has an augmentation  $\varepsilon : A \rightarrow \mathbb{k}$ . Then  $\mathbb{k}$  is a  $A$ -bimodule. The (*reduced*) *Bar construction*  $B(A)$  is just then the Hochschild chain complex  $\mathcal{C}_*(A, \mathbb{k})$  and  $\text{Ext}_A^*(\mathbb{k}, \mathbb{k}) = HH^*(A, \mathbb{k})$ .

Dually, let  $C$  be a coalgebra with diagonal  $\Delta_C : C \rightarrow C \otimes C$ . Let  $N$  be a  $C$ -bicomodule with left  $C$ -coaction  $\Delta_N^l : N \rightarrow C \otimes N$  and right  $C$ -coaction  $\Delta_N^r : N \rightarrow N \otimes C$ . The Hochschild cochain complex  $\mathcal{C}_{\text{coalg}}^*(C, N)$  ([12, p. 57] or [1, 30.3]) is the cochain complex  $\mathcal{C}^n(C, N) = \text{Hom}(N, C^n)$  with differential  $d : \mathcal{C}^n(C, N) \rightarrow \mathcal{C}^{n+1}(C, N)$  given by

$$d(\varphi) = (C \otimes \varphi) \circ \Delta_N^l + \sum_{i=1}^n (-1)^i (C^{\otimes i-1} \otimes \Delta_C \otimes C^{\otimes n-i}) \circ \varphi + (-1)^{n+1} (\varphi \otimes C) \circ \Delta_N^r.$$

The *Hochschild coalgebra cohomology*  $HH_{\text{coalg}}^*(C, N)$  is the homology of  $\mathcal{C}_{\text{coalg}}^*(C, N)$ . Suppose that  $C$  has a coaugmentation  $\eta : \mathbb{k} \rightarrow C$ . Then  $\mathbb{k}$  is a  $C$ -bicomodule. The (*reduced*) *cobar construction*  $\Omega(C)$  [20, p. 432] is just  $\mathcal{C}_{\text{coalg}}^*(C, \mathbb{k})$  and  $\text{Cotor}_C^*(\mathbb{k}, \mathbb{k}) = HH_{\text{coalg}}^*(C, \mathbb{k})$ .

### 3. OPERADS WITH MULTIPLICATION

A *Gerstenhaber algebra* is a commutative graded algebra  $A = \{A^i\}_{i \in \mathbb{Z}}$  equipped with a bracket of degree  $-1$

$$\{-, -\} : A^i \otimes A^j \rightarrow A^{i+j-1}, \quad x \otimes y \mapsto \{x, y\}$$

such that the product and the Lie bracket satisfy the Poisson rule: for any  $c \in A^k$  the adjunction map  $\{-, c\} : A^i \rightarrow A^{i+k-1}$ ,  $a \mapsto \{a, c\}$  is a  $(k-1)$ -derivation: ie. for  $a, b, c \in A$ ,  $\{ab, c\} = \{a, c\}b + (-1)^{|a|(|c|-1)} a\{b, c\}$ .

In this paper, every Gerstenhaber algebra comes from a (linear) operad with multiplication using the following general theorem:

**Theorem 1.** [12, 13, 30] *a) Each operad with multiplication  $O$  is a cosimplicial module (See 5). Denote by  $\mathcal{C}^*(O)$  the associated cochain complex.*

*b) Its homology  $H(\mathcal{C}^*(O))$  is a Gerstenhaber algebra.*

Let us first recall what is a (linear) operad and a (linear) operad with multiplication.

In this paper, *operad* means non- $\Sigma$ -operad in the category of  $\mathbb{k}$ -modules. That is: a sequence of modules  $\{O(n)\}_{n \geq 0}$ , an identity element  $id \in O(1)$  and structure maps

$$\begin{aligned} \gamma &: O(n) \otimes O(i_1) \otimes \cdots \otimes O(i_n) \rightarrow O(i_1 + \cdots + i_n) \\ f \otimes g_1 \otimes \cdots \otimes g_n &\mapsto \gamma(f; g_1, \dots, g_n) \end{aligned}$$

satisfying associativity and unit [28].

Hereafter we use mainly the composition operations  $\circ_i : O(m) \otimes O(n) \rightarrow O(m + n - 1)$ ,  $f \otimes g \mapsto f \circ_i g$  defined for  $m \in \mathbb{N}^*$ ,  $n \in \mathbb{N}$  and  $1 \leq i \leq m$  by  $f \circ_i g := \gamma(f; id, \dots, g, id, \dots, id)$  where  $g$  is the  $i$ -th element after the semicolon.

*Example 2.* Let  $(\mathcal{C}, \otimes, \mathbb{k})$  be a monoidal category. Suppose that  $\mathcal{C}$  is enriched over the category of  $\mathbb{k}$ -modules [27, I.8] and that

$$\otimes : \text{Hom}_{\mathcal{C}}(V_1, W_1) \times \text{Hom}_{\mathcal{C}}(V_2, W_2) \rightarrow \text{Hom}_{\mathcal{C}}(V_1 \otimes V_2, W_1 \otimes W_2),$$

mapping  $(g_1, g_2)$  to  $g_1 \otimes g_2$ , is  $\mathbb{k}$ -bilinear (we say that  $\mathcal{C}$  is a  $\mathbb{k}$ -linear monoidal category). Let  $V$  be a object of  $\mathcal{C}$ . The *endomorphism operad* of  $V$  in  $\mathcal{C}$  [28, p. 43] is the operad  $\mathcal{E}nd_{\mathcal{C}}(V)$  defined by

$$\mathcal{E}nd_{\mathcal{C}}(V)(n) := \text{Hom}_{\mathcal{C}}(V^{\otimes n}, V).$$

The structure maps  $\gamma$

$$\text{Hom}_{\mathcal{C}}(V^{\otimes n}, V) \otimes \text{Hom}_{\mathcal{C}}(V^{\otimes i_1}, V) \otimes \cdots \otimes \text{Hom}_{\mathcal{C}}(V^{\otimes i_n}, V) \rightarrow \text{Hom}_{\mathcal{C}}(V^{\otimes i_1 + \cdots + i_n}, V)$$

are given by  $\gamma(f; g_1, \dots, g_n) = f \circ (g_1 \otimes \cdots \otimes g_n)$ . The identity element of  $\mathcal{E}nd_{\mathcal{C}}(V)$  is the identity map  $id_V : V \rightarrow V$ .

*Example 3.* The *coendomorphism operad* of  $V$  in  $\mathcal{C}$ , denoted  $\text{Co}\mathcal{E}nd_{\mathcal{C}}(V)$ , is by definition the endomorphism operad of  $V$  in the opposite category  $\mathcal{C}^{op}$ ,  $\mathcal{E}nd_{\mathcal{C}^{op}}(V)$ . Explicitly [28, p. 43-4]  $\text{Co}\mathcal{E}nd_{\mathcal{C}}(V)$  is the operad given by

$$\text{Co}\mathcal{E}nd_{\mathcal{C}}(V)(n) := \text{Hom}_{\mathcal{C}}(V, V^{\otimes n}).$$

The structure maps  $\gamma$

$$\text{Hom}_{\mathcal{C}}(V, V^{\otimes n}) \otimes \text{Hom}_{\mathcal{C}}(V, V^{\otimes i_1}) \otimes \cdots \otimes \text{Hom}_{\mathcal{C}}(V, V^{\otimes i_n}) \rightarrow \text{Hom}_{\mathcal{C}}(V, V^{\otimes i_1 + \cdots + i_n})$$

are given by  $\gamma(f; g_1, \dots, g_n) = (g_1 \otimes \cdots \otimes g_n) \circ f$ . The identity element of  $\mathcal{E}nd_{\mathcal{C}}(V)$  is again  $id$ .

**Definition 4.** An *operad with multiplication* is an operad equipped with an element  $\mu \in O(2)$  called the multiplication and an element  $e \in O(0)$  such that  $\mu \circ_1 \mu = \mu \circ_2 \mu$  and  $\mu \circ_1 e = id = \mu \circ_2 e$ .

Let  $\underline{Ass}$  be the (non- $\Sigma$ ) associative operad [28]:  $\underline{Ass}(n) := \mathbb{k}$ . An operad  $O$  is an operad with multiplication if and only if  $O$  is equipped with a morphism of operads  $\underline{Ass} \rightarrow O$ .

*Sketch of proof of 1.* a) The coface maps  $\delta_i : O(n) \rightarrow O(n+1)$  and codegeneracy maps  $\sigma_i : O(n) \rightarrow O(n-1)$  are defined [30] by

$$(5) \quad \delta_0 f = \mu \circ_2 f, \quad \delta_i f = f \circ_i \mu, \quad \delta_{n+1} f = \mu \circ_1 f, \quad \sigma_{i-1} f = f \circ_i e \text{ for } 1 \leq i \leq n.$$

b) The associated cochain complex  $\mathcal{C}^*(O)$  is the cochain complex whose differential  $d$  is given by

$$d := \sum_{i=0}^{n+1} (-1)^i \delta_i : O(n) \rightarrow O(n+1).$$

The linear maps  $\cup : O(m) \otimes O(n) \rightarrow O(m+n)$  defined by

$$(6) \quad f \cup g := (\mu \circ_1 f) \circ_{m+1} g = (\mu \circ_2 g) \circ_1 f$$

gives  $\mathcal{C}^*(O)$  a structure of differential graded algebra. The linear maps of degree  $-1$

$$\bar{\circ}, \{-, -\} : O(m) \otimes O(n) \rightarrow O(m+n-1)$$

are defined by

$$(7) \quad f \bar{\circ} g := (-1)^{(m-1)(n-1)} \sum_{i=1}^m (-1)^{(n-1)(i-1)} f \circ_i g$$

and

$$\{f, g\} := f \bar{\circ} g - (-1)^{(m-1)(n-1)} g \bar{\circ} f.$$

The bracket  $\{-, -\}$  defines a structure of differential graded Lie algebra of degree  $-1$  on  $\mathcal{C}^*(O)$ . After passing to cohomology, the cup product  $\cup$  and the bracket  $\{-, -\}$  satisfy the Poisson rule.  $\square$

*Remark 8.* As pointed by Turchin in [43], the Gerstenhaber algebra  $H(\mathcal{C}^*(\mathcal{O}))$  has Dyer-Lashof operations. In particular [12, p. 63], if  $n$  is even or if  $2 = 0$  in  $\mathbb{k}$ , a Steenrod or Dyer-lashof (non additive) operation  $Sq^{n-1} : H^n(\mathcal{C}^*(\mathcal{O})) \rightarrow H^{2n-1}(\mathcal{C}^*(\mathcal{O}))$  is defined by  $Sq^{n-1}(f) = f \bar{\circ} f$  for  $f \in \mathcal{O}(n)$ .

*Remark 9.* Let  $\mathcal{O}$  be an operad. Then  $\mathcal{O}(1)$  equipped with  $\circ_1 : \mathcal{O}(1) \otimes \mathcal{O}(1) \rightarrow \mathcal{O}(1)$  and  $id : \mathbb{k} \rightarrow \mathcal{O}(1)$  is an algebra. By (7), the Lie algebra  $\mathcal{C}^1(O)$  is just  $\mathcal{O}(1)$  equipped with the Lie bracket given by  $\{f, g\} := f \circ_1 g - g \circ_1 f$ .

CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

*Example 10.* Let  $A$  be a monoid in  $\mathcal{C}$ , i. e. an object of  $\mathcal{C}$  equipped with an associative multiplication  $\mu : A \otimes A \rightarrow A$  and an unit  $e : \mathbb{k} \rightarrow A$ . Then the endomorphism operad  $\mathcal{E}nd_{\mathcal{C}}(A)$  of  $A$  equipped with  $\mu \in \text{Hom}_{\mathcal{C}}(A^{\otimes 2}, A) = \mathcal{E}nd_{\mathcal{C}}(A)(2)$  and  $e \in \text{Hom}_{\mathcal{C}}(A^{\otimes 0}, A) = \mathcal{E}nd_{\mathcal{C}}(A)(0)$  is an operad with multiplication. The associated cosimplicial module is the cosimplicial module  $\{\text{Hom}_{\mathcal{C}}(A^{\otimes n}, A)\}_{n \in \mathbb{N}}$ . The coface maps  $\delta_i : \text{Hom}_{\mathcal{C}}(A^{\otimes n}, A) \rightarrow \text{Hom}_{\mathcal{C}}(A^{\otimes n+1}, A)$  and the codegeneracy map  $\sigma_i : \text{Hom}_{\mathcal{C}}(A^{\otimes n}, A) \rightarrow \text{Hom}_{\mathcal{C}}(A^{\otimes n-1}, A)$  are given by [31, (2.5)]

(11)

$$\delta_0 f = \mu \circ (id \otimes f), \quad \delta_i f = f \circ (id^{\otimes i-1} \otimes \mu \otimes id^{\otimes n-i}), \quad \delta_{n+1} f = \mu \circ (f \otimes id),$$

and  $\sigma_{i-1} f = f \circ (id^{\otimes i-1} \otimes e \otimes id^{\otimes n-i})$  for  $1 \leq i \leq n$ .

If  $\mathcal{C}$  is the category of  $\mathbb{k}$ -modules,  $A$  is an algebra and the cochain complex  $\mathcal{C}^*(\mathcal{E}nd_{\mathcal{C}}(A))$  associated to this cosimplicial module is the Hochschild cochain complex of  $A$ , denoted  $\mathcal{C}^*(A, A)$ . This is why Turchin in his work on knots [41, 42] always call the cochain complex associated to a linear operad with multiplication, the Hochschild cochain complex of the operad with multiplication.

*Property 12.* Let  $\mathcal{C}$  and  $\mathcal{D}$  be two  $\mathbb{k}$ -linear monoidal categories. Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a monoidal functor (in the sense of [27, p. 255]). Let  $\psi : F(V) \otimes F(W) \rightarrow F(V \otimes W)$  be the associated associative unital natural transformation. Suppose that  $F : \text{Hom}_{\mathcal{C}}(V, W) \rightarrow \text{Hom}_{\mathcal{D}}(F(V), F(W))$  is  $\mathbb{k}$ -linear. Let  $A$  be a monoid in  $\mathcal{C}$ . Then  $F(A)$  is a monoid in  $\mathcal{D}$  and the map  $\Gamma$  from  $\mathcal{E}nd_{\mathcal{C}}(A)$  to  $\mathcal{E}nd_{\mathcal{D}}(F(A))$ , mapping  $f : A^{\otimes n} \rightarrow A$  to the composite  $F(f) \circ \psi : F(A)^{\otimes n} \rightarrow F(A)$ , is a morphism of operads with multiplication.

*Example 13.* Dually, let  $C$  be a comonoid in  $\mathcal{C}$ , i. e. an object of  $\mathcal{C}$  equipped with a coassociative diagonal  $\Delta : C \rightarrow C \otimes C$  and a counit  $\varepsilon : C \rightarrow \mathbb{k}$ . Since  $C$  is a monoid in  $\mathcal{C}^{op}$ , the coendomorphism operad of  $C$ ,  $\text{Co}\mathcal{E}nd_{\mathcal{C}}(C)$  equipped with  $\Delta \in \text{Hom}_{\mathcal{C}}(C, C^{\otimes 2}) = \text{Co}\mathcal{E}nd_{\mathcal{C}}(C)(2)$  and  $\varepsilon \in \text{Hom}_{\mathcal{C}}(C, C^{\otimes 0}) = \text{Co}\mathcal{E}nd_{\mathcal{C}}(C)(0)$  is also an operad with multiplication. The associated cosimplicial module is the cosimplicial module  $\{\text{Hom}_{\mathcal{C}}(C, C^{\otimes n})\}_{n \in \mathbb{N}}$ . The coface maps  $\delta_i : \text{Hom}_{\mathcal{C}}(C, C^{\otimes n}) \rightarrow \text{Hom}_{\mathcal{C}}(C, C^{\otimes n+1})$  and the codegeneracy map  $\sigma_i : \text{Hom}_{\mathcal{C}}(C, C^{\otimes n}) \rightarrow \text{Hom}_{\mathcal{C}}(C, C^{\otimes n-1})$  are given by

(14)

$$\delta_0 f = (id \otimes f) \circ \Delta, \quad \delta_i f = (id^{\otimes i-1} \otimes \Delta \otimes id^{\otimes n-i}) \circ f, \quad \delta_{n+1} f = (f \otimes id) \circ \Delta,$$

and  $\sigma_{i-1} f = (id^{\otimes i-1} \otimes \varepsilon \otimes id^{\otimes n-i}) \circ f$  for  $1 \leq i \leq n$ .

If  $\mathcal{C}$  is the category of  $\mathbb{k}$ -modules,  $C$  is a coalgebra and the cochain complex  $\mathcal{C}^*(\text{Co}\mathcal{E}nd_{\mathcal{C}}(C))$  associated to this cosimplicial module is the Hochschild cochain complex of the coalgebra  $C$ , denoted  $\mathcal{C}_{coalg}^*(C, C)$ .

More generally, let  $A$  be  $\mathbb{k}$ -algebra. Let  $\mathcal{C}$  be the category of  $A$ -bimodules. Let  $C$  be a  $A$ -coring, i. e. a comonoid in  $\mathcal{C}$  ([19, 4.2] or [1, 17.1]). The cochain complex  $\mathcal{C}^*(\text{CoEnd}_{\mathcal{C}}(C))$  associated to this cosimplicial module is the Cartier cochain complex of  $C$  with coefficients in  $C$ , denoted  $C_{Ca}(C, C)$ . Therefore, without any calculations, we have obtained that  $C_{Ca}(C, C)$  is an operad with multiplication [1, 30.8]. This is again an example of our leitmotiv in this paper:

“Every operad with multiplication should be the endomorphism operad of a monoid in a appropriate monoidal category  $\mathcal{C}$ ”.

#### 4. GERSTENHABER ALGEBRA STRUCTURE ON $\text{EXT}_A^*(\mathbb{k}, \mathbb{k})$

Let  $C$  be a bialgebra. The Cobar construction of  $C$  is the cosimplicial module associated to a specific linear operad with multiplication [12, p. 65]. Therefore its cohomology  $\text{Cotor}_C^*(\mathbb{k}, \mathbb{k})$  has a Gerstenhaber algebra structure. In the following, we show that this operad with multiplication is just the endomorphism operad of a monoid in an appropriate monoidal category and we show:

**Theorem 15.** *Let  $C$  be a bialgebra. Then  $\text{Cotor}_C^*(\mathbb{k}, \mathbb{k})$  is a sub Gerstenhaber algebra of the Hochschild cohomology of the coalgebra  $C$ ,  $HH_{\text{coalg}}^*(C, C)$ .*

By Property 17, this Lie bracket of degree  $-1$  on the cotorsion product of a bialgebra is an extension of the well-known Lie bracket on the primitive elements of a bialgebra. Dually, we prove

**Theorem 16.** *Let  $A$  be a bialgebra. Then  $\text{Ext}_A^*(\mathbb{k}, \mathbb{k})$  is a sub Gerstenhaber algebra of the Hochschild cohomology of the algebra  $A$ ,  $HH^*(A, A)$ .*

When  $A$  is a Hopf algebra, this theorem was proved by Farinati and Solotar [9]. But as we would like to emphasize, antipodes are not needed for the Gerstenhaber algebra structure. As we explain in Theorem 49, antipodes are needed only to have a Batalin-Vilkovisky algebra structure.

By Property 20, this inclusion of Gerstenhaber algebras is in degree 1 the inclusion of the Lie algebra of “differentiations” into the Lie algebra of derivations, well known in algebraic groups.

In Proposition 22, we prove that when the bialgebra  $C$  is  $\mathbb{k}$ -free of finite type, Theorem 16 is the dual of Theorem 15. This duality will be later extended in Corollary 48.

In Conjectures 23 and 25, we explain that if the bialgebra  $A$  is braided, the Lie bracket of degree  $-1$  given by Theorem 16 on  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  should vanish and be replaced by a Lie bracket of degree  $-2$ . This is related to a conjecture of Kontsevich.

In Corollary 26, we explain that the homology of a double loop space  $H_*(\Omega^2 X)$  is always a sub Gerstenhaber algebra of Hochschild cohomology if  $X$  is 2-connected.

*Proof of Theorem 15.* The category of left  $C$ -modules,  $C\text{-mod}$ , is a monoidal category. Let  $M$  be a comonoid in this monoidal category, i. e.  $M$  is a  $C$ -module coalgebra [20, Definition IX.2.1]. The coendomorphism operad associated to  $M$  is the operad  $\{\text{Hom}_{C\text{-mod}}(M, M^{\otimes n})\}_{n \in \mathbb{N}}$  with multiplication  $\Delta : M \rightarrow M \otimes M \in \text{Hom}_{C\text{-mod}}(M, M^{\otimes 2})$  and  $\varepsilon : M \rightarrow \mathbb{k} \in \text{Hom}_{C\text{-mod}}(M, M^{\otimes 0})$ . The inclusion maps  $i_C : \text{Hom}_{C\text{-mod}}(M, M^{\otimes n}) \hookrightarrow \text{Hom}_{\mathbb{k}\text{-mod}}(M, M^{\otimes n})$  defines obviously a morphism of linear operads with multiplication.

The underlying coalgebra  $C$  is an example of  $C$ -module coalgebra. Therefore we can take in particular  $M = C$ . The linear morphism  $ev : \text{Hom}_{C\text{-mod}}(C, C^{\otimes n}) \xrightarrow{\cong} C^{\otimes n}$ , mapping  $f : C \rightarrow C^{\otimes n}$  to  $f(1)$  is an isomorphism. The inverse is the linear map  $ext : C^{\otimes n} \xrightarrow{\cong} \text{Hom}_{C\text{-mod}}(C, C^{\otimes n})$ , mapping  $c_1 \otimes \cdots \otimes c_n$  to  $f : C \rightarrow C^{\otimes n}$  defined by  $f(c) = c^{(1)}c_1 \otimes \cdots \otimes c^{(n)}c_n$ . Here we have denoted by  $c^{(1)} \otimes \cdots \otimes c^{(n)}$  the iterated diagonal of  $c$ ,  $\Delta^{n-1}(c)$ . Consider the associated cosimplicial set  $\{\text{Hom}_{C\text{-mod}}(C, C^{\otimes n})\}_{n \in \mathbb{N}}$ . The coface maps  $\delta_i$  and codegeneracy maps  $\sigma_i$  are given by equations (14). Therefore for  $1 \leq i \leq n$ ,

$$\begin{aligned} ev \circ \delta_0 \circ ext(c_1 \otimes \cdots \otimes c_n) &= 1 \otimes c_1 \otimes \cdots \otimes c_n, \\ ev \circ \delta_i \circ ext(c_1 \otimes \cdots \otimes c_n) &= c_1 \otimes \cdots \otimes \Delta(c_i) \otimes \cdots \otimes c_n, \\ ev \circ \delta_{n+1} \circ ext(c_1 \otimes \cdots \otimes c_n) &= c_1 \otimes \cdots \otimes c_n \otimes 1 \text{ and} \\ ev \circ \sigma_{i-1} \circ ext(c_1 \otimes \cdots \otimes c_n) &= c_1 \otimes \cdots \otimes \varepsilon(c_i) \otimes \cdots \otimes c_n. \end{aligned}$$

So  $ext : C^{\otimes n} \xrightarrow{\cong} \text{Hom}_{C\text{-mod}}(C, C^{\otimes n})$  is an isomorphism of cosimplicial modules between the Cobar construction of  $C$ ,  $\Omega C$ , and the cosimplicial module associated to the operad with multiplication  $\mathcal{CoEnd}_{C\text{-mod}}(C)$ . Therefore  $\text{Cotor}_C^*(\mathbb{k}, \mathbb{k}) := H^*(\Omega C)$  is a Gerstenhaber algebra. The composite

$$C^{\otimes n} \xrightarrow{ext} \text{Hom}_{C\text{-mod}}(C, C^{\otimes n}) \subset \text{Hom}_{\mathbb{k}\text{-mod}}(C, C^{\otimes n})$$

admits the morphism of differential graded algebras

$$\mathcal{C}^*(C, \eta) : \mathcal{C}_{\text{coalg}}^*(C, C) \rightarrow \mathcal{C}_{\text{coalg}}^*(C, \mathbb{k}) = \Omega C$$

mapping  $f : C \rightarrow C^{\otimes n}$  to  $f(1)$  as retract. Passing to cohomology, we obtain an injective morphism of Gerstenhaber algebras

$$\text{Cotor}_C^*(\mathbb{k}, \mathbb{k}) \hookrightarrow HH_{\text{coalg}}^*(C, C)$$

which admits the morphism of graded algebras

$$HH^*(C, \eta) : HH_{\text{coalg}}^*(C, C) \rightarrow \text{Cotor}_C^*(\mathbb{k}, \mathbb{k})$$

as retract.  $\square$

*Property 17.* The Lie algebra structure on  $\text{Cotor}_C^1(\mathbb{k}, \mathbb{k})$  given by Theorem 15 coincides with the Lie algebra of primitive elements  $P(C)$  of the bialgebra  $C$ .

*Proof.* Consider the isomorphisms  $ext$  and  $ev$  given in the proof of Theorem 15. We have:

$$(18) \quad ev \circ \circ_i \circ (ext \otimes ext)(a_1 \otimes \cdots \otimes a_m \otimes b_1 \otimes \cdots \otimes b_n) = \\ a_1 \otimes \cdots \otimes a_{i-1} \otimes a_i^{(1)} b_1 \otimes \cdots \otimes a_i^{(n)} b_n \otimes a_{i+1} \otimes \cdots \otimes a_m.$$

$ev(id_C) = 1_C \in C$ ,  $ev(\varepsilon) = 1_{\mathbb{k}} \in \mathbb{k}$  and  $ev(\Delta) = 1_C \otimes 1_C \in C \otimes C$ . Therefore  $ev : \text{Hom}_{C\text{-mod}}(C, C^{\otimes n}) \xrightarrow{\cong} C^{\otimes n}$  is an isomorphism of linear operads with multiplication between  $\mathcal{CoEnd}_{C\text{-mod}}(C)$  and the operad with multiplication  $\mathcal{O}$  of [31, Proof of Corollary 2.9], first considered by Gerstenhaber and Schack [12, p. 65] (See also [30, Example 3.5]). In particular  $\circ_1 : \mathcal{O}(1) \otimes \mathcal{O}(1) \rightarrow \mathcal{O}(1)$  is the multiplication of  $C$ ,  $\mu : C \otimes C \rightarrow C$ . Therefore, by (9), the Lie algebra  $\text{Cotor}_C^1(\mathbb{k}, \mathbb{k})$  coincides with the Lie algebra of primitive elements of  $C$ , denoted  $P(C)$ .  $\square$

In order to check that the Gerstenhaber algebra structure given by Theorem 16 coincides with the Gerstenhaber algebra structure on  $\text{Ext}_A^*(\mathbb{k}, \mathbb{k})$  given by Farinati and Solotar [9], we give the proof of Theorem 16.

*Property 19.* Let  $C$  be a coalgebra. Let  $\varepsilon : C \rightarrow \mathbb{k}$  be its counit. Let  $N$  be a left  $C$ -comodule. Then the linear morphism

$$proj : \text{Hom}_{C\text{-comod}}(N, C) \xrightarrow{\cong} N^\vee, \quad F \mapsto \varepsilon \circ F,$$

is an isomorphism. Its inverse is the linear map  $lift : N^\vee \xrightarrow{\cong} \text{Hom}_{C\text{-comod}}(N, C)$  mapping  $f : N \rightarrow \mathbb{k}$  to the composite  $N \xrightarrow{\Delta_N} C \otimes N \xrightarrow{C \otimes f} C \otimes \mathbb{k} = C$ .

*Proof of Theorem 16.* The category of left  $A$ -comodules,  $A\text{-comod}$ , is a monoidal category. Let  $M$  be a monoid in this monoidal category, i. e.  $M$  is a  $A$ -comodule algebra [20, Definition III.7.1]. The endomorphism operad associated to  $M$  is the operad  $\{\text{Hom}_{A\text{-comod}}(M^{\otimes n}, M)\}_{n \in \mathbb{N}}$  with multiplication  $\mu : M \otimes M \rightarrow M \in \text{Hom}_{A\text{-comod}}(M^{\otimes 2}, M)$  and  $\eta : \mathbb{k} \rightarrow M \in \text{Hom}_{A\text{-comod}}(M^{\otimes 0}, M)$ . The coaction of  $A$  on  $M^{\otimes n}$ ,  $\Delta_{M^{\otimes n}}$ , is the composite  $M^{\otimes n} \xrightarrow{\Delta_M^{\otimes n}} (A \otimes M)^{\otimes n} \xrightarrow{\tau} A^{\otimes n} \otimes M^{\otimes n} \xrightarrow{\mu_A \otimes M^{\otimes n}} A \otimes M^{\otimes n}$  where  $\tau$  is the exchange isomorphism. Explicitly  $\Delta_{M^{\otimes n}}(a_1 \otimes \cdots \otimes a_n) = a_1^{(1)} \cdots a_n^{(1)} \otimes (a_1^{(2)} \otimes \cdots \otimes a_n^{(2)})$  where  $\Delta_M a_i = a_i^{(1)} \otimes a_i^{(2)}$ .

CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

We now take  $M = A$ . Using Property 19 with  $C = A$  and  $N = A^{\otimes n}$ , we obtain that

$$proj : \text{Hom}_{A\text{-comod}}(A^{\otimes n}, A) \xrightarrow{\cong} (A^{\otimes n})^\vee, \quad F \mapsto \varepsilon \circ F,$$

is an isomorphism. Its inverse is the linear map  $lift : (A^{\otimes n})^\vee \xrightarrow{\cong} \text{Hom}_{A\text{-comod}}(A^{\otimes n}, A)$  mapping  $f : A^{\otimes n} \rightarrow \mathbb{k}$  to  $F : A^{\otimes n} \rightarrow A$  defined by  $F(a_1 \otimes \cdots \otimes a_n) = a_1^{(1)} \cdots a_n^{(1)} f(a_1^{(2)} \otimes \cdots \otimes a_n^{(2)})$ .

Therefore the composite

$$(A^{\otimes n})^\vee \xrightarrow{lift} \text{Hom}_{A\text{-comod}}(A^{\otimes n}, A) \subset \text{Hom}_{\mathbb{k}\text{-mod}}(A^{\otimes n}, A)$$

coincides with the section of  $\mathcal{C}^*(A, \varepsilon) : \mathcal{C}^*(A, A) \rightarrow BA^\vee$  defined by Farinati and Solotar [9, p. 2862].

Consider the associated cosimplicial set  $\{\text{Hom}_{A\text{-comod}}(A^{\otimes n}, A)\}_{n \in \mathbb{N}}$ . The coface maps  $\delta_i$  and codegeneracy maps  $\sigma_i$  are given by equations (11). Therefore for  $1 \leq i \leq n$ ,

$$\begin{aligned} proj \circ \delta_0 \circ lift(f)(a_1 \otimes \cdots \otimes a_{n+1}) &= \varepsilon(a_1) f(a_2 \otimes \cdots \otimes a_{n+1}), \\ proj \circ \delta_i \circ lift(f)(a_1 \otimes \cdots \otimes a_{n+1}) &= f(a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_{n+1}), \\ proj \circ \delta_{n+1} \circ lift(f)(a_1 \otimes \cdots \otimes a_{n+1}) &= f(a_1 \otimes \cdots \otimes a_n) \varepsilon(a_{n+1}) \text{ and} \\ proj \circ \sigma_{i-1} \circ lift(f)(a_1 \otimes \cdots \otimes a_{n-1}) &= f(a_1 \otimes \cdots \otimes a_{i-1} \otimes 1_A \otimes a_i \otimes \cdots \otimes a_n). \end{aligned}$$

So  $lift : (A^{\otimes n})^\vee \xrightarrow{\cong} \text{Hom}_{A\text{-comod}}(A^{\otimes n}, A)$  is an isomorphism of cosimplicial modules between the dual of the bar construction of  $A$ ,  $BA^\vee$ , and the cosimplicial module associated to the operad with multiplication  $\mathcal{E}nd_{A\text{-comod}}(A)$ . Therefore  $\text{Ext}_A^*(\mathbb{k}, \mathbb{k}) := H^*(BA^\vee)$  is a Gerstenhaber algebra.  $\square$

Let  $A$  be an algebra and  $M$  be a  $A$ -bimodule. The cocycles of degree 1 of the Hochschild complex  $\mathcal{C}^*(A, M)$  are exactly the module of derivations  $\text{Der}(A, M)$ . A linear map  $f : A \rightarrow M$  is a *derivation* if and only if  $\forall a, b \in A, f(ab) = f(a)b + af(b)$ . The boundaries of degree 1 of  $\mathcal{C}^*(A, M)$  are the inner derivations, i. e. the linear maps  $f : A \rightarrow M, a \mapsto am - ma$ , where  $m$  is a given element of  $M$ . The degree 1 component of Hochschild cohomology,  $HH^1(A, M)$ , can be identified with the quotient  $\text{Der}(A, M)/\{\text{inner derivations}\}$  [25, 1.5.2]. In particular, suppose that  $A$  has an augmentation  $\varepsilon : A \rightarrow \mathbb{k}$ . Then  $\text{Ext}_A^1(\mathbb{k}, \mathbb{k}) = HH^1(A, \mathbb{k}) = \text{Der}(A, \mathbb{k})$ .

*Property 20.* Let  $A$  be a bialgebra. The inclusion of Lie algebra  $\text{Ext}_A^1(\mathbb{k}, \mathbb{k}) \hookrightarrow HH^1(A, A)$  given by Theorem 16 can be identified with the following composite of Lie algebra morphisms

$$\text{Der}(A, \mathbb{k}) \xrightarrow{i} \text{Der}(A, A) \xrightarrow{q} \text{Der}(A, A)/\{\text{innerderivations}\}.$$

Here  $q$  is the obvious quotient map and  $i$  is the inclusion of the Lie algebra of “differentiations” of  $A$  into the Lie algebra of derivations of  $A$  given by [16, p. 36].

Let  $G$  be an affine algebraic group. Then the *algebra of polynomial functions* on  $G$ ,  $\mathcal{P}(G)$ , is a commutative Hopf algebra. By definition [16, p. 36], the Lie algebra of  $G$  is  $\text{Ext}_{\mathcal{P}(G)}^1(\mathbb{k}, \mathbb{k}) = \text{Der}(\mathcal{P}(G), \mathbb{k})$ .

Let  $G$  be a Lie group. The algebra of smooth maps on  $G$ ,  $C^\infty(G)$ , is a module algebra over the group ring  $\mathbb{R}[G]$ , but is not a bialgebra (except when  $G$  is finite and discrete). However there is still an analogue of the inclusion  $i$ : the composite

$$T_e(G) \xrightarrow[\text{lift}]{\cong} \text{Hom}_{\text{mod-}\mathbb{R}[G]}(C^\infty(G), C^\infty(G)) \cap \text{Der}(C^\infty(G)) \subset \text{Der}(C^\infty(G)).$$

Here *lift* is the isomorphism between the tangent space and the right invariant vector fields on  $G$ .

*Proof.* Consider the inverse isomorphisms  $\text{proj} : \text{Hom}_{A\text{-comod}}(A^n, A) \xrightarrow{\cong} (A^{\otimes n})^\vee$  and  $\text{lift} : (A^{\otimes n})^\vee \xrightarrow{\cong} \text{Hom}_{A\text{-comod}}(A^n, A)$  given in the proof of Theorem 16. Let  $\mathcal{O}$  denote the linear operad with multiplication such that  $\text{proj} : \mathcal{E}nd_{A\text{-comod}}(A) \xrightarrow{\cong} \mathcal{O}$  is an isomorphism of linear operads with multiplication. Explicitly, for  $f \in \mathcal{O}(m) = (A^{\otimes m})^\vee$  and  $g \in \mathcal{O}(n) = (A^{\otimes n})^\vee$ ,  $f \circ_i g$  is given by

$$f \circ_i g(a_1 \otimes \cdots \otimes a_{m+n-1}) = f(a_1 \otimes \cdots \otimes a_{i-1} \otimes a_i^{(1)} \cdots a_{i+n-1}^{(1)} g(a_i^{(2)} \otimes \cdots \otimes a_{i+n-1}^{(2)}) \otimes a_{i+n} \otimes \cdots \otimes a_{m+n-1})$$

where  $\Delta a_j = a_j^{(1)} \otimes a_j^{(2)}$ . The identity element of  $\mathcal{O}$  is the counit of  $A$ ,  $\varepsilon \in A^\vee = \mathcal{O}(1)$ . The multiplication of  $\mathcal{O}$  is the composite  $\varepsilon \circ \mu \in (A \otimes A)^\vee = \mathcal{O}(2)$  and the unit is  $\text{id}_{\mathbb{k}} \in (A^{\otimes 0})^\vee = \mathcal{O}(0)$ . In particular,  $\circ_1 : \mathcal{O}(1) \otimes \mathcal{O}(1) \rightarrow \mathcal{O}(1)$  is the multiplication of  $A^\vee$ ,  $\mu_{A^\vee} : A^\vee \otimes A^\vee \rightarrow A^\vee$  obtained by dualizing the diagonal of  $A$ . Therefore, by (9), the Lie algebra  $\mathcal{C}^1(\mathcal{O})$  is just the Lie algebra associated to the associative algebra  $A^\vee$ . The composite  $\mathcal{O} \xrightarrow[\cong]{\text{lift}} \mathcal{E}nd_{A\text{-comod}}(A) \subset \mathcal{E}nd_{\mathbb{k}\text{-mod}}(A)$  is an injective morphism of linear operads with multiplication. Therefore this composite  $\mathcal{C}^*(\mathcal{O}) \xrightarrow[\cong]{\text{lift}} \mathcal{C}^*(\mathcal{E}nd_{A\text{-comod}}(A)) \subset \mathcal{C}^*(\mathcal{E}nd_{\mathbb{k}\text{-mod}}(A))$  is an injective morphism of differential graded Lie algebras. In degree 1, this composite  $\mathcal{O}(1) \hookrightarrow \mathcal{E}nd_{\mathbb{k}\text{-mod}}(A)(1) = \text{Hom}_{\mathbb{k}\text{-mod}}(A, A)$  is the injective morphism of (associative) algebras, mapping  $f : A \rightarrow \mathbb{k}$  to  $(A \otimes f) \circ \Delta_A$ , given by [16, I.Proposition 2.1]. Restricted at the cycles in degree 1, this composite gives the injective morphism of Lie algebras  $\text{Der}(A, \mathbb{k}) \xrightarrow{i} \text{Der}(A, A)$  considered in [16, p. 36].  $\square$

Let us prove that Theorem 16 is the dual of Theorem 15.

**Lemma 21.** *Let  $C$  be a coalgebra with coaugmentation  $\eta : \mathbb{k} \rightarrow C$ . Let  $A = C^\vee$  be the dual algebra with augmentation  $\varepsilon : A \rightarrow \mathbb{k}$ . Then*

*i) the linear map  $\Gamma : \text{CoEnd}_{\mathbb{k}\text{-mod}}(C) \rightarrow \text{End}_{\mathbb{k}\text{-mod}}(A)$ , mapping  $f : C \rightarrow C^{\otimes n}$  to the composite  $A^{\otimes n} \rightarrow (C^{\otimes n})^\vee \xrightarrow{f^\vee} A$ , is a morphism of linear operads with multiplication,*

*ii) the linear map  $\phi : \Omega C \rightarrow (BA)^\vee$ , such that  $\phi(c_1 \otimes \cdots \otimes c_n)$  is the form on  $A^{\otimes n}$ , mapping  $\varphi_1 \otimes \cdots \otimes \varphi_n$  to the product  $\varphi_1(c_1) \cdots \varphi_n(c_n)$ , is a morphism of differential graded algebras.*

*iii) We have the commutative diagram of differential graded algebras*

$$\begin{array}{ccc} \mathcal{C}_{\text{coalg}}^*(C, C) & \xrightarrow{\mathcal{C}_{\text{coalg}}^*(C, \eta)} & \Omega C \\ \Gamma \downarrow & & \downarrow \phi \\ \mathcal{C}^*(A, A) & \xrightarrow{\mathcal{C}^*(A, \varepsilon)} & (BA)^\vee \end{array}$$

*If  $C$  is  $\mathbb{k}$ -free of finite type then both  $\Gamma$  and  $\phi$  are isomorphisms.*

*Proof.* Let  $\psi : V^\vee \otimes W^\vee \rightarrow (V \otimes W)^\vee$  be the linear map, mapping the tensor product  $\varphi_1 \otimes \varphi_2$  of a form on  $V$  and of a form on  $W$ , to the form on  $V \otimes W$ , also denoted  $\varphi_1 \otimes \varphi_2$ , mapping  $v \otimes w$  to the product  $\varphi_1(v)\varphi_2(w)$ . The functor  ${}^\vee$  from the opposite category of  $\mathbb{k}$ -modules to the category of  $\mathbb{k}$ -modules, mapping a  $\mathbb{k}$ -module  $V$ , to its dual  $V^\vee := \text{Hom}(V, \mathbb{k})$  is a monoidal functor. Therefore by applying Property 12, we obtain i). ii) is well-known and iii) is easy to check.  $\square$

Note that in [11], together with Felix and Thomas, we gave a different proof that  $H^*(\Gamma) : HH_{\text{coalg}}^*(C, C) \rightarrow HH^*(A, A)$  is a morphism of Gerstenhaber algebras.

**Proposition 22.** *Let  $C$  be a bialgebra  $\mathbb{k}$ -free of finite type. Let  $A$  be the dual bialgebra. Then the inclusions of Gerstenhaber algebras given by Theorems 15 and 16 fit into the commutative diagram of Gerstenhaber algebras.*

$$\begin{array}{ccc} \text{Cotor}_C^*(\mathbb{k}, \mathbb{k}) & \longrightarrow & HH_{\text{coalg}}^*(C, C) \\ H^*(\phi) \downarrow \cong & & \cong \downarrow H^*(\Gamma) \\ \text{Ext}_A^*(\mathbb{k}, \mathbb{k}) & \longrightarrow & HH^*(A, A) \end{array}$$

*Proof.* Since  $C$  is an algebra  $\mathbb{k}$ -free of finite type, the dualizing functor  ${}^\vee$ , defined in the proof of Lemma 21, restrict to a functor  $F$  from the opposite category of left  $C$ -modules to the category of left  $A$ -comodules. If

$M$  and  $N$  are left  $C$ -modules,  $\psi : M^\vee \otimes N^\vee \rightarrow (M \otimes N)^\vee$  is a morphism of left  $A$ -comodules. Therefore by Property 12, we obtain the morphism of linear operads with multiplication  $\Gamma_F : \mathcal{CoEnd}_{C\text{-mod}}(C) \rightarrow \mathcal{End}_{A\text{-comod}}(A)$ . Consider the two commutatives squares

$$\begin{array}{ccccc} \mathcal{CoEnd}_{C\text{-mod}}(C) & \xrightarrow{i_C} & \mathcal{CoEnd}_{\mathbb{k}\text{-mod}}(C) & \xrightarrow{\mathcal{C}_{\text{coalg}}^*(C,\eta)} & \Omega C \\ \Gamma_F \downarrow \cong & & \Gamma \downarrow \cong & & \cong \downarrow \phi \\ \mathcal{End}_{A\text{-comod}}(A) & \xrightarrow{i_A} & \mathcal{End}_{\mathbb{k}\text{-mod}}(A) & \xrightarrow{\mathcal{C}^*(A,\varepsilon)} & (BA)^\vee \end{array}$$

The left square commutes by definition of  $\Gamma_F$  since the two horizontal maps  $i_C$  and  $i_A$  are just the inclusions. Part iii) of Lemma 21 says that the right square commutes. The composite  $\mathcal{CoEnd}_{C\text{-mod}}(C) \xrightarrow{i_C} \mathcal{CoEnd}_{\mathbb{k}\text{-mod}}(C) \xrightarrow{\mathcal{C}_{\text{coalg}}^*(C,\eta)} \Omega C$  is the isomorphism  $ev$  considered in the proof of Theorem 15. The composite  $\mathcal{End}_{A\text{-comod}}(A) \xrightarrow{i_A} \mathcal{End}_{\mathbb{k}\text{-mod}}(A) \xrightarrow{\mathcal{C}^*(A,\varepsilon)} (BA)^\vee$  is the isomorphism  $proj$  considered in the proof of Theorem 16. Therefore, we have the commutative square of linear operads with multiplication

$$\begin{array}{ccc} \Omega C & \xrightarrow{i_C \circ ev^{-1}} & \mathcal{C}_{\text{coalg}}^*(C, C) \\ \phi \downarrow \cong & & \cong \downarrow \Gamma \\ (BA)^\vee & \xrightarrow{i_A \circ proj^{-1}} & \mathcal{C}^*(A, A) \end{array}$$

Applying homology, we obtain the Proposition.  $\square$

Let  $H$  be a finite dimensional Hopf algebra. Let  $D(H)$  be the Drinfeld double of  $H$ . Then Taillefer [40] proved that the Gerstenhaber-Schack cohomology of  $H$ ,  $H_{GS}(H, H)$  is isomorphic as graded algebras to  $\text{Ext}_{D(H)^{op}}(\mathbb{F}, \mathbb{F})$ . Since  $D(H)$  is a Hopf algebra, by Theorem 16, Farinati and Solotar [9] have obtained a Gerstenhaber algebra structure on  $\text{Ext}_{D(H)^{op}}(\mathbb{F}, \mathbb{F}) = H_{GS}(H, H)$ . But Taillefer using a braiding [40, Beginning of Section 5] shows that the Lie bracket in this Gerstenhaber algebra structure is trivial. The Drinfeld double  $D(H)$  is a braided Hopf algebra. Therefore, following the proof of Taillefer, it should be easy to prove

**Conjecture 23.** *Let  $A$  be braided bialgebra. Then the Lie algebra of the Gerstenhaber algebra  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  given by Theorem 16 is trivial.*

*Proof when  $A$  is a cocommutative Hopf algebra.* Let  $A$  be a cocommutative Hopf algebra. Since  $A$  is cocommutative, the antipode  $S$  is involutive. Therefore by Theorem 49,  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  is a Batalin-Vilkovisky algebra. By [22, Theorem 4.1], the operator  $B$  of this Batalin-Vilkovisky algebra is trivial. Therefore by (29), the Lie bracket is null.  $\square$

In [36], Shoikhet mentions the following conjecture of Kontsevich.

**Conjecture 24.** (*Kontsevich*) *Let  $H$  be a bialgebra. Then  $H_{GS}(H, H)$  is a 3-algebra, i. e. [28, Theorem p. 26-7] an algebra over the homology of the little 3-cubes operad,  $\mathcal{C}_3$ .*

Shoikhet [36, Corollary 0.3] has announced that the proof of this conjecture when  $H$  is a Hopf algebra. We formulate the following related conjecture:

**Conjecture 25.** *Let  $A$  be braided bialgebra. Then  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  is a 3-algebra.*

If  $A$  is cocommutative, again this Lie algebra bracket (of degree  $-2$  this time) should vanish: modulo  $p$ , only the Steenrod or Dyer-Lashoff operations on  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  (Remark 8 and [29, Theorem 11.8]) should be non trivial.

As an algebraic topologist, we find the following Corollary of Theorem 15, highly interesting.

**Corollary 26.** *Let  $X$  be a 2-connected pointed topological space. Denote by  $\Omega_M X$  the pointed Moore loops on  $X$ . Then the homology of the double loop spaces on  $X$ ,  $H_*(\Omega^2 X)$ , equipped with the Pontryagin product, is a sub Gerstenhaber algebra of  $HH_{\text{coalg}}^*(S_*(\Omega_M X), S_*(\Omega_M X))$ , the Hochschild cohomology of the coalgebra  $S_*(\Omega_M X)$ .*

*Proof.* The bialgebra  $C$  in Theorem 15 can be differential graded. Since  $\Omega_M X$  is a topological monoid, the (reduced normalized) singular chains on  $\Omega_M X$  form a differential graded bialgebra  $C = S_*(\Omega_M X)$ . Therefore, by Theorem 15,  $\text{Cotor}_{S_*(\Omega_M X)}(\mathbb{k}, \mathbb{k})$  is a sub Gerstenhaber algebra of  $HH_{\text{coalg}}^*(S_*(\Omega_M X), S_*(\Omega_M X))$ . By Adams cobar equivalence, there is an isomorphism of graded algebras  $\text{Cotor}_{S_*(\Omega_M X)}(\mathbb{k}, \mathbb{k}) \cong H_*(\Omega_M \Omega_M X)$ . The inclusion of the (ordinary) pointed loops into the Moore loops  $\Omega X \xrightarrow{\cong} \Omega_M X$  is both a homotopy equivalence [46, p. 112-3] and a morphism of  $H$ -spaces. So as graded algebras,  $H_*(\Omega_M \Omega_M X)$  is isomorphic to  $H_*(\Omega^2 X)$ .  $\square$

Corollary 26 gives in particular a Gerstenhaber algebra structure on  $H_*(\Omega^2 X)$  extending the Pontryagin product. Of course, we believe that

this Gerstenhaber algebra structure coincides with the usual one given by Cohen in [4]:

**Conjecture 27.** *Let  $X$  be a 2-connected pointed topological space. There is an isomorphism of Gerstenhaber algebras between the Gerstenhaber algebra  $\text{Cotor}_{S^*(\Omega_M X)}(\mathbb{k}, \mathbb{k})$  given by Theorem 15 and the Gerstenhaber algebra  $H_*(\Omega^2 X)$  given by Cohen in [4].*

Recall that the Gerstenhaber algebra on  $H_*(\Omega^2 X)$  is usually defined as follows: the little 2-cube operad  $\mathcal{C}_2$  acts on the double loop space,  $\Omega^2 X$ . So its homology  $H_*(\Omega^2 X)$  is an algebra over the homology of  $\mathcal{C}_2$ , i. e. is a Gerstenhaber algebra by Cohen [4].

## 5. BATALIN-VILKOVISKY ALGEBRAS

**28.** A *Batalin-Vilkovisky algebra* is a Gerstenhaber algebra  $A$  equipped with a degree  $-1$  linear map  $B : A^i \rightarrow A^{i-1}$  such that  $B \circ B = 0$  and

$$(29) \quad \{a, b\} = (-1)^{|a|} (B(a \cup b) - (Ba) \cup b - (-1)^{|a|} a \cup (Bb))$$

for  $a$  and  $b \in A$ .

**Definition 30.** A *cyclic operad* is a non- $\Sigma$ -operad  $\mathcal{O}$  equipped with linear maps  $\tau_n : \mathcal{O}(n) \rightarrow \mathcal{O}(n)$  for  $n \in \mathbb{N}$  such that

$$(31) \quad \forall n \in \mathbb{N}, \quad \tau_n^{n+1} = id_{\mathcal{O}(n)},$$

$$(32) \quad \forall m \geq 1, n \geq 1, \quad \tau_{m+n-1}(f \circ_1 g) = \tau_n g \circ_n \tau_m f,$$

$$(33) \quad \forall m \geq 2, n \geq 0, 2 \leq i \leq m, \quad \tau_{m+n-1}(f \circ_i g) = \tau_m f \circ_{i-1} g,$$

for each  $f \in \mathcal{O}(m)$  and  $g \in \mathcal{O}(n)$ . In particular, we have  $\tau_1 id = id$ .

**Definition 34.** [31] A *cyclic operad with multiplication* is an operad which is both an operad with multiplication and a cyclic operad such that

$$\tau_2 \mu = \mu.$$

**Theorem 35.** [31] *If  $\mathcal{O}$  is a cyclic operad with a multiplication then*

a) *the structure of cosimplicial module on  $\mathcal{O}$  extends to a structure of cocyclic module and*

b) *the Connes coboundary map  $B$  on  $\mathcal{C}^*(\mathcal{O})$  induces a natural structure of Batalin-Vilkovisky algebra on the Gerstenhaber algebra  $H^*(\mathcal{C}^*(\mathcal{O}))$ .*

**Theorem 36.** *Let  $\mathcal{O}$  be a linear cyclic operad with multiplication. Consider the associated cocyclic module. Then the cyclic cochains  $\mathcal{C}_\lambda(\mathcal{O})$  forms a subcomplex of  $\mathcal{C}^*(\mathcal{O})$ , stable under the Lie bracket of degree  $-1$ . In particular, the cyclic cohomology  $HC_\lambda^*(\mathcal{C}^*(\mathcal{O}))$  has naturally a graded Lie algebra structure of degree  $-1$ .*

CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

*Proof.* Let  $\mathcal{O}$  be a linear cyclic operad. Let  $f \in \mathcal{C}_\lambda^m(\mathcal{O})$  and  $g \in \mathcal{C}_\lambda^n(\mathcal{O})$ . Using (32), (33) and the change of variable  $i' = i - 1$  for the first equation and using  $\tau_m(f) = (-1)^m f$  and  $\tau_m(g) = (-1)^n g$  for the second equation, we have

$$\begin{aligned} \tau_{m+n-1}(f\bar{\circ}g) &= (-1)^{(m-1)(n-1)} \left( \tau_n g \circ_n \tau_m f + \sum_{i'=1}^{m-1} (-1)^{(n-1)i'} \tau_m f \circ_{i'} g \right) \\ &= (-1)^{(m-1)(n-1)} \left( (-1)^{m+n} g \circ_n f + (-1)^{m+n-1} \sum_{i=1}^{m-1} (-1)^{(n-1)(i-1)} f \circ_i g \right). \end{aligned}$$

By symmetry

$$(-1)^{(m-1)(n-1)} \tau_{m+n-1}(g\bar{\circ}f) = (-1)^{m+n} f \circ_m g + (-1)^{m+n-1} \sum_{i=1}^{n-1} (-1)^{(m-1)(i-1)} g \circ_i f.$$

Therefore  $\tau_{m+n-1}\{f, g\} = (-1)^{m+n-1}\{f, g\}$ .

Suppose now that  $\mathcal{O}$  has an associative multiplication  $\mu$  such that  $\tau_2\mu = \mu$ . Since  $\mu \in \mathcal{C}_\lambda^2(\mathcal{O})$ , we have just proved above that for any  $g \in \mathcal{C}_\lambda^n(\mathcal{O})$ , the differential of  $g$ ,  $d(g) = \{\mu, g\} \in \mathcal{C}_\lambda(\mathcal{O})$ .  $\square$

*Remark 37.* Let  $\mathcal{O}$  be a cyclic operad. Then  $\tau_1 : \mathcal{O}(1) \rightarrow \mathcal{O}(1)$  is an involutive morphism of anti-algebras. And  $\mathcal{C}_\lambda^1(\mathcal{O}) = \text{Ker}(\tau_1 + \text{Id} : \mathcal{O}(1) \rightarrow \mathcal{O}(1))$  is a sub Lie algebra of the Lie algebra associated to associative algebra  $\mathcal{O}(1)$  (Compare with Remark 9).

*Remark 38.* In [31, Corollary 1.5], motivated by applications to string topology [2], we proved that the negative cyclic cohomology of a cyclic operad  $\mathcal{O}$  with multiplication,  $HC_-^*(\mathcal{C}^*(\mathcal{O}))$ , has a Lie bracket of degree  $-2$ .

*Remark 39.* The (ordinary) cyclic cohomology of  $\mathcal{O}$ ,  $HC^*(\mathcal{C}^*(\mathcal{O}))$ , has also a Lie bracket of degree  $-1$ . This was stated only in the case of the cyclic cohomology of the group ring  $\mathbb{k}[G]$  of a finite group  $G$  [3, Theorem 67 a)]. But the proof of [3, Theorem 67 a)] works for any cyclic operad with multiplication.

*Remark 40.* The proof of Theorem 36 is a lot more simple than the proofs of remarks 38 and 39. Indeed, the proofs of remarks 38 and 39 use that  $H^*(\mathcal{C}^*(\mathcal{O}))$  is a Batalin-Vilkovisky algebra (Theorem 35). On the contrary, in the proof of Theorem 36, we don't even use that  $H^*(\mathcal{C}^*(\mathcal{O}))$  is a Gerstenhaber algebra: we use only the Lie algebra on  $\mathcal{C}^*(\mathcal{O})$ .

If our ground ring  $\mathbb{k}$  contains  $\mathbb{Q}$ , there is a natural isomorphism [25, p. 72]

$$HC_\lambda^n(\mathcal{C}^*(\mathcal{O})) \xrightarrow{\cong} HC^n(\mathcal{C}^*(\mathcal{O})).$$

This isomorphism obviously should be compatible with the brackets.

Recall the following well-known result in string topology.

**Corollary 41.** ([44],[31, Theorem 1.6]) *Let  $A$  be a symmetric Frobenius algebra (Definition 54). Then its Hochschild cohomology  $HH^*(A, A)$  is a Batalin-Vilkovisky algebra.*

We need to sketch our proof given in [31].

*Proof.* Let  $\Theta : A \xrightarrow{\cong} A^\vee$  be an isomorphism of  $A$ -bimodules given by the symmetric Frobenius algebra structure on  $A$ . Then  $\mathcal{C}^*(A, \Theta) : \mathcal{C}^*(A, A) \xrightarrow{\cong} \mathcal{C}^*(A, A^\vee)$  is an isomorphism of cosimplicial modules. Let  $Ad : \mathcal{C}^*(A, A^\vee) \xrightarrow{\cong} \mathcal{C}_*(A, A)^\vee$  be the adjunction map [31, (4.1)] which associates to any  $g \in \text{Hom}(A^n, A^\vee)$ , the linear map  $Ad(g) : A \otimes A^{\otimes n} \rightarrow \mathbb{k}$  given by  $Ad(g)(a_0 \otimes \cdots \otimes a_n) = g(a_1, \dots, a_n)(a_0)$ . Then  $Ad : \mathcal{C}^*(A, A^\vee) \xrightarrow{\cong} \mathcal{C}_*(A, A)^\vee$  is an isomorphism of cosimplicial modules. By [31, Proof of Theorem 1.6]

$$\mathcal{C}^*(A, A) \xrightarrow[\cong]{\mathcal{C}^*(A, \Theta)} \mathcal{C}^*(A, A^\vee) \xrightarrow[\cong]{Ad} \mathcal{C}_*(A, A)^\vee$$

equipped with the  $\tau_n$  [31, (4.2)] is a cyclic operad with multiplication. Using Theorem 35,  $HH^*(A, A) \xrightarrow{\cong} HH^*(A, A^\vee)$  is a Batalin-Vilkovisky algebra.  $\square$

If instead of using Theorem 35, we apply Theorem 36 in the previous proof, we obtain the following Corollary:

**Corollary 42.** *Let  $A$  be a symmetric Frobenius algebra. Then its cyclic cohomology  $HC_\lambda^*(A)$  (in the sense of [25, 2.4.2]), is a graded Lie algebra of degree  $-1$ .*

We wonder if this Corollary is not a particular simple case of [15, Proposition 2.11]?

In [31], our main objective was the following result

**Corollary 43.** [31, Theorem 1.1] *Let  $H$  be a Hopf algebra endowed with a modular pair in involution  $(\delta, 1)$  (Definition 47). Then the Connes-Moscovici cocyclic on the Cobar construction on  $H$ , defines a Batalin-Vilkovisky algebra structure on  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$ .*

*Proof.* A computation [31, Section 5] shows that the operad with multiplication  $\mathcal{CoEnd}_{H\text{-mod}}(H) \cong \Omega H$  considered in the proof of Theorem 15 equipped with the  $\tau_n$  defined by Connes and Moscovici, is cyclic. Therefore, by Theorem 35, its homology  $\text{Cotor}_H^*(\mathbb{k}, \mathbb{k})$  is a Batalin-Vilkovisky algebra.  $\square$

If instead of using Theorem 35, we apply Theorem 36 in the previous proof, we obtain the following Corollary:

**Corollary 44.** *Let  $H$  be a Hopf algebra endowed with a modular pair in involution  $(\delta, 1)$  (Definition 47). Then its cyclic cohomology,  $HC_{(\delta,1)}^*(H)$ , is a graded Lie algebra of degree  $-1$ .*

## 6. BATALIN-VILKOVISKY ALGEBRA STRUCTURE ON $\text{EXT}_H^*(\mathbb{k}, \mathbb{k})$

Everybody is more familiar with an algebra  $A$  than with a coalgebra  $C$ . And therefore, one usually prefers the Exterior product  $\text{Ext}_A^*(\mathbb{k}, \mathbb{k})$  instead of the Cotorsion product  $\text{Cotor}_C^*(\mathbb{k}, \mathbb{k})$ . The goal of this section is to give the duals of Corollaries 43 and 44, Theorem 49 below. Taillefer [39], Khalkhali and Rangipour [22] developed a theory dual to Connes-Moscovici cyclic cohomology of Hopf algebras. First, we are going to explain this duality.

**Proposition 45.** *Let  $K$  be a finite dimensional Hopf algebra with a modular pair in involution  $(\delta, \sigma)$  in the sense of Khalkhali-Rangipour [22, (1)]. Then i) its dual  $K^\vee$  is a Hopf algebra equipped with a modular pair in involution  $(\text{ev}_\sigma, \delta)$  in the sense of Connes-Moscovici where  $\text{ev}_\sigma : K^\vee \rightarrow \mathbb{F}$  is defined by  $\text{ev}_\sigma(\varphi) = \varphi(\sigma)$ .*

*Let  $\psi_n : (K^\vee)^{\otimes n} \xrightarrow{\cong} (K^{\otimes n})^\vee$  be the linear map mapping the tensor product  $\varphi_1 \otimes \cdots \otimes \varphi_n$  of  $n$  forms on  $K$  to the form on  $K^{\otimes n}$ , also denoted  $\varphi_1 \otimes \cdots \otimes \varphi_n$ , mapping  $k_1 \otimes \cdots \otimes k_n$  to the product  $\varphi_1(k_1) \cdots \varphi_n(k_n)$ . Then ii)  $\psi_*$  is an isomorphism of cocyclic modules between the cocyclic modules  $\Omega(K^\vee)_{(\text{ev}_\sigma, \delta)}$  introduced by Connes-Moscovici and the dual of the cyclic module  $B(K)^{(\delta, \sigma)}$  introduced by Khalkhali-Rangipour [22, Theorem 2.1] and Taillefer [39].*

*iii) In particular,  $\psi_*$  induces an isomorphism between Connes-Moscovici cyclic cohomology of  $K^\vee$ ,  $HC_{(\text{ev}_\sigma, \delta)}^*(K^\vee)$  and the dual of Khalkhali-Rangipour-Taillefer cyclic homology of  $K$ ,  $\widetilde{HC}_*^{(\delta, \sigma)}(K)^\vee$ .*

The cocyclic module  $\Omega(H)_{(\delta, \sigma)}$  is denoted  $H_{(\delta, \sigma)}^\natural$  in [6, Theorem 1]. The cyclic module  $B(K)^{(\delta, \sigma)}$  is denoted  $\widetilde{K}^{(\delta, \sigma)}$  in [22, Theorem 2.1] and  $C_*^{(\sigma, \varepsilon, \delta)}(K)$  in [39, 2.1].

**Definition 46.** A *modular pair* is a couple  $(\delta, \sigma)$  when  $\delta$  is a character and  $\sigma$  is a group-like element such that  $\delta(\sigma) = 1$ .

*Proof of Proposition 45.* i) An element  $\sigma$  is a *group-like* element of  $K$  by definition if and only if  $\Delta\sigma = \sigma \otimes \sigma$  and  $\varepsilon(\sigma) = 1$ . This means that the linear map that we denoted again  $\sigma : \mathbb{F} \rightarrow K$ , mapping 1 to  $\sigma$  is a morphism of coalgebras. Therefore its dual  $ev_\sigma = \sigma^\vee : K^\vee \rightarrow \mathbb{F}$  is a morphism of algebras, i. e. a *character* of  $K^\vee$ . Let  $\delta : K \rightarrow \mathbb{F}$  be a character of  $K$ , i. e. a morphism of algebras. Its dual  $\delta^\vee : \mathbb{F} \rightarrow K^\vee$ , mapping 1 to  $\delta$ , is a morphism of coalgebras, i.e  $\delta$  is a group-like element of  $K^\vee$ . By definition,  $ev_\sigma(\delta) = \delta(\sigma)$ . Therefore  $(\delta, \sigma)$  is a modular pair on  $K$  if and only if  $(ev_\sigma, \delta)$  is a modular pair in  $K^\vee$ .

Let  $(\delta, \sigma)$  be a modular pair on  $H$ . The *twisted antipode*  $\tilde{S}$  associated to  $(\delta, \sigma)$  (in the sense of Connes-Moscovici) is by definition the convolution product  $(\eta \circ \delta) \star S$  in  $\text{Hom}(H, H)$ . Explicitly, for  $h \in H$ ,  $\tilde{S}(h) = \delta(h^1)S(h^2)$ , where  $\Delta h = h^1 \otimes h^2$ . Consider the map  $\tau_n : H^{\otimes n} \rightarrow H^{\otimes n}$  defined by

$$\tau_n(h_1 \otimes \cdots \otimes h_n) := \mu_{H^{\otimes n}} \left( \Delta^{n-1} \tilde{S}(h_1) \otimes (h_2 \otimes \cdots \otimes h_n \otimes \sigma) \right)$$

Here  $\mu_{H^{\otimes n}} : H^{\otimes n} \otimes H^{\otimes n} \rightarrow H^{\otimes n}$  is the product in  $H^{\otimes n}$  and  $\Delta^{n-1} : H \rightarrow H^{\otimes n}$  is the iterated diagonal on  $H$ . In particular,  $\tau_1(h) = \tilde{S}(h)\sigma$ .

**Definition 47.** A modular pair  $(\delta, \sigma)$  (Definition 46) is *in involution* in the sense of Connes-Moscovici if and only if  $\tau_1^2 = id_H$ , i. e.  $\forall h \in H$ ,  $\tilde{S}^2(h) = \sigma h \sigma^{-1}$ .

Let  $(\Omega H)_{(\delta, \sigma)}$  be the usual cosimplicial module defining the Cobar construction, except that  $\delta_{n+1} : H^{\otimes n} \rightarrow H^{\otimes n+1}$  is given by [6, (2.9)]  $\delta_{n+1}(h_1 \otimes \cdots \otimes h_n) = h_1 \otimes \cdots \otimes h_n \otimes \sigma$ . Connes and Moscovici have shown that if  $\tau_1^2 = id_H$ , then  $(\Omega H)_{(\delta, \sigma)}$  (equipped with the  $\tau_n$ ) is a cocyclic module.

Let  $(\delta, \sigma)$  be a modular pair on  $K$ . Let  $t_n : K^{\otimes n} \rightarrow K^{\otimes n}$  defined by ([22, Theorem 2.1] or [39, 2.1] which generalizes [25, (7.3.3.1)])

$$t_n(k_1 \otimes \cdots \otimes k_n) = \sigma S(k_1^{(1)} \dots k_n^{(1)}) \otimes k_1^{(2)} \otimes \cdots \otimes k_{n-1}^{(2)} \delta(k_n^{(2)})$$

where  $\Delta(k_i) = k_i^{(1)} \otimes k_i^{(2)}$ . In particular,  $t_1$  is equal to  $\sigma(S \star \eta \circ \delta)$ , the left multiplication by  $\sigma$  of the convolution product  $\star$  of  $S$  and  $\eta \circ \delta$ . By definition, the couple  $(\delta, \sigma)$  is a *modular pair in involution* in the sense of Khalkhali-Rangipour [22, (1)] if and only if  $t_1^2 = id_K$ .

Therefore to prove part i) of this Proposition, it suffices to show that  $\tau_1 = t_1^\vee$ . This will be proved in the proof of ii) below.

(Denote by  $K^{op, cop}$  the Hopf algebra with the opposite multiplication, the opposite diagonal and the same antipode [8, Remark 4.2.10], since

CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

the convolution product  $\star$  on  $\text{Hom}(K^{op,cop}, K^{op,cop})$  is the opposite of the convolution product  $\star$  on  $\text{Hom}(K, K)$ , note that a modular pair in involution for  $K$  in the sense of Khalkhali-Rangipour is the same as a modular pair in involution for  $K^{op,cop}$  in the sense of Connes-Moscovici.)

ii) Let  $B(K)^{(\delta,\sigma)}$  be the usual simplicial module defining the Bar construction except that  $d_{n+1} : K^{\otimes n+1} \rightarrow K^{\otimes n}$  is given by ([22, Theorem 2.1] or [39, 2.1])  $d_{n+1}(k_1 \otimes \cdots \otimes k_{n+1}) = k_1 \otimes \cdots \otimes k_n \delta(k_{n+1})$ . It is well known [20, Lemma XVIII.7.3] that  $\psi_*$  is an isomorphism of cosimplicial modules from the usual Cobar construction on  $K^\vee$ ,  $\Omega(K^\vee)_{(ev_\sigma, \varepsilon)}$ , to the dual of the usual Bar construction on  $K$ ,  $(B(K)^{(\varepsilon, \sigma)})^\vee$ . Obviously,  $\psi_{n+1} \circ \delta_{n+1} = d_{n+1}^\vee \circ \psi_n$ . Therefore  $\psi_* : \Omega(K^\vee)_{(ev_\sigma, \delta)} \xrightarrow{\cong} (B(K)^{(\delta, \sigma)})^\vee$  is an isomorphism of cosimplicial modules even if  $\delta \neq \varepsilon$ .

Denote by  $\sigma S : K \rightarrow K$  the linear map defined by  $(\sigma S)(k) = \sigma S(k)$ ,  $k \in K$ . The transposition map  $\text{Hom}(K, K) \rightarrow \text{Hom}(K^\vee, K^\vee)$ , mapping a linear map  $f : K \rightarrow K$  to its dual  $f^\vee : K^\vee \rightarrow K^\vee$  is a morphism of algebras with respect to the convolution products  $\star$ . Since  $\sigma S$  can be written as the convolution product  $(\sigma \circ \varepsilon) \star S$  of the composite  $K \xrightarrow{\varepsilon} \mathbb{F} \xrightarrow{\sigma} K$  and of the antipode  $S$ , its dual  $(\sigma S)^\vee$  is equal to  $(\varepsilon^\vee \circ \sigma^\vee) \star S^\vee = (\varepsilon \circ ev_\sigma) \star S^\vee$  which is the twisted antipode  $\tilde{S}$  on  $K^\vee$  associated to the modular pair  $(ev_\sigma, \delta)$ .

The cyclic operator  $t_n : K^{\otimes n} \rightarrow K^{\otimes n}$  can be written as the composite

$$K^{\otimes n} \xrightarrow{\Delta_{K^{\otimes n}}} K^{\otimes n} \otimes K^{\otimes n} \xrightarrow{\mu^{(n-1)} \otimes K^{\otimes n}} K \otimes K^{\otimes n} \xrightarrow{\sigma S \otimes K^{\otimes n-1} \otimes \delta} K \otimes K^{\otimes n-1} \otimes \mathbb{F}.$$

Here  $\mu^{(n-1)} : K^{\otimes n} \rightarrow K$  is the iterated product on  $K$  and  $\Delta_{K^{\otimes n}}$  is the diagonal of  $K^{\otimes n}$ . The cocyclic operator  $\tau_n : H^{\otimes n} \rightarrow H^{\otimes n}$  can be written as the composite

$$H \otimes H^{\otimes n-1} \otimes \mathbb{F} \xrightarrow{\tilde{S} \otimes H^{\otimes n-1} \otimes \sigma} H \otimes H^{\otimes n} \xrightarrow{\Delta^{(n-1)} \otimes H^{\otimes n}} H^{\otimes n} \otimes H^{\otimes n} \xrightarrow{\mu_{H^{\otimes n}}} H^{\otimes n}$$

Here  $\Delta^{(n-1)} : H \rightarrow H^{\otimes n}$  is the iterated diagonal on  $H$  and  $\mu_{H^{\otimes n}}$  is the multiplication of  $H^{\otimes n}$ . We saw that the twisted antipode  $\tilde{S}$  on  $K^\vee$  associated to  $(ev_\sigma, \delta)$  was  $(\sigma S)^\vee$ , the dual of  $\sigma S$ . Therefore  $\psi_n \circ \tau_n = t_n^\vee \circ \phi_n$ . In particular when  $n = 1$ , since  $\psi_1$  is the identity,  $\tau_1 = t_1^\vee$ . So finally,  $\psi_* : \Omega(K^\vee)_{(ev_\sigma, \delta)} \xrightarrow{\cong} (B(K)^{(\delta, \sigma)})^\vee$  is an isomorphism of cocyclic modules.  $\square$

**Corollary 48.** *Let  $K$  be a finite dimensional Hopf algebra equipped with a group-like element  $\sigma$  such that  $\forall k \in K, S \circ S(k) = \sigma^{-1}k\sigma$ . Then  $\psi_* : \Omega(K^\vee)_{(ev_\sigma, \varepsilon)} \xrightarrow{\cong} (B(K)^{(\varepsilon, \sigma)})^\vee$  is an isomorphism of cyclic operads with multiplication. In particular,  $H^*(\psi_*) : \text{Cotor}_{K^\vee}^*(\mathbb{F}, \mathbb{F}) \xrightarrow{\cong}$*

$Ext_K^*(\mathbb{F}, \mathbb{F})$  is an isomorphism of Batalin-Vilkovisky algebras and  $\psi_*$  induces an isomorphism of graded Lie algebras  $HC_{(ev_\sigma, \varepsilon)}^*(K^\vee) \xrightarrow{\cong} \widetilde{HC}_{(\varepsilon, \sigma)}^*(K)$ .

*Proof.* The canonical injection of  $K$  into its bidual  $K^{\vee\vee}$ ,  $\nu : K \hookrightarrow K^{\vee\vee}$ , is an isomorphism of bialgebras. Let  $C := K^\vee$  be the dual bialgebra. In the proof of Proposition 22, we saw that  $\phi : \Omega C \xrightarrow{\cong} B(C^\vee)^\vee$  is an isomorphism of linear operads with multiplication. Therefore the composite  $\Omega(K^\vee) \xrightarrow{\phi} B(K^{\vee\vee})^\vee \xrightarrow{B(\nu)^\vee} B(K)^\vee$  is also an isomorphism of linear operads with multiplication. But this composite coincides with the isomorphism of cocyclic modules  $\psi_* : \Omega(K^\vee)_{(ev_\sigma, \varepsilon)} \xrightarrow{\cong} (B(K)^{(\varepsilon, \sigma)})^\vee$  given by part ii) of Proposition 45.  $\square$

**Theorem 49.** *Let  $K$  be a Hopf algebra equipped with a group-like element  $\sigma$  such that for all  $k \in K$ ,  $S^2(k) = \sigma^{-1}k\sigma$ . Let  $t_n : K^{\otimes n} \rightarrow K^{\otimes n}$  be the linear map defined by*

$$t_n(k_1 \otimes \cdots \otimes k_n) = \sigma S(k_1^{(1)} \cdots k_{n-1}^{(1)} k_n) \otimes k_1^{(2)} \otimes \cdots \otimes k_{n-1}^{(2)}.$$

*The dual of the Bar construction on  $K$ ,  $B(K)^\vee$  is a cyclic operad with multiplication. In particular, the Gerstenhaber algebra given by Theorem 16,  $Ext_K^*(\mathbb{k}, \mathbb{k})$ , is in fact a Batalin-Vilkovisky algebra and the cyclic cohomology of  $K$ ,  $\widetilde{HC}_{(\varepsilon, \sigma)}^*(K)$  has a Lie bracket of degree  $-1$ .*

*Proof.* Corollary 48 explains in details that this Theorem is the dual of Corollaries 43 and 44. Therefore, the computations dual to [31, Proof of Theorem 1.1] show that the operad with multiplication  $B(K)^\vee$  given in the proof of Theorem 16 together with the cyclic operators  $t_n$  defines a cyclic operad with multiplication. Using Theorems 35 and 36, we conclude.  $\square$

## 7. CHARACTERISTIC MAPS

**Lemma 50.** *Let  $H$  be a bialgebra. Let  $A$  be a left module algebra over  $H$  (in the sense of [20, Definition V.6.1]). Then the application  $\Phi : H^{\otimes n} \rightarrow Hom_{\mathbb{k}\text{-mod}}(A^{\otimes n}, A)$  mapping  $h_1 \otimes \cdots \otimes h_n$  to  $f : A^{\otimes n} \rightarrow A$  defined by  $f(a_1 \otimes \cdots \otimes a_n) = (h_1 \cdot a_1) \cdots (h_n \cdot a_n)$  defines a morphism of linear operads with multiplication from the coendomorphism operad of  $H$ ,  $CoEnd_{H\text{-mod}}(H)$ , to the endomorphism operad of  $A$ ,  $End_{\mathbb{k}\text{-mod}}(A)$ . In particular,  $\Phi$  induces a morphism of Gerstenhaber algebras  $H^*(\Phi) : Cotor_H^*(\mathbb{k}, \mathbb{k}) \rightarrow HH^*(A, A)$ .*

*Proof.* Since  $1_H \cdot a_1 = a_1$ ,  $\Phi(1_H) = id_A$ . Let  $h_1 \otimes \cdots \otimes h_m \in H^{\otimes m}$ ,  $k_1 \otimes \cdots \otimes k_m \in H^{\otimes n}$  and  $a_1 \otimes \cdots \otimes a_{m+n-1} \in A^{\otimes m+n-1}$ . Using (18), we

have that  $\Phi[(h_1 \otimes \cdots \otimes h_m) \circ_i (k_1 \otimes \cdots \otimes k_n)]$  evaluated on  $a_1 \otimes \cdots \otimes a_{m+n-1}$  is equal to the product

$$(h_1.a_1) \cdots (h_{i-1}.a_{i-1})(h_i^{(1)}k_1.a_i) \cdots (h_i^{(n)}k_n.a_{i+n-1})(h_{i+1}.a_{i+n}) \cdots (h_m.a_{m+n-1}).$$

On the other hand, using example 2,  $\Phi(h_1 \otimes \cdots \otimes h_m) \circ_i \Phi(k_1 \otimes \cdots \otimes k_n)$  evaluated on  $a_1 \otimes \cdots \otimes a_{m+n-1}$  is equal to the product

$$(h_1.a_1) \cdots (h_{i-1}.a_{i-1}) (h_i.[(k_1.a_i) \cdots (k_n.a_{i+n-1})]) (h_{i+1}.a_{i+n}) \cdots (h_m.a_{m+n-1}).$$

Since for any  $h \in H$ ,  $a$  and  $b \in A$ ,  $h.(ab) = (h^{(1)}.a)(h^{(2)}.a)$ , the previous two products are equal. So  $\Phi$  is a morphism of operads. Now  $\Phi$  is a morphism of operads with multiplication, since  $\Phi(1_{\mathbb{k}})$  is the unit map  $\eta : \mathbb{k} \rightarrow A$  and since  $\Phi(1_H \otimes 1_H)$  is the multiplication  $\mu : A \otimes A \rightarrow A$  of  $A$ .  $\square$

The following Lemma is a variant of the previous lemma if  $H$  is finite dimensional, since in this case,  $A$  is a left module algebra over  $H$  if and only if  $A$  be a right comodule algebra over the dual of  $H$ ,  $H^\vee$ .

**Lemma 51.** *Let  $H$  be a bialgebra. Let  $A$  be a right comodule algebra over  $H$  (in the sense of [20, Definition III.7.1]). Then the application  $\Phi : (H^{\otimes n})^\vee \rightarrow \text{Hom}_{\mathbb{k}\text{-mod}}(A^{\otimes n}, A)$  mapping  $f : H^{\otimes n} \rightarrow \mathbb{k}$  to  $F : A^{\otimes n} \rightarrow A$  defined by  $F(a_1 \otimes \cdots \otimes a_n) = a_1^{(1)} \cdots a_n^{(1)} f(a_1^{(2)} \otimes \cdots \otimes a_n^{(2)})$ . defines a morphism of linear operads with multiplication from the endomorphism operad of  $H$ ,  $\mathcal{E}nd_{H\text{-comod}}(H)$ , to the endomorphism operad of  $A$ ,  $\mathcal{E}nd_{\mathbb{k}\text{-mod}}(A)$ . In particular,  $\Phi$  induces a morphism of Gerstenhaber algebras  $H^*(\Phi) : \text{Ext}_H^*(\mathbb{k}, \mathbb{k}) \rightarrow HH^*(A, A)$ .*

Note that in the case  $A = H$ ,  $H^*(\Phi)$  coincides with the inclusion of Gerstenhaber algebras given by Theorem 16. The proof of Lemma 51 is a computation similar to the proof of Lemma 50.

**Theorem 52.** *Let  $H$  be a Hopf algebra equipped with a group-like element  $\sigma \in H$  such that  $\forall h \in H$ ,  $S^2(h) = \sigma^{-1}h\sigma$ . Let  $A$  be a right comodule algebra over  $H$ . Let  $\tau : A \rightarrow \mathbb{k}$  be a non degenerate 1-trace, i. e. the morphism of left  $A$ -modules  $\Theta : A \xrightarrow{\cong} A^\vee$ , mapping  $b \in A$  to  $\varphi : A \rightarrow \mathbb{k}$  given by  $\varphi(a) = \tau(ab)$ , is an isomorphism of  $A$ -bimodules. Suppose that  $\tau$  is  $\sigma$ -invariant in the sense of [22, Definition 3.1]:  $\forall a, b \in A$ ,  $\tau(a^{(1)}a^{(2)}) = \tau(a)\sigma$ . Then*

1) *the morphism  $H^*(\Phi) : \text{Ext}_H^*(\mathbb{k}, \mathbb{k}) \rightarrow HH^*(A, A)$  given by Lemma 50, is a morphism of Batalin-Vilkovisky algebras,*

2) *the characteristic map defined by Khalkhali-Rangipour [22, (10)]  $\gamma^* : \widetilde{HC}_{(\varepsilon, \sigma)}^*(H) \rightarrow HC_\lambda^*(A)$  is a morphism of graded Lie algebras.*

*Proof.* Recall from the proof of Corollary 41 that

$$\mathcal{C}^*(A, A) \xrightarrow[\cong]{\mathcal{C}^*(A, \Theta)} \mathcal{C}^*(A, A^\vee) \xrightarrow[\cong]{Ad} \mathcal{C}_*(A, A)^\vee$$

is a cyclic operad with multiplication. By Lemma 51,  $\Phi : B(H)^\vee \rightarrow \mathcal{C}^*(A, A)$  is a morphism of linear operads with multiplication. Let  $\gamma : \mathcal{C}_*(A, A) \rightarrow B(H)$  be the morphism of cyclic modules defined by [22, Proposition 3.1]

$$\gamma(a_0 \otimes a_1 \otimes \cdots \otimes a_n) = \tau(a_0 a_1^{(1)} \cdots a_n^{(1)})(a_1^{(2)} \otimes \cdots \otimes a_n^{(2)}).$$

Here the coaction of  $a_i$ ,  $\Delta a_i = a_i^{(1)} \otimes a_i^{(2)}$ . A straightforward calculation shows that the composite

$$B(H)^\vee \xrightarrow{\Phi} \mathcal{C}^*(A, A) \xrightarrow[\cong]{\mathcal{C}^*(A, \Theta)} \mathcal{C}^*(A, A^\vee) \xrightarrow[\cong]{Ad} \mathcal{C}_*(A, A)^\vee$$

is the dual of  $\gamma$ ,  $\gamma^\vee : \mathcal{C}_*(A, A)^\vee \rightarrow B(H)^\vee$ . Since  $\gamma^\vee$  is a morphism of cocyclic modules,  $\Phi : B(H)^\vee \rightarrow \mathcal{C}^*(A, A)$  is a morphism of linear cyclic operads with multiplication. By applying Theorem 35,  $H(\Phi)$  is a morphism of Batalin-Vilkovisky algebras between the Batalin-Vilkovisky algebras given by Theorem 49 and Corollary 41. This is 1). By applying Theorem 36, we obtain 2).  $\square$

Using this time, Lemma 50 and the cocyclic map  $\gamma$  defined by Connes and Moscovici [6, Theorem 6], we obtain easily the following variant of the previous Theorem.

**Theorem 53.** *Let  $H$  be a Hopf algebra endowed with a modular pair in involution  $(\delta, 1)$ . Let  $A$  be a module algebra over  $H$ . Let  $\tau : A \rightarrow \mathbb{k}$  be a non degenerate 1-trace, i. e. the morphism of left  $A$ -modules  $\Theta : A \xrightarrow{\cong} A^\vee$ , mapping  $b \in A$  to  $\varphi : A \rightarrow \mathbb{k}$  given by  $\varphi(a) = \tau(ab)$ , is an isomorphism of  $A$ -bimodules. Suppose that  $\tau$  is  $\delta$ -invariant. Then*

1) *the morphism  $H^*(\Phi) : \text{Cotor}_H^*(\mathbb{k}, \mathbb{k}) \rightarrow HH^*(A, A)$  given by Lemma 50, is a morphism of Batalin-Vilkovisky algebras,*

2) *the characteristic map defined by Connes and Moscovici  $\chi_\tau : HC_{(\delta, 1)}^*(H) \rightarrow HC_\lambda^*(A)$  is a morphism of graded Lie algebras.*

In [6, Theorem 6] or [37, Section 4.4], Connes and Moscovici have defined more generally a characteristic map  $\chi_\tau : HC_{(\delta, \sigma)}^*(H) \rightarrow HC_\lambda^*(A)$  without assuming that

- i) the group-like element  $\sigma$  is the unit 1 of  $H$ ,
- and without assuming that
- ii) the  $\sigma$ -trace  $\tau$  is non degenerated.

But we need i) to have a Lie bracket on  $HC_{(\delta, \sigma)}^*(H)$  (Corollary 44) and we need i) and ii) to have a Lie bracket on  $HC_\lambda^*(A)$  (Corollary 42).

However, note that in their first construction of the characteristic map in [5], Connes and Moscovici were assuming i) like us. We believe that ii) can be weakened, since the Batalin-Vilkovisky algebra on  $HH^*(A, A^\vee)$  can be defined for non counital symmetric Frobenius algebras, i. e. “unital associative algebras with an invariant co-inner product” [45, p. 61-2]. In particular, as Tradler explained us,  $A$  does not need to be finite dimensional.

### 8. HOPF ALGEBRAS THAT ARE SYMMETRIC FROBENIUS

In this section, we work over an arbitrary field  $\mathbb{F}$ . We want to consider in Theorem 52, the case where the the comodule algebra  $A$  over  $H$  is the Hopf algebra  $H$  itself. We remark that for a finite dimensional Hopf algebra  $H$ , there is a close relationship between being a symmetric Frobenius algebra and being equipped with a modular pair in involution of the form  $(\varepsilon, u)$  (Theorem 60). Therefore (Theorem 62), for Hopf algebras which are symmetric Frobenius algebras, often we have an inclusion of Batalin-Vilkovisky algebras  $\text{Ext}_H^*(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(H, H)$  and in some cases, the characteristic map  $\widetilde{HC}_{(\varepsilon, \sigma)}^*(H) \rightarrow HC_\lambda^*(H)$  is injective.

First, we recall the notion of (symmetric) Frobenius algebra and that the Nakayama automorphisms of a symmetric Frobenius algebra are all inner automorphisms. Then we recall that an augmented symmetric Frobenius algebra is always unimodular. Specializing to Hopf algebras, we recall that finite dimensional Hopf algebras are always Frobenius algebras and that the square  $S \circ S$  of the antipode of an unimodular Hopf algebra is a particular Nakayama automorphism. Finally, we can recall Theorem 60 due to Oberst and Schneider [34], which explains when a Hopf algebra is a symmetric Frobenius algebra. In the proof of Theorem 60, we recall the construction of a non-degenerated trace  $\tau$  on  $H$ . Checking that the diagonal of  $H$  is compatible with this trace  $\tau$ , we obtain Theorem 62.

**8.1. Frobenius algebras.** Let  $A$  be an algebra. The morphism of right  $A$ -modules  $\Theta : A \rightarrow A^\vee$ , mapping 1 to the form  $\phi$ , is an isomorphism (of  $A$ -bimodules) if and only if  $A$  is finite dimensional and the bilinear form  $\langle \ , \ \rangle : A \otimes A \rightarrow \mathbb{F}$  defined by  $\langle a, b \rangle := \phi(ab)$  is non degenerate (and symmetric).

**Definition 54.** An algebra  $A$  is a (symmetric) Frobenius algebra if there exists an isomorphism  $\Theta : A \xrightarrow{\cong} A^\vee$  of right  $A$ -modules (respectively of  $A$ -bimodules). We call  $\phi := \Theta(1)$  a Frobenius form.

*Example 55.* Let  $G$  be a finite group then its group algebra  $\mathbb{F}[G]$  is a non commutative symmetric Frobenius algebra. By definition, the group ring  $\mathbb{F}[G]$  admits the set  $\{g \in G\}$  as a basis. Denote by  $\delta_g$  the dual basis in  $\mathbb{F}[G]^\vee$ . The linear isomorphism  $\Theta : \mathbb{F}[G] \rightarrow \mathbb{F}[G]^\vee$ , sending  $g$  to  $\delta_{g^{-1}}$  is an isomorphism of  $\mathbb{F}[G]$ -bimodules.

Let  $A$  be a Frobenius algebra with Frobenius form  $\phi$ . By definition [23, (16.42)], the *Nakayama automorphism* of  $\phi$  is the unique automorphism of algebras  $\sigma : A \xrightarrow{\cong} A$  such that for all  $a$  and  $b \in A$ ,  $\phi(ab) = \phi(\sigma(b)a)$ . Let  $\sigma$  and  $\sigma'$  be two Nakayama automorphisms of a Frobenius algebra  $A$ . Then, by [23, (16.43)], there exists an invertible element  $u \in A$  such that for all  $x \in A$ ,  $\sigma'(x) = u\sigma(x)u^{-1}$ . In particular, if  $A$  is a symmetric Frobenius algebra, the identity map of  $A$ ,  $id_A : A \rightarrow A$  is a particular Nakayama automorphism of  $A$ . And all the other Nakayama automorphisms are inner automorphisms [26, p. 483 Lemma (b)].

**Definition 56.** Let  $(A, \mu, \eta, \varepsilon)$  be an augmented algebra. A left (respectively right) *integral* in  $A$  is an element  $l$  of  $A$  such that  $\forall h \in A$ ,  $h \times l = \varepsilon(h)l$  (respectively  $l \times h = \varepsilon(h)l$ ). The augmented algebra  $A$  is *unimodular* if the set of left integrals in  $A$  coincides with the set of right integrals in  $A$ .

*Remark 57.* An element  $l$  of  $A$  is a right integral in  $A$  such that  $\varepsilon(l) = 1$  if and only if  $1_A - l$  is a left unit in  $\overline{A}$ , the augmentation ideal of  $A$ . Suppose that there exists a right integral  $l$  in  $A$  such that  $\varepsilon(l) = 1$ . Then  $l$  defines a morphism of right  $A$ -modules  $s : \mathbb{F} \rightarrow A$  such that  $\varepsilon \circ s = id_{\mathbb{F}}$ . Therefore  $\mathbb{F}$  is a right projective  $A$ -module and  $\text{Ext}_A^*(\mathbb{F}, \mathbb{F})$  is concentrated in degree 0 (Compare with [7, Proof of Proposition 5.4]).

The set of right (respectively left) integrals in an augmented Frobenius algebra is a  $\mathbb{F}$ -vector space of dimension 1 [19, Proposition 6.1]. Let  $A$  be an augmented algebra and let  $\Theta : A \xrightarrow{\cong} A^\vee$  be an isomorphism of left (respectively right)  $A$ -modules. Then  $\Theta^{-1}(\varepsilon)$  is non-zero left (respectively right) integral in  $A$  [19, just above Proposition 6.1]. In particular, if  $\Theta : A \xrightarrow{\cong} A^\vee$  is an isomorphism of  $A$ -bimodules,  $\Theta^{-1}(\varepsilon)$  is both a non-zero left et right integral in  $A$ . Therefore a symmetric Frobenius algebra with an augmentation is always unimodular.

Let  $A$  be a Frobenius algebra with an augmentation. Let  $t$  be any non-zero left integral in  $A$ . The *distinguished group-like element* or left modular function [19, (6.2)] in  $A^\vee$  is the unique morphism of algebras  $\alpha : A \rightarrow \mathbb{F}$  such that for all  $h \in A$ ,  $t \times h = \alpha(h)t$  ([33, 2.2.3] or [35, p. 590]). Note that  $A$  is unimodular if and only if the distinguished group-like element in  $A^\vee$  is  $\varepsilon$  the augmentation of  $A$ .

**8.2. Hopf algebras.** Let  $(H, \mu, \eta, \Delta, \varepsilon, S)$  be a finite dimensional Hopf algebra. Its dual is also a Hopf algebra  $(H^\vee, \Delta^\vee, \varepsilon^\vee, \mu^\vee, \eta^\vee, S^\vee)$ . In particular, a form  $\lambda$  on  $H$  is a left (respectively right) integral in  $H^\vee$  if and only if for every  $\varphi \in H^\vee$  and  $k \in H$ ,  $\sum \varphi(k^{(1)})\lambda(k^{(2)}) = \varphi(1_H)\lambda(k)$  (respectively  $\sum \lambda(k^{(1)})\varphi(k^{(2)}) = \varphi(1_H)\lambda(k)$ ). Here  $\Delta k = \sum k^{(1)} \otimes k^{(2)}$ .

*Example 58.* [33, 2.1.2] If  $G$  is a finite group,  $\sum_{g \in G} g$  is both a left and right integral in the group algebra  $\mathbb{F}[G]$ . And  $\delta_1$ , the form mapping  $g \in G$  to 1 if  $g = 1$  and 0 otherwise, is both a left and right integral in  $\mathbb{F}[G]^\vee$ . Since  $\delta_1(1) = 1$ , by Remark 57,  $\text{Cotor}_{\mathbb{F}[G]}^*(\mathbb{F}, \mathbb{F})$  and  $\text{Ext}_{\mathbb{F}[G]^\vee}^*(\mathbb{F}, \mathbb{F})$  are both concentrated in degree 0 (Note that here the product on  $G$  is not used and that  $G$  does not need to be finite [21, 4. p. 97]).

The set of left (respectively right) integrals in the dual Hopf algebra  $H^\vee$  is a  $\mathbb{F}$ -vector space of dimension 1 [38, Corollary 5.1.6 2)]. So let  $\lambda$  be any non-zero left (respectively right) integral in  $H^\vee$ . The morphism of left (respectively right)  $H$ -modules,  $H \xrightarrow{\cong} H^\vee$ , sending  $g$  to the form, denoted [38, p. 95]  $g \mapsto \lambda$ , mapping  $h$  to  $\lambda(hg)$  (respectively to the form mapping  $h$  to  $\lambda(gh)$ ), is an isomorphism [38, Proof of Corollary 5.1.6 2)]. So a finite dimensional Hopf algebra is always a Frobenius algebra, but not always a symmetric Frobenius algebra as Theorem 60 will show.

**Lemma 59.** ([35, Theorem 3(a)] or [19, (6.8)]) *Let  $H$  be a finite dimensional Hopf algebra. Let  $\lambda$  be a non-zero right integral in  $H^\vee$ . Let  $\alpha$  be the distinguished group-like element in  $H^\vee$ . Then for all  $a$  and  $b \in H$ ,*

*i)  $\lambda(ab) = \lambda(S^2(b \leftarrow \alpha)a)$  where  $b \leftarrow \alpha = \sum \alpha(b^{(1)})b^{(2)}$  ([38, p. 95], [35, p. 585] or [19, p. 55]),*

*ii) In the case  $H$  is unimodular,  $\lambda(ab) = \lambda(S^2(b)a)$  [24, Proposition 8].*

We have seen that if  $H$  is unimodular, then  $\alpha = \varepsilon$ . Therefore ii) follows from i). Note that Kadison's distinguished group-like element [19, (6.2) or p. 57]  $m$  in  $H^\vee$  is  $\alpha^{-1} = \alpha \circ S$ , the inverse of ours ([33, 2.2.3] or [19, p. 57]), since he uses right integrals to define it and we use left integrals. Lemma 59 means that the Nakayama automorphism  $\sigma$  of any non-zero right integral  $\lambda$  in  $H^\vee$  is given by  $\sigma(b) = S^2(b \leftarrow \alpha)$  for any  $b \in H$ .

**Theorem 60.** [34, 10, 26, 17] *A finite dimensional Hopf algebra  $H$  is a symmetric Frobenius algebra if and only if  $H$  is unimodular and its antipode  $S$  satisfies  $S^2$  is an inner automorphism of  $H$ .*

*Proof.* Suppose that  $H$  is a symmetric Frobenius algebra. Then we saw that  $H$  is unimodular and that all its Nakayama automorphisms

are inner automorphisms. By ii) of Lemma 59,  $S^2$  is a Nakayama automorphism of  $H$ .

Conversely, assume that  $H$  is unimodular and that  $S^2$  is an inner automorphism of  $H$ . Let  $u$  be an invertible element of  $H$  such that  $\forall h \in H, S^2(h) = uhu^{-1}$ . Let  $\lambda$  be any non-zero right integral in  $H^\vee$ . We saw above that  $\lambda(ab)$  is a non-degenerate bilinear form on  $H$ . By ii) of Lemma 59,  $\lambda(ab) = \lambda(S^2(b)a) = \lambda(ubu^{-1}a)$ . Therefore  $\beta(h, k) := \lambda(uhk)$  is a non-degenerate symmetric bilinear form [26, p. 487 proof of Proposition].  $\square$

*Example 61.* Let  $G$  be a finite group. Since  $S^2 = Id$  and since  $\delta_1$  is a right integral for  $\mathbb{F}[G]^\vee$ , we recover that the linear isomorphism  $\mathbb{F}[G] \rightarrow \mathbb{F}[G]^\vee$ , sending  $g$  to  $\delta_1(g-) = \delta_{g^{-1}}$  is an isomorphism of  $\mathbb{F}[G]$ -bimodules.

The Sweedler algebra is an example of non unimodular Hopf algebra over any field of characteristic different from 2 [33, 2.1.2]. Notice that a cocommutative Hopf algebra over a field of characteristic different from 0 can be non unimodular [26, p. 487-8, Remark and Examples (1) and (4)].

The square of the antipode of every quasi-cocommutative Hopf algebra with bijective antipode is an inner automorphism ([20, Proposition VIII.4.1] or [33, 10.1.4]). Therefore by Theorem 60, every braided (also called quasitriangular) unimodular finite dimensional Hopf algebra is a symmetric Frobenius algebra. In particular, the Drinfeld double  $D(H)$  of any finite dimensional Hopf algebra is a symmetric Frobenius algebra [19, Theorem 6.10].

**Theorem 62.** *Let  $H$  be a finite dimensional unimodular (Definition 56) Hopf algebra equipped with a group-like element  $\sigma$  such that  $\forall h \in H, S^2(h) = \sigma^{-1}h\sigma$ . Then 1)  $H^*(\Phi) : Ext_H^*(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(H, H)$  is an inclusion of Batalin-Vilkovisky algebras.*

*2) Suppose moreover that  $H$  is cosemisimple. Then  $\gamma^* : \widetilde{HC}_{(\varepsilon, \sigma)}^*(H) \rightarrow HC_\lambda^*(H)$  is an inclusion of graded Lie algebras.*

Remark that by [8, Exercice 5.5.10], a finite dimensional cosemisimple Hopf algebra is always unimodular.

*Proof.* By Theorem 16,  $H^*(\Phi) : Ext_H^*(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(H, H)$  is an inclusion of Gerstenhaber algebras. By Theorem 49 (or Corollary 48),  $Ext_H^*(\mathbb{F}, \mathbb{F})$  is a Batalin-Vilkovisky algebra and  $\widetilde{HC}_{(\varepsilon, \sigma)}^*(H)$  has a Lie bracket of degree  $-1$ . By Theorem 60,  $H$  is a symmetric Frobenius algebra. Therefore by Corollary 41,  $HH^*(H, H)$  is a Batalin-Vilkovisky algebra. And by Corollary 42,  $HC_\lambda^*(H)$  has a Lie bracket of degree  $-1$ .

More precisely, let  $\lambda$  be any non-zero right integral in  $H^\vee$ . Let  $\tau : H \rightarrow \mathbb{F}$  given by  $\tau(a) = \lambda(\sigma^{-1}a)$  for all  $a \in H$ . In the proof of Theorem 60, we saw that  $\tau$  is a non degenerate 1-trace. Since  $\lambda$  is right integral in  $H^\vee$ , using the canonical injection of  $H$  into its bidual, for every  $k \in H$ ,  $\lambda(k^{(1)})k^{(2)} = \lambda(k)1_H$ . Here  $\Delta k = k^{(1)} \otimes k^{(2)}$ . By taking  $k = \sigma^{-1}a$ , since  $\sigma^{-1}$  is a group-like element, for all  $a \in H$ ,  $\tau(a^{(1)})\sigma^{-1}a^{(2)} = \lambda(\sigma^{-1}a^{(1)})\sigma^{-1}a^{(2)} = \lambda(\sigma^{-1}a)1_H = \tau(a)1_H$ . This means that  $\tau$  is  $\sigma$ -invariant in the sense of [22, Definition 3.1]. Therefore by applying part 1) of Theorem 52 in the case  $A = H$ , we obtain that  $H^*(\Phi) : \text{Ext}_H(\mathbb{F}, \mathbb{F}) \hookrightarrow HH^*(H, H)$  is a morphism of Batalin-Vilkovisky algebras. This is 1).

By [33, 2.4.6] or [8, Exercice 5.5.9],  $H$  is cosemisimple means that there exists a right integral  $t$  in  $H^\vee$  such that  $t(1) = 1$ . Since the set of right integrals in  $H^\vee$  is a  $\mathbb{F}$ -vector space of dimension 1, any non-zero right integral  $\lambda$  in  $H^\vee$  satisfies  $\lambda(1) \neq 0$ . Since  $\tau(\sigma) = \lambda(\sigma^{-1}\sigma) = \lambda(1)$  is different from zero, by [22, Theorem 3.1], the morphism of graded Lie algebras given by part 2) of Theorem 52,  $\gamma^* : \widetilde{HC}_{(\varepsilon, \sigma)}^*(H) \rightarrow HC_\lambda^*(H)$  is injective. So 2) is proved.  $\square$

Note that by Theorem 60, any Hopf algebra satisfying the hypotheses of Theorem 62 is a symmetric Frobenius algebra. On the contrary, any Hopf algebra which is also a symmetric Frobenius algebra does not necessarily satisfies the hypotheses of Theorem 62. Indeed, in a symmetric Frobenius Hopf algebra,  $S^2$  is an inner automorphism, not necessarily given by a group-like element  $\sigma$ . But in order, to apply Connes-Moscovici (or more precisely its dual Khalkhali-Rangipour-Taillefer) Hopf cyclic cohomology, we have to suppose that  $\sigma$  is a group-like element.

#### REFERENCES

- [1] Tomasz Brzezinski and Robert Wisbauer, *Corings and comodules*, London Mathematical Society Lecture Note Series, vol. 309, Cambridge University Press, Cambridge, 2003.
- [2] M. Chas and D. Sullivan, *String topology*, preprint: math.GT/991159, 1999.
- [3] D. Chataur and L. Menichi, *String topology of classifying spaces*, preprint: math.AT/0801.0174v3, 2008.
- [4] F. Cohen, T. Lada, and J. May, *The homology of iterated loop spaces*, Lecture Notes in Mathematics, vol. 533, Springer-Verlag, 1976.
- [5] A. Connes and H. Moscovici, *Hopf algebras, cyclic cohomology and the transverse index theorem*, Comm. Math. Phys. **198** (1998), no. 1, 199–246.
- [6] Alain Connes and Henri Moscovici, *Cyclic cohomology and Hopf algebra symmetry*, Lett. Math. Phys. **52** (2000), no. 1, 1–28, Conference Moshé Flato 1999 (Dijon).

- [7] Marius Crainic, *Cyclic cohomology of Hopf algebras*, J. Pure Appl. Algebra **166** (2002), no. 1-2, 29–66.
- [8] Sorin Dăscălescu, Constantin Năstăsescu, and Şerban Raianu, *Hopf algebras*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 235, Marcel Dekker Inc., New York, 2001, An introduction.
- [9] Marco A. Farinati and Andrea L. Solotar, *G-structure on the cohomology of Hopf algebras*, Proc. Amer. Math. Soc. **132** (2004), no. 10, 2859–2865.
- [10] Rolf Farnsteiner, *On Frobenius extensions defined by Hopf algebras*, J. Algebra **166** (1994), no. 1, 130–141.
- [11] Y. Félix, L. Menichi, and J.-C. Thomas, *Gerstenhaber duality in Hochschild cohomology*, J. Pure Appl. Algebra **199** (2005), no. 1-3, 43–59.
- [12] Murray Gerstenhaber and Samuel D. Schack, *Algebras, bialgebras, quantum groups, and algebraic deformations*, Contemp. Math., vol. 134, Amer. Math. Soc., 1992, pp. 51–92.
- [13] Murray Gerstenhaber and Alexander A. Voronov, *Homotopy G-algebras and moduli space operad*, Internat. Math. Res. Notices (1995), no. 3, 141–153.
- [14] E. Getzler, *Batalin-Vilkovisky algebras and two-dimensional topological field theories*, Comm. Math. Phys. **159** (1994), no. 2, 265–285.
- [15] Alastair Hamilton and Andrey Lazarev, *Symplectic  $A_\infty$ -algebras and string topology operations*, Amer. Math. Soc. Transl. Ser. 2, vol. 224, Amer. Math. Soc., 2008, pp. 147–157.
- [16] Gerhard P. Hochschild, *Basic theory of algebraic groups and Lie algebras*, Graduate Texts in Mathematics, vol. 75, Springer-Verlag, New York, 1981.
- [17] J. E. Humphreys, *Symmetry for finite dimensional Hopf algebras*, Proc. Amer. Math. Soc. **68** (1978), no. 2, 143–146.
- [18] T. Kadeishvili, *On the cobar construction of a bialgebra*, Homology Homotopy Appl. **7** (2005), no. 2, 109–122 (electronic).
- [19] Lars Kadison, *New examples of Frobenius extensions*, University Lecture Series, vol. 14, American Mathematical Society, Providence, RI, 1999.
- [20] Christian Kassel, *Quantum groups*, Graduate Texts in Mathematics, vol. 155, Springer-Verlag, New York, 1995.
- [21] M. Khalkhali, *Lectures on noncommutative geometry*, preprint: math/0702140 v2, 2007.
- [22] M. Khalkhali and B. Rangipour, *A new cyclic module for Hopf algebras*, K-Theory **27** (2002), no. 2, 111–131.
- [23] T. Y. Lam, *Lectures on modules and rings*, Graduate Texts in Mathematics, vol. 189, Springer-Verlag, New York, 1999.
- [24] Richard Gustavus Larson and Moss Eisenberg Sweedler, *An associative orthogonal bilinear form for Hopf algebras*, Amer. J. Math. **91** (1969), 75–94.
- [25] J. Loday, *Cyclic homology*, second ed., Grundlehren der Mathematischen Wissenschaften, vol. 301, Springer-Verlag, Berlin, 1998.
- [26] Martin Lorenz, *Representations of finite-dimensional Hopf algebras*, J. Algebra **188** (1997), no. 2, 476–505.
- [27] S. Mac Lane, *Categories for the working mathematician*, Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York-Berlin, 1971.
- [28] M. Markl, S. Shnider, and J. Stasheff, *Operads in algebra, topology and physics*, Mathematical Surveys and Monographs, vol. 96, Amer. Math. Soc., 2002.

CONNES-MOSCOVICI CHARACTERISTIC MAP IS A LIE ALGEBRA MORPHISM

- [29] J. Peter May, *A general algebraic approach to Steenrod operations*, Lecture Notes in Mathematics, Vol. 168, Springer, Berlin, 1970, pp. 153–231.
- [30] J. McClure and J. Smith, *A solution of Deligne’s hochschild cohomology conjecture*, Contemp. Math., vol. 293, pp. 153–193, Amer. Math. Soc., 2002.
- [31] L. Menichi, *Batalin-Vilkovisky algebras and cyclic cohomology of Hopf algebras*, *K-Theory* **32** (2004), no. 3, 231–251.
- [32] ———, *Van den Bergh isomorphisms in string topology*, To appear in *J. Non-commut. Geom.*, arXiv:0907.2105, 2009.
- [33] Susan Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics, vol. 82, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993.
- [34] Ulrich Oberst and Hans-Jürgen Schneider, *Über Untergruppen endlicher algebraischer Gruppen*, *Manuscripta Math.* **8** (1973), 217–241.
- [35] David E. Radford, *The trace function and Hopf algebras*, *J. Algebra* **163** (1994), no. 3, 583–622.
- [36] Boris Shoikhet, *Hopf algebras, tetramodules, and n-fold monoidal categories*, 2009.
- [37] Georges Skandalis, *Géométrie non commutative, opérateur de signature transverse et algèbres de Hopf (d’après A. Connes et H. Moscovici)*, *Astérisque* (2002), no. 282, Exp. No. 892, ix, 345–364, Séminaire Bourbaki, Vol. 2000/2001.
- [38] Moss E. Sweedler, *Hopf algebras*, Mathematics Lecture Note Series, W. A. Benjamin, Inc., New York, 1969.
- [39] Rachel Taillefer, *Cyclic homology of Hopf algebras*, *K-Theory* **24** (2001), no. 1, 69–85.
- [40] ———, *Injective Hopf bimodules, cohomologies of infinite dimensional Hopf algebras and graded-commutativity of the Yoneda product*, *J. Algebra* **276** (2004), no. 1, 259–279.
- [41] V. Tourtchine, *On the homology of the spaces of long knots*, *Advances in topological quantum field theory*, NATO Sci. Ser. II Math. Phys. Chem., vol. 179, Kluwer Acad. Publ., Dordrecht, 2004, pp. 23–52.
- [42] ———, *On the other side of the bialgebra of chord diagrams*, *J. Knot Theory Ramifications* **16** (2007), no. 5, 575–629.
- [43] Victor Tourtchine, *Dyer-Lashof-Cohen operations in Hochschild cohomology*, *Algebr. Geom. Topol.* **6** (2006), 875–894 (electronic).
- [44] T. Tradler, *The BV algebra on Hochschild cohomology induced by infinity inner products*, preprint: math.QA/0210150v1, 2002.
- [45] Thomas Tradler and Mahmoud Zeinalian, *Algebraic string operations*, *K-Theory* **38** (2007), no. 1, 59–82.
- [46] G. Whitehead, *Elements of homotopy theory*, Graduate Texts in Mathematics, vol. 61, Springer-Verlag, 1978.

UMR 6093 ASSOCIÉE AU CNRS, UNIVERSITÉ D’ANGERS, FACULTÉ DES SCIENCES, 2 BOULEVARD LAVOISIER, 49045 ANGERS, FRANCE  
*E-mail address:* `firstname.lastname at univ-angers.fr`