



HAL
open science

Practical use of variational principles for modeling water waves

Didier Clamond, Denys Dutykh

► **To cite this version:**

Didier Clamond, Denys Dutykh. Practical use of variational principles for modeling water waves. 2010. hal-00456891v1

HAL Id: hal-00456891

<https://hal.science/hal-00456891v1>

Preprint submitted on 15 Feb 2010 (v1), last revised 8 Dec 2015 (v6)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Practical use of variational principles for modeling water waves

Didier Clamond ^{a,*}

^a*Laboratoire J.-A. Dieudonné, Parc Valrose, 06108 Nice, France*

Denys Dutykh ^b

^b*LAMA, UMR 5127 CNRS, Université de Savoie, 73376 Le Bourget-du-Lac Cedex, France*

Abstract

This paper describes a method for deriving approximate equations for water waves. The method is based on a ‘relaxed’ variational principle, i.e., on a Lagrangian involving as many variables as possible. This formulation is particularly suitable for the construction of approximate water wave models, since it allows more freedom while preserving the variational structure. The advantages of this relaxed formulation are illustrated with various examples in shallow and deep waters, as well as arbitrary depths.

Using subordinate constraints (e.g., irrotationality and free surface impermeability) in various combinations, several model equations are derived, some being well-known, other being new. The models obtained are studied analytically and exact travelling wave solutions are constructed when possible.

Key words: water waves, variational principles, Lagrangian, Hamiltonian, shallow water, deep water

1 Introduction

The water wave problem in fluid mechanics has been known since more than two hundreds years (Craik 2004). The classical mathematical formulation of

* Corresponding author.

Email addresses: diderc@unice.fr (Didier Clamond),
Denys.Dutykh@univ-savoie.fr (Denys Dutykh).

surface gravity waves involves five equations: the irrotationality of the fluid flow, the fluid incompressibility, the bottom and the surface impermeabilities, and the surface isobaricity (Mei 1989). This system of equations cannot be generally solved exactly and, historically, the water wave theory has been developed by constructing various approximations. In shallow water, we have the equations of Korteweg and de Vries (1895), Boussinesq (1871), Benjamin *et al.* (1972), Serre (1953), Green and Naghdi (1976), Camassa and Holm (1993), Degasperis–Procesi (1999), and many other model equations. On finite depth and deep water, there is the celebrated nonlinear Schrödinger equation (Mei 1989) and the equations of Dysthe (1979), Trulsen *et al.* (2000), Kraenkel *et al.* (2005), among others. These equations are most often derived via some perturbation techniques and are thus valid for waves of small amplitude. Moreover, these equations are generally valid for a very limited range of the ratio *wavelength/water depth* and for narrow-banded spectra. However, for many applications it is necessary to use models uniformly valid for all depths and which are accurate for large amplitudes. It is well-known in theoretical physics that variational formulations are tools of choice to derive such approximations when small parameter expansions are inefficient. Variational principles have also the big advantage of ensuring that one can build approximations with optimal ‘fit’ among all the equations defining the problem at hand.

There are two variational formulations for surface waves that are commonly used, namely the Lagrangian of Luke (1967) and the Hamiltonian of Zakharov (1968). Details on the variational formulations for surface waves can be found in review papers, e.g., Radder (1999), Salmon (1988), Zakharov and Kuznetsov (1997). Luke’s Lagrangian assumes that the flow is exactly irrotational, i.e., the Lagrangian involves a velocity potential but not explicitly the velocity components. If in addition the fluid incompressibility and the bottom impermeability are satisfied identically, the equations at the surface can be derived from Zakharov’s Hamiltonian. Thus, both principles naturally assume that the flow is exactly irrotational, as it is the case of the water wave problem formulation, but Zakharov’s Hamiltonian is more constrained than Luke’s Lagrangian. Luke’s and Zakharov’s variational formulations require that part or all of the equations in the bulk of the fluid and at the bottom are satisfied identically, while the remaining relations must be approximated (Craig & Sulem 1993). It is because the irrotationality and incompressibility are mathematically easy to fulfill, that they are chosen to be satisfied identically. Beside simplicity, there are generally no reasons to fulfill irrotationality and/or incompressibility instead of the impermeability or the isobaricity of the free surface, for example.

Variational formulations involving as few dependent variables as possible are often regarded as simpler (Yahalom & Lynden-Bell 2008). It is understandably tempting to solve exactly (i.e., analytically) as many equations as possible in order to ‘improve’ the solution accuracy. This is not always a good idea, however. Indeed, numerical analysis and scientific computing know many ex-

amples when efficient and most used algorithms do exactly the opposite. These so-called *relaxation methods* — e.g., pseudo-compressibility for incompressible fluid flows (Kameyama *et al.* 2005) — have proven to be very effective for stiff problems. The same idea may also apply to analytical approximations. When solving a system of equations, the exact resolution of a few equations does not necessarily ensure that the overall error is reduced: what really matters is that the global error is minimized. Since for water waves it is possible to use a variational formulation, approximations derived from the latter are guaranteed to be optima.

In this study we would like to elucidate the benefit of using relaxed variational methods for the water wave problem. In other words, we illustrate the advantage of using a variational principle involving as many dependent variables as possible. The paper is organized as follows.

In Section 2, Luke’s Lagrangian is *relaxed* to incorporate explicitly more degrees of freedom. This modification yields the Hamilton principle in its most general form. The advantage of this formulation is subsequently illustrated with examples over a fixed horizontal bottom, for the sake of simplicity. We begin in Section 3 with shallow water models, where some well-known and a few novel models are derived from the same Lagrangian, but with different subordinate constraints. In Section 4, similar model equations are derived in the deep water limit. In particular, a generalization of the Klein–Gordon equation and a remarkably simple accurate approximation for traveling waves are derived. A generalized ansatz, including the shallow and deep waters as limiting particular cases is presented in Section 5. Further generalizations are discussed in Section 6 and their advantage is illustrated with a variant of Serre’s equations. Finally, conclusions and perspectives for future studies are outlined in Section 7.

2 Generalized variational formulation

Water wave problem possesses several variational structures (Whitham 1965; Luke 1967; Zakharov 1968). In the present study we will extensively exploit the Lagrangian variational formalism. Surface gravity wave equations, for a potential flow with an impermeable bottom and an impermeable free surface where the pressure is constant (taken to be zero), can be derived minimizing the following functional (Luke 1967):

$$\mathcal{L} = \int_{t_1}^{t_2} \int_{\Omega} \mathcal{L} \rho \, d^2\mathbf{x} \, dt, \quad \mathcal{L} = - \int_{-d}^{\eta} \left[gy + \phi_t + \frac{1}{2}(\nabla \phi)^2 + \frac{1}{2}\phi_y^2 \right] dy, \quad (1)$$

with $\mathbf{x} = (x_1, x_2)$ the horizontal Cartesian coordinates, y the upward vertical coordinate, t the time, ∇ the horizontal gradient, Ω the horizontal domain, ϕ

the velocity potential and $g > 0$ the acceleration due to gravity; $y = \eta(\mathbf{x}, t)$, $y = 0$ and $y = -d(\mathbf{x}, t)$ being, respectively, the equations of the free surface, of the still water level and of the bottom. For the sake of simplicity, the surface tension effect is neglected and the fluid density ρ is constant (and can thus be set to unity without any loss of generality), but this is not a limitation for the purpose of this paper.

Integrating by parts, then neglecting the terms at the horizontal and temporal boundaries because they do not contribute to the minimization (this will be done repeatedly below without explicit mention), Luke's variational formulation (1) can be rewritten with the following Lagrangian density:

$$\mathcal{L} = \tilde{\phi} \eta_t + \check{\phi} d_t - \frac{1}{2} g \eta^2 + \frac{1}{2} g d^2 - \int_{-d}^{\eta} \left[\frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} \phi_y^2 \right] dy, \quad (2)$$

where the over 'tildes' and 'wedges' denote, respectively, the quantities written at the free surface $y = \eta$ and at the bottom $y = -d$. We shall also denote with 'bars' the quantities averaged over the water depth, e.g.

$$\bar{\mathbf{u}}(\mathbf{x}, t) \equiv \frac{1}{\eta(\mathbf{x}, t) + d(\mathbf{x}, t)} \int_{-d(\mathbf{x}, t)}^{\eta(\mathbf{x}, t)} \mathbf{u}(\mathbf{x}, y, t) dy.$$

The variational formulations (1) and (2) impose that any approximation is exactly irrotational, i.e., the choice of an ansatz for ϕ necessarily implies an irrotational motion. Note that the term $\frac{1}{2} g d^2$ in (2) can be omitted because, d being prescribed, it does not contribute to the minimization process.

To give us more freedom while keeping an exact formulation, the variational principle is modified (relaxed) by introducing explicitly the horizontal velocity $\mathbf{u} = \nabla \phi$ and the vertical one $v = \phi_y$. The variational formulation can thus be reformulated with the Lagrangian density

$$\mathcal{L} = \tilde{\phi} \eta_t + \check{\phi} d_t - \frac{1}{2} g \eta^2 - \int_{-d}^{\eta} \left[\frac{1}{2} (\mathbf{u}^2 + v^2) + \boldsymbol{\mu} \cdot (\nabla \phi - \mathbf{u}) + \nu (\phi_y - v) \right] dy, \quad (3)$$

where the Lagrange multipliers $\boldsymbol{\mu}$ and ν have to be determined. By variations with respect of \mathbf{u} and v , one finds at once that $\boldsymbol{\mu} = \mathbf{u}$ and $\nu = v$, implying that (3) becomes

$$\mathcal{L} = \tilde{\phi} \eta_t + \check{\phi} d_t - \frac{1}{2} g \eta^2 + \int_{-d}^{\eta} \left[\frac{1}{2} \mathbf{u}^2 + \frac{1}{2} v^2 - \mathbf{u} \cdot \nabla \phi - v \phi_y \right] dy. \quad (4)$$

The Lagrangian density (3) involves six variables $\{\eta, \phi, \mathbf{u}, v, \boldsymbol{\mu}, \nu\}$, while the simplified Lagrangian (4) involves only four variables $\{\eta, \phi, \mathbf{u}, v\}$ and the original Lagrangian (2) only two (η and ϕ). These additional variables introduce additional freedom in the construction of approximations, thus allowing more subordinate relations to be fulfilled. The Lagrangian density (4) was used by

Kim *et al.* (2001) to derive the ‘irrotational’ Green–Naghdi equations for long waves in shallow water. The more general Lagrangian density (3) provides more flexibility to derive model equations, as illustrated below.

The connection of (3) with the variational formulation of the classical mechanics can be seen applying Green’s theorem to (3) that yields another equivalent variational formulation involving the Lagrangian density

$$\begin{aligned} \mathcal{L} = & (\eta_t + \tilde{\boldsymbol{\mu}} \cdot \nabla \eta - \tilde{\nu}) \tilde{\phi} + (d_t + \check{\boldsymbol{\mu}} \cdot \nabla d + \check{\nu}) \check{\phi} - \frac{1}{2} g \eta^2 \\ & + \int_{-d}^{\eta} \left[\boldsymbol{\mu} \cdot \mathbf{u} - \frac{1}{2} \mathbf{u}^2 + \nu v - \frac{1}{2} v^2 + (\nabla \cdot \boldsymbol{\mu} + \nu_y) \phi \right] dy, \end{aligned} \quad (5)$$

which in deep water limit ($d \rightarrow \infty$) becomes (if ϕ decay faster than y^{-1} as $y \rightarrow -\infty$)

$$\begin{aligned} \mathcal{L} = & (\eta_t + \tilde{\boldsymbol{\mu}} \cdot \nabla \eta - \tilde{\nu}) \tilde{\phi} - \frac{1}{2} g \eta^2 \\ & + \int_{-\infty}^{\eta} \left[\boldsymbol{\mu} \cdot \mathbf{u} - \frac{1}{2} \mathbf{u}^2 + \nu v - \frac{1}{2} v^2 + (\nabla \cdot \boldsymbol{\mu} + \nu_y) \phi \right] dy. \end{aligned}$$

Thus, the Hamilton principle of classical mechanics is recovered in Eulerian description of motion, i.e., the variational principle involves the kinetic energy minus the potential energy plus some constraints for the fluid incompressibility, for the flow irrotationality and for the bottom and surface impermeabilities. In other words, the Lagrangian density (5) is the Hamilton principle in its most general form for irrotational surface gravity waves.

Note that, via the integration by parts, the term $\eta_t \tilde{\phi}$, for example, can be replaced by $-\eta \tilde{\phi}_t$ in all the Lagrangian densities given above, without loss (nor gain) of generality. Note also that the relaxed variational formulations involving (3) and (5) being strictly equivalent, one should use the more convenient one depending on the problem under consideration. Note finally that extensions of (3) and (5) including, e.g., obstacles, surface tensions and stratifications in several homogeneous layers are straightforward generalizations. For instance, to include the surface tension it is sufficient to add the term $-\sigma(\sqrt{1 + (\nabla \eta)^2} - 1)$ into the definition of the Lagrangian density (5), σ being the surface tension coefficient.

The goal in this paper is to illustrate the power of the relaxed variational principle via some simple examples. We shall thus consider, for simplicity, a fixed horizontal bottom and we shall derive various approximate equations for shallow and deep waters. Possibilities for arbitrary depths and some other generalizations will also be discussed.

3 Examples in shallow water

Let consider first the shallow water case, with constant depth for the clarity of exposition. We introduce a realistic ansatz for these waves and then apply several constraints to derive various approximations, some of them being well-known, others being new.

3.1 Choice of a simple ansatz

For a long wave in shallow water, in potential motion on a horizontal impermeable sea bed at $y = -d$, it has long been noticed that the velocity field can be well approximated truncating the following expansion (Lagrange 1781)

$$\mathbf{u} = \tilde{\mathbf{u}} - \frac{1}{2}(y+d)^2 \nabla^2 \tilde{\mathbf{u}} + \frac{1}{24}(y+d)^4 \nabla^4 \tilde{\mathbf{u}} + \dots$$

All Lagrange's followers (e.g., Airy, Boussinesq, Rayleigh and many others) used this type of expansions to derive their respective approximations (Craig 2004). Reviews on shallow water approximations can be found in Bona *et al.* (2002, 2004), Kirby (1997), Madsen and Schäffer (1999), Wu (2001), Dougalis & Mitsotakis (2008), among others.

We consider here a simple ansatz of polynomial type, that is a zeroth-order polynomial in y for ϕ and for \mathbf{u} , and a first-order one for v , i.e., we approximate flows that are nearly uniform along the vertical direction. Our ansatz thus reads

$$\phi \approx \bar{\phi}(\mathbf{x}, t), \quad \mathbf{u} \approx \bar{\mathbf{u}}(\mathbf{x}, t), \quad v \approx (y+d)(\eta+d)^{-1} \tilde{v}(\mathbf{x}, t). \quad (6)$$

Such ansatz are the basis of most shallow water approximations. We have also to introduce suitable ansatz for the Lagrange multiplier $\boldsymbol{\mu}$ and ν . Since $\boldsymbol{\mu} = \mathbf{u}$ and $\nu = v$ for the exact solution, the natural ansatz for the multipliers are

$$\boldsymbol{\mu} \approx \bar{\boldsymbol{\mu}}(\mathbf{x}, t), \quad \nu \approx (y+d)(\eta+d)^{-1} \tilde{\nu}(\mathbf{x}, t). \quad (7)$$

With the ansatz (6) and (7), the Lagrangian density (5) becomes

$$\begin{aligned} \mathcal{L} = & (\eta_t + \bar{\boldsymbol{\mu}} \cdot \nabla \eta) \bar{\phi} - \frac{1}{2} g \eta^2 \\ & + (\eta+d) \left[\bar{\boldsymbol{\mu}} \cdot \bar{\mathbf{u}} - \frac{1}{2} \bar{\mathbf{u}}^2 + \frac{1}{3} \tilde{\nu} \tilde{v} - \frac{1}{6} \tilde{v}^2 + \bar{\phi} \nabla \cdot \bar{\boldsymbol{\mu}} \right]. \end{aligned} \quad (8)$$

Using the Green formula, the variational problem can also be written such that the Lagrangian density is in the following simpler form

$$\mathcal{L} = \bar{\phi} \eta_t - \frac{1}{2} g \eta^2 + (\eta+d) \left[\bar{\boldsymbol{\mu}} \cdot \bar{\mathbf{u}} - \frac{1}{2} \bar{\mathbf{u}}^2 + \frac{1}{3} \tilde{\nu} \tilde{v} - \frac{1}{6} \tilde{v}^2 - \bar{\boldsymbol{\mu}} \cdot \nabla \bar{\phi} \right]. \quad (9)$$

The two Lagrangian densities (8) and (9) differing by a divergence term, they yield exactly the same equations. Thus, depending on the constraints, we use the Lagrangian density leading to the simpler expression. We now investigate equations led by this shallow water model under various subordinate relations.

3.2 Unconstrained approximation

Without further constraints, the minimization of (9) yields

$$\delta \bar{\mathbf{u}} : 0 = \bar{\boldsymbol{\mu}} - \bar{\mathbf{u}}, \quad (10)$$

$$\delta \tilde{\nu} : 0 = \tilde{\nu} - \bar{\nu}, \quad (11)$$

$$\delta \bar{\boldsymbol{\mu}} : 0 = \bar{\mathbf{u}} - \nabla \bar{\phi}, \quad (12)$$

$$\delta \tilde{\nu} : 0 = \tilde{\nu}, \quad (13)$$

$$\delta \bar{\phi} : 0 = \eta_t + \nabla \cdot [(\eta + d) \bar{\boldsymbol{\mu}}], \quad (14)$$

$$\delta \eta : 0 = \bar{\boldsymbol{\mu}} \cdot \bar{\mathbf{u}} - \frac{1}{2} \bar{\mathbf{u}}^2 + \frac{1}{3} \tilde{\nu} \bar{\nu} - \frac{1}{6} \bar{\nu}^2 - \bar{\boldsymbol{\mu}} \cdot \nabla \bar{\phi} - \bar{\phi}_t - g\eta. \quad (15)$$

The relations (10) – (13) imply that the flow is exactly potential, but the fluid incompressibility is not satisfied identically. With these four relations, the last two equations can be rewritten in this form:

$$h_t + \nabla \cdot [h \bar{\mathbf{u}}] = 0, \quad (16)$$

$$\bar{\mathbf{u}}_t + (\bar{\mathbf{u}} \cdot \nabla) \bar{\mathbf{u}} + g \nabla h = 0, \quad (17)$$

where $h = \eta + d$ is the total water depth. Equations (16) – (17) are the very well-known nonlinear shallow water equations, also known as Airy or Saint-Venant equations (Wehausen & Laitone 1960, §28). They are sometimes called *non-dispersive fully-nonlinear approximation* because their classical derivation assumes long waves without the extra hypothesis of small amplitudes.

The Saint-Venant equations do not admit smooth progressive wave solutions. They are nevertheless widely used because they can be solved analytically by the method of characteristics (Stoker 1957). Moreover, numerous efficient finite volumes type schemes have been proposed (Zhou 2002).

3.3 Constraining with free surface impermeability

We now constrain the ansatz (6) imposing that the impermeability of the free surface is satisfied identically. Since the surface impermeability is expressed through the velocity $(\boldsymbol{\mu}, \nu)$ in (5), we substitute

$$\tilde{\nu} = \eta_t + \bar{\boldsymbol{\mu}} \cdot \nabla \eta, \quad (18)$$

into the Lagrangian density (8), and the minimization procedure gives

$$\delta \bar{\mathbf{u}} : 0 = \bar{\boldsymbol{\mu}} - \bar{\mathbf{u}}, \quad (19)$$

$$\delta \tilde{v} : 0 = \eta_t + \bar{\boldsymbol{\mu}} \cdot \nabla \eta - \tilde{v}, \quad (20)$$

$$\delta \bar{\boldsymbol{\mu}} : 0 = \bar{\mathbf{u}} + \frac{1}{3} \tilde{v} \nabla \eta - \nabla \bar{\phi}, \quad (21)$$

$$\delta \bar{\phi} : 0 = \eta_t + \nabla \cdot [(\eta + d) \bar{\boldsymbol{\mu}}], \quad (22)$$

$$\begin{aligned} \delta \eta : 0 = & \bar{\boldsymbol{\mu}} \cdot \bar{\mathbf{u}} - \frac{1}{2} \bar{\mathbf{u}}^2 - \frac{1}{6} \tilde{v}^2 - \bar{\boldsymbol{\mu}} \cdot \nabla \bar{\phi} - \bar{\phi}_t - g \eta \\ & - \frac{1}{3} (\eta + d) [\tilde{v}_t + \bar{\boldsymbol{\mu}} \cdot \nabla \tilde{v} + \tilde{v} \nabla \cdot \bar{\boldsymbol{\mu}}]. \end{aligned} \quad (23)$$

The relations (19) and (21) link the velocity potential and the horizontal velocity as $\nabla \bar{\phi} \neq \bar{\mathbf{u}} = \bar{\boldsymbol{\mu}}$ and, therefore, equations (19) – (23) cannot be derived from Luke’s variational principle. Nevertheless, the relaxed variational principle ensures that the vorticity is minimum. Relations (19) and (22) provide the mass conservation and hence, with (18), the approximation (19) – (23) implies that the fluid incompressibility is fulfilled identically.

Eliminating $\bar{\phi}$, $\bar{\boldsymbol{\mu}}$ and \tilde{v} from the horizontal gradient of (23), the system (19) – (23) becomes

$$h_t + \nabla \cdot [h \bar{\mathbf{u}}] = 0, \quad (24)$$

$$\begin{aligned} \bar{\mathbf{u}}_t + \bar{\mathbf{u}} \cdot \nabla \bar{\mathbf{u}} + g \nabla h + \frac{1}{3} h^{-1} \nabla [h^2 \tilde{\gamma}] = & (\bar{\mathbf{u}} \cdot \nabla h) \nabla (h \nabla \cdot \bar{\mathbf{u}}) \\ & - [\bar{\mathbf{u}} \cdot \nabla (h \nabla \cdot \bar{\mathbf{u}})] \nabla h, \end{aligned} \quad (25)$$

with $h = \eta + d$ and where

$$\tilde{\gamma} = \tilde{v}_t + \bar{\mathbf{u}} \cdot \nabla \tilde{v} = h \left\{ (\nabla \cdot \bar{\mathbf{u}})^2 - \nabla \cdot \bar{\mathbf{u}}_t - \bar{\mathbf{u}} \cdot \nabla [\nabla \cdot \bar{\mathbf{u}}] \right\}, \quad (26)$$

is the fluid vertical acceleration at the free surface.

In the two-dimensional case (one horizontal dimension) the right-hand side of (25) vanishes and the system (24), (25) reduces to the equations first derived by Serre (1953), independently rediscovered by Su and Gardner (1969) and again by Green, Laws and Naghdi (1974). It is sometimes called *weakly-dispersive fully-nonlinear approximation* (Wu 2001). These equations admit a traveling solitary wave solution

$$\eta = a \operatorname{sech}^2 \frac{1}{2} \varkappa (x_1 - ct), \quad c^2 = g(d + a), \quad (\varkappa d)^2 = 3a(d + a)^{-1},$$

which is linearly stable (Li 2002). Note that this solution does not impose any limitation on the wave amplitude, meaning that Serre’s equations are inconsistent for the highest waves. Note also that the Serre equations have a *non-canonical* Hamiltonian structure (Holm 1988; Li 2002).

In three dimensions, equations (24), (25) were called by Kim *et al.* (2001) ‘irrotational’ Green–Naghdi equations. If the right-hand side of (25) is neglected,

we recover the classical Green–Naghdi equations (Green & Naghdi 1976).

3.4 Constraining with incompressibility and partial potential flow I

Here, we restrict the freedom imposing that the pseudo-velocity is related to the horizontal velocity as $\mathbf{u} = \nabla\phi$, and that the fluid incompressibility $\nabla \cdot \mathbf{u} + v_y = 0$ is fulfilled together with the relations $\boldsymbol{\mu} = \mathbf{u}$ and $\nu = v$, i.e., we take

$$\bar{\boldsymbol{\mu}} = \bar{\mathbf{u}}, \quad \tilde{\nu} = \tilde{v}, \quad \bar{\mathbf{u}} = \nabla\bar{\phi}, \quad \tilde{v} = -(\eta + d) \nabla^2\bar{\phi}.$$

These constraints do not impose exact irrotationality because $v \neq \phi_y$. Obviously, we shall derive an approximation which lies “between” the Saint-Venant and Serre equations.

Thus, the Lagrangian density (9) becomes

$$\mathcal{L} = \bar{\phi}\eta_t - \frac{1}{2}g\eta^2 - \frac{1}{2}(\eta + d) (\nabla\bar{\phi})^2 + \frac{1}{6}(\eta + d)^3 (\nabla^2\bar{\phi})^2,$$

and its minimization yields

$$\begin{aligned} \delta\bar{\phi}: 0 &= \eta_t + \nabla \cdot [(\eta + d) \nabla\bar{\phi}] + \frac{1}{3} \nabla^2 [(\eta + d)^3 (\nabla^2\bar{\phi})], \\ \delta\eta: 0 &= \bar{\phi}_t + g\eta + \frac{1}{2} (\nabla\bar{\phi})^2 - \frac{1}{2}(\eta + d)^2 (\nabla^2\bar{\phi})^2. \end{aligned}$$

It seems that these equations have never appeared before in the literature. They are a generalization of the so-called Kaup–Boussinesq (or canonical Boussinesq) equations (Kaup 1975; Kupershmidt 1985) and are thus referred to as the gKB equations. This can be seen noticing that the gKB equations can be derived from the canonical Hamiltonian

$$\int_{\Omega} \left\{ \frac{1}{2}g\eta^2 + \frac{1}{2}(\eta + d) (\nabla\bar{\phi})^2 - \frac{1}{6}(\eta + d)^3 (\nabla^2\bar{\phi})^2 \right\} d^2\mathbf{x}, \quad (27)$$

while the classical Kaup–Boussinesq (cKB) equations are obtained replacing $(\eta+d)^3$ by d^3 in (27) and restricting the resulting Hamiltonian to one horizontal dimension.

The linearized gKB and cKB systems admit the special traveling wave solution

$$\eta = a \cos k(x_1 - ct), \quad c^2 = gd \left(1 - \frac{1}{3}k^2d^2\right), \quad (28)$$

implying that these equations are linearly ill-conditioned ($c^2 < 0$ for $kd > \sqrt{3}$). Moreover, the Hamiltonian (27) is not always positive. However, if, like the cKB, the gKB equations are integrable, they may be a somewhat interesting model for gravity waves in shallow water.

3.5 Constraining with incompressibility and partial potential flow II

So far, all the approximations derived turned out to be such that $\boldsymbol{\mu} = \mathbf{u}$ and $\nu = v$. We propose here a novel approximation that does not satisfies one of these identities and that is an interesting variant of the previous model.

We impose a partially potential flow such that $\boldsymbol{\mu} = \nabla\phi$ and $\nu = \phi_y$, together with the incompressibility condition $\nabla \cdot \mathbf{u} + v_y = 0$ and the condition $\mathbf{u} = \boldsymbol{\mu}$. Thus, substituting the constraints

$$\bar{\boldsymbol{\mu}} = \bar{\mathbf{u}} = \nabla\bar{\phi}, \quad \tilde{\nu} = 0, \quad \tilde{v} = -(\eta + d) \nabla^2\bar{\phi},$$

into the Lagrangian density (9) yields

$$\mathcal{L} = \bar{\phi}\eta_t - \frac{1}{2}g\eta^2 - \frac{1}{2}(\eta + d) (\nabla\bar{\phi})^2 - \frac{1}{6}(\eta + d)^3 (\nabla^2\bar{\phi})^2,$$

and the minimization procedure gives the equations

$$\delta\bar{\phi}: 0 = \eta_t + \nabla \cdot [(\eta + d) \nabla\bar{\phi}] - \frac{1}{3}\nabla^2 [(\eta + d)^3 (\nabla^2\bar{\phi})], \quad (29)$$

$$\delta\eta: 0 = \bar{\phi}_t + g\eta + \frac{1}{2}(\nabla\bar{\phi})^2 + \frac{1}{2}(\eta + d)^2 (\nabla^2\bar{\phi})^2. \quad (30)$$

These equations can be derived from the canonical Hamiltonian

$$\int_{\Omega} \left\{ \frac{1}{2}g\eta^2 + \frac{1}{2}(\eta + d) (\nabla\bar{\phi})^2 + \frac{1}{6}(\eta + d)^3 (\nabla^2\bar{\phi})^2 \right\} d^2\mathbf{x},$$

which is always positive (an interesting feature for modeling water waves). To the linear approximation, equations (29), (30) have the progressive wave solution

$$\eta = a \cos k(x_1 - ct), \quad c^2 = gd(1 + \frac{1}{3}k^2d^2), \quad (31)$$

which is well-behaved (i.e., c^2 is never negative). Comparisons with the gKB equations suggest to call equations (29), (30) *regularized general Kaup-Boussinesq* (rgKB). However, the linear dispersion relation (31) does not correspond to a Taylor expansion around $kd = 0$ of the exact dispersion relation of linear waves (i.e., $c^2 = g \tanh(kd)/k$), while (28) does. Therefore, the rgKB equations are not very interesting for modeling water waves. However, these equations may be of mathematical interest or to model some physical processes other than water waves.

3.6 Other constraints and generalizations

3.6.1 Constraining with incompressibility and potential flow I

In the previous example, we have constructed an approximation such that $\boldsymbol{\mu} = \mathbf{u}$ but $\nu \neq v$. Now, we release the constrain $\boldsymbol{\mu} = \mathbf{u}$ and keep other remaining constraints. Thus, we impose

$$\bar{\boldsymbol{\mu}} = \nabla \bar{\phi}, \quad \tilde{v} = 0, \quad \tilde{v} = -(\eta + d) \nabla \cdot \bar{\mathbf{u}},$$

so that the pseudo velocity field $(\boldsymbol{\mu}, \nu)$ is irrotational while the velocity field (\mathbf{u}, v) is incompressible. After some elementary algebra, the Lagrangian density becomes

$$\mathcal{L} = \bar{\phi} h_t - \frac{1}{2} g \eta^2 + h \bar{\mathbf{u}} \cdot \nabla \bar{\phi} - \frac{1}{2} h \bar{\mathbf{u}}^2 - \frac{1}{6} h^3 (\nabla \cdot \bar{\mathbf{u}})^2 - h (\nabla \bar{\phi})^2,$$

where $h = \eta + d$. The minimization procedure yields

$$\begin{aligned} \delta \bar{\mathbf{u}} : 0 &= h \nabla \bar{\phi} - h \bar{\mathbf{u}} + \frac{1}{3} \nabla [h^3 \nabla \cdot \bar{\mathbf{u}}], \\ \delta \bar{\phi} : 0 &= h_t - \nabla \cdot [h \bar{\mathbf{u}}] + 2 \nabla \cdot [h \nabla \bar{\phi}], \\ \delta \eta : 0 &= \bar{\phi}_t + g \eta + \frac{1}{2} \bar{\mathbf{u}}^2 + (\nabla \bar{\phi})^2 - \bar{\mathbf{u}} \cdot \nabla \bar{\phi} + \frac{1}{2} h^2 (\nabla \cdot \bar{\mathbf{u}})^2. \end{aligned}$$

The linearization of this system of equations have a $(2\pi/k)$ -periodic sinusoidal traveling wave solution with the dispersion relation

$$c^2 = gd(1 + \frac{2}{3}k^2d^2)(1 + \frac{1}{3}k^2d^2)^{-1} = gd(1 + \frac{1}{3}k^2d^2) + O(k^4),$$

which, like the previous example, is not satisfactory for water waves. However, these equations may be of interest in other contexts than water waves.

3.6.2 Constraining with incompressibility and potential flow II

We now assume that the pseudo velocity field $(\boldsymbol{\mu}, \nu)$ is divergence free, while the velocity field (\mathbf{u}, v) is irrotational, i.e., we impose the constraints

$$\bar{\mathbf{u}} = \nabla \bar{\phi}, \quad \tilde{v} = 0, \quad \tilde{v} = -(\eta + d) \nabla \cdot \bar{\boldsymbol{\mu}}.$$

The Lagrangian density becomes

$$\mathcal{L} = \bar{\phi} \eta_t - \frac{1}{2} g \eta^2 - \frac{1}{2} (\nabla \bar{\phi})^2,$$

which, after minimization, yields the Saint-Venant equations. Thus, these constraints do not bring anything new. It should be emphasized that this is the

case for the special shallow water ansatz we are considering here, but this is not necessarily the case for other ansatz.

3.6.3 Further possibilities

The constraints of Sections (3.4) to (3.6.2) can be unified into a single formalism considering combinations. Indeed, the velocity field (\mathbf{u}, v) being not more (nor less) physical than the pseudo-velocity field $(\boldsymbol{\mu}, \nu)$ and the potential velocity field $(\nabla\phi, \phi_y)$, the constraints can be imposed by combinations of these three fields. For instance, we could impose the irrotationality for the field

$$\{ c_1\mathbf{u} + c_2\boldsymbol{\mu} + (1 - c_1 - c_2)\nabla\phi ; c_1v + c_2\nu + (1 - c_1 - c_2)\phi_y \},$$

the fluid incompressibility for the field

$$\{ c_3\mathbf{u} + c_4\boldsymbol{\mu} + (1 - c_3 - c_4)\nabla\phi ; c_3v + c_4\nu + (1 - c_3 - c_4)\phi_y \},$$

and so on for any constraint we may think of. The c_n are parameters at our disposal. We can choose them in a convenient way based on some mathematical and physical considerations. For example, imposing that the approximate equations derived must be linearly well-posed and/or have better dispersion relation properties.

In the examples above, only some kinematic constraints (irrotationality, incompressibility, impermeability) were used. We could have also considered dynamical constraints based on, e.g., the Bernoulli equation, or other relevant dynamical equations.

The relaxed variational principle provided a common platform for deriving several shallow water equations from the same ansatz in changing only the constraints. Beside the ansatz, no further approximation were made and the derivations required only some elementary algebra. Using more general ansatz — i.e., involving more free functions and parameters — one can introduce more constraints, if desired, and derives more accurate approximations. A simple example is given in Section 6 below.

4 Examples in deep water

We illustrate here the advantages of the relaxed variational principle in the opposite limiting case of deep water.

4.1 Choice of an ansatz

For progressive waves in deep water, the Stokes expansion shows that the velocity field varies nearly exponentially along the vertical (Appendix A). Even for very large unsteady waves (including breaking waves), accurate numerical simulations and experiments have shown that the vertical variation of the velocity field is indeed very close to an exponential (Grue *et al.* 2003; Jensen *et al.* 2007). Thus, this property is exploited here to derive simple approximations for waves in deep water.

Let κ be a characteristic wavenumber corresponding, for example, to the carrier wave of a modulated wave group or to the peak frequency of a JONSWAP spectrum. Following the discussion above, it is natural to seek approximations in the form

$$\{\phi; \mathbf{u}; v; \boldsymbol{\mu}; \nu\} \approx \{\tilde{\phi}; \tilde{\mathbf{u}}; \tilde{v}; \tilde{\boldsymbol{\mu}}; \tilde{\nu}\} e^{\kappa(y-\eta)}, \quad (32)$$

where $\tilde{\phi}$, $\tilde{\mathbf{u}}$, \tilde{v} , $\tilde{\boldsymbol{\mu}}$ and $\tilde{\nu}$ are functions of \mathbf{x} and t that will be determined using the variational principle. The ansatz (32) is certainly the simplest possible that is consistent with experimental evidences. This ansatz has already been used by Kraenkel *et al.* (2005) for building their approximation. Possible generalizations are discussed in Section 6.

For the sake of simplicity, we introduce the constraints $\tilde{\boldsymbol{\mu}} = \tilde{\mathbf{u}}$ and $\tilde{\nu} = \tilde{v}$. Thus, the ansatz (32) substituted into the Lagrangian density (4) yields

$$2\kappa \mathcal{L} = 2\kappa \tilde{\phi} \eta_t - g\kappa \eta^2 + \frac{1}{2} \tilde{\mathbf{u}}^2 + \frac{1}{2} \tilde{v}^2 - \tilde{\mathbf{u}} \cdot (\nabla \tilde{\phi} - \kappa \tilde{\phi} \nabla \eta) - \kappa \tilde{v} \tilde{\phi}. \quad (33)$$

With (or without) subordinate relations, this Lagrangian gives various equations. We investigate two cases here.

4.2 Unconstrained approximation

Without further constraints, the minimization procedure yields

$$\begin{aligned} \delta \tilde{\mathbf{u}} : 0 &= \tilde{\mathbf{u}} - \nabla \tilde{\phi} + \kappa \tilde{\phi} \nabla \eta, \\ \delta \tilde{v} : 0 &= \tilde{v} - \kappa \tilde{\phi}, \\ \delta \tilde{\phi} : 0 &= 2\kappa \eta_t + \nabla \cdot \tilde{\mathbf{u}} - \kappa \tilde{v} + \kappa \tilde{\mathbf{u}} \cdot \nabla \eta, \\ \delta \eta : 0 &= 2g\kappa \eta + 2\kappa \tilde{\phi}_t + \kappa \nabla \cdot (\tilde{\phi} \tilde{\mathbf{u}}). \end{aligned}$$

The two first relations imply that this approximation is exactly irrotational and their use in the last two equations gives

$$\eta_t + \frac{1}{2}\kappa^{-1}\nabla^2\tilde{\phi} - \frac{1}{2}\kappa\tilde{\phi} = \frac{1}{2}\tilde{\phi}\left[\nabla^2\eta + \kappa(\nabla\eta)^2\right], \quad (34)$$

$$\tilde{\phi}_t + g\eta = -\frac{1}{2}\nabla\cdot\left[\tilde{\phi}\nabla\tilde{\phi} - \kappa\tilde{\phi}^2\nabla\eta\right]. \quad (35)$$

Since equations (34), (35) derive from an irrotational motion, they can also be obtained from Luke's Lagrangian (2) under ansatz (32). Equations (34), (35) are a deep water counterpart of Saint-Venant equations for shallow water waves; this claim will appear clearer in Section 5. They can also be derived from the canonical Hamiltonian

$$\int_{\Omega}\left\{\frac{1}{2}g\eta^2 + \frac{1}{4}\kappa^{-1}\left[\nabla\tilde{\phi} - \kappa\tilde{\phi}\nabla\eta\right]^2 + \frac{1}{4}\kappa\tilde{\phi}^2\right\}d^2\mathbf{x}. \quad (36)$$

This 'simple' Hamiltonian is quartic in nonlinearities and involves only first-order derivatives. It has to be compared with Zakharov's quartic Hamiltonian (B.2) which involves second-order derivatives and Hilbert transforms. However, Zakharov's quartic Hamiltonian is valid for broad spectra. Note that the Hamiltonian (36) cannot be derived from the exact one (B.1), since the latter assumes that irrotationality and incompressibility are both satisfied identically in the bulk, while the incompressibility is not fulfilled by equations (34), (35).

To the linear approximation, after elimination of $\tilde{\phi}$, equations (34), (35) yield

$$\eta_{tt} - \frac{1}{2}(g/\kappa)\nabla^2\eta + \frac{1}{2}g\kappa\eta = 0, \quad (37)$$

that is a Klein–Gordon equation. For this reason, equations (34), (35) will be referred here as *generalized Klein–Gordon* (gKG). The Klein–Gordon equation is prominent in mathematical physics and appears, e.g., as a relativistic generalization of the Schrödinger equation. The Klein–Gordon equation (37) admits a special $(2\pi/k)$ -periodic traveling wave solution

$$\eta = a \cos k(x_1 - ct), \quad c^2 = \frac{1}{2}g(k^2 + \kappa^2)(\kappa k^2)^{-1}.$$

Therefore, if $k = \kappa$ the exact dispersion relation of linear waves (i.e., $c^2 = g/k$) is recovered, as it should be. This means, in particular, that the gKG model is valid for narrow-banded spectra.

We focus now on $(2\pi/\kappa)$ -periodic progressive waves solution of the gKG equations, i.e., we seek for solutions depending only on the variable $\theta = \kappa(x_1 - ct)$. We were not able to find an exact analytic solution but a Stokes-like expansion gives some interesting insights. To the seventh-order, we have

$$\begin{aligned}
\kappa \eta &= \alpha \cos \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{25}{12} \alpha^2 + \frac{1675}{192} \alpha^4 \right) \cos 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{99}{16} \alpha^2 + \frac{11807}{320} \alpha^4 \right) \cos 3\theta + \frac{1}{3} \alpha^4 \left(1 + \frac{64}{5} \alpha^2 \right) \cos 4\theta \\
&\quad + \frac{125}{384} \alpha^5 \left(1 + \frac{6797}{300} \alpha^2 \right) \cos 5\theta + \frac{27}{80} \alpha^6 \cos 6\theta + \frac{16807}{46080} \alpha^7 \cos 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{3}{2}} \tilde{\phi} &= \alpha \left(1 - \frac{1}{4} \alpha^2 - \frac{59}{96} \alpha^4 - \frac{4741}{1536} \alpha^6 \right) \sin \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{11}{12} \alpha^2 + \frac{547}{192} \alpha^4 \right) \sin 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{163}{48} \alpha^2 + \frac{221}{15} \alpha^4 \right) \sin 3\theta + \frac{1}{3} \alpha^4 \left(1 + \frac{149}{20} \alpha^2 \right) \sin 4\theta \\
&\quad + \frac{125}{384} \alpha^5 \left(1 + \frac{5057}{375} \alpha^2 \right) \sin 5\theta + \frac{27}{80} \alpha^6 \sin 6\theta + \frac{16807}{46080} \alpha^7 \sin 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{1}{2}} c &= 1 + \frac{1}{2} \alpha^2 + \frac{1}{2} \alpha^4 + \frac{899}{384} \alpha^6 + O(\alpha^8).
\end{aligned}$$

The expansions of η and $\tilde{\phi}$ match the exact Stokes wave (c.f. Appendix A) up to the third-order (non-matching coefficients are displayed bold). This is not surprising since the gKG equations are cubic in nonlinearities. A bit more surprising is that the phase velocity c is correct up to the fifth-order. But the most interesting is that, to the leading order, the n^{th} Fourier coefficient is (for all n up to infinity)

$$\frac{n^{n-2} \alpha^n}{2^{n-1} (n-1)!}, \quad (38)$$

which is also the case for the exact Stokes wave (Appendix A).

In comparison, for the cubic Zakharov equations (B.3), (B.4) the phase velocity is correct only up to the third-order and the Fourier coefficients do not verify the asymptotic behavior (38) (see Appendix B). Truncating Zakharov's Hamiltonian at the order $N + 1$ in nonlinearities, the corresponding Stokes double series is correct up to the order N in the expansion parameter. But none of these higher approximations have the exact asymptotic behavior (38) for their Fourier coefficients because they involve expansions around $\eta = 0$, while the gKG does not.

4.3 Constraining with the free surface impermeability

In order to satisfy the free surface impermeability identically, we take

$$\tilde{v} = \eta_t + \tilde{\mathbf{u}} \cdot \nabla \eta,$$

and the Lagrangian density (33) becomes

$$2\kappa \mathcal{L} = \tilde{\phi} (\kappa \eta_t + \nabla \cdot \tilde{\mathbf{u}}) - g\kappa \eta^2 + \frac{1}{2} \tilde{\mathbf{u}}^2 + \frac{1}{2} (\eta_t + \tilde{\mathbf{u}} \cdot \nabla \eta)^2, \quad (39)$$

while the minimization procedure yields the equations

$$\delta \tilde{\mathbf{u}} : 0 = \tilde{\mathbf{u}} + (\eta_t + \tilde{\mathbf{u}} \cdot \nabla \eta) \nabla \eta - \nabla \tilde{\phi}, \quad (40)$$

$$\delta \tilde{\phi} : 0 = \kappa \eta_t + \nabla \cdot \tilde{\mathbf{u}}, \quad (41)$$

$$\delta \eta : 0 = 2g\kappa\eta + \kappa\tilde{\phi}_t + \eta_{tt} + (\tilde{\mathbf{u}} \cdot \nabla \eta)_t + \nabla \cdot (\tilde{\mathbf{u}}\eta_t) + \nabla \cdot [(\tilde{\mathbf{u}} \cdot \nabla \eta)\tilde{\mathbf{u}}]. \quad (42)$$

The relation (41) implying that $\nabla \cdot \mathbf{u} + v_y = 0$, the solution satisfies the incompressibility identically. On the other hand, the irrotationality being not verified identically, equations (40) – (42) cannot be derived from Luke’s variational formulation. Note that (40) yields $\nabla \tilde{\phi} = \tilde{\mathbf{u}} + \tilde{v} \nabla \eta$ that is exact for potential flows, meaning that this ‘incompressible’ approximation is “potential at the free surface”.

As for the shallow water case, the potential $\tilde{\phi}$ can be eliminated from equations (40), thus yielding a deep water analog of the Serre’s and Green–Naghdi equations.

To the linear approximation, relations (40) – (42) can be combined into a single equation for the elevation of the free surface:

$$(\nabla^2 - \kappa^2) \eta_{tt} + 2g\kappa \nabla^2 \eta = 0,$$

which admits the special $(2\pi/k)$ -periodic solution

$$\eta = a \cos k(x_1 - ct), \quad c^2 = 2g\kappa(k^2 + \kappa^2)^{-1}.$$

Therefore, if $k = \kappa$ the exact linear approximation is recovered, as it should be. Again, this means that this model is valid for narrow-banded spectra. How narrow will be investigated now.

4.3.1 Two-dimensional progressive waves

We seek now exact solutions depending only on $\xi \equiv x_1 - ct$ (two-dimensional progressive waves). The equations (40) and (41) yield respectively

$$\tilde{\phi}_\xi = \tilde{u} + (\tilde{u} - c) \eta_\xi^2, \quad \tilde{u} = c\kappa\eta + cK_1,$$

where K_1 is an integration constant. Substituting these relations into (42) and multiplying the result by η_ξ , after one integration and some algebra, we obtain

$$\left(\frac{d\eta}{d\xi} \right)^2 = \frac{K_0^2 - (\kappa\eta - K_1 \hat{\varepsilon}^{-2})^2}{\hat{\varepsilon}^{-2} (\kappa\eta + K_1 - 1)^2}, \quad (43)$$

with

$$\hat{\varepsilon}^2 \equiv 2g\kappa^{-1}c^{-2} - 1,$$

and where K_0 is another integration constant. Assuming that $\eta_\xi = 0$ at the wave crest where $\eta = a$ and at the trough where $\eta = -b$ (hence $a+b$ is the total wave height), we find

$$K_0 = \frac{1}{2} \kappa (a + b), \quad K_1 = \frac{1}{2} \kappa (a - b) \hat{\varepsilon}^2.$$

Solving equation (43), together with the condition of zero mean free surface elevation, we obtain an exact $(2\pi/k)$ -periodic solution in the parametric form (Geniet 2003):

$$k \xi(\tau) = \tau - \varepsilon \hat{\varepsilon}^{-1} \sin \tau, \quad k \eta(\tau) = \frac{1}{2} \varepsilon^2 \hat{\varepsilon}^{-1} + \varepsilon \cos \tau, \quad (44)$$

with

$$\varepsilon = \frac{1}{2} k (a + b), \quad k \kappa^{-1} = \hat{\varepsilon} + \frac{1}{2} (\hat{\varepsilon}^{-1} + \hat{\varepsilon}) \varepsilon^2,$$

ε being a wave steepness and τ being a parametric variable such that a crest is at $\tau = 0$ and the closest troughs at $\tau = \pm\pi$. This remarkably simple solution describes surface waves as trochoids and is similar (but not identical) to the well-known Gerstner wave (Wehausen & Laitone 1960, §34- β). Smooth surface profiles are obtained for $0 \leq \varepsilon < \hat{\varepsilon}$, $\varepsilon \ll \hat{\varepsilon}$ corresponding to quasi-sinusoidal solutions (infinitesimal waves). For the limiting case $\varepsilon = \hat{\varepsilon}$, in the vicinity of the crest $k(a-\eta) \sim \frac{1}{2} \varepsilon (6k\xi)^{\frac{2}{3}}$ so the solution involves a sharp angle forming a 0° inner angle (i.e., a cusp), while the exact angle should be 120° .

So far, κ is a free parameter at our disposal. We shall now investigate various choices.

4.3.2 Simple approximation

A ‘natural’ choice is to take $\kappa = k$, yielding the steepness $\hat{\varepsilon} \approx 0.596$ for the limiting wave. With this peculiar choice of κ , a Stokes-like expansion of the solution of (44) is

$$\begin{aligned} k \eta &= \alpha \cos k\xi + \frac{1}{2} \alpha^2 \left(1 + \frac{13}{12} \alpha^2 + \frac{395}{192} \alpha^4 \right) \cos 2k\xi \\ &\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{35}{16} \alpha^2 + \frac{1727}{320} \alpha^4 \right) \cos 3k\xi + \frac{1}{3} \alpha^4 \left(1 + \frac{33}{10} \alpha^2 \right) \cos 4k\xi \\ &\quad + \frac{125}{384} \alpha^5 \left(1 + \frac{53}{12} \alpha^2 \right) \cos 5k\xi + \frac{27}{80} \alpha^6 \cos 6k\xi + \frac{16807}{46080} \alpha^7 \cos 7k\xi + O(\alpha^8), \\ \sqrt{k/gc} &= 1 + \frac{1}{2} \alpha^2 + \frac{1}{2} \alpha^4 + \frac{611}{384} \alpha^6 + O(\alpha^8). \end{aligned}$$

As for the gKB approximation, the Fourier coefficients of this Stokes-like expansion satisfy the asymptotic expression (38) and the expansion of c is exact

up to the fifth-order. Therefore, taking $\kappa = k$ leads to a quite accurate approximation. However, the optimum value of κ leads to an even more interesting approximation.

4.3.3 Optimum parameter

With the trochoidal solution and the relation between the parameters obtained above, the Lagrangian density (39) integrated over one wavelength yields the Lagrangian

$$\mathcal{L} = \frac{\pi g \varepsilon^4}{4 k^3} = \frac{\pi g}{k^3} \left[-1 + \frac{\kappa c^2}{2g} + \sqrt{\frac{k^2 c^2}{2g\kappa} - \frac{k^2 c^4}{4g^2}} \right]^2.$$

Thus, the Lagrangian is minimum when the steepness is minimum or, equivalently, the wavelength is maximum if the wave height is kept constant. The variational principle being defined with fixed horizontal and temporal boundaries, the optimum parameter κ is obtained minimizing \mathcal{L} keeping k and c constant. After some algebra, the equation $d\mathcal{L}/d\kappa = 0$ gives two solutions for κ :

$$\kappa_{\pm} = \sqrt{2} k \left[1 + 2\varepsilon^2 \pm \sqrt{1 - 4\varepsilon^2} \right]^{-\frac{1}{2}},$$

provided that $\varepsilon \leq 1/2$. Both solutions correspond to an extremum of \mathcal{L} . However, a solution is stable only if the optimal κ is a minimum of \mathcal{L} , i.e., if

$$\left. \frac{d^2 \mathcal{L}}{d\kappa^2} \right|_{\kappa=\kappa_{\pm}} > 0,$$

which, after some algebra, yields the condition

$$\sqrt{3} \varepsilon \kappa_{\pm} < k. \quad (45)$$

The first solution

$$\kappa_+ / k = \sqrt{2} \left[1 + 2\varepsilon^2 + \sqrt{1 - 4\varepsilon^2} \right]^{-\frac{1}{2}} = 1 + \frac{1}{2} \varepsilon^4 + \mathcal{O}(\varepsilon^6),$$

is very close to the wavenumber k when the steepness ε is small. When the steepness increases from 0 to 1/2, the dimensionless parameters κ/k , kc^2/g and $\sqrt{3}\varepsilon\kappa/k$ increase monotonically. The highest wave is obtained for $\varepsilon = 1/2$, where $\kappa/k = 2/\sqrt{3} \approx 1.15$, $kc^2/g = 3\sqrt{3}/4 \approx 1.3$ and $\sqrt{3}\varepsilon\kappa/k = 1$. All these waves are smooth and stable because the condition (45) is fulfilled.

The second solution

$$\kappa_- / k = \sqrt{2} \left[1 + 2\varepsilon^2 - \sqrt{1 - 4\varepsilon^2} \right]^{-\frac{1}{2}},$$

is admissible (i.e., real) if $1/2 \geq \varepsilon \geq \sqrt{\sqrt{5} - 2} \approx 0.486$. As the steepness decays from $1/2$ to $\sqrt{\sqrt{5} - 2}$, the crest sharpens, the limiting value $\varepsilon = \hat{\varepsilon} = \sqrt{\sqrt{5} - 2}$ corresponding to a cusp at the crest. Thus the sharp-crested wave is not the highest one. All these waves are unstable because the condition (45) is violated.

4.3.4 Remarks

Taking the optimum parameter κ has improved the accuracy of the approximation. The main gain is qualitative, however. Indeed, we found that waves are unstable before a sharp crest is formed. Such behavior is not predicted by low-order perturbation expansions. This simple example is a remarkable illustration of the power of the variational method. The trochoidal wave described here is probably the approximation with the highest ratio *accuracy/complexity* ever derived for a traveling wave in deep water. This approximation has first been derived by Geniet (2003) via a different route.

5 Arbitrary depth

A general ansatz, for waves in finite constant depth and satisfying identically the bottom impermeability, is suggested by the linear theory of water waves:

$$\begin{aligned} \phi &\approx \frac{\cosh \kappa Y}{\cosh \kappa h} \tilde{\phi}(\mathbf{x}, t), & \mathbf{u} &\approx \frac{\cosh \kappa Y}{\cosh \kappa h} \tilde{\mathbf{u}}(\mathbf{x}, t), & v &\approx \frac{\sinh \kappa Y}{\sinh \kappa h} \tilde{v}(\mathbf{x}, t), \\ & & \boldsymbol{\mu} &\approx \frac{\cosh \kappa Y}{\cosh \kappa h} \tilde{\boldsymbol{\mu}}(\mathbf{x}, t), & \nu &\approx \frac{\sinh \kappa Y}{\sinh \kappa h} \tilde{\nu}(\mathbf{x}, t), \end{aligned} \quad (46)$$

where $Y = y + d$ and $h = \eta + d$. The parameter κ is a characteristic wave number to be made precise *a posteriori*. This ansatz is uniformly valid for all depths because it yields the shallow water one (6) as $\kappa \rightarrow 0$, and the deep water one (32) as $d \rightarrow \infty$. Obviously, the ansatz (46) is valid for wave fields with wavenumber spectra that are narrow-banded around κ . Substituting the ansatz (46) into the relaxed variational principle (5), we obtain

$$\begin{aligned} \mathcal{L} &= [\eta_t + \tilde{\boldsymbol{\mu}} \cdot \nabla \eta] \tilde{\phi} - \frac{1}{2} g \eta^2 + [\tilde{\nu} \tilde{v} - \frac{1}{2} \tilde{v}^2] \frac{\sinh(2\kappa h) - 2\kappa h}{2\kappa \cosh(2\kappa h) - 2\kappa} \\ &+ [\tilde{\boldsymbol{\mu}} \cdot \tilde{\mathbf{u}} - \frac{1}{2} \tilde{\mathbf{u}}^2 + \tilde{\phi} \nabla \cdot \tilde{\boldsymbol{\mu}} - \kappa \tanh(\kappa h) \tilde{\phi} \tilde{\boldsymbol{\mu}} \cdot \nabla \eta] \frac{\sinh(2\kappa h) + 2\kappa h}{2\kappa \cosh(2\kappa h) + 2\kappa} \\ &+ \frac{1}{2} \tilde{\phi} \tilde{\nu} \left[\frac{2\kappa h}{\sinh(2\kappa h)} - 1 \right]. \end{aligned}$$

Applying various constraints, one obtains generalized equations including the ones derived in Sections 3 and 4 as limiting cases. In particular, we can derive

arbitrary depth generalizations of the Saint-Venant and ‘irrotational’ Green–Naghdi equations. Thus, the relaxed variational principle leads simple derivations of model equations for finite amplitude waves in arbitrary depth, which is not the case with the classical perturbations techniques. These developments are left to future investigations.

6 Generalizations

The ansatz (46) can be generalized in many relevant ways, depending on the problem under consideration. Natural generalizations can be based on the velocity vertical variations given by, e.g., higher-order deep and shallow water theories, or obtained fitting some experimental data. In this section we propose a possible generalization based on an ansatz of the form

$$\phi \approx \left[\frac{\cosh \kappa Y}{\cosh \kappa h} \right]^\lambda \tilde{\phi}(\mathbf{x}, t), \quad \mathbf{u} \approx \left[\frac{\cosh \kappa Y}{\cosh \kappa h} \right]^\lambda \tilde{\mathbf{u}}(\mathbf{x}, t), \quad v \approx \left[\frac{\sinh \kappa Y}{\sinh \kappa h} \right]^\lambda \tilde{v}(\mathbf{x}, t), \quad (47)$$

where λ is a parameter at our disposal. If $\lambda = 1$, the ansatz (46) is recovered, but the case $\lambda \neq 1$ does not correspond to the vertical profile predicted by any theory based on perturbation expansions. Still, this type of ansatz is of some interest, as we shall illustrate it below.

Note first that in the deep water limit $d \rightarrow \infty$, the ansatz (47) becomes

$$\phi \approx e^{\lambda\kappa(y-\eta)} \tilde{\phi}(\mathbf{x}, t), \quad \mathbf{u} \approx e^{\lambda\kappa(y-\eta)} \tilde{\mathbf{u}}(\mathbf{x}, t), \quad v \approx e^{\lambda\kappa(y-\eta)} \tilde{v}(\mathbf{x}, t),$$

and thus, via the change of parameter $\lambda\kappa \mapsto \kappa$, the ansatz (32) is recovered. This means that (47) is not more general than (46) in deep water. On the contrary, these two ansatz are very different in finite depth. We illustrate this claim in the simple case of shallow water ($\kappa h \rightarrow 0$) when (47) becomes

$$\phi \approx \bar{\phi}(\mathbf{x}, t), \quad \mathbf{u} \approx \bar{\mathbf{u}}(\mathbf{x}, t), \quad v \approx \left[\frac{y+d}{\eta+d} \right]^\lambda \tilde{v}(\mathbf{x}, t),$$

where we have replaced $\tilde{\phi}$ and $\tilde{\mathbf{u}}$ by $\bar{\phi}$ and $\bar{\mathbf{u}}$, respectively, since they are equal in this limiting case.

6.1 Modified Serre’s equations

For the sake of simplicity, we consider here only one horizontal dimension, say x_1 , and we set $x_1 = x$ and $u_1 = u$, for brevity. We also consider the special

case $\mu = u$ and $\nu = v$ together with the constraint $\tilde{v} = \eta_t + \tilde{u}\eta_x$ (free surface's impermeability). Thus, the Lagrangian density (5) yields

$$\mathcal{L} = (\eta_t + [(\eta + d)\bar{u}]_x) \tilde{\phi} - \frac{1}{2}g\eta^2 + \frac{1}{2}(\eta + d)\bar{u}^2 + \frac{1}{2}\beta(\eta + d) [\eta_t + \bar{u}\eta_x]^2, \quad (48)$$

where $\beta = (2\lambda + 1)^{-1}$. After some algebra, the minimization procedure leads to the following equations

$$h_t + [h\bar{u}]_x = 0, \quad (49)$$

$$\bar{u}_t + \bar{u}\bar{u}_x + g h_x + \beta h^{-1} [h^2 \tilde{\gamma}]_x = 0, \quad (50)$$

where $\tilde{\gamma}$ is defined in (26). If $\beta = \frac{1}{3}$ the classical Serre equations are recovered.

Equations (49), (50) admit a $(2\pi/k)$ -periodic cnoidal traveling wave solution:

$$\bar{u} = \frac{c\eta}{d + \eta}, \quad (51)$$

$$\eta = a \frac{\operatorname{dn}^2\left(\frac{1}{2}\varkappa(x - ct)|m\right) - E/K}{1 - E/K} = a - H \operatorname{sn}^2\left(\frac{1}{2}\varkappa(x - ct)|m\right), \quad (52)$$

dn and sn being elliptic functions of Jacobi of parameter m ($0 \leq m \leq 1$), and where $K = K(m)$ and $E = E(m)$ are the complete elliptic integrals of the first and second kinds, respectively (Abramowitz & Stegun 1965, #17.3). The parameter \varkappa is a sort of wavenumber, a is the wave amplitude (mean level to crest elevation), H is the total wave height (trough to crest elevation) and c is the wave phase velocity observed in the frame of reference without mean flow. The wave parameters are related via the relations

$$k = \frac{\pi \varkappa}{2K}, \quad H = \frac{m a K}{K - E}, \quad (\varkappa d)^2 = \frac{g H}{m \beta c^2}, \quad (53)$$

$$m = \frac{g H (d + a) (d + a - H)}{g (d + a)^2 (d + a - H) - d^2 c^2}. \quad (54)$$

In the limiting case $m \rightarrow 1$, we have $K \rightarrow \infty$, $E/K \rightarrow 0$, $k \rightarrow 0$, $H \rightarrow a$ and hence, the classical solitary wave solution is recovered

$$\eta = a \operatorname{sech}^2 \frac{1}{2} \varkappa (x - ct), \quad c^2 = g(d + a), \quad \frac{a}{d} = \frac{\beta (\varkappa d)^2}{1 - \beta (\varkappa d)^2}.$$

At this stage, β is still a free parameter. An optimum expression for this parameter can be obtained substituting the solution (51), (52) into the Lagrangian density (48), integrating \mathcal{L} over one wavelength, then solving $d\mathcal{L}/d\beta = 0$

keeping k and c constant (as well as g and d), the other parameters varying according to relations (53), (54). Thus, after some cumbersome algebra, we found that $\beta = 0$ is the optimum value for this parameter, which is not very interesting for practical applications.

Nonetheless, the main objective of this section is to show that one can derive sensible equations from ansatz not deriving from some classical approximation procedure. This is another illustration of the power of the variational procedure.

7 Discussion

In this study, we have illustrated the advantage of using a variational principle with as many variables as possible. We call it the *relaxed* variational principle, since the Lagrangian density (5) involves more degrees of freedom (i.e., the variables η , ϕ , \mathbf{u} , v , $\boldsymbol{\mu}$ and ν) compared to the two degrees of freedom (η and ϕ) in the classical case. In particular, these extra variables can be used to impose various constraints such that incompressibility, irrotationality, impermeability, etc. The practical use of the relaxed formulation was illustrated on numerous examples in shallow, deep and intermediate waters. Thus, we obtained several approximations, some well-known, some new to our knowledge.

In the shallow water regime, we have first obtained the classical nonlinear shallow water (or Saint-Venant) equations (16), (17). Then, with the same ansatz (6) but imposing the constraint of the free surface impermeability, we have derived the irrotational Green–Naghdi equations (24), (25). Applying the incompressibility constraint and choosing differently the pseudo-velocity field, we have obtained two kinds of *generalized Kaup–Boussinesq* equations. Several ways of further generalizations were also outlined.

In deep water, two models were considered. Namely, we derived deep water counterparts of the celebrated Saint-Venant and Serre equations. The former has a canonical Hamiltonian formulation and degenerates to the Klein–Gordon equation in the linear approximation; we thus called the new system (34), (35) *generalized Klein–Gordon* equations. The latter could be solved analytically for a two-dimensional traveling wave. This solution is a striking illustration of the power of the variational formulation compared to asymptotic expansions, specially for large amplitudes. In addition, both equations were shown to possess excellent asymptotic properties with respect to Stokes-like expansions.

The case of arbitrary depth has also been briefly considered. In particular, it has been shown how easily one can introduce an ansatz valid for all depths. Indeed, the vertical variation of the velocity field suggested by the linear theory

provides at once such a general ansatz which degenerates to previous cases when the water is shallow ($\kappa d \rightarrow 0$) or deep ($d \rightarrow \infty$). This simplicity and flexibility of the variational principle is quite remarkable.

Finally, we considered a generalized ansatz and we illustrated its consequence in the limiting case of shallow water. In this way, we derived the *modified Serre equations* and subsequently obtained exact cnoidal and solitary wave solutions. The main purpose of this example was to illustrate the fact that one can introduce an ansatz which is not inspired by any asymptotic expansion and nevertheless lead to good approximations.

In the present paper, some further possibilities for generalizations are also mentioned. However, we have to emphasize that not all ansatz and constraints will necessarily lead to physically relevant and tractable approximations (the same is true for models derived from asymptotic expansions.) Nonetheless, the relaxed variational formulation is sufficiently versatile to allow easy derivations of physically sound models. We have illustrated this claim, in particular, by showing how it is simple to obtain approximate equations valid for all depths.

In order to derive approximate models, variational formulations are more efficient than asymptotic expansions. However, both approaches can be also combined. Indeed, once the variational principle has been applied to an ansatz, asymptotic expansions can be further applied to obtain simpler models. For instance, one could consider ‘unidirectionalized’ approximations (Olver 1984, 1988) to derive variants of Korteweg and de Vries (1895), Dysthe (1979), Camassa and Holm (1993), Degasperis and Procesi (1999), Kraenkel *et al.* (2005), and other equations. This possibility will be investigated in future works.

For the sake of simplicity, we have considered only gravity waves propagating at the surface of a single layer of a homogeneous fluid with a horizontal bottom. It is trivial to introduce a relaxed variational formulation including, e.g., surface tension, stratifications in several homogeneous layers and obstacles. Such a general variational formulation, together with relevant ansatz and well chosen constraints, will easily lead to interesting models. For perfect fluids, variational formulations can also be obtained for rotational motions (Eckart 1960; Luke 1967; Salmon 1988; Morrison 1998; Constantin *et al.* 2006). A relaxed version of such variational principles will facilitate the derivation of approximate models.

The numerical models for simulating water waves are undergoing constant improvements. The state of the art can be found in recent reviews (e.g., Fenton 1999; Dias & Bridges 2006; Ma 2010). Certainly, the variational principle is the tool of choice to derive efficient approximations. By efficient, we mean models that capture most of the relevant physics and which, in the same time, can be easily and rapidly solved numerically. One interesting direction for future

researches is the development of numerical schemes preserving hamiltonian structure at the discrete level. Another interesting application is the derivation of new improved models with uneven bathymetry for coastal hydrodynamics and tsunami wave modeling (Synolakis & Bernard 2006).

The use of a variational principle for modeling surface waves is by no mean new. However, its power has not yet been fully exploited. The present paper is a further contribution in this direction.

Acknowledgement

The second author would like to acknowledge the support from ANR MathOcéan (Project n° ANR-08-BLAN-0301-01) and from the program “Risques gravitaires, séismes” of Cluster Environnement and the research network VOR.

A Exact Stokes wave

In deep water, a seventh-order Stokes expansion (for the exact equations) is

$$\begin{aligned}
\kappa \eta &= \alpha \cos \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{17}{12} \alpha^2 + \frac{233}{64} \alpha^4 \right) \cos 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{51}{16} \alpha^2 + \frac{3463}{320} \alpha^4 \right) \cos 3\theta + \frac{1}{3} \alpha^4 \left(1 + \frac{307}{60} \alpha^2 \right) \cos 4\theta \\
&\quad + \frac{125}{384} \alpha^5 \left(1 + \frac{10697}{1500} \alpha^2 \right) \cos 5\theta + \frac{27}{80} \alpha^6 \cos 6\theta + \frac{16807}{46080} \alpha^7 \cos 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{3}{2}} \tilde{\phi} &= \alpha \left(1 - \frac{1}{4} \alpha^2 - \frac{43}{96} \alpha^4 - \frac{2261}{1536} \alpha^6 \right) \sin \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{7}{12} \alpha^2 + \frac{81}{64} \alpha^4 \right) \sin 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{281}{144} \alpha^2 + \frac{5813}{1080} \alpha^4 \right) \sin 3\theta + \frac{1}{3} \alpha^4 \left(1 + \frac{431}{120} \alpha^2 \right) \sin 4\theta \\
&\quad + \frac{125}{384} \alpha^5 \left(1 + \frac{3369}{625} \alpha^2 \right) \sin 5\theta + \frac{27}{80} \alpha^6 \sin 6\theta + \frac{16807}{46080} \alpha^7 \sin 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{1}{2}} c &= 1 + \frac{1}{2} \alpha^2 + \frac{1}{2} \alpha^4 + \frac{707}{384} \alpha^6 + O(\alpha^8),
\end{aligned}$$

where $\theta = x - ct$. Note that, to the leading order, the n -th Fourier coefficient is $2^{1-n} n^{n-2} \alpha^n / (n-1)!$ (this is also true for all $n > 7$). In the bulk of the fluid, the velocity potential is

$$\begin{aligned}
g^{-\frac{1}{2}} \kappa^{\frac{3}{2}} \phi &= \alpha \left(1 - \frac{1}{8} \alpha^2 - \frac{7}{12} \alpha^4 - \frac{14761}{9216} \alpha^6 \right) e^{\kappa y} \sin \theta + \frac{1}{2} \alpha^4 \left(1 + \frac{11}{6} \alpha^2 \right) e^{2\kappa y} \sin 2\theta \\
&\quad + \frac{1}{12} \alpha^5 \left(1 + \frac{191}{24} \alpha^2 \right) e^{3\kappa y} \sin 3\theta + \frac{1}{72} \alpha^6 e^{4\kappa y} \sin 4\theta \\
&\quad + \frac{1}{480} \alpha^7 e^{5\kappa y} \sin 5\theta + O(\alpha^8),
\end{aligned}$$

meaning that harmonics appear at the fourth-order only, thus justifying the ansatz (32). Note that, to the leading order, the n -th Fourier coefficient is $\alpha^{n+2}/n!(n-1)$ for all $n > 1$.

B Cubic Zakharov's equations

Satisfying exactly the Laplace equation and the bottom impermeability, the gravity waves variational formulation (Luke 1967) yields the Hamiltonian (Zakharov 1968)

$$\mathcal{H} = \frac{1}{2} \int \left\{ g \eta^2 + \tilde{\phi} V \right\} d^2 \mathbf{x}, \quad V = [\phi_y - \nabla \eta \cdot \nabla \phi]_{y=\eta}. \quad (\text{B.1})$$

Introducing a Dirichlet–Neumann operator G , such that $V = G(\eta)\tilde{\phi}$ (Craig & Sulem 1993), expanding G around $\eta = 0$ and neglecting the terms beyond the quartic nonlinearities, the Hamiltonian (B.1) becomes

$$\mathcal{H} = \frac{1}{2} \int \left\{ g \eta^2 + \tilde{\phi} \left[\mathfrak{d} \tilde{\phi} - \mathfrak{d}(\eta \mathfrak{d} \tilde{\phi}) - \nabla \cdot (\eta \nabla \tilde{\phi}) + \frac{1}{2} \mathfrak{d}(\eta^2 \nabla^2 \tilde{\phi}) + \mathfrak{d}(\eta \mathfrak{d}(\eta \mathfrak{d} \tilde{\phi})) + \frac{1}{2} \nabla^2(\eta^2 \mathfrak{d} \tilde{\phi}) \right] \right\} d^2 \mathbf{x}, \quad (\text{B.2})$$

with the pseudo-differential operator $\mathfrak{d} = (-\nabla^2)^{\frac{1}{2}} \tanh[(-\nabla^2)^{\frac{1}{2}} d]$. (For one horizontal dimension in infinite depth $\mathfrak{d}f = -\mathfrak{H}(f_x)$, \mathfrak{H} the Hilbert transform.) Thus, the cubic Zakharov's equations (CZE) are

$$\eta_t - \mathfrak{d} \tilde{\phi} = -\nabla \cdot (\eta \nabla \tilde{\phi}) - \mathfrak{d}(\eta \mathfrak{d} \tilde{\phi}) + \frac{1}{2} \nabla^2(\eta^2 \mathfrak{d} \tilde{\phi}) + \mathfrak{d}(\eta \mathfrak{d}(\eta \mathfrak{d} \tilde{\phi})) + \frac{1}{2} \mathfrak{d}(\eta^2 \nabla^2 \tilde{\phi}), \quad (\text{B.3})$$

$$\tilde{\phi}_t + g \eta = \frac{1}{2} (\mathfrak{d} \tilde{\phi})^2 - \frac{1}{2} (\nabla \tilde{\phi})^2 - (\eta \mathfrak{d} \tilde{\phi}) \nabla^2 \tilde{\phi} - (\mathfrak{d} \tilde{\phi}) \mathfrak{d}(\eta \mathfrak{d} \tilde{\phi}). \quad (\text{B.4})$$

For progressive $(2\pi/\kappa)$ -periodic solutions in infinite depth, a seventh-order Stokes expansion is

$$\begin{aligned}
\kappa \eta &= \alpha \cos \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{3}{2} \alpha^2 + \frac{445}{96} \alpha^4 \right) \cos 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{41}{12} \alpha^2 + \frac{5213}{384} \alpha^4 \right) \cos 3\theta + \frac{7}{24} \alpha^4 \left(1 + \frac{263}{42} \alpha^2 \right) \cos 4\theta \\
&\quad + \frac{67}{384} \alpha^5 \left(1 + \frac{2569}{201} \alpha^2 \right) \cos 5\theta - \frac{9}{320} \alpha^6 \cos 6\theta - \frac{16751}{46080} \alpha^7 \cos 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{3}{2}} \tilde{\phi} &= \alpha \left(1 - \frac{1}{4} \alpha^2 - \frac{31}{64} \alpha^4 - \frac{465}{256} \alpha^6 \right) \sin \theta + \frac{1}{2} \alpha^2 \left(1 + \frac{3}{4} \alpha^2 + \frac{123}{64} \alpha^4 \right) \sin 2\theta \\
&\quad + \frac{3}{8} \alpha^3 \left(1 + \frac{89}{36} \alpha^2 + \frac{27271}{3456} \alpha^4 \right) \sin 3\theta + \frac{7}{24} \alpha^4 \left(1 + \frac{1795}{336} \alpha^2 \right) \sin 4\theta \\
&\quad + \frac{67}{384} \alpha^5 \left(1 + \frac{24769}{2010} \alpha^2 \right) \sin 5\theta - \frac{9}{320} \alpha^6 \sin 6\theta - \frac{16751}{46080} \alpha^7 \sin 7\theta + O(\alpha^8), \\
g^{-\frac{1}{2}} \kappa^{\frac{1}{2}} c &= 1 + \frac{1}{2} \alpha^2 + \frac{41}{64} \alpha^4 + \frac{913}{384} \alpha^6 + O(\alpha^8),
\end{aligned}$$

where the incorrect (compared to the exact expansion) coefficients and signs are displayed in bold face. Thus, the CZE match the exact Stokes wave up to the third-order only. Truncating the Hamiltonian at the order $N + 1$ in nonlinearities, the corresponding Stokes double series is correct up to the order N in the expansion parameter. None of these approximations have the exact asymptotic behavior (38) for their Fourier coefficients.

References

- [1] ABRAMOWITZ, M. & STEGUN, I. A. 1965. *Handbook of Mathematical Functions*. Dover.
- [2] BENJAMIN, T.B., BONA, J. L., MAHONY, J. J. 1972. Model equations for long waves in nonlinear dispersive systems. *Philos. Trans. Royal Soc. London Ser. A* **272**, 47–78.
- [3] BONA, J. L., CHEN, M. & SAUT, J.-C. 2002. Boussinesq equations and other systems for small-amplitude long waves in nonlinear dispersive media. I: Derivation and linear theory. *J. Nonlinear Sci.* **12**, **4**, 283–318.
- [4] BONA, J. L., CHEN, M. & SAUT, J.-C. 2004. Boussinesq equations and other systems for small-amplitude long waves in nonlinear dispersive media. II: The nonlinear theory. *Nonlinearity* **17**, **3**, 925–952.
- [5] BOUSSINESQ, J. 1871. Théorie de l’intumescence liquide appelée onde solitaire ou de translation se propageant dans un canal rectangulaire. *C. R. Acad. Sci. Paris Sér. A-B* **72**, 755–759.
- [6] CONSTANTIN, A., SATTINGER, D. H. & STRAUSS, W. 2006. Variational formulations for steady water waves with vorticity. *J. Fluid Mech.* **548**, 151–163.
- [7] CRAIG, W. & SULEM, C. 1993. Numerical simulation of gravity waves. *J. Comp. Phys.* **108**, 73–83.
- [8] CRAIK, A. 2004. The Origins of Water Wave Theory. *Annu. Rev. Fluid Mech.* **36**, 1–28.

- [9] DEGASPERIS, A. & PROCESI, M. 1999. Asymptotic integrability, ed. A. Degasperis & G. Gaeta, *Symmetry and Perturbation Theory*, World Scientific, 23–37.
- [10] DIAS, F. & BRIDGES, T. J. 2006. The numerical computation of freely propagating time-dependent irrotational water waves. *Fluid Dyn. Res.* **38**, 803–830.
- [11] DOUGALIS, V. A. & MITSOTAKIS, D. E. 2008. Theory and numerical analysis of Boussinesq systems: A review. In *Effective Computational Methods in Wave Propagation*, 63–110, Ed. N. A. Kampanis, V. A. Dougalis & J. A. Ekaterinaris.
- [12] DYSTHE, K. B. 1979. Note on a modification to the nonlinear Schrödinger equation for application to deep water. *Proc. R. Soc. Lond. A* **369**, 105–114.
- [13] ECKART, C. 1960. Variation principles of hydrodynamics. *Phys. Fluids* **3**, 5, 421–427.
- [14] FENTON, J. D. 1999. Numerical methods for nonlinear waves. *Adv. Coastal Ocean Engng* **5**, 241–324. World Scientific.
- [15] GENIET, F. 2003. Large amplitude surface waves in infinite depth, a variational approach. Preprint LPMT 2003 PM/03-25.
- [16] GREEN, A. E., LAWS, N. & NAGHDI, P. M. 1974. On the theory of water waves. *Proc. Roy. Soc. Lond. A* **338**, 43–55.
- [17] GREEN, A. & NAGHDI, P. M. 1976. A derivation of equations for wave propagation in water of variable depth. *J. Fluid Mech.* **78**, 237–246.
- [18] GRUE, J., CLAMOND, D., HUSEBY, M. & JENSEN, A. 2003. Kinematics of extreme water waves. *Appl. Ocean Res.* **25**, 6, 355–366.
- [19] JENSEN, A., CLAMOND, D., HUSEBY, M. & GRUE, J. 2007. On local and convective accelerations in steep wave events. *Ocean Engin.* **34**, 426–435.
- [20] KAMEYAMA, M., KAGEYAMA, A. & SATO, T. 2005. Multigrid iterative algorithm using pseudo-compressibility for three-dimensional mantle convection with strongly variable viscosity. *J. Comput. Phys.* **206**, 162–181.
- [21] KAUP, D. J. 1975. Higher-order water-wave equation and method for solving it. *Prog. Theor. Phys.* **54**, 396–408.
- [22] KIM, J.W., BAI, K. J., ERTEKIN, R. C. & WEBSTER, W. C. 2001. A derivation of the Green–Naghdi equations for irrotational flows. *J. Engineering Math.* **40**, 1, 17–42.
- [23] KIRBY, J. T. 1997. Nonlinear, dispersive long waves in water of variable depth. *Gravity Waves in Water of Finite Depth*, ed. J. N. Hunt, *Adv. Fluid Mech.* **10**, 55–125, Computational Mechanics Publications.
- [24] KORTEWEG, D. & DE VRIES, G. 1895. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Phil. Mag.* **39**, 422–443.

- [25] KRAENKEL, R. A., LEON, J. & MANNA, M. A. 2005. Theory of small aspect ratio waves in deep water. *Physica D* **211**, 377–390.
- [26] KUPERSHMIDT, B. A. 1985. Mathematics of dispersive water waves. *Commun. Math. Phys.* **99**, 51–73.
- [27] LAGRANGE, J.-L. 1781. Mémoire sur la théorie du mouvement des fluides. *Nouv. Mém. Acad. Berlin*, **196**. (in French)
- [28] LI, Y. A. 2002. Hamiltonian structure and linear stability of solitary waves of the Green-Naghdi equations. *J. Nonlin. Math. Phys.* **9**, **1**, 99–105.
- [29] LUKE, J. C. 1967. A variational principle for a fluid with a free surface. *J. Fluid Mech.* **27**, 395–397.
- [30] MA, Q. 2010. Advances in numerical simulation of nonlinear water waves. *Adv. Coastal Ocean Engng* **11**. World Scientific.
- [31] MADSEN, P. A. & SCHÄFFER, H. A. 1999. A review of Boussinesq-type equations for surface gravity waves. *Adv. Coastal & Ocean Engin.* **5**, World Scientific, 1–94.
- [32] MEI, C. C. 1989. *The applied dynamics of water waves*. World Scientific.
- [33] MORRISON, P. J. 1998. Hamiltonian description of the ideal fluid. *Rev. Mod. Phys.* **70**, **2**, 467–521.
- [34] OLVER, P. J. 1984. Hamiltonian perturbation theory and water waves. *Contemp. Math.* **28**, 231–249.
- [35] OLVER, P. J. 1988. Unidirectionalization of hamiltonian waves. *Phys. Lett. A* **126**, **8-9**, 501–506.
- [36] RADDER, A. C. 1999. Hamiltonian dynamics of water waves. *Adv. Coast. Ocean Engng.* **4**, 21–59.
- [37] SALMON, R. 1988. Hamiltonian fluid mechanics. *Ann. Rev. Fluid Mech.* **20**, 225–256.
- [38] SERRE, F. 1953. Contribution à l'étude des écoulements permanents et variables dans les canaux. *La Houille blanche* **8**, 374–388 & 830–872. (in French)
- [39] STOKER, J. J. 1957. *Water waves: The mathematical theory with applications*. Wiley.
- [40] SU, C. H. & GARDNER, C. S. 1969. Korteweg–deVries equation and generalizations. III. Derivation of the Korteweg–deVries equation and Burger equation. *J. Math. Phys.* **10**, **3**, 536–539.
- [41] SYNOLAKIS, C. & BERNARD, E. 2006. Tsunami science before and beyond Boxing Day 2004. *Phil. Trans. R. Soc. A* **364**, 2231–2265.
- [42] TRULSEN, K., KLIAKHANDLER, I., DYSTHE, K. B. & VELARDE, M. G. 2000. On weakly nonlinear modulation of waves on deep water. *Phys. Fluids* **12**, **10**, 2432–2437.

- [43] WEHAUSEN, J. V. & LAITONE, E. V. 1960. Surface waves. *Handbuch der Physik* **9**, **3**, 446–778. Springer-Verlag.
- [44] WHITHAM, G. B. 1965. A general approach to linear and non-linear dispersive waves using a Lagrangian. *J. Fluid Mech.* **22**, 273–283.
- [45] WU, T. Y. 2001 A unified theory for modeling water waves. *Adv. App. Mech.* **37**, 1–88.
- [46] YAHALOM, A. & LYNDEN-BELL, D. 2008. Simplified variational principles for barotropic magnetohydrodynamics. *J. Fluid Mech.* **607**, 235–265.
- [47] ZAKHAROV, V. E. 1968. Stability of periodic wave of finite amplitude on the surface of a deep fluid. *J. Appl. Mech. Phys.*, Engl. Transl. **2**, 190–198.
- [48] ZAKHAROV, V. E. & KUZNETSOV, E. A. 1997. Hamiltonian formalism for nonlinear waves. *Usp. Fiz. Nauk* **167** (11), 1137–1168.
- [49] ZHOU, J. G., CAUSON, D. M., INGRAM, D. M. & MINGHAM, C. G. 2002. Numerical solutions of the shallow water equations with discontinuous bed topography. *Int. J. Numer. Meth. Fluids* **38**, 769–788.