



HAL
open science

Sums of large global solutions to the incompressible Navier-Stokes equations

Jean-Yves Chemin, Isabelle Gallagher, Ping Zhang

► **To cite this version:**

Jean-Yves Chemin, Isabelle Gallagher, Ping Zhang. Sums of large global solutions to the incompressible Navier-Stokes equations. 2010. hal-00459759v1

HAL Id: hal-00459759

<https://hal.science/hal-00459759v1>

Preprint submitted on 25 Feb 2010 (v1), last revised 1 Oct 2010 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SUMS OF LARGE GLOBAL SOLUTIONS TO THE INCOMPRESSIBLE NAVIER-STOKES EQUATIONS

JEAN-YVES CHEMIN, ISABELLE GALLAGHER, AND PING ZHANG

ABSTRACT. Let \mathcal{G} be the (open) set of $\dot{H}^{\frac{1}{2}}$ divergence free vector fields generating a global smooth solution to the three dimensional incompressible Navier-Stokes equations. We prove that any element of \mathcal{G} can be perturbed by an arbitrarily large, smooth divergence free vector field which varies slowly in one direction, and the resulting vector field (which remains arbitrarily large) is an element of \mathcal{G} if the variation is slow enough. This result implies that through any point in \mathcal{G} passes an uncountable number of arbitrarily long segments included in \mathcal{G} .

1. INTRODUCTION

1.1. Setting of the problem and statement of the result. Let us first recall the classical Navier-Stokes system for incompressible fluids in three space dimensions:

$$(NS) \quad \begin{cases} \partial_t u + u \cdot \nabla u - \Delta u = -\nabla p \\ \operatorname{div} u = 0 \\ u|_{t=0} = u_0 \end{cases}$$

where $u(t, x)$ denotes the fluid velocity and $p(t, x)$ the pressure. In this paper the space variable x is chosen in \mathbb{R}^3 .

To start with, let us recall the history of results concerning small data. The first one states that if the initial data u_0 is such that $\|u_0\|_{L^2} \|\nabla u_0\|_{L^2}$ is small enough, a global regular solution exists; this was proved by J. Leray in his seminal paper [12]. Then, starting with the paper by H. Fujita and T. Kato (see [5]), the following approach was developed: let us denote by \mathbb{B} the bilinear operator defined by

$$\begin{cases} \partial_t \mathbb{B}(v, w) - \Delta \mathbb{B}(v, w) = \frac{1}{2} \mathbb{P} \operatorname{div}(v \otimes w + w \otimes v) \\ \mathbb{B}(v, w)|_{t=0} = 0 \end{cases}$$

where \mathbb{P} denotes the Leray projection onto divergence free vector fields. Then, it is easily checked that u is a solution of (NS) if and only if

$$u = e^{t\Delta} u_0 + \mathbb{B}(u, u)$$

which is something like Duhamel's formula. Then the theory of small initial data reduces to finding a Banach space X of time-dependent divergence free vector fields on $\mathbb{R}^+ \times \mathbb{R}^3$ such that \mathbb{B} is a bilinear map from $X \times X$ to X . An elementary abstract fixed point theorem claims that if X is a Banach space of time-dependent divergence free vector fields on $\mathbb{R}^+ \times \mathbb{R}^3$ such that

$$\|\mathbb{B}(v, w)\|_X \leq C \|v\|_X \|w\|_X$$

(X will be called from now on an adapted space), a solution of (NS) exists in X and is global as soon as

$$\|e^{t\Delta} u_0\|_X \leq (4C)^{-1}.$$

The search of the largest possible adapted space X is a long story. It started in 1964 with the paper [5] where the space X is defined by the norm

$$\|u\|_X \stackrel{\text{def}}{=} \sup_{t \geq 0} t^{\frac{1}{4}} \|\nabla u(t)\|_{L^2}.$$

After a number of important steps (see in particular [6], [9], [17] and [1]), the problem of finding the largest adapted space was achieved by H. Koch and D. Tataru. They proved in [10] that the space of time-dependent divergence free vector fields on $\mathbb{R}^+ \times \mathbb{R}^3$ such that

$$\|u\|_{X_{KT}} \stackrel{\text{def}}{=} \sup_{t \geq 0} t^{\frac{1}{2}} \|u(t)\|_{L^\infty} + \sup_{\substack{x \in \mathbb{R}^3 \\ R > 0}} \frac{1}{R^{\frac{3}{2}}} \left(\int_{P(x,R)} |u(t,y)|^2 dy dt \right)^{\frac{1}{2}} < \infty$$

where $P(x, R)$ is the parabolic ball $[0, R^2] \times B(x, R)$, is an adapted space.

Now let us observe that the incompressible Navier-Stokes system is translation and scaling invariant: if u is a solution of (NS) on $[0, T] \times \mathbb{R}^3$ then, for any positive λ and for any x_0 in \mathbb{R}^3 , the vector field u_{λ, x_0} defined by

$$u_{\lambda, x_0}(t, x) \stackrel{\text{def}}{=} \lambda u(\lambda^2 t, \lambda(x - x_0))$$

is also a solution of (NS) on $[0, \lambda^{-2}T] \times \mathbb{R}^3$. Thus, an adapted space must be translation and scaling invariant in the following sense: a constant C exists such that, for any positive λ and for any x_0 in \mathbb{R}^3 ,

$$C^{-1} \|u\|_X \leq \|u_{\lambda, x_0}\|_X \leq C \|u\|_X.$$

The second term appearing in the norm $\|\cdot\|_{X_{KT}}$ above comes from the fact that the solution of (NS) should be locally in L^2 in order to be able to define the product as a locally L^1 function. The relevant norm on the initial data is $\|e^{t\Delta} u_0\|_X$. In the case of the Koch and Tataru theorem, this norm turns out to be equivalent to the norm of the space ∂BMO of derivatives of BMO functions. Of course, the space of initial data which measures the size of the initial data must be translation and scaling invariant. A remark due to Y. Meyer (see [14]) is that the norm in such a space is always greater than the norm in the Besov space $\dot{B}_{\infty, \infty}^{-1}$ defined by

$$\|u\|_{\dot{B}_{\infty, \infty}^{-1}} \stackrel{\text{def}}{=} \sup_{t \geq 0} t^{\frac{1}{2}} \|u(t)\|_{L^\infty}.$$

This leads to the following definition of a large initial data for the incompressible Navier-Stokes equation.

Definition 1.1. *A divergence free vector field u_0 is a large initial data for the incompressible Navier-Stokes system if its $\dot{B}_{\infty, \infty}^{-1}$ norm is large.*

Let us point out that this approach using Duhamel's formula does not use the very special structure of the incompressible Navier-Stokes system. A family of results does use the special structure of (NS) : in those cases some geometrical invariance on the initial data is preserved by the flow of (NS) and this leads to some unexpected conservation of quantities, which makes the problem subcritical and thus prevents blow up. We refer for instance to [11], [13], [15], or [16], where special symmetries (like helicoidal, or axisymmetric without swirl) allow to prove global wellposedness for any data.

Some years ago, the first two authors investigated the possible existence of large initial data (in the sense of Definition 1.1) which have no preserved geometrical invariance and which nevertheless generate global regular solution. The first result in this direction was proved in [2] where such a family of large initial data was constructed, with strong oscillations in one direction. The main point of the proof is that for any element of this family, the first iterate $\mathbb{B}(e^{t\Delta} u_0, e^{t\Delta} u_0)$ is exponentially small with respect to the large initial data u_0 in

some appropriate norm. Let us notice that this result does use the fine structure of the non linear term of (NS) : M. Paicu and the second author proved in [8] that for a modified incompressible Navier-Stokes system, this family of initial data generates solutions that blow up at finite time.

In [3], the first two authors constructed another class of examples, in which the initial data has slow variations in one direction. The proof of global regularity uses the fact that the 2D Navier-Stokes equations are globally wellposed. The initial data presented in the next theorem will be referred to in the following as “quasi-2D” (see [3]).

Theorem 1. *Let $v_0^h = (v_0^1, v_0^2)$ be a horizontal, smooth divergence free vector field on \mathbb{R}^3 (i.e. v_0^h is in $L^2(\mathbb{R}^3)$ as well as all its derivatives), belonging, as well as all its derivatives, to $L^2(\mathbb{R}_{x_3}; \dot{H}^{-1}(\mathbb{R}^2))$; let w_0 be a smooth divergence free vector field on \mathbb{R}^3 . Then there exists a positive ε_0 such that if $\varepsilon \leq \varepsilon_0$, the initial data*

$$u_{0,\varepsilon}(x) \stackrel{\text{def}}{=} (v_0^h + \varepsilon w_0^h, w_0^3)(x_1, x_2, \varepsilon x_3)$$

generates a unique, global solution u^ε of (NS) .

Remark 1.2. It is clear from the proof of [3] that the dependence of the parameter ε_0 on the profiles v_0^h and w_0 is only through their norms.

Note that such an initial data may be arbitrarily large in the sense of Definition 1.1 (see [3]). We recall for the convenience of the reader the result proved in [3].

Proposition 1.3. *Let (f, g) be in $\mathcal{S}(\mathbb{R}^2) \times \mathcal{S}(\mathbb{R})$ and define $h_\varepsilon(x_h, x_3) \stackrel{\text{def}}{=} f(x_h)g(\varepsilon x_3)$. We have, if ε is small enough,*

$$\|h_\varepsilon\|_{\dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^3)} \geq \frac{1}{4} \|f\|_{\dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^2)} \|g\|_{L^\infty(\mathbb{R})}.$$

In this paper we consider the global wellposedness of the Navier-Stokes equations with data which is the sum of an initial data (which may be large) giving rise to a global solution, and a quasi-2D initial data as presented above (which may also be large). The theorem is the following.

Theorem 2. *Let u_0, v_0^h and w_0 be three smooth divergence free vector fields defined on \mathbb{R}^3 , satisfying*

- u_0 belongs to $\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$ and generates a unique global solution to the Navier-Stokes equations;
- $v_0^h = (v_0^1, v_0^2)$ is a horizontal, smooth divergence free vector field on \mathbb{R}^3 belonging, as well as all its derivatives, to the space $L^2(\mathbb{R}_{x_3}; \dot{H}^{-1}(\mathbb{R}^2))$;
- $v_0^h(x_1, x_2, 0) = w_0^3(x_1, x_2, 0) = 0$ for all $(x_1, x_2) \in \mathbb{R}^2$.

Then there exists a positive number ε_0 depending on u_0 and on norms of v_0^h and w_0 such that for any $\varepsilon \in (0, \varepsilon_0]$, there is a unique, global solution to the Navier-Stokes equations with initial data

$$u_{0,\varepsilon}(x) \stackrel{\text{def}}{=} u_0(x) + (v_0^h + \varepsilon w_0^h, w_0^3)(x_1, x_2, \varepsilon x_3).$$

Remark 1.4. Let u_0 be any element of the (open) set \mathcal{G} of $\dot{H}^{\frac{1}{2}}$ divergence free vector fields generating a global smooth solution to (NS) , and let N be an arbitrarily large number. Then for any smooth divergence free vector field f^h (over \mathbb{R}^2) and g (over \mathbb{R}) satisfying $\|f^h\|_{\dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^2)} \|g\|_{L^\infty(\mathbb{R})} \geq 4N$, and such that $g(0) = 0$, Theorem 2 implies that there is ε_N depending on u_0 and on norms of f^h and g such that $u_0 + (f^h \otimes g, 0)(x_1, x_2, \varepsilon_N x_3)$ belongs to \mathcal{G} . Since ε_N only depends on norms of f^h and g , that implies that for any $\lambda \in [-1, 1]$, the initial data $u_0 + \lambda(f^h \otimes g, 0)(x_1, x_2, \varepsilon_N x_3)$ also belongs to \mathcal{G} . Using Proposition 1.3 one

concludes that through u_0 passes an uncountable number of segments of length N included in \mathcal{G} .

Remark 1.5. With the notation of Theorem 2, the data $u_0(x) + (v_0^h + \varepsilon w_0^h, w_0^3)(x_1, x_2, \varepsilon x_3)$ belongs to \mathcal{G} as soon as ε is small enough, so one can add to that initial data any vector field of the type $(v_0^{h(1)} + \varepsilon_1 w_0^{h(1)}, w_0^{3(1)})(x_1, x_2, \varepsilon_1 x_3)$ and if ε_1 is small enough (depending on u_0 , on ε , and on norms of $v_0^h, w_0, v_0^{h(1)}$ and $w_0^{(1)}$), then the resulting vector field belongs to \mathcal{G} . One thus immediately constructs by induction superpositions of the type

$$u_0(x) + \sum_{j=0}^J (v_0^{h(j)} + \varepsilon_j w_0^{h(j)}, w_0^{3(j)})(x_1, x_2, \varepsilon_j x_3)$$

which belong to \mathcal{G} for small enough ε_j 's, depending on u_0 , on the norms of the profiles $v_0^{h(j)}$ and $w_0^{(j)}$, and on $(\varepsilon_k)_{k < j}$.

Finally notice that one can also require the slow variation on the profiles to hold on another coordinate than x_3 , up to obvious modifications of the assumptions of the theorem.

Remark 1.6. In [4], an even larger initial data than the one of Theorem 1 is constructed (the size of the solution blows up when ε tends to 0); this is a strong obstacle to the use of a perturbative argument such as the one we use here.

1.2. Scheme of the proof and organization of the paper. Let us start by introducing some notation. In the following we shall denote, for any point $x = (x_1, x_2, x_3) \in \mathbb{R}^3$, its horizontal coordinates by $x_h \stackrel{\text{def}}{=} (x_1, x_2)$. Similarly the horizontal components of any vector field $u = (u^1, u^2, u^3)$ will be denoted by $u^h \stackrel{\text{def}}{=} (u^1, u^2)$ and the horizontal divergence will be defined by $\text{div}_h u^h \stackrel{\text{def}}{=} \nabla^h \cdot u^h$, where $\nabla^h \stackrel{\text{def}}{=} (\partial_1, \partial_2)$. Finally we shall define the horizontal Laplacian by $\Delta_h \stackrel{\text{def}}{=} \partial_1^2 + \partial_2^2$. We shall often use the following shorthand notation for slowly varying functions: for any function f defined on \mathbb{R}^3 , we write

$$(1.1) \quad [f]_\varepsilon(x_h, x_3) \stackrel{\text{def}}{=} f(x_h, \varepsilon x_3).$$

In order to prove Theorem 2, we look for the solution (which exists and is smooth for a short time depending on ε , due to classical existence theory) under the form

$$(1.2) \quad u_\varepsilon \stackrel{\text{def}}{=} u_\varepsilon^{app} + R_\varepsilon$$

where the approximate solution u_ε^{app} is defined by the sum of the global solution associated with u_0 and the quasi-2D approximation:

$$(1.3) \quad u_\varepsilon^{app} \stackrel{\text{def}}{=} u + [v_\varepsilon^{(2D)}]_\varepsilon \quad \text{with} \quad v_\varepsilon^{(2D)} \stackrel{\text{def}}{=} (v^h, 0) + (\varepsilon w_\varepsilon^h, w_\varepsilon^3)$$

while

- u is the global smooth solution of (NS) associated with the initial data u_0 ;
- v^h is the global smooth solution of the two dimensional Navier-Stokes equation (with parameter y_3 in \mathbb{R}) with pressure p_0 and data $v_0^h(\cdot, y_3)$

$$(NS2D_3) \quad \begin{cases} \partial_t v^h + v^h \cdot \nabla^h v^h - \Delta_h v^h = -\nabla^h p_0 \\ \text{div}_h v^h = 0 \\ v^h|_{t=0} = v_0^h(x_h, y_3); \end{cases}$$

- w_ε solves the linear equation with data w_0 (and pressure $p_{\varepsilon,1}$)

$$(T_v^\varepsilon) \quad \begin{cases} \partial_t w_\varepsilon + v^h \cdot \nabla^h w_\varepsilon - \Delta_h w_\varepsilon - \varepsilon^2 \partial_3^2 w_\varepsilon = -(\nabla^h p_{\varepsilon,1}, \varepsilon^2 \partial_3 p_{\varepsilon,1}) \\ \text{div} w_\varepsilon = 0 \\ w_\varepsilon|_{t=0} = w_0. \end{cases}$$

We will also define the approximate pressure

$$(1.4) \quad p_\varepsilon^{app} \stackrel{\text{def}}{=} p + [p_0 + \varepsilon p_{\varepsilon,1}]_\varepsilon.$$

The stability of this approximate solution is described by the following proposition, which is proved in [3], Lemma 2.1. As in the rest of this paper, we have used the following notation: if X (resp. Y) is a function space over \mathbb{R}^2 (resp. \mathbb{R}), then we write X_h for $X(\mathbb{R}^2)$ and Y_v for $Y(\mathbb{R})$. We also denote the space $Y(\mathbb{R}; X(\mathbb{R}^2))$ by $Y_v X_h$.

Proposition 1.7 ([3]). *For any positive ε_0 , the family $(u_\varepsilon^{app})_{\varepsilon \leq \varepsilon_0}$ of approximate solutions is uniformly bounded in $L^2(\mathbb{R}^+; L^\infty(\mathbb{R}^3))$ and the family $(\nabla u_\varepsilon^{app})_{\varepsilon \leq \varepsilon_0}$ is uniformly bounded in $L^2(\mathbb{R}^+; L_v^\infty(L_h^2))$.*

The size of the error term E_ε (this denomination will become apparent in the next section) defined by

$$(1.5) \quad E_\varepsilon \stackrel{\text{def}}{=} (\partial_t - \Delta)u_\varepsilon^{app} + u_\varepsilon^{app} \cdot \nabla u_\varepsilon^{app} + (0, \partial_3[p_0])_\varepsilon$$

can be estimated as follows.

Proposition 1.8. *The family $(E_\varepsilon)_{\varepsilon \leq \varepsilon_0}$ of error terms satisfies*

$$\lim_{\varepsilon \rightarrow 0} \|E_\varepsilon\|_{L^2(\mathbb{R}^+; \dot{H}^{-\frac{1}{2}})} = 0.$$

The structure of this article is the following:

- the second section is devoted the proof of Theorem 2 using the above two propositions;
- the third section consists in proving Proposition 1.8 using estimates on the product in anisotropic spaces.

2. PROOF OF THEOREM 2

Assuming Proposition 1.8, the proof of Theorem 2 follows the same lines as the proof of Theorem 3 of [3]; we recall it for the reader's convenience. Using the definition of the approximate solution $(u_\varepsilon^{app}, p_\varepsilon^{app})$ given in (1.3, 1.4), and the error term E_ε given in (1.5), we find that the remainder R_ε satisfies the following modified three-dimensional Navier-Stokes equation

$$(MNS_\varepsilon) \quad \begin{cases} \partial_t R_\varepsilon + R_\varepsilon \cdot \nabla R_\varepsilon - \Delta R_\varepsilon + u_\varepsilon^{app} \cdot \nabla R_\varepsilon + R_\varepsilon \cdot \nabla u_\varepsilon^{app} = -E_\varepsilon - \nabla q_\varepsilon \\ \operatorname{div} R_\varepsilon = 0 \quad \text{and} \quad R_\varepsilon|_{t=0} = 0, \end{cases}$$

with $q_\varepsilon \stackrel{\text{def}}{=} p_\varepsilon - p_\varepsilon^{app}$. The proof of the theorem reduces to the proof that (MNS_ε) is globally wellposed. In order to do so, let us define

$$R_\varepsilon^\lambda(t) \stackrel{\text{def}}{=} R_\varepsilon(t) \exp\left(-\lambda \int_0^t V_\varepsilon(t') dt'\right) \quad \text{with} \quad V_\varepsilon(t) \stackrel{\text{def}}{=} \|u_\varepsilon^{app}(t)\|_{L^\infty}^2 + \|\nabla u_\varepsilon^{app}(t)\|_{L_v^\infty L_h^2}^2.$$

Note that Proposition 1.7 implies that $\int_{\mathbb{R}^+} V_\varepsilon(t) dt$ is uniformly bounded, by a constant denoted by U in the following. Writing also $E_\varepsilon^\lambda(t) \stackrel{\text{def}}{=} E_\varepsilon(t) \exp\left(-\lambda \int_0^t V_\varepsilon(t') dt'\right)$, an $\dot{H}^{\frac{1}{2}}$ energy estimate on (MNS_ε) implies

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 &= -2\lambda V_\varepsilon(t) \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 - (E_\varepsilon^\lambda | R_\varepsilon^\lambda)_{\dot{H}^{\frac{1}{2}}}(t) \\ &\quad - \left(\exp\left(\lambda \int_0^t V_\varepsilon(t') dt'\right) R_\varepsilon^\lambda \cdot \nabla R_\varepsilon^\lambda + u_\varepsilon^{app} \cdot \nabla R_\varepsilon^\lambda + R_\varepsilon^\lambda \cdot \nabla u_\varepsilon^{app} | R_\varepsilon^\lambda \right)_{\dot{H}^{\frac{1}{2}}}(t). \end{aligned}$$

A law of product in Sobolev spaces and Proposition 1.7 imply that

$$\exp\left(\lambda \int_0^t V_\varepsilon(t') dt'\right) (R_\varepsilon^\lambda \cdot \nabla R_\varepsilon^\lambda | R_\varepsilon^\lambda)_{\dot{H}^{\frac{1}{2}}} \leq C e^{\lambda U} \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}} \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2.$$

Lemma 2.3 of [3] claims that

$$(2.1) \quad (b \cdot \nabla a + a \cdot \nabla b | b)_{\dot{H}^{\frac{1}{2}}} \leq C (\|a\|_{L^\infty} + \|\nabla a\|_{L_v^\infty L_h^2}) \|b\|_{\dot{H}^{\frac{1}{2}}} \|\nabla b\|_{\dot{H}^{\frac{1}{2}}},$$

so by definition of V_ε we get

$$(u_\varepsilon^{app} \cdot \nabla R_\varepsilon^\lambda + R_\varepsilon^\lambda \cdot \nabla u_\varepsilon^{app} | R_\varepsilon^\lambda)_{\dot{H}^{\frac{1}{2}}} \leq \frac{1}{4} \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + C V_\varepsilon(t) \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2.$$

Let us choose $\lambda = C/2$. Then using the fact that

$$|(E_\varepsilon^\lambda | R_\varepsilon^\lambda)_{\dot{H}^{\frac{1}{2}}}(t)| \leq \frac{1}{4} \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + C \|E_\varepsilon^\lambda(t)\|_{\dot{H}^{-\frac{1}{2}}}^2$$

we obtain

$$\frac{d}{dt} \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + \frac{1}{2} \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 \leq C \|E_\varepsilon(t)\|_{\dot{H}^{-\frac{1}{2}}}^2 + C e^{CU} \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}} \|\nabla R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2.$$

Since $R_\varepsilon|_{t=0} = 0$ and $\lim_{\varepsilon \rightarrow 0} \|E_\varepsilon\|_{L^2(\mathbb{R}^+; \dot{H}^{-\frac{1}{2}})} = 0$ by Proposition 1.8, we deduce that as long as $\|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}$ is smaller than $1/4C e^{-CU}$, then for any $\eta > 0$ there is ε_0 such that

$$\forall \varepsilon \leq \varepsilon_0, \quad \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + \frac{1}{4} \int_0^t \|\nabla R_\varepsilon^\lambda(t')\|_{\dot{H}^{\frac{1}{2}}}^2 dt' \leq \eta,$$

which in turn implies that

$$\forall \varepsilon \leq \varepsilon_0, \quad \forall t \in \mathbb{R}^+, \quad \|R_\varepsilon^\lambda(t)\|_{\dot{H}^{\frac{1}{2}}}^2 + \frac{1}{4} \int_0^t \|\nabla R_\varepsilon^\lambda(t')\|_{\dot{H}^{\frac{1}{2}}}^2 dt' \leq \eta.$$

That concludes the proof of the theorem. \square

3. THE ESTIMATE OF THE ERROR TERM

In this section, we shall prove Proposition 1.8. Let us first remark that the error term E_ε can be decomposed as

$$E_\varepsilon = E_\varepsilon^1 + E_\varepsilon^2 \quad \text{with} \quad E_\varepsilon^2 \stackrel{\text{def}}{=} u \cdot \nabla [v_\varepsilon^{(2D)}]_\varepsilon + [v_\varepsilon^{(2D)}]_\varepsilon \cdot \nabla u.$$

Thus the term E_ε^1 is exactly the error term which appears in [3], and Lemmas 4.1, 4.2 and 4.3 of [3] imply that

$$(3.2) \quad \|E_\varepsilon^1\|_{L^2(\mathbb{R}^+; \dot{H}^{-\frac{1}{2}})} \leq C_0 \varepsilon^{\frac{1}{3}}.$$

In order to estimate the term E_ε^2 , let us first observe that Lemmas 3.1 and 3.2 of [3] imply the following proposition.

Proposition 3.1 ([3]). *For any s greater than -1 , for any $\alpha \in \mathbb{N}^3$ and for any positive t , we have*

$$\|\partial^\alpha v_\varepsilon^{(2D)}(t)\|_{L_v^2 \dot{H}_h^s}^2 + \int_0^t \|\partial^\alpha \nabla^h v_\varepsilon^{(2D)}(t')\|_{L_v^2 \dot{H}_h^s}^2 dt' \leq C_0.$$

We shall also be using the following result, whose proof is postponed to the end of this paragraph.

Proposition 3.2. *The vector field $v_\varepsilon^{(2D)}$ satisfies*

$$\|v_\varepsilon^{(2D)}(\cdot, 0)\|_{L^\infty(\mathbb{R}^+; L_h^2)} + \|\nabla^h v_\varepsilon^{(2D)}(\cdot, 0)\|_{L^2(\mathbb{R}^+; L_h^2)} \leq C \varepsilon^{\frac{1}{2}}.$$

Assuming this result, let us prove Proposition 1.8. The stability theorem of [7] claims in particular that

$$\lim_{t \rightarrow \infty} \|u(t)\|_{\dot{H}^{\frac{1}{2}}} = 0.$$

As the set of smooth compactly supported divergence free vector field is dense in the space of $\dot{H}^{\frac{1}{2}}$ divergence free vector fields, this allows to construct for any positive η , a family $(t_j)_{1 \leq j \leq N}$ of positive real numbers and a family $(\phi_j)_{1 \leq j \leq N}$ of smooth compactly supported divergence free vector fields such that (with $t_0 = 0$)

$$(3.3) \quad \|\underline{u}_\eta\|_{L^\infty(\mathbb{R}^+; \dot{H}^{\frac{1}{2}})} \leq \eta \quad \text{with} \quad \underline{u}_\eta \stackrel{\text{def}}{=} u - \tilde{u}_\eta \quad \text{and} \quad \tilde{u}_\eta(t, x) \stackrel{\text{def}}{=} \sum_{j=1}^N \mathbf{1}_{[t_{j-1}, t_j]}(t) \phi_j(x).$$

Then, for any positive η , let us decompose E_ε^2 as

$$(3.4) \quad E_\varepsilon^2 = \underline{E}_{\varepsilon, \eta} + \tilde{E}_{\varepsilon, \eta} \quad \text{with} \quad \underline{E}_{\varepsilon, \eta} \stackrel{\text{def}}{=} \underline{u}_\eta \cdot \nabla [v_\varepsilon^{(2D)}]_\varepsilon + [v_\varepsilon^{(2D)}]_\varepsilon \cdot \nabla \underline{u}_\eta.$$

The term $\underline{E}_{\varepsilon, \eta}$ will be estimated thanks to the following lemma which is a generalization of (2.1).

Lemma 3.3. *Let a and b be two smooth functions. We have*

$$\|ab\|_{\dot{H}^{\frac{1}{2}}} \leq C \|a\|_{\dot{H}^{\frac{1}{2}}} (\|\nabla^h b\|_{L_v^\infty(L_h^2)} + \|b\|_{L^\infty} + \|\partial_3 b\|_{L_v^2(\dot{H}_h^{\frac{1}{2}})}).$$

Proof. Let us first notice that, for any function f in $\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$

$$(3.5) \quad \|f\|_{\dot{H}^{\frac{1}{2}}} \leq \|f\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} + \|f\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}}.$$

Now we observe that by two-dimensional product laws, one has for any x_3 in \mathbb{R}

$$\|a(\cdot, x_3)b(\cdot, x_3)\|_{\dot{H}_h^{\frac{1}{2}}} \leq C (\|a(\cdot, x_3)\|_{\dot{H}_h^{\frac{1}{2}}} \|\nabla^h b(\cdot, x_3)\|_{L_h^2} + \|a(\cdot, x_3)\|_{\dot{H}_h^{\frac{1}{2}}} \|b(\cdot, x_3)\|_{L_h^\infty}).$$

One has of course

$$(3.6) \quad s \leq 0 \implies \|a\|_{\dot{H}^s} \leq \|a\|_{L_v^2(\dot{H}_h^s)} \quad \text{and} \quad s \geq 0 \implies \|a\|_{L_v^2(\dot{H}_h^s)} \leq \|a\|_{\dot{H}^s}$$

so taking $s = 1/2$ gives

$$(3.7) \quad \begin{aligned} \|ab\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}} &\leq C \|a\|_{L_v^2(\dot{H}_h^{\frac{1}{2}})} (\|\nabla^h b\|_{L_v^\infty(L_h^2)} + \|b\|_{L^\infty}) \\ &\leq \|a\|_{\dot{H}^{\frac{1}{2}}} (\|\nabla^h b\|_{L_v^\infty(L_h^2)} + \|b\|_{L^\infty}). \end{aligned}$$

Now let us estimate $\|ab\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}}$. A law of product in the vertical variable implies that for any x_h in \mathbb{R}^2

$$\|a(x_h, \cdot)b(x_h, \cdot)\|_{\dot{H}_v^{\frac{1}{2}}} \leq C (\|a(x_h, \cdot)\|_{\dot{H}_v^{\frac{1}{2}}} \|b(x_h, \cdot)\|_{L_v^\infty} + \|a(x_h, \cdot)\|_{L_v^2} \|\partial_3 b(x_h, \cdot)\|_{L_v^2}).$$

Taking the L^2 norm in the horizontal variable gives

$$\|ab\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} \leq C (\|a\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} \|b\|_{L^\infty} + \|a\|_{L_h^4 L_v^4} \|\partial_3 b\|_{L_h^4 L_v^4}).$$

Using Minkowski's inequality, we get that

$$\|ab\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} \leq C (\|a\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} \|b\|_{L^\infty} + \|a\|_{L_v^2 L_h^4} \|\partial_3 b\|_{L_v^2 L_h^4}).$$

Then using the Sobolev embedding $L_h^4 \hookrightarrow \dot{H}_h^{\frac{1}{2}}$ and again (3.6), we infer

$$\begin{aligned} \|ab\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} &\leq C(\|a\|_{L_h^2 \dot{H}_v^{\frac{1}{2}}} \|b\|_{L^\infty} + \|a\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}} \|\partial_3 b\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}}) \\ &\leq C\|a\|_{\dot{H}^{\frac{1}{2}}} (\|b\|_{L^\infty} + \|\partial_3 b\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}}). \end{aligned}$$

Together with (3.5) and (3.7), this proves Lemma 3.3. \square

That lemma allows to obtain the required estimate for $\underline{E}_{\varepsilon, \eta}$. Using the divergence free condition, we indeed have that

$$\underline{E}_{\varepsilon, \eta} = \operatorname{div}(\underline{u}_\eta \otimes [v_\varepsilon^{(2D)}]_\varepsilon + [v_\varepsilon^{(2D)}]_\varepsilon \otimes \underline{u}_\eta).$$

So the above lemma implies that for any positive time t

$$\begin{aligned} \|\underline{E}_{\varepsilon, \eta}(t)\|_{\dot{H}^{-\frac{1}{2}}} &\leq C\|\underline{u}_\eta \otimes [v_\varepsilon^{(2D)}]_\varepsilon + [v_\varepsilon^{(2D)}]_\varepsilon \otimes \underline{u}_\eta\|_{\dot{H}^{\frac{1}{2}}}(t) \\ &\leq C\|\underline{u}_\eta\|_{\dot{H}^{\frac{1}{2}}} (\|\nabla^h [v_\varepsilon^{(2D)}]_\varepsilon\|_{L_v^\infty L_h^2} + \|[v_\varepsilon^{(2D)}]_\varepsilon\|_{L^\infty} + \|\partial_3 [v_\varepsilon^{(2D)}]_\varepsilon\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}})(t). \end{aligned}$$

By definition of $[\cdot]_\varepsilon$ and using (3.3), we get

$$\|\underline{E}_{\varepsilon, \eta}(t)\|_{\dot{H}^{-\frac{1}{2}}} \leq C\eta (\|\nabla^h v_\varepsilon^{(2D)}\|_{L_v^\infty L_h^2} + \|v_\varepsilon^{(2D)}\|_{L^\infty} + \varepsilon^{\frac{1}{2}} \|\partial_3 v_\varepsilon^{(2D)}\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}}).$$

Proposition 3.1 gives finally

$$(3.8) \quad \|\underline{E}_{\varepsilon, \eta}\|_{L^2(\mathbb{R}^+, \dot{H}^{-\frac{1}{2}})} \leq C_0 \eta.$$

In order to estimate the term $\tilde{E}_{\varepsilon, \eta}$ let us observe that thanks to the divergence free condition, we have

$$(3.9) \quad \tilde{E}_{\varepsilon, \eta} = \tilde{u}_\eta^h \cdot \nabla^h [v_\varepsilon^{(2D)}]_\varepsilon + \varepsilon \tilde{u}_\eta^3 [\partial_3 v_\varepsilon^{(2D)}]_\varepsilon + [v_\varepsilon^{(2D)}]_\varepsilon \cdot \nabla \tilde{u}_\eta.$$

Using the 3D law of product gives

$$\|\varepsilon \tilde{u}_\eta^3 [\partial_3 v_\varepsilon^{(2D)}]_\varepsilon\|_{\dot{H}^{-\frac{1}{2}}} \leq C\varepsilon \|\tilde{u}_\eta\|_{\dot{H}^{\frac{1}{2}}} \|[\partial_3 v_\varepsilon^{(2D)}]_\varepsilon\|_{\dot{H}^{\frac{1}{2}}}$$

Using (3.6), this gives

$$(3.10) \quad \|\varepsilon \tilde{u}_\eta^3 [\partial_3 v_\varepsilon^{(2D)}]_\varepsilon\|_{\dot{H}^{-\frac{1}{2}}} \leq \varepsilon^{\frac{1}{2}} \|\tilde{u}_\eta\|_{\dot{H}^{\frac{1}{2}}} \|v_\varepsilon^{(2D)}\|_{\dot{H}^{\frac{3}{2}}}.$$

The two other terms of (3.9) are estimated using the following lemma.

Lemma 3.4. *Let a and b be two smooth functions. We have*

$$\|ab\|_{\dot{H}^{-\frac{1}{2}}} \leq C\|a\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}} \|b(\cdot, 0)\|_{L_h^2} + C\|x_3 a\|_{L^2} \|\partial_3 b\|_{L_v^\infty \dot{H}_h^{\frac{1}{2}}}.$$

Proof. Let us decompose b in the following way:

$$(3.11) \quad b(x_h, x_3) = b(x_h, 0) + \int_0^{x_3} \partial_3 b(x_h, y_3) dy_3.$$

Laws of product for Sobolev spaces on \mathbb{R}^2 together with Assertion (3.6) gives

$$\begin{aligned} \|a(b|_{x_3=0})\|_{\dot{H}^{-\frac{1}{2}}} &\leq \|a(b|_{x_3=0})\|_{L_v^2 \dot{H}_h^{-\frac{1}{2}}} \\ &\leq \left(\int_{\mathbb{R}} \|a(\cdot, x_3) b(\cdot, 0)\|_{\dot{H}_h^{-\frac{1}{2}}}^2 dx_3 \right)^{\frac{1}{2}} \\ &\leq C\|b(\cdot, 0)\|_{L^2} \left(\int_{\mathbb{R}} \|a(\cdot, x_3)\|_{\dot{H}_h^{\frac{1}{2}}}^2 dx_3 \right)^{\frac{1}{2}} \\ (3.12) \quad &\leq C\|a\|_{L_v^2 \dot{H}_h^{\frac{1}{2}}} \|b(\cdot, 0)\|_{L_h^2}. \end{aligned}$$

In order to use (3.11), let us observe that for any x_3 , two-dimensional product laws give

$$\begin{aligned} \left\| a(\cdot, x_3) \int_0^{x_3} \partial_3 b(\cdot, y_3) dy_3 \right\|_{\dot{H}_h^{-\frac{1}{2}}} &\leq C \|a(\cdot, x_3)\|_{L_h^2} \left\| \int_0^{x_3} \|\partial_3 b(\cdot, y_3)\|_{\dot{H}_h^{\frac{1}{2}}} dy_3 \right\| \\ &\leq C |x_3| \|a(\cdot, x_3)\|_{L_h^2} \|\partial_3 b\|_{L_v^\infty \dot{H}_h^{\frac{1}{2}}}. \end{aligned}$$

The above estimate integrated in x_3 together with (3.11) and (3.12) gives the result. \square

Now let us apply this lemma to estimate $\tilde{u}_\eta^h \cdot \nabla^h [v_\varepsilon^{(2D)}]_\varepsilon$ and $[v_\varepsilon^{(2D)}]_\varepsilon \cdot \nabla \tilde{u}_\eta$. We get

$$\begin{aligned} \|\tilde{u}_\eta^h \cdot \nabla^h [v_\varepsilon^{(2D)}]_\varepsilon(t)\|_{\dot{H}^{-\frac{1}{2}}} &\leq C \|\tilde{u}_\eta^h(t, \cdot)\|_{L_v^2(\dot{H}_h^{\frac{1}{2}})} \|\nabla^h v_\varepsilon^{(2D)}(t, \cdot, 0)\|_{L_h^2} \\ &\quad + \varepsilon \|x_3 \tilde{u}_\eta^h(t)\|_{L^2} \|\partial_3 \nabla^h v_\varepsilon^{(2D)}(t, \cdot)\|_{L_v^\infty(\dot{H}_h^{\frac{1}{2}})} \end{aligned}$$

and

$$\begin{aligned} \|[v_\varepsilon^{(2D)}]_\varepsilon \cdot \nabla \tilde{u}_\eta(t)\|_{\dot{H}^{-\frac{1}{2}}} &\leq C \|\nabla \tilde{u}_\eta(t, \cdot)\|_{L_v^2(\dot{H}_h^{\frac{1}{2}})} \|v_\varepsilon^{(2D)}(t, \cdot, 0)\|_{L_h^2} \\ &\quad + \varepsilon \|x_3 \nabla \tilde{u}_\eta(t, \cdot)\|_{L^2} \|\partial_3 v_\varepsilon^{(2D)}(t, \cdot)\|_{L_v^\infty(\dot{H}_h^{\frac{1}{2}})}. \end{aligned}$$

By construction of \tilde{u}_η and by Proposition 3.1 and 3.2, together with (3.9) and (3.10), we infer that

$$(3.13) \quad \|\tilde{E}_{\varepsilon, \eta}\|_{L^2(\mathbb{R}^+; \dot{H}^{-\frac{1}{2}})} \leq C_\eta \varepsilon^{\frac{1}{2}}$$

and putting (3.2), (3.8) and (3.13) together proves Proposition 1.8, up to the proof of Proposition 3.2.

So let us finally prove Proposition 3.2. We recall that $v_\varepsilon^{(2D)} = (v^h, 0) + (\varepsilon w_\varepsilon^h, w_\varepsilon^3)$, and due to the form of $(NS2D_3)$ it is clear that $v^h(t, x_h, 0) = 0$ for any (t, x_h) in $\mathbb{R}^+ \times \mathbb{R}^2$. So it remains to estimate $(\varepsilon w_\varepsilon^h, w_\varepsilon^3)$. We first notice that due to Proposition 3.1

$$\|\varepsilon w_\varepsilon^h(\cdot, 0)\|_{L^\infty(\mathbb{R}^+; L_h^2)} + \|\varepsilon \nabla^h w_\varepsilon^h(\cdot, 0)\|_{L^2(\mathbb{R}^+; L_h^2)} \leq C\varepsilon,$$

so we are left with the computation of $w_\varepsilon^3(t, \cdot, 0)$. By definition of w_ε we have

$$\begin{cases} \partial_t w_\varepsilon^3 + v^h \cdot \nabla^h w_\varepsilon^3 - \Delta_h w_\varepsilon^3 = \varepsilon^2 F_\varepsilon & \text{with } F_\varepsilon \stackrel{\text{def}}{=} \partial_3^2 w_\varepsilon^3 - \partial_3 p_{\varepsilon, 1}, \\ w_\varepsilon^3|_{t=0} = w_0^3 \end{cases}$$

We shall start by writing an $\dot{H}_h^{\frac{1}{2}}$ energy estimate (with y_3 seen as a parameter) which will imply that $w_\varepsilon^3(t, \cdot, 0)$ is smaller than $C\varepsilon$ in $L^\infty(\mathbb{R}^+; \dot{H}_h^{\frac{1}{2}}) \cap L^2(\mathbb{R}^+; \dot{H}_h^{\frac{3}{2}})$. The result in the space $L^\infty(\mathbb{R}^+; L_h^2) \cap L^2(\mathbb{R}^+; \dot{H}_h^1)$ will follow by interpolation with a bound in a negative order Sobolev space, given by Proposition 3.1.

Let us start by the $\dot{H}_h^{\frac{1}{2}}$ energy estimate. We claim that there is a constant C_0 such that for any $\varepsilon \leq \varepsilon_0$,

$$(3.14) \quad \varepsilon \|F_\varepsilon\|_{L^2(\mathbb{R}^+; L_v^\infty \dot{H}_h^{-\frac{1}{2}})} \leq C_0.$$

Assuming (3.14), an $\dot{H}_h^{\frac{1}{2}}$ energy estimate (joint with the fact that $w_\varepsilon^3|_{t=0}(\cdot, 0) = 0$) gives directly that

$$\|w_\varepsilon^3(\cdot, 0)\|_{L^\infty(\mathbb{R}^+; \dot{H}_h^{\frac{1}{2}})} + \|\nabla^h w_\varepsilon^3(\cdot, 0)\|_{L^2(\mathbb{R}^+; \dot{H}_h^{\frac{1}{2}})} \leq C\varepsilon.$$

But by Proposition 3.1 we know that w_ε^3 is uniformly bounded, say in $L^\infty(\mathbb{R}^+; L_v^\infty \dot{H}_h^{-\frac{1}{2}})$ and $\nabla^h w_\varepsilon^3$ is uniformly bounded in $L^2(\mathbb{R}^+; L_v^\infty \dot{H}_h^{-\frac{1}{2}})$, so we get by interpolation that

$$\|w_\varepsilon^3(\cdot, 0)\|_{L^\infty(\mathbb{R}^+; L_h^2)} + \|\nabla^h w_\varepsilon^3(\cdot, 0)\|_{L^2(\mathbb{R}^+; L_h^2)} \leq C\varepsilon^{\frac{1}{2}}.$$

It remains to prove the claim (3.14). On the one hand, Proposition 3.1 and Lemma 3.2 of [3] imply that

$$(3.15) \quad \|\partial_3^2 w_\varepsilon^3\|_{L^2(\mathbb{R}^+; L_v^\infty \dot{H}_h^{-\frac{1}{2}})} \leq C_0.$$

The estimate on the pressure seems slightly more delicate, but we notice as in [3] that

$$-(\varepsilon^2 \partial_3^2 + \Delta_h) p_{\varepsilon,1} = \operatorname{div}_h (v^h \cdot \nabla_h w_\varepsilon^h + \partial_3 (w_\varepsilon^3 v^h)).$$

Since $\varepsilon \partial_3 \operatorname{div}_h (\varepsilon^2 \partial_3^2 + \Delta_h)^{-1}$ is a uniformly bounded Fourier multiplier, this implies easily using the estimates of Proposition 3.1 and Sobolev embeddings that that

$$(3.16) \quad \|\varepsilon \partial_3 p_{\varepsilon,1}\|_{L^2(\mathbb{R}^+; L_v^\infty \dot{H}_h^{-\frac{1}{2}})} \leq C \|\varepsilon \partial_3 p_{\varepsilon,1}\|_{L^2(\mathbb{R}^+; H_v^1 \dot{H}_h^{-\frac{1}{2}})} \leq C_0.$$

The combination of (3.15) and (3.16) proves the claim, hence Proposition 3.2.

Acknowledgments. Part of this work was done when P. Zhang visited Laboratoire J.-L. Lions of Université Pierre et Marie Curie, as well as when J.-Y. Chemin visited the Academy of Mathematics & Systems Science, CAS. They would like to thank these two institutions for their hospitality. I. Gallagher is partially supported by the French Ministry of Research grant ANR-08-BLAN-0301-01. P. Zhang is partially supported by NSF of China under Grant 10131050 and 10276036, National 973 project and the innovation grant from the Chinese Academy of Sciences.

REFERENCES

- [1] M. Cannone, Y. Meyer and F. Planchon, Solutions autosimilaires des équations de Navier-Stokes, *Séminaire "Équations aux Dérivées Partielles de l'École Polytechnique*, 1993–1994.
- [2] J.-Y. Chemin and I. Gallagher, On the global wellposedness of the 3D Navier-Stokes equations with large data, *Annales de l'École Normale Supérieure*, **39**, 2006, pages 679–698.
- [3] J.-Y. Chemin and I. Gallagher, Large, global solutions to the Navier-Stokes equations, slowly varying in one direction, *to appear, Transactions of the American Mathematical Society*.
- [4] J.-Y. Chemin, I. Gallagher and M. Paicu, Global regularity for some classes of large solutions to the Navier-Stokes equations, *to appear, Annals of Mathematics*.
- [5] H. Fujita and T. Kato, On the Navier-Stokes initial value problem I, *Archive for Rational Mechanic Analysis*, **16**, 1964, pages 269–315.
- [6] T. Kato, Nonstationary flows of viscous and ideal fluids in \mathbb{R}^3 , *Journal of Functional Analysis*, **9**, 1972, pages 296–305.
- [7] I. Gallagher, D. Iftimie and F. Planchon, Asymptotics and stability for global solutions to the Navier-Stokes equations, *Annales de l'Institut Fourier*, **53**, 2003, pages 1387–1424.
- [8] I. Gallagher and M. Paicu, Remarks on the blow-up of solutions to a toy model for the Navier-Stokes equations, *Proceedings of the American Mathematical Society*, **137**, 2009, pages 2075–2083.
- [9] Y. Giga and T. Miyakawa, Solutions in L^r of the Navier-Stokes initial value problem, *Archiv for Rational Mechanics and Analysis*, **89**, 1985, pages 267–281.
- [10] H. Koch and D. Tataru, Well-posedness for the Navier-Stokes equations, *Advances in Mathematics*, **157**, 2001, pages 22–35.
- [11] O. Ladyzhenskaya, *The mathematical theory of viscous incompressible flow*. Second English edition, revised and enlarged. Mathematics and its Applications, Vol. 2 Gordon and Breach, Science Publishers, New York-London-Paris 1969 xviii+224 pp.
- [12] J. Leray, Essai sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Mathematica*, **63**, 1933, pages 193–248.
- [13] A. Mahalov, E. Titi and S. Leibovich, Invariant helical subspaces for the Navier-Stokes equations. *Archiv for Rational Mechanics and Analysis*, **112**, 1990, pages 193–222.

- [14] Y. Meyer, *Wavelets, Paraproducts and Navier–Stokes*. Current Developments in Mathematics, International Press, Cambridge, Massachusetts, 1996.
- [15] G. Ponce, R. Racke, T. Sideris and E. Titi, Global stability of large solutions to the 3D Navier-Stokes equations, *Communications in Mathematical Physics*, **159**, 1994, pages 329–341.
- [16] M. Ukhovskii and V. Iudovich, Axially symmetric flows of ideal and viscous fluids filling the whole space. *Prikl. Mat. Meh.* 32 59–69 (Russian); translated as *Journal of Applied Mathematics and Mechanics*, **32**, 1968, pages 52–61.
- [17] F. Weissler, The Navier-Stokes Initial Value Problem in L^p , *Archiv for Rational Mechanics and Analysis*, **74**, 1980, pages 219–230.

(J.-Y. Chemin) LABORATOIRE J.-L. LIONS, UMR 7598, UNIVERSITÉ PIERRE ET MARIE CURIE, 75230 PARIS CEDEX 05, FRANCE

E-mail address: `chemin@ann.jussieu.fr`

(I. Gallagher) INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PARIS DIDEROT, 75251 PARIS CEDEX 05, FRANCE

E-mail address: `gallagher@math.jussieu.fr`

(P. Zhang) ACADEMY OF MATHEMATICS & SYSTEMS SCIENCE, CAS, BEIJING 100080, CHINA, AND HUA LOO-KENG KEY LABORATORY OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES.

E-mail address: `zp@amss.ac.cn`