

Characterization and cartography of topsoil hydraulic properties in a French mountainous peri-urban catchment

E. Gonzalez Sosa, Isabelle Braud, J. Dehotin, F. Branger, M. Lagouy

▶ To cite this version:

E. Gonzalez Sosa, Isabelle Braud, J. Dehotin, F. Branger, M. Lagouy. Characterization and cartography of topsoil hydraulic properties in a French mountainous peri-urban catchment. Geophysical Research Abstracts, 2009, 11, p. EGU2009-2220-2 - p. hal-00471243

HAL Id: hal-00471243 https://hal.science/hal-00471243

Submitted on 7 Apr 2010 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Geophysical Research Abstracts, Vol. 11, EGU2009-2220-2, 2009 EGU General Assembly 2009 © Author(s) 2009



Characterization and cartography of topsoil hydraulic properties in a French mountainous peri-urban catchment

E. Gonzalez-Sosa (1), I. Braud (1), E. Gonzalez-Sosa (2), J. Dehotin (1), F. Branger (1), and M. Lagouy (1)

(1) Cemagref, UR HHLY, Lyon Cedex 9, France (isabelle.braud@cemagref.fr), (2) DEFI-CIAQ. University of Querétaro. CU Cerro de las Campanas. S/N. Centro. 6900 Queretaro, Qro. Mexico

Due to the increase of urbanization and modification of agricultural practices, peri-urban areas experiment a quick change in land use. The impact of such change on the catchment hydrological cycle must be quantified. To achieve this goal, distributed hydrological models offer the ability to take into account land use change, and more specifically its effect on surface infiltration capacity. A distributed assessment of infiltration properties and their variability at the catchment scale is thus of great importance if accurate simulation of the water balance are expected on such catchments.

This paper presents a field campaign conducted in a 7 km2 peri-urban catchment, located in the "Mont du Lyonnais" area, close to the city of Lyon (France) in order to document the topsoil hydraulic properties. The sampling strategy was set up in order to sample the largest number of soil/land use combinations. The locations were chosen from a GIS analysis based on the overlapping of the pedology and land use maps, and accessibility consideration. At each location, two types of infiltration tests were performed: infiltration tests under suction using mini-disk infiltrometers and single ring infiltration tests under positive head. Three replicates were performed for each method. Particle size data and organic matter analysis were also conducted at each location.

Results will be discussed in terms of soil hydraulic properties and particle size data statistics. Relationship with external factors such as pedological unit, land use, slope, texture will be explored. Preliminary results show that forest and pasture soils exhibit the highest hydraulic conductivity and sorptivity.

In order to provide models with values at the modelling unit scale (field and/or sub-catchment scale), existing pedotransfer function will be assessed and if necessary calibrated using the local measurements. Finally a methodology for the cartography of the soil hydraulic properties will be proposed.