



HAL
open science

Baire class one colorings and a dichotomy for countable unions of F_σ rectangles

Dominique Lecomte

► **To cite this version:**

Dominique Lecomte. Baire class one colorings and a dichotomy for countable unions of F_σ rectangles. 2010. hal-00472830

HAL Id: hal-00472830

<https://hal.science/hal-00472830v1>

Preprint submitted on 13 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Baire class one colorings and a dichotomy for countable unions of F_σ rectangles

Dominique LECOMTE

March 2010

- Université Paris 6, Institut de Mathématiques de Jussieu, Projet Analyse Fonctionnelle
Tour 46-0, boîte 186, 4, place Jussieu, 75 252 Paris Cedex 05, France.
dominique.lecomte@upmc.fr
- Université de Picardie, I.U.T. de l'Oise, site de Creil,
13, allée de la faïencerie, 60 107 Creil, France.

Abstract. We study the Baire class one countable colorings, i.e., the countable partitions into F_σ sets. Such a partition gives a covering of the diagonal into countably many F_σ squares. This leads to the study of countable unions of F_σ rectangles. We give a Hurewicz-like dichotomy for such countable unions.

2010 Mathematics Subject Classification. Primary: 03E15, Secondary: 54H05

Keywords and phrases. Baire class one, Borel chromatic number, Borel class, coloring, dichotomy, Hurewicz, partition, product

1 Introduction

The reader should see [K] for the standard descriptive set theoretic notation used in this paper. We study a definable coloring problem. We will need some more notation:

Notation. The letters X, Y will refer to some sets. We set $\Delta(X) := \{(x_0, x_1) \in X^2 \mid x_0 = x_1\}$.

Definition 1.1 (1) Let $A \subseteq X^2$. We say that A is a digraph if $A \cap \Delta(X) = \emptyset$.

(2) Let A be a digraph. A countable coloring of (X, A) is a map $c : X \rightarrow \omega$ such that A does not meet $(c \times c)^{-1}(\Delta(\omega))$.

In [K-S-T], the authors characterize the analytic digraphs of having a Borel countable coloring. The characterization is given in terms of the following notion of comparison between relations.

Notation. Let X, Y be Polish spaces, A (resp., B) a relation on X (resp., Y), and Γ a class of sets.

$$(X, A) \preceq_{\Gamma} (Y, B) \Leftrightarrow \exists f : X \rightarrow Y \text{ } \Gamma\text{-measurable with } A \subseteq (f \times f)^{-1}(B).$$

In this case, we say that f is a Γ -measurable homomorphism from (X, A) into (Y, B) . This notion essentially makes sense for digraphs (we can take f to be constant if B is not a digraph).

We also have to introduce a minimum digraph without Borel countable coloring:

- Let $\psi : \omega \rightarrow 2^{<\omega}$ be the natural bijection. More specifically, $\psi(0) := \emptyset$ is the sequence of length 0, $\psi(1) := 0$, $\psi(2) := 1$ are the sequences of length 1, and so on. Note that $|\psi(n)| \leq n$ if $n \in \omega$. Let $n \in \omega$. As $|\psi(n)| \leq n$, we can define $s_n := \psi(n)0^{n-|\psi(n)|}$. The crucial properties of the sequence $(s_n)_{n \in \omega}$ are the following:

- For each $s \in 2^{<\omega}$, there is $n \in \omega$ such that $s \subseteq s_n$ (we say that $(s_n)_{n \in \omega}$ is dense in $2^{<\omega}$).
- $|s_n| = n$.

- We put $\mathbb{G}_0 := \{(s_n 0 \gamma, s_n 1 \gamma) \mid n \in \omega \text{ and } \gamma \in 2^\omega\} \subseteq 2^\omega \times 2^\omega$. Note that \mathbb{G}_0 is analytic since the map $(n, \gamma) \mapsto (s_n 0 \gamma, s_n 1 \gamma)$ is continuous.

The previous definitions were given, when $\Gamma = \Delta_1^1$, in [K-S-T], where the following is proved:

Theorem 1.2 (Kechris, Solecki, Todorćević) Let X be a Polish space, and A an analytic relation on X . Then exactly one of the following holds:

- (a) There is a Borel countable coloring of (X, A) , i.e., $(X, A) \preceq_{\Delta_1^1} (\omega, \neg \Delta(\omega))$,
- (b) $(2^\omega, \mathbb{G}_0) \preceq_{\Sigma_1^0} (X, A)$.

This result had several developments during the last years:

- We can characterize the potentially closed sets via a Hurewicz-like test, and in finite dimension it is a consequence of the previous result. Let us specify this. The following definition can be found in [Lo2] (see Definition 3.3).

Definition 1.3 (Louveau) Let X, Y be Polish spaces, A a Borel subset of $X \times Y$, and Γ a Borel class. We say that A is potentially in Γ (denoted $A \in \text{pot}(\Gamma)$) iff we can find a finer Polish topology σ (resp., τ) on X (resp., Y) such that $A \in \Gamma((X, \sigma) \times (Y, \tau))$.

In particular, the potentially open sets are exactly the countable unions of Borel rectangles. A consequence of this is that the Borel hierarchy build on the Borel rectangles is exactly the hierarchy of the classes of the sets potentially in some Borel class.

The good notion of comparison to study the $\text{pot}(\Gamma)$ sets is as follows. Let X_0, X_1, Y_0, Y_1 be Polish spaces, and $A_0^\varepsilon, A_1^\varepsilon$ disjoint analytic subsets of $X_\varepsilon \times Y_\varepsilon$. Then we set

$$(X_0, Y_0, A_0^0, A_1^0) \leq (X_1, Y_1, A_0^1, A_1^1) \Leftrightarrow \\ \exists f: X_0 \rightarrow X_1 \exists g: Y_0 \rightarrow Y_1 \text{ continuous with } \forall \varepsilon \in 2 \ A_\varepsilon^0 \subseteq (f \times g)^{-1}(A_\varepsilon^1),$$

The following theorem is proved in [L1], and is a consequence of Theorem 1.2:

Theorem 1.4 Let X, Y be Polish spaces, and A_0, A_1 disjoint analytic subsets of $X \times Y$. Then exactly one of the following holds:

- (a) The set A_0 can be separated from A_1 by a $\text{pot}(\Sigma_1^0)$ set,
- (b) $(2^\omega, 2^\omega, \Delta(2^\omega), \mathbb{G}_0) \leq (X, Y, A_0, A_1)$.

In [L1], it is also proved that we cannot have f one-to-one in Theorem 1.2.(b) in general. It is easy to check that Theorem 1.2 is also an easy consequence of Theorem 1.4. This means that the study of the Borel countable colorings is highly related to the study of countable unions of Borel rectangles.

- We can extend Theorem 1.2 to any finite dimension, and also in infinite dimension if we change the space in which lives the infinite dimensional version of \mathbb{G}_0 (see [L2]).

- B. Miller recently developed some techniques to recover many dichotomy results of descriptive set theory, but without using effective descriptive set theory. He replaces it with some versions of Theorem 1.2. In particular, he can prove Theorem 1.2 without effective descriptive set theory.

When A is Borel, it is natural to ask about the relation between the Borel class of A and that of the coloring f when Theorem 1.2.(a) holds. This leads to consider Δ_ξ^0 -measurable countable colorings (or equivalently Σ_ξ^0 -measurable countable colorings). We have the following conjecture:

Conjecture 1 Let $1 \leq \xi < \omega_1$. Then there are

- a 0-dimensional Polish space \mathbb{X}_ξ ,
- an analytic relation \mathbb{A}_ξ on \mathbb{X}_ξ

such that for any 0-dimensional Polish space X , and for any analytic relation A on X , exactly one of the following holds:

- (a) $(X, A) \preceq_{\Delta_\xi^0} (\omega, \neg\Delta(\omega))$,
- (b) $(\mathbb{X}_\xi, \mathbb{A}_\xi) \preceq_{\Sigma_1^0} (X, A)$.

We will prove it when $1 \leq \xi \leq 2$, and in these cases we do not have to assume that A is analytic. A sequence $s \in 3^{<\omega}$ will be said to be *suitable* if $s = \emptyset$ or $s(|s| - 1) = 2$. We will have $\mathbb{X}_2 := 3^\omega$ and $\mathbb{A}_2 := \{(s0\alpha, s1\beta) \mid s \text{ suitable} \wedge \alpha, \beta \in 2^\omega\}$.

We saw that the study of the Borel countable colorings is highly related to the study of countable unions of Borel rectangles, and gave some motivation for studying Σ_ξ^0 -measurable countable colorings. This motivates the study of countable unions of Σ_ξ^0 rectangles. Another motivation is that $(X, A) \preceq_{\Delta_\xi^0} (\omega, \neg\Delta(\omega))$ is equivalent to the fact that $\Delta(X)$ can be separated from A by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set, by the generalized reduction property for the class Σ_ξ^0 (see 22.16 in [K]).

Conjecture 2 *Let $1 \leq \xi < \omega_1$. Then there are*

- 0-dimensional Polish spaces $\mathbb{X}_\xi^0, \mathbb{X}_\xi^1$,

- disjoint analytic subsets A_ξ^0, A_ξ^1 of $\mathbb{X}_\xi^0 \times \mathbb{X}_\xi^1$

such that for any Polish spaces X, Y , and for any pair A_0, A_1 of disjoint analytic subsets of $X \times Y$, exactly one of the following holds:

(a) *The set A_0 can be separated from A_1 by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set,*

(b) $(\mathbb{X}_\xi^0, \mathbb{X}_\xi^1, A_\xi^0, A_\xi^1) \leq (X, Y, A_0, A_1)$.

It is easy to prove this when $\xi = 1$. Our main result is that Conjecture 2 holds when $\xi = 2$. We now describe our minimum example $(\mathbb{X}_2^0, \mathbb{X}_2^1, A_2^0, A_2^1)$.

Notation. We put $\mathbb{X}_2^0 := 3^\omega \setminus \{s1\beta \mid s \text{ suitable} \wedge \beta \in 2^\omega\}$, $\mathbb{X}_2^1 := 3^\omega \setminus \{s0\alpha \mid s \text{ suitable} \wedge \alpha \in 2^\omega\}$, $A_2^0 := \Delta(\mathbb{X}_2^0 \cap \mathbb{X}_2^1)$ and $A_2^1 := \Delta_2 := \{(s0\alpha, s1\beta) \mid s \text{ suitable} \wedge \alpha, \beta \in 2^\omega\}$.

We use effective descriptive set theory, and give effective strengthenings of our results. The reader should see [M] for basic notions of effective descriptive set theory. In particular, we will see that to test whether an analytic relation has a Σ_ξ^0 -measurable countable coloring, it is enough to test countably many partitions instead of continuum many. We will use the topology T_2 generated by the $\Sigma_1^1 \cap \Pi_1^0$ subsets of a recursively presented Polish space (introduced in [Lo1]). Our main result can be strengthened as follows (see [L3]).

Theorem 1.5 *Let X, Y be recursively presented Polish spaces, and A_0, A_1 disjoint Σ_1^1 subsets of $X \times Y$. The following are equivalent:*

(a) *The set A_0 cannot be separated from A_1 by a $(\Sigma_2^0 \times \Sigma_2^0)_\sigma$ set.*

(b) *The set A_0 cannot be separated from A_1 by a $\Delta_1^1 \cap (\Sigma_2^0 \times \Sigma_2^0)_\sigma$ set.*

(c) *The set A_0 cannot be separated from A_1 by a $\Sigma_1^0(T_2 \times T_2)$ set.*

(d) $A_0 \cap \overline{A_1}^{T_2 \times T_2} \neq \emptyset$.

(e) $(\mathbb{X}_2^0, \mathbb{X}_2^1, A_2^0, A_2^1) \leq (X, Y, A_0, A_1)$.

2 Some general effective facts

One can hope for an effective strengthening of Conjecture 1:

Effective conjecture 1 *Let $1 \leq \xi < \omega_1$. Then there are*

- a 0-dimensional Polish space \mathbb{X}_ξ ,

- an analytic relation \mathbb{A}_ξ on \mathbb{X}_ξ

such that $(\mathbb{X}_\xi, \mathbb{A}_\xi) \not\preceq_{\Delta_\xi^0} (\omega, \neg\Delta(\omega))$ and for any $\alpha \in \omega^\omega$ with $1 \leq \xi < \omega_1^\alpha$, for any 0-dimensional recursively in α presented Polish space X , and for any $\Sigma_1^1(\alpha)$ relation A on X , one of the following holds:

(a) $(X, A) \preceq_{\Delta_1^1(\alpha) \cap \Delta_\xi^0} (\omega, \neg\Delta(\omega))$,

(b) $(\mathbb{X}_\xi, \mathbb{A}_\xi) \preceq_{\Sigma_1^0} (X, A)$.

We will see that this effective conjecture is true when $1 \leq \xi \leq 2$. The following statement is a corollary of this effective conjecture, and is in fact a theorem:

Theorem 2.1 *Let $1 \leq \xi < \omega_1^{CK}$, X a 0-dimensional recursively presented Polish space, and A a Σ_1^1 relation on X . We assume that $(X, A) \preceq_{\Delta_\xi^0} (\omega, \neg\Delta(\omega))$. Then $(X, A) \preceq_{\Delta_1^1 \cap \Delta_\xi^0} (\omega, \neg\Delta(\omega))$.*

A consequence of this is that to test whether an analytic relation has a Σ_ξ^0 -measurable countable coloring, it is enough to test countably many partitions instead of continuum many. Another consequence is the equivalence between Conjecture 1 and the Effective conjecture 1. We have in fact preliminary results that will help us to prove also the equivalence between (a)-(d) in Theorem 1.5, in the general case.

Lemma 2.2 *Let $1 \leq \xi < \omega_1^{CK}$, X, Y recursively presented Polish spaces, and $A \in \Sigma_1^1(X) \cap \Sigma_\xi^0$, $B \in \Sigma_1^1(Y) \cap \Sigma_\xi^0$ and $C \in \Sigma_1^1(X \times Y)$ disjoint from $A \times B$. Then there are $A', B' \in \Delta_1^1 \cap \Sigma_\xi^0$ such that $A' \times B'$ separates $A \times B$ from C .*

Proof. Note that A and $\{x \in X \mid \exists y \in B (x, y) \in C\}$ are disjoint Σ_1^1 sets, separable by a Σ_ξ^0 subset of X . By Theorems 1.A and 1.B in [Lo1], there is $A' \in \Delta_1^1 \cap \Sigma_\xi^0$ separating these two sets. Similarly, B and $\{y \in Y \mid \exists x \in A' (x, y) \in C\}$ are disjoint Σ_1^1 sets, and there is $B' \in \Delta_1^1 \cap \Sigma_\xi^0$ separating these two sets. \square

Theorem 2.3 *Let $1 \leq \xi < \omega_1^{CK}$, X, Y recursively presented Polish spaces, and A_0, A_1 disjoint Σ_1^1 subsets of $X \times Y$. We assume that A_0 is separable from A_1 by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set. Then A_0 is separable from A_1 by a $\Delta_1^1 \cap ((\Delta_1^1 \cap \Sigma_\xi^0) \times (\Delta_1^1 \cap \Sigma_\xi^0))_\sigma$ set.*

Proof. By Example 2 of Chapter 3 in [Lo2], the family $(N(n, X))_{n \in \omega}$ is regular without parameter. By Corollary 2.10 in [Lo2], $\Pi_\xi^0(X)$, as well as $\Sigma_\xi^0(X) = (\bigcup_{\eta < \xi} \Pi_\eta^0(X))_\sigma$, are regular without parameter. By Theorem 2.12 in [Lo2], $\Sigma_\xi^0(X) \times \Sigma_\xi^0(Y)$ is also regular without parameter. By Theorem 2.8 in [Lo2], the family $\Phi := (\Sigma_\xi^0(X) \times \Sigma_\xi^0(Y))_\sigma$ is separating which imply the existence of $S \in \Delta_1^1 \cap \Phi$ separating A_0 from A_1 .

With the notation of [Lo2], let n be an integer with $(0^\infty, n) \in W$ and $C_{0^\infty, n} = S$. Then $(0^\infty, n)$ is in W_Φ , which by Theorem 2.8.(ii) in [Lo2] is

$$\left\{ (\alpha, n) \in W \mid \exists \beta \in \Delta_1^1(\alpha) \ \forall m \in \omega \ (\alpha, \beta(m)) \in W_{\Sigma_\xi^0(X) \times \Sigma_\xi^0(Y)} \wedge C_{\alpha, n} = \bigcup_{m \in \omega} C_{\alpha, \beta(m)} \right\}.$$

This implies that $S \in \Delta_1^1 \cap (\Delta_1^1 \cap (\Sigma_\xi^0 \times \Sigma_\xi^0))_\sigma$. It remains to check that $\Delta_1^1 \cap (\Sigma_\xi^0 \times \Sigma_\xi^0) = (\Delta_1^1 \cap \Sigma_\xi^0) \times (\Delta_1^1 \cap \Sigma_\xi^0)$. The second set is clearly a subset of the first one. So assume that $R = A \times B \in \Delta_1^1 \cap (\Sigma_\xi^0 \times \Sigma_\xi^0)$. We may assume that R is not empty. Then the projections A, B are Σ_1^1 since $R \in \Delta_1^1$. Lemma 2.2 gives $A', B' \in \Delta_1^1 \cap \Sigma_\xi^0$ with $A \times B \subseteq A' \times B' \subseteq R = A \times B$. \square

Recall that if A is a relation on X and $D \subseteq X$, then D is A -discrete if $A \cap D^2 = \emptyset$.

Proof of Theorem 2.1. We apply Theorem 2.3 to $Y := X$, $A_0 := \Delta(X)$ and $A_1 := A$. As

$$(X, A) \preceq_{\Delta_\xi^0} (\omega, \neg \Delta(\omega)),$$

$\Delta(X)$ is separable from A by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set. Theorem 2.3 gives $C_n, D_n \in \Delta_1^1 \cap \Sigma_\xi^0$ such that $S := \bigcup_{n \in \omega} C_n \times D_n \in \Delta_1^1$ separates $\Delta(X)$ from A . As the set of codes for $\Delta_1^1 \cap \Sigma_\xi^0$ subsets of X is Π_1^1 (see Proposition 1.4 in [Lo1]), the Δ_1^1 -selection theorem and the separation theorem imply that we may assume that the sequences (C_n) and (D_n) are Δ_1^1 . Note that $(C_n \cap D_n)$ is a Δ_1^1 covering of X into A -discrete $\Delta_1^1 \cap \Sigma_\xi^0$ sets. As X is 0-dimensional we can reduce this covering into a Δ_1^1 covering (Δ_n) of X into $\Delta_1^1 \cap \Sigma_\xi^0$ sets, which are in fact Δ_ξ^0 . This gives the desired partition. \square

Notation. Following [Lo1], we define the following topologies on a 0-dimensional recursively in α presented Polish space X , for any $\alpha \in \omega^\omega$. Let $T_1(\alpha)$ be the usual topology on X , and for $2 \leq \xi < \omega_1$, $T_\xi(\alpha)$ be the topology generated by the $\Sigma_1^1(\alpha) \cap \Pi_{<\xi}^0$ subsets of X . The next proposition gives a reformulation of the inequality $(X, A) \preceq_{\Delta_1^1(\alpha) \cap \Delta_\xi^0} (\omega, \neg \Delta(\omega))$ of the Effective conjecture 1.

Proposition 2.4 *Let $1 \leq \xi < \omega_1^{CK}$, X a 0-dimensional recursively presented Polish space, and A a Σ_1^1 relation on X . Then $(X, A) \preceq_{\Delta_1^1 \cap \Delta_\xi^0} (\omega, \neg \Delta(\omega))$ is equivalent to $\Delta(X) \cap \overline{A}^{T_\xi \times T_\xi} = \emptyset$.*

Proof. Assume first that $(X, A) \preceq_{\Delta_1^1 \cap \Delta_\xi^0} (\omega, \neg \Delta(\omega))$. Then there is a partition (B_n) of X into A -discrete $\Delta_1^1 \cap \Delta_\xi^0$ sets. In particular, Theorem 1.A in [Lo1] implies that B_n is a countable union of $\Delta_1^1 \cap \Pi_{<\xi}^0$ sets if $\xi \geq 2$. In particular, B_n is T_ξ -open and $\Delta(X)$ is disjoint from $\overline{A}^{T_\xi \times T_\xi}$.

Conversely, assume that $\Delta(X) \cap \overline{A}^{T_\xi \times T_\xi} = \emptyset$. Then each element x of X is contained in a A -discrete $\Sigma_1^1 \cap \Pi_{<\xi}^0$ set (basic clopen set if $\xi = 1$). Lemma 2.2 implies that each element x of X is in fact contained in a A -discrete $\Delta_1^1 \cap \Pi_{<\xi}^0$ set if $\xi \geq 2$. It remains to apply Proposition 1.4 in [Lo1] and the Δ_1^1 -selection theorem to get the desired partition. \square

One can also hope for an effective strengthening of Conjecture 2 generalizing Theorem 1.5:

Effective conjecture 2 Let $1 \leq \xi < \omega_1$. Then there are

- 0-dimensional Polish spaces $\mathbb{X}_\xi^0, \mathbb{X}_\xi^1$,

- disjoint analytic subsets $\mathbb{A}_\xi^0, \mathbb{A}_\xi^1$ of the space $\mathbb{X}_\xi^0 \times \mathbb{X}_\xi^1$, not separable by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set,

such that for any $\alpha \in \omega^\omega$ such that $1 \leq \xi < \omega_1^\alpha$, for any recursively in α presented Polish spaces X, Y , and for any pair A_0, A_1 of disjoint $\Sigma_1^1(\alpha)$ subsets of $X \times Y$, the following are equivalent:

- (a) The set A_0 cannot be separated from A_1 by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set.
- (b) The set A_0 cannot be separated from A_1 by a $\Delta_1^1(\alpha) \cap (\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set.
- (c) The set A_0 cannot be separated from A_1 by a $\Sigma_1^0(T_\xi(\alpha) \times T_\xi(\alpha))$ set.
- (d) $A_0 \cap \overline{A_1}^{T_\xi(\alpha) \times T_\xi(\alpha)} \neq \emptyset$.
- (e) $(\mathbb{X}_\xi^0, \mathbb{X}_\xi^1, \mathbb{A}_\xi^0, \mathbb{A}_\xi^1) \leq (X, Y, A_0, A_1)$.

In fact, the statements (a)-(d) are indeed equivalent:

Theorem 2.5 Let $1 \leq \xi < \omega_1^{CK}$, X, Y recursively presented Polish spaces, and A_0, A_1 disjoint Σ_1^1 subsets of $X \times Y$. The following are equivalent:

- (a) The set A_0 cannot be separated from A_1 by a $(\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set.
- (b) The set A_0 cannot be separated from A_1 by a $\Delta_1^1 \cap (\Sigma_\xi^0 \times \Sigma_\xi^0)_\sigma$ set.
- (c) The set A_0 cannot be separated from A_1 by a $\Sigma_1^0(T_\xi \times T_\xi)$ set.
- (d) $A_0 \cap \overline{A_1}^{T_\xi \times T_\xi} \neq \emptyset$.

Proof. Theorem 2.3 implies that (a) is indeed equivalent to (b). It also implies, using the proof of Proposition 2.4, that (c) implies (a), and the converse is clear. It is also clear that (c) and (d) are equivalent. \square

A consequence of this is that Conjecture 2 and the Effective conjecture 2 are equivalent.

3 The case $\xi = 1$

We set $\mathbb{X}_1 := 2^\omega$ and $\mathbb{A}_1 := \{(0^{2k+1}1\alpha, 0^{2k}1\beta) \mid k \in \omega \wedge \alpha, \beta \in 2^\omega\}$.

Lemma 3.1 The space \mathbb{X}_1 is a 0-dimensional metrizable compact space, \mathbb{A}_1 is a Σ_1^0 relation on \mathbb{X}_1 , and $(\mathbb{X}_1, \mathbb{A}_1) \not\leq_{\Delta_1^0} (\omega, \neg\Delta(\omega))$.

Proof. The first two assertions are clear. We argue by contradiction for the last assertion, which gives $f: \mathbb{X}_1 \rightarrow \omega$ continuous with $f(\alpha) \neq f(\beta)$ if $(\alpha, \beta) \in \mathbb{A}_1$. We set $C_n := f^{-1}(\{n\})$, so that $(C_n)_{n \in \omega}$ is a partition of \mathbb{X}_1 into \mathbb{A}_1 -discrete Δ_1^0 sets. Choose n with $0^\infty \in C_n$. Then $0^i\alpha \in C_n$ if i is big enough. This gives an integer k with $0^{2k+1}1^\infty, 0^{2k}1^\infty \in C_n$, and $(0^{2k+1}1^\infty, 0^{2k}1^\infty) \in \mathbb{A}_1 \cap C_n^2$, which is absurd. \square

Theorem 3.2 *Let X be a 0-dimensional Polish space, and A a relation on X . Then exactly one of the following holds:*

- (a) $(X, A) \preceq_{\Delta_1^0} (\omega, \neg\Delta(\omega))$,
- (b) $(\mathbb{X}_1, \mathbb{A}_1) \preceq_{\Sigma_1^0} (X, A)$.

Moreover, this is not true, even if A is analytic, if X is not 0-dimensional, and we cannot have f one-to-one in (b) (with this couple $(\mathbb{X}_1, \mathbb{A}_1)$ or any other).

Proof. Note first that (a) and (b) cannot hold simultaneously, by Lemma 3.1. We enumerate a basis $(N(n, X))_{n \in \omega}$ for the topology of X made of clopen sets. Assume that (a) does not hold. We build

- an increasing sequence of integers $(n_k)_{k \in \omega}$,
- a sequence $(x_p)_{p \in \omega}$ of points of X .

We want these objects to satisfy the following conditions:

- (1) $(x_{2k}, x_{2k+1}) \in A \cap N(n_k, X)^2$
- (2) $N(n_{k+1}, X) \subseteq N(n_k, X)$
- (3) $\text{diam}(N(n_k, X)) \leq 2^{-k}$
- (4) There is no covering of $N(n_k, X)$ into A -discrete clopen subsets of X

- Assume that this is done. Then we can define a point x of X by $\{x\} = \bigcap_{k \in \omega} N(n_k, X)$. Note that (x_p) tends to x . We define $f: \mathbb{X}_1 \rightarrow X$ by $f(0^\infty) := x$, $f(0^{2k+1}1\alpha) := x_{2k}$ and $f(0^{2k}1\beta) := x_{2k+1}$. Note that f is continuous. Moreover, $(f(0^{2k+1}1\alpha), f(0^{2k}1\beta)) = (x_{2k}, x_{2k+1}) \in A$, so that (b) holds.

- Let us prove that the construction is possible. We set $N(n_{-1}, X) := X$. Assume that $(n_k)_{k < l}$ and $(x_{2k}, x_{2k+1})_{k < l}$ satisfying (1)-(4) have been constructed, which is the case for $l = 0$. We choose a covering of $N(n_{l-1}, X)$ with basic clopen sets of diameter at most 2^{-l} , contained in $N(n_{l-1}, X)$. Then one of these basic sets, say $N(n_l, X)$, satisfies (4). It remains to choose (x_{2l}, x_{2l+1}) in the set $A \cap N(n_l, X)^2$.

- Consider now $X := \mathbb{R}$ and $A := \{(0, 1)\}$. Then (a) does not hold since \mathbb{R} is connected. If (b) holds, then we must have $f(0^{2k+1}1\alpha) = 0$ and $f(0^{2k}1\beta) = 1$. By continuity of f , we get $f(0^\infty) = 0 = 1$.

This would be the same with any $(\mathbb{X}_1, \mathbb{A}_1)$. Indeed, as $(\mathbb{X}_1, \mathbb{A}_1) \not\preceq_{\Delta_1^0} (\omega, \neg\Delta(\omega))$, we have $\overline{\Pi_0[\mathbb{A}_1]} \cap \overline{\Pi_1[\mathbb{A}_1]} \neq \emptyset$, since otherwise there would be a clopen subset C of \mathbb{X}_1 separating $\overline{\Pi_0[\mathbb{A}_1]}$ from $\overline{\Pi_1[\mathbb{A}_1]}$, and we would have $\Delta(\mathbb{X}_1) \subseteq C^2 \cup (\neg C)^2 \subseteq \neg\mathbb{A}_1$. So we can choose $x \in \overline{\Pi_0[\mathbb{A}_1]} \cap \overline{\Pi_1[\mathbb{A}_1]}$, $x_{2k} \in \Pi_0[\mathbb{A}_1]$ such that (x_{2k}) tends to x , $y_{2k+1} \in \Pi_1[\mathbb{A}_1]$ such that (y_{2k+1}) tends to x , y_{2k} with $(x_{2k}, y_{2k}) \in \mathbb{A}_1$, and x_{2k+1} with $(x_{2k+1}, y_{2k+1}) \in \mathbb{A}_1$. Then $f(x_{2k}) = 0$, $f(y_{2k+1}) = 1$ and we conclude as before.

- Consider $X := 2^\omega$ and $A := \{0^\infty\} \times (2^\omega \setminus \{0^\infty\})$. Then (a) does not hold since if a clopen subset C of 2^ω contains 0^∞ , then it contains also $\alpha \neq 0^\infty$, so that $(0^\infty, \alpha) \in A \cap C^2$. If (b) holds, then $f(0^{2k+1}1\alpha) = 0^\infty$ for each integer k and f is not one-to-one.

This argument works as soon as $\Pi_0[\mathbb{A}_1]$ has at least two elements. If we argue in the other factor, then we see that an example $(\mathbb{X}_1, \mathbb{A}_1)$ with injectivity must satisfy that \mathbb{A}_1 is a singleton $\{(\alpha, \beta)\}$. As $(\mathbb{X}_1, \mathbb{A}_1) \preceq_{\Sigma_1^0} (2^\omega, \mathbb{G}_0)$, $\alpha \neq \beta$. So take a clopen subset C of \mathbb{X}_1 containing α but not β . Then $\Delta(\mathbb{X}_1) \subseteq C^2 \cup (\neg C)^2 \subseteq \neg\mathbb{A}_1$. □

Proposition 3.3 *Conjecture 2 holds for $\xi = 1$.*

Proof. We set $\mathbb{X}_1^\varepsilon := \mathbb{X}_1$, $\mathbb{A}_1^0 := \{(0^\infty, 0^\infty)\}$ and $\mathbb{A}_1^1 := \mathbb{A}_1$. If $(x, y) \in A_0 \cap \overline{A_1}$, then choose (x_k, y_k) in A_1 tending to (x, y) , and set $f(0^\infty) := x$, $g(0^\infty) := y$, $f(0^{2k+1}1\alpha) := f(0^{2k}1\beta) := x_k$, $g(0^{2k+1}1\alpha) := g(0^{2k}1\beta) := y_k$. \square

4 The case $\xi = 2$

Lemma 4.1 *The space \mathbb{X}_2 is a 0-dimensional metrizable compact space, \mathbb{A}_2 is a Σ_2^0 relation on \mathbb{X}_2 , and $(\mathbb{X}_2, \mathbb{A}_2) \not\preceq_{\Delta_2^0} (\omega, \neg\Delta(\omega))$.*

Proof. The first two assertions are clear. We argue by contradiction for the last assertion, which gives $f: \mathbb{X}_2 \rightarrow \omega$ Δ_2^0 -measurable with $f(\alpha) \neq f(\beta)$ if $(\alpha, \beta) \in \mathbb{A}_2$. We set $C_n := f^{-1}(\{n\})$, so that $(C_n)_{n \in \omega}$ is a partition of \mathbb{X}_2 into \mathbb{A}_2 -discrete Δ_2^0 sets. By Baire's theorem, there are an integer n and $s \in 2^{<\omega}$ such that C_n contains the basic clopen set N_s . Then $(s20^\infty, s21^\infty) \in \mathbb{A}_2 \cap C_n^2$, which is absurd. \square

We have a stronger result than Conjecture 1, in the sense that we do not need any regularity assumption on A , neither that X is 0-dimensional:

Theorem 4.2 (Lecomte-Zelený) *Let X be a Polish space, and A a relation on X . Then exactly one of the following holds:*

- (a) $(X, A) \preceq_{\Delta_2^0} (\omega, \neg\Delta(\omega))$,
- (b) $(\mathbb{X}_2, \mathbb{A}_2) \preceq_{\Sigma_1^0} (X, A)$.

Proof. Note first that (a) and (b) cannot hold simultaneously, by Lemma 4.1. If A is not a digraph, then choose x with $(x, x) \in A$, and put $f(\alpha) := x$. So we may assume that A is a digraph. We set

$$U := \bigcup \left\{ V \in \Sigma_1^0(X) \mid \exists D \in \Sigma_2^0(\omega \times X) \ V \subseteq \bigcup_{p \in \omega} D_p \wedge \forall p \in \omega \ A \cap (D_p \times D_p) = \emptyset \right\}.$$

Case 1. $U = X$.

There is a countable covering of X into A -discrete Σ_2^0 sets. We just have to reduce them to get a partition showing that (a) holds.

Case 2. $U \neq X$.

Then $Y := X \setminus U$ is a nonempty closed subset of X .

Claim *If $\emptyset \neq W \in \Sigma_1^0(Y)$, then there is no Σ_2^0 subset of $\omega \times X$ whose sections are A -discrete and cover W . In particular, W is not A -discrete.*

We argue by contradiction. Let $y \in W$, and Z an open subset of X with $Z \cap Y = W$. As $Z \cap U$ can be covered with some $\bigcup_{p \in \omega} D_p$'s, so is Z . Thus $Z \subseteq U$, so that $y \in Z \cap Y \subseteq U \setminus U = \emptyset$, which is the desired contradiction. \diamond

We construct a sequence $(V_s)_{s \in 3^{<\omega}}$ of open subsets of Y , and a sequence $(x_s)_{s \in 3^{<\omega}}$ of points of Y . We want these objects to satisfy the following conditions:

- (1) $x_s \in V_s$
- (2) $\overline{V_{s\varepsilon}} \subseteq V_s$
- (3) $\text{diam}(V_s) \leq 2^{-|s|}$
- (4) $(x_{s0}, x_{s1}) \in A$ if s is suitable
- (5) $x_{s\varepsilon} = x_s$ if $\varepsilon = 2 \vee s$ is not suitable

• Assume that this is done. We define $f: 3^\omega \rightarrow Y \subseteq X$ by $\{f(\alpha)\} := \bigcap_{k \in \omega} \overline{V_{\alpha|k}} = \bigcap_{k \in \omega} V_{\alpha|k}$, so that f is continuous. Note that $f(\alpha)$ is the limit of $x_{\alpha|k}$, and that

$$x_{s\varepsilon} = x_{s\varepsilon(\alpha|1)} = \dots = x_{s\varepsilon(\alpha|(q+1))}$$

for each $(s, \varepsilon, \alpha) \in 3^{<\omega} \times 2 \times 2^\omega$. Thus $f(s\varepsilon\alpha) = \lim_{q \rightarrow \infty} x_{s\varepsilon(\alpha|q)} = x_{s\varepsilon}$ and

$$(f(s0\alpha), f(s1\beta)) = (x_{s0}, x_{s1}) \in A.$$

So (b) holds.

• Let us prove that the construction is possible. We choose $x_\emptyset \in Y$ and an open neighborhood V_\emptyset of x_\emptyset in Y , of diameter at most 1. Assume that $(V_s)_{s \in 3^{\leq l}}$ and $(x_s)_{s \in 3^{\leq l}}$ satisfying (1)-(5) have been constructed, which is the case for $l = 0$.

An application of the Claim gives $(x_{s0}, x_{s1}) \in A \cap V_s^2$ if s is suitable. We satisfy (5), so that the definition of the x_s 's is complete. Note that $x_s \in V_{s|l}$ if $s \in 3^{l+1}$.

We choose an open neighborhood V_s of x_s in Y , of diameter at most 2^{-l-1} , ensuring the inclusion $\overline{V_s} \subseteq V_{s|l}$. This finishes the proof. \square

Remark. We cannot replace $(\mathbb{X}_2, \mathbb{A}_2)$ with $(2^\omega, \{(s0\alpha, s1\beta) \mid s \in 2^{<\omega} \wedge \alpha, \beta \in 2^\omega\})$. Indeed, otherwise we get $f: 2^\omega \rightarrow 3^\omega$ continuous with

$$\{(s0\alpha, s1\beta) \mid s \in 2^{<\omega} \wedge \alpha, \beta \in 2^\omega\} \subseteq (f \times f)^{-1}(\{(s0\alpha, s1\beta) \mid s \text{ suitable} \wedge \alpha, \beta \in 2^\omega\}).$$

Thus $(f(0^\infty), f(0^k 1^\infty)) = (s_k 0 \alpha_k, s_k 1 \beta_k) = (s_0 0 \alpha_0, s_0 1 \beta_0)$. But $f(0^\infty) = s_0 0 \alpha_0$ is the limit of $f(0^k 1^\infty) = s_0 1 \beta_k$, which cannot be. This shows that it is useful to take 3 instead of 2.

Now we come to the proof of our main theorem.

Lemma 4.3 *The spaces $\mathbb{X}_2^0, \mathbb{X}_2^1$ are 0-dimensional Polish spaces, $\mathbb{A}_2^0, \mathbb{A}_2^1$ are disjoint analytic subsets of $\mathbb{X}_2^0 \times \mathbb{X}_2^1$, and are not separable by a $(\Sigma_2^0 \times \Sigma_2^0)_\sigma$ set.*

Proof. The first two assertions are clear since $\mathbb{X}_2^0, \mathbb{X}_2^1$ are G_δ subsets of 3^ω , $\mathbb{A}_2^0, \mathbb{A}_2^1$ have disjoint projections, $\mathbb{A}_2^0 = \Delta(3^\omega) \cap (\mathbb{X}_2^0 \times \mathbb{X}_2^1)$ is closed and \mathbb{A}_2^1 is Σ_2^0 . We argue by contradiction for the last assertion, which gives $C_n \in \mathbf{\Pi}_1^0(\mathbb{X}_2^0)$ and $D_n \in \mathbf{\Pi}_1^0(\mathbb{X}_2^1)$ with $\mathbb{A}_2^0 \subseteq \bigcup_{n \in \omega} (C_n \times D_n) \subseteq \neg \mathbb{A}_2^1$. In particular, $\mathbb{X}_2^0 \cap \mathbb{X}_2^1 = \bigcup_{n \in \omega} C_n \cap D_n$, and Baire's theorem gives n and $s \in 3^{<\omega}$ such that the inclusion $N_s \cap \mathbb{X}_2^0 \cap \mathbb{X}_2^1 \subseteq C_n \cap D_n$ holds. Note that $N_s \cap \mathbb{X}_2^0 \subseteq C_n$ and $N_s \cap \mathbb{X}_2^1 \subseteq D_n$. Then $(s20^\infty, s21^\infty) \in \mathbb{A}_2^1 \cap (C_n \times D_n)$, which is absurd. \square

Remark. This proof shows that the spaces $\mathbb{X}_2^0, \mathbb{X}_2^1$ of Conjecture 2 cannot be both compact, which is quite unusual in this kind of dichotomy (even if it was already the case in [L2]). Indeed, our example shows that $\mathbb{A}_2^0, \mathbb{A}_2^1$ must be separable by a closed set C , and C, \mathbb{A}_2^1 must have disjoint projections. If $\mathbb{X}_2^0, \mathbb{X}_2^1$ are compact, then C and its projections are compact too. The product of these compact projections is a $(\Sigma_2^0 \times \Sigma_2^0)_\sigma$ set separating \mathbb{A}_2^0 from \mathbb{A}_2^1 , which cannot be. This fact implies that we cannot extend the continuous maps of Theorem 1.5.(e) to 3^ω in general.

Notation. We now recall some facts about the Gandy-Harrington topology (see [L2]). Let Z be a recursively presented Polish space. The *Gandy-Harrington topology* on Z is generated by the Σ_1^1 subsets of Z . We set $\Omega := \{z \in Z \mid \omega_1^z = \omega_1^{\text{CK}}\}$. Then Ω is Σ_1^1 , dense in (Z, GH) , and $W \cap \Omega$ is a clopen subset of (Ω, Σ_Z) for each $W \in \Sigma_1^1(Z)$. Moreover, (Ω, GH) is a 0-dimensional Polish space. So we fix a complete compatible metric d_{GH} on (Ω, GH) .

Proof of Theorem 1.5. We already saw that (a)-(d) are equivalent at the end of Section 2. Lemma 4.3 shows that (e) implies (a). So it is enough to show that (d) implies (e). We set $N := A_0 \cap \overline{A_1}^{T_2 \times T_2}$, which is not empty. Lemma 2.2 implies that

$$\begin{aligned} (x, y) \notin \overline{A_1}^{T_2 \times T_2} &\Leftrightarrow \exists C, D \in \Sigma_1^1 \cap \Pi_1^0 \ (x, y) \in C \times D \subseteq \neg A_1 \\ &\Leftrightarrow \exists C, D \in \Delta_1^1 \cap \Sigma_2^0 \ (x, y) \in C \times D \subseteq \neg A_1. \end{aligned}$$

This and Proposition 1.4 in [Lo1] show that N is Σ_1^1 .

• Note that s is not suitable if and only if it is of the form $u\varepsilon v$, where u is suitable, $\varepsilon \in 2$ and $v \in 2^{<\omega}$. If $\emptyset \neq s$ is suitable, then we set $s^- := s \mid \max\{l < |s| \mid s \upharpoonright l \text{ is suitable}\}$. We construct

- a sequence $(x_s)_{s \in 3^{<\omega}}$ of points of X ,
- a sequence $(y_s)_{s \in 3^{<\omega}}$ of points of Y ,
- a sequence $(U_s)_{s \in 3^{<\omega}}$ of Σ_1^0 subsets of X ,
- a sequence $(V_s)_{s \in 3^{<\omega}}$ of Σ_1^0 subsets of Y ,
- a sequence $(W_s)_{s \in 3^{<\omega}}$ suitable of Σ_1^1 subsets of $X \times Y$.

We want these objects to satisfy the following conditions:

- (1) $(x_s, y_s) \in U_s \times V_s$
- (2) $(x_s, y_s) \in W_s \subseteq N \cap \Omega$ if s is suitable
- (3) $\overline{U_{s\varepsilon}} \subseteq U_s$ if s is suitable or $s = u0v$, and $\overline{U_{u1v2}} \subseteq U_u$
- (4) $\overline{V_{s\varepsilon}} \subseteq V_s$ if s is suitable or $s = u1v$, and $\overline{V_{u0v2}} \subseteq V_u$
- (5) $W_s \subseteq W_{s^-}$ if $\emptyset \neq s$ is suitable
- (6) $\text{diam}(U_s), \text{diam}(V_s) \leq 2^{-|s|}$
- (7) $\text{diam}_{\text{GH}}(W_s) \leq 2^{-|s|}$ if s is suitable
- (8) $(x_{u0}, y_{u1}) \in (\overline{\Pi_0[(U_u \times V_u) \cap W_u]} \times \overline{\Pi_1[(U_u \times V_u) \cap W_u]}) \cap A_1$
- (9) $(x_{u0v}, y_{u1v}) = (x_{u0}, y_{u1})$

• Assume that this is done. Let $\alpha \in \mathbb{X}_2^0$. Then the increasing sequence (p_k) of integers such that $\alpha|p_k$ is suitable or of the form $u0v$ is infinite. Condition (3) implies that $(\overline{U_{\alpha|p_k}})_{k \in \omega}$ is non-increasing. Moreover, $(\overline{U_{\alpha|p_k}})_{k \in \omega}$ is a sequence of nonempty closed subsets of X whose diameters tend to 0, so that we can define $\{f(\alpha)\} := \bigcap_{k \in \omega} \overline{U_{\alpha|p_k}} = \bigcap_{k \in \omega} U_{\alpha|p_k}$. This defines a continuous map $f: \mathbb{X}_2^0 \rightarrow X$ with $f(\alpha) = \lim_{k \rightarrow \infty} x_{\alpha|p_k}$. Similarly, we define $g: \mathbb{X}_2^1 \rightarrow Y$ continuous with $g(\beta) = \lim_{k \rightarrow \infty} y_{\beta|q_k}$.

If $\alpha \in \mathbb{X}_2^0 \cap \mathbb{X}_2^1$, then the sequence (k_j) of integers such that $\alpha|p_{k_j}$ is suitable is infinite. Note that $(W_{\alpha|p_{k_j}})_{j \in \omega}$ is a non-increasing sequence of nonempty closed subsets of Ω whose GH-diameters tend to 0, so that we can define $F(\alpha)$ by $\{F(\alpha)\} := \bigcap_{j \in \omega} W_{\alpha|p_{k_j}} \subseteq N \subseteq A_0$. As $F(\alpha)$ is the limit (in $(X \times Y, \text{GH})$, and thus in $X \times Y$) of $(x_{\alpha|p_{k_j}}, y_{\alpha|p_{k_j}})_{j \in \omega}$, we get $F(\alpha) = (f(\alpha), g(\alpha))$. Thus $\mathbb{A}_2^0 \subseteq (f \times g)^{-1}(A_0)$.

Note that $x_{s\varepsilon} = x_{s\varepsilon(\alpha|1)} = \dots = x_{s\varepsilon(\alpha|(q+1))}$ for each $(s, \varepsilon, \alpha) \in 3^{<\omega} \times 2 \times 2^\omega$. This implies that $f(s0\alpha) = \lim_{q \rightarrow \infty} x_{s0(\alpha|q)} = x_{s0}$. Similarly, $g(s1\beta) = y_{s1}$ and $(f(s0\alpha), g(s1\beta)) = (x_{s0}, y_{s1}) \in A_1$. Thus $\mathbb{A}_2^1 \subseteq (f \times g)^{-1}(A_1)$.

• Let us prove that the construction is possible. As N is not empty, we can choose $(x_\emptyset, y_\emptyset) \in N \cap \Omega$, a Σ_1^1 subset W_\emptyset of $X \times Y$ with $(x_\emptyset, y_\emptyset) \in W_\emptyset \subseteq N \cap \Omega$ of GH-diameter at most 1, and a Σ_1^0 neighborhood U_\emptyset (resp., V_\emptyset) of x_\emptyset (resp., y_\emptyset) of diameter at most 1. Assume that $(x_s)_{s \in 3^{\leq l}}$, $(y_s)_{s \in 3^{\leq l}}$, $(U_s)_{s \in 3^{\leq l}}$, $(V_s)_{s \in 3^{\leq l}}$ and $(W_s)_{s \in 3^{\leq l}}$ satisfying (1)-(9) have been constructed, which is the case for $l=0$.

Note that $(x_u, y_u) \in (U_u \times V_u) \cap W_u \subseteq \overline{A_1}^{-T_2 \times T_2}$ since u is suitable. We choose $U, V \in \Sigma_1^0$ with $(x_u, y_u) \in U \times V \subseteq \overline{U} \times \overline{V} \subseteq U_u \times V_u$. As $\Pi_\varepsilon[(U \times V) \cap W_u]$ is Σ_1^1 , $\overline{\Pi_\varepsilon[(U \times V) \cap W_u]}$ is $\Sigma_1^1 \cap \Pi_1^0$. In particular, $\overline{\Pi_\varepsilon[(U \times V) \cap W_u]}$ is T_2 -open. This shows the existence of

$$(x_{u0}, y_{u1}) \in (\overline{\Pi_0[(U \times V) \cap W_u]} \times \overline{\Pi_1[(U \times V) \cap W_u]}) \cap A_1.$$

Note that $(x_{u0}, y_{u1}) \in \overline{U} \times \overline{V} \subseteq U_u \times V_u$. We set $x_{u1} := x_u, y_{u0} := y_u$. We defined x_s, y_s when $s \in 3^{l+1}$ is not suitable but $s|l$ is suitable.

Assume now that s is suitable, but not $s|l$. This gives (u, ε, v) such that $s = u\varepsilon v2$. Assume first that $\varepsilon=0$. Note that $x_{u0v} = x_{u0} \in U_{u0v} \cap \Pi_0[(U_u \times V_u) \cap W_u]$. This gives $x_s \in U_{u0v} \cap \Pi_0[(U_u \times V_u) \cap W_u]$, and also y_s with $(x_s, y_s) \in ((U_u \cap U_{u0v}) \times V_u) \cap W_u = (U_{u0v} \times V_u) \cap W_u$. If $\varepsilon=1$, then similarly we get $(x_s, y_s) \in (U_u \times V_{u1v}) \cap W_u$.

If s and $s|l$ are both suitable, or both non suitable, then we set $(x_s, y_s) := (x_{s|l}, y_{s|l})$. So we defined x_s, y_s in any case. Note that Conditions (8) and (9) are fulfilled, and that $(x_s, y_s) \in W_{s^-}$ if s is suitable. Moreover, $x_s \in U_{s|l}$ if $s|l$ is suitable or $s|l = u0v$, and $x_s \in U_u$ if $s = u1v2$, and similarly in Y . We choose Σ_1^0 sets U_s, V_s of diameter at most 2^{-l-1} with

$$(x_s, y_s) \in U_s \times V_s \subseteq \overline{U_s} \times \overline{V_s} \subseteq \begin{cases} U_{s|l} \times V_{s|l} & \text{if } s \text{ is not suitable or } s|l \text{ is suitable,} \\ U_{s|l} \times V_u & \text{if } s = u0v2, \\ U_u \times V_{s|l} & \text{if } s = u1v2. \end{cases}$$

It remains to choose, when s is suitable, $W_s \in \Sigma_1^1(X \times Y)$ of GH-diameter at most 2^{-l-1} with $(x_s, y_s) \in W_s \subseteq W_{s^-}$. \square

5 References

- [K] A. S. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, 1995
- [K-S-T] A. S. Kechris, S. Solecki and S. Todorcević, Borel chromatic numbers, *Adv. Math.* 141 (1999), 1-44
- [L1] D. Lecomte, On minimal non potentially closed subsets of the plane, *Topology Appl.* 154, 1 (2007) 241-262
- [L2] D. Lecomte, A dichotomy characterizing analytic graphs of uncountable Borel chromatic number in any dimension, *Trans. Amer. Math. Soc.* 361 (2009), 4181-4193
- [L3] D. Lecomte, How can we recognize potentially Π^0_ξ subsets of the plane?, *to appear in J. Math. Log.* (see *arXiv*)
- [Lo1] A. Louveau, A separation theorem for Σ^1_1 sets, *Trans. Amer. Math. Soc.* 260 (1980), 363-378
- [Lo2] A. Louveau, Ensembles analytiques et boréliens dans les espaces produit, *Astérisque (S. M. F.)* 78 (1980)
- [M] Y. N. Moschovakis, *Descriptive set theory*, North-Holland, 1980

Acknowledgements. This work started last summer, when I was invited at the University of Prague by Miroslav Zelený. I am very grateful to him for that, and pleased that we could prove Theorem 4.2 together.