



HAL
open science

An Exact Connection between two Solvable SDEs and a Non Linear Utility Stochastic PDEs

Nicole El Karoui, Mohamed M'Rad

► **To cite this version:**

Nicole El Karoui, Mohamed M'Rad. An Exact Connection between two Solvable SDEs and a Non Linear Utility Stochastic PDEs. 2010. hal-00477381v1

HAL Id: hal-00477381

<https://hal.science/hal-00477381v1>

Preprint submitted on 29 Apr 2010 (v1), last revised 13 Feb 2013 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An Exact Connection between two Solvable SDEs and a Non Linear Utility Stochastic PDEs ^{*†}

El Karoui Nicole and M'rad Mohamed

Paris VI/CMAP

Ecole Polytechnique,

nicole.elkaroui@cmap.polytechnique.fr, mrad@cmap.polytechnique.fr

Abstract

The paper proposes a new approach to consistent stochastic utilities, also called forward dynamic utility, recently introduced by M. Musiela and T. Zariphopoulou [27]. These utilities satisfy a property of consistency with a given incomplete financial market which gives them properties similar to the function values of classical portfolio optimization. First, we derive a non linear stochastic PDEs that satisfy consistent stochastic utilities processes of Itô type and their dual convex conjugates. Then, under some assumptions of regularity and monotony on the stochastic flow associated with the optimal wealth as function of the initial capital, and on the optimal state price dual process, we characterize all consistent utilities for a given increasing optimal wealth process from the composition of the dual optimal process and the inverse of the optimal wealth. This allows us to reduce the resolution of fully nonlinear second order utility SPDE to the existence of monotone solutions of two stochastic differential equations. We also, express the volatility of consistent utilities as an operator of the first and the second order derivatives of the utility in terms of the optimal primal and dual policies.

*With the financial support of the "Fondation du Risque" and the Fédération des banques Françaises.

†**Keywords:** forward utility, performance criteria, horizon-unbiased utility, consistent utility, progressive utility, portfolio optimization, optimal portfolio, duality, minimal martingale measure, Stochastic flows, Stochastic partial differential equation

Introduction

Recently, the concept of forward dynamic utilities has been introduced by M. Musiela and T. Zariphopoulou (2004-2007) [27, 26, 28, 30, 32, 34], to model possible changes over the time of individual preferences of an agent. Such a concept has also been studied by F. Berrier, M. Tehranchi and C. Rogers (2009) [12] and G. Zitkovic [45]. Further works are related to this problem from T. Choulli, C. Stricker and L. Jia (2007) [3], V. Henderson and D. Hobson (2007) [6].

The economic agent will adjust its preferences based on the information that is revealed over time and is represented by the filtration $(\mathcal{F}_t, t \geq 0)$ defined on the probability space $(\Omega, \mathbb{P}, (\mathcal{F}_t, t \geq 0))$. In contrast to the classical literature, there is no pre-specified trading horizon at the end of which the utility datum is assigned. Rather, the agent starts with today's specification of its utility, $U(0, x) = u(x)$, and then builds the process $U(t, x)$ for $t > 0$ in relation to the information flow given by $(\mathcal{F}_t, t \geq 0)$. This, together with the choice of a initial utility, distinguishes the forward dynamic utility from the recursive utility for which the aggregator can be specified exogenously and the value function is recovered backward in time.

Working with positive wealth processes X^π in an incomplete market, we define a consistent stochastic utility as a progressive non negative stochastic utility $U(t, x)$, for which $U(t, X_t^\pi)$ is a supermartingale, and a martingale for one optimal wealth. However we restrict our study to forward utilities which are Itô-semimartingales with spatial parameter x , whose the local characteristics (β, γ) are such that

$$dU(t, x) = \beta(t, x)dt + \gamma(t, x).dW_t.$$

The key tool is the Itô-Ventzel's formula we recall at the beginning of Section 2. As in the classical Hamilton-Jacobi-Bellman framework, we proceed by verification to establish the dynamics of consistent utilities. Assuming a sufficient constraint on the drift β of HJB type, we get the utility stochastic PDE that we investigate in this paper. In particular, we study the role of the utility risk premium defined by $\eta^U(t, x) = \gamma_x/U_x(t, x)$. Paragraph 3 goes into details on the question of duality and gives a characterization of the non linear SPDE satisfied by the progressive convex conjugate \tilde{U} of U . This allows us to obtain a complete interpretation of the volatility γ . Unlike the backward case, we do not give a positive answer on the question of existence and uniqueness of solutions of the primal and dual Hamilton-Jacobi-Bellman equations established in this work, but show the important role of the volatility γ of the \mathcal{X} -Consistent utility U and the strong analogy between the primal and dual problem. The obstacles in the analysis come from the fact that the HJB

equations are forward in time and then without maximum principle. Therefore existing results of existence, uniqueness and regularity of weak (viscosity) solutions are not directly applicable. An additional difficulty comes from the fact that the volatility coefficient may depend on higher order derivatives of U , in which case the SPDE cannot be turned into a regular PDE with random coefficients, using the method of stochastic characteristics. Moreover, the concavity property cannot be derived directly from the dynamics; that is still an open question in general, which we try to answer in Section 4. In paragraph 3.3, we show the stability of the notion of consistent utility by change of numeraire and then, without loss of generality, we can consider the martingale market where the portfolios are simple local martingales and the stochastic PDE's are easier to deal with.

The main contribution of this paper is a new approach to consistent dynamic utilities, by using stochastic flows, which we introduce in Section 4. The idea is very simple and natural: Suppose that the optimal portfolio denoted by $(\mathcal{X}(t, x))$ is strictly increasing with respect to the initial capital, and denote by $(\mathcal{X}(t, x))$ the adapted inverse process, defined by $X(t, \mathcal{X}(t, x)) = x$. Then, using the dual identity $U_x(t, X_t^*(x)) = Y_t^*(u_x(x))$, we can find $U_x(t, x)$ from $U_x(t, x) = \mathcal{Y}(t, \mathcal{X}(t, x))$ where $(Y_t^*(y))$ is the optimal state price density process. Finally we get U by integration. So, we are able to generate all the consistent utilities with given optimal portfolio.

The problem of recovering the utility function coherent with a given optimal portfolio is known in the financial literature as the “inverse“ Merton problem; it has been considered by many authors in the past in particular by H.He and C.Huang (1992), [14]. In the classical expected optimization problem, there are restrictions to put so that the portfolio is consistent with a deterministic utility criterion at some fixed time horizon. These difficulties disappear when the criterion is a progressive utility as we show in this paper.

The study provides a fine analysis of the utility volatility vector and its derivative in terms of optimal allocation policy and optimal choice of state price density. In fact, given these optimal policies, the volatility vector γ is interpreted as an operator $\Upsilon(x, U', U'')$ which is linear on U_{xx} and the dependence on U_x (resp. x) is identical to how volatility of flow \mathcal{Y} (resp. X^*) depends on \mathcal{Y} (resp. X^*).

To the best of our knowledge, the utilities fully non linear stochastic partial differential equations established in this paper and satisfied by forward utilities and their dual have not been established in a general way. In [12] and [35] the authors study the case where the volatility vector of the utility is zero. In [32], the authors derive a stochastic PDE and study examples where the volatility of the utility is constant, proportional to U (case of change of probability) and the case where the volatility is proportional to xU_x which correspond to a change of numeraire .

Furthermore, as far as we know, there is no general consistent utilities construction proposed in the literature, except the case of power or exponential type, or decreasing utilities.

Another main contribution of this paper is a connection between two solvable SDEs and the utility SPDEs early established. In particular, given a volatility vector γ such that $\gamma_x(t, x) = -xU_{xx}\kappa^*(t, x) + \nu^*(t, U_x(t, x))$, we show the existence and uniqueness of a solution to the fully nonlinear second order SPDE from that of a pair of SDE's. In any case this represents an interesting result in the theory of stochastic partial differential equations.

The paper is organized as follows, we give the definition of consistent dynamic utilities. Then, in order to study the HJB Stochastic PDE, we give more precisions on the market model and introduce the useful Itô-Ventzel formula. In Section 2, we provide the dynamics of consistent utilities and a closed form for the optimal policy. In Section 3, we study the dual process and establish a duality identity. In Section 3.3, we show the stability of the notion of consistent utility by a change of numeraire and we provide an example of consistent utility obtained by combining a standard utility function with some positive processes. In Section 4, we present our new approach and the main results of this work.

1 Consistent Stochastic Utilities

We start by introducing the concept of a forward utility consistent with a given family of portfolios. All stochastic processes are defined on a filtered probability space $(\Omega, \mathcal{F}_{t \geq 0}, \mathbb{P})$ with complete filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. In general, \mathcal{F}_0 is assumed to be trivial. In the Itô framework considered afterwards, $(\mathcal{F}_t)_{t \geq 0}$ will be usual Brownian augmented filtration.

1.1 Definition of Consistent Stochastic Utilities

A *progressive utility* U is a *positive* adapted continuous random field $U(t, x)$, such that $t \geq 0$, $x > 0 \mapsto U(t, x)$ is an increasing concave function, (in short utility function).

Obviously, this very general definition has to be compelled to represent more realistically the individual preferences of an investor in a given financial market, possibly changing over time. The idea is to calibrate these utilities with regard to some convex class (in particular vector space) of wealth processes, denoted by \mathcal{X} , on which utilities may have more properties.

As classical utility function, a progressive stochastic utility measures the relative satisfaction of any portfolio and gives a selection criterion which allows to identify an optimal choice of investment at any time. In general, we will impose below the uniqueness of the optimal

process, to be as close as possible to the usual expectations of investors. Furthermore, the satisfaction for the optimal choice is maximum and will be preserved at all futures times which explains the martingale property in the definition below. On the other hand if the strategy in \mathcal{X} fails to be optimal then it is better not to make investment. The fact of making a bad investment choice can be seen as a loss, compared with what he could won if he had followed the optimal policy. From this, we will suppose that the utility of any strategy is a supermartingale. The optimum represents the reference (benchmark) for the investor.

The class \mathcal{X} is a test-class which only allows us to specify the stochastic utility. Once his utility is defined, an investor can then turn to a portfolio optimization problem on the general financial market to establish his optimal policy or to calculate indifference prices.

Now we are able to define the \mathcal{X} -consistent stochastic utility as follows.

Definition 1.1 (*\mathcal{X} -consistent Utility*). *A \mathcal{X} -consistent stochastic utility process $U(t, x)$ is a positive progressive utility, ($t \geq 0, x > 0 \mapsto U(t, x)$ is an increasing, strictly concave function) with the following properties:*

- **Consistency with the test-class:** *For any admissible wealth process $X \in \mathcal{X}$, $\mathbb{E}(U(t, X_t)) < +\infty$ and*

$$\mathbb{E}(U(t, X_t)/\mathcal{F}_s) \leq U(s, X_s), \quad \forall s \leq t \quad .a.s.$$

- **Existence of optimal wealth:** *For any initial wealth $x > 0$, there exists an optimal wealth process $X^* \in \mathcal{X}$, such that $X_0^* = x$, and $U(s, X_s^*) = \mathbb{E}(U(t, X_t^*)/\mathcal{F}_s) \quad \forall s \leq t$.*

In short for any admissible wealth $X \in \mathcal{X}$, $U(t, X_t)$ is a positive supermartingale and a martingale for the optimal-benchmark wealth X^ .*

Our definition of consistent dynamic utilities differs from the one introduced by Musiela and Zariphopoulou [27, 26, 28, 30, 32] or Barrier and al. [12] by the fact that we do not require that the wealth processes X are discounted. This variation offers more options and allows us to study the invariance of the class of stochastic utilities by change of numéraire. In any case, there is no fixed horizon.

Remark A deterministic utility u is a \mathcal{X} -consistent utility only when the test-portfolios are local martingales. In this case, the optimal strategy is to do nothing.

The Market Model We consider a securities market which consists of $d + 1$ assets, one riskless asset, with price S^0 given by $dS_t^0 = S_t^0 r_t dt$ and d risky assets. We model the price of the d risky assets as a locally bounded positive semimartingale $S^i, \quad i = 1, \dots, d$ defined on the filtered probability space $(\Omega, \mathcal{F}_{t \geq 0}, \mathbb{P})$.

A (self-financing) portfolio is defined as a pair (x, ϕ) , where the constant x is the initial value of the portfolio and the column vector $\phi = (\phi^i)_{1 \leq i \leq d}$ is a predictable S -integrable process specifying the amount of each asset held in the portfolio. The value process, also called wealth process, $X^\phi = (X_t^\phi)_{t \geq 0}$ of such portfolio ϕ is given by

$$\frac{X_t^\phi}{S_t^0} = \frac{x}{S_0} + \int_0^t \frac{\phi_\alpha}{S_\alpha^0} \cdot d\left(\frac{S_\alpha}{S_\alpha^0}\right), \quad t \geq 0. \quad (1)$$

Let us denote by \mathbb{X}^+ the set of positive wealth processes. To facilitate the exposition we only consider wealth processes in \mathbb{X}^+ . This naturally leads us to characterize portfolios by means of relative weights π in place of the amounts ϕ . The relation between these two notions is easy since $\phi_t = (\pi_t^1 X_t^\phi(x), \dots, \pi_t^d X_t^\phi(x))^T$, where the transpose operator is denoted by T . The advantage of the second formulation is that the assumption of positive wealth is automatically satisfied, since the previous equation becomes with the notation X^π in place of X^ϕ ,

$$\frac{dX_t^\pi}{X_t^\pi} = r_t dt + \pi_t \cdot \left(\frac{dS_t}{S_t} - r_t \mathbf{1} dt \right), \quad t \geq 0 \quad (2)$$

where the d -dimensional vector such all components are equal to 1 is denoted by $\mathbf{1}$. Let us now recall that a probability measure $\mathbb{Q} \sim \mathbb{P}$ is called an equivalent local martingale measure if, for any $X \in \mathbb{X}^+$, $\frac{X}{S_0}$ is a local martingale under \mathbb{Q} . To ensure the absence of arbitrage opportunities, we postulate that the family of equivalent local martingale measures is not empty, (see [9], [7] for a precise statement and references). We stress that no assumption concerning completeness is made and in particular, many equivalent martingale measures may exist.

Itô's Market: Let $W = (W_1, W_2, \dots, W_n)^T$ be a n -standard Brownian motion ($n \geq d$), defined on the filtered probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The filtration $(\mathcal{F}_t)_{t \geq 0}$ is the \mathbb{P} -augmented filtration generated by the Brownian motion W .

The risky asset prices are continuous Itô's semimartingales with the dynamics:

$$\frac{dS_t^i}{S_t^i} = b_t^i dt + \sigma_t^i \cdot dW_t, \quad \text{for } 0 \leq i \leq d \quad (3)$$

where the inner scalar product is denoted by " \cdot ". The coefficient b^i represents the appreciation rate by time unit of the asset i and σ^i its volatility vector in \mathbb{R}^n , considered as a $n \times 1$ matrix. Denote by b the appreciation rate column vector $n \times 1$ $(b^i)_{i=1, \dots, d}$, and by σ_t the volatility matrix $n \times d$ (n lines d columns), whose i^{th} column is the vector σ_t^i for $i = 1, \dots, d$. The processes b , σ and r are \mathcal{F}_t non-anticipating processes and satisfy some minimal appropriate integrability conditions. Using vector and matrix notation, we have $dS_t = S_t(b_t dt + \sigma_t^T dW_t)$. Moreover, equation (2) may be rewritten as, $dX_t^\pi = X_t^\pi [(r_t + \pi_t \cdot (b_t - r_t) \mathbf{1}) dt + \sigma_t \pi_t \cdot dW_t]$.

As usual, the matrix $(\sigma\sigma^T)(t, \omega)$ is assumed to be **non singular**. This assumption is equivalent to suppose that, for any $i \in 1..d$, the asset S^i cannot be replicated by an admissible portfolio.

The existence of an equivalent local martingale measure is equivalent in this framework, to the fact that the excess of return vector belongs to the range of volatility matrix: in other words, there exists a \mathcal{F} -progressively measurable process $\eta \in \mathbb{R}^n$ such that $b_t - r_t \mathbf{1} = \sigma_t^T \eta_t$. Additional integrability assumptions are necessary to ensure that the exponential martingale generated by $\eta \cdot W$ is the density of some probability measure.

We get that the dynamics of the portfolio becomes $dX_t^\pi = X_t^\pi [r_t dt + \sigma_t \pi_t \cdot (dW_t + \eta_t dt)]$. The key role is played by the volatility vector $\sigma\pi$. For this and in order to facilitate the exposition, we denote it by $\kappa := \sigma\pi$. To fix the notation, we denote by $\mathcal{R}_t^\sigma \subset \mathbb{R}^n$ the range of σ_t , and by $\mathcal{R}_t^{\sigma, \perp}$ the orthogonal vector subspace. By assumption, κ_t is required to lie at any time t in \mathcal{R}_t^σ . Replacing X_t^π by X_t^κ , the above equation becomes

$$dX_t^\kappa = X_t^\kappa [r_t dt + \kappa_t \cdot (dW_t + \eta_t dt)], \quad \kappa_t \in \mathcal{R}_t^\sigma. \quad (4)$$

Algebraic Notations The following short notation will be used extensively. Let \mathcal{R}^σ be a vector subspace of \mathbb{R}^n . For any $\alpha \in \mathbb{R}^n$, we denote by α^σ the orthogonal projection of the vector α onto \mathcal{R}^σ and by α^\perp the orthogonal projection onto $\mathcal{R}^{\sigma, \perp}$. Then, the following decomposition : $\alpha = \alpha^\sigma + \alpha^\perp$ holds.

To close this section, let us introduce as Musiela and Zariphopoulou ([32]), the generalized inverse of σ known as the Moore-Penrose inverse σ^+ and recall that σ^+ is the unique matrix $d \times N$ satisfying the following four Penrose equations:

$$\sigma\sigma^+ = (\sigma\sigma^+)^T, \quad \sigma^+\sigma = (\sigma^+\sigma)^T, \quad \sigma\sigma^+\sigma = \sigma, \quad \sigma^+\sigma\sigma^+ = \sigma^+ \quad (5)$$

Then $\sigma\sigma^+$ is the orthogonal projection matrix onto \mathcal{R}^σ , and $\alpha^\sigma = \sigma\sigma^+\alpha$. Moreover, denoting by I_n the n -dimensional identity matrix, $I_n - \sigma\sigma^+$ is the orthogonal projection matrix onto $\mathcal{R}^{\sigma, \perp}$. Moreover, under market assumptions ($\sigma^T\sigma$ non singular) there exists a unique vector π such that $\kappa = \sigma\pi$ which is $\pi = \sigma^+\kappa$.

Minimal Risk premium The market incompleteness is described through the family of risk premium η . Since for any $\kappa \in \mathcal{R}^\sigma$, $\kappa \cdot \eta = \kappa \cdot \eta^\sigma$, we assume throughout this paper and without further mention that $\eta = \eta^\sigma \in \mathcal{R}^\sigma$. η^σ is often referred to as the minimal risk premium.

2 Stochastic Partial Differential Equation

In this section, under some additional regularity assumptions we will focus on the Hamilton-Jacobi-Bellman stochastic PDE satisfied by a \mathcal{X} -consistent stochastic utility using essentially Itô-Ventzel's formula and techniques of dynamic programming established and developed in the classical theory of utility maximization (see for example H. Pham [15]). Additional regularity assumptions are necessary to advance in the study. From now, \mathcal{X} -consistent stochastic utilities $U(t, x)$ are described as Itô's semimartingales with spatial parameter $x > 0$; in other words, $U(t, x)$ is a continuous random field with dynamics,

$$dU(t, x) = \beta(t, x)dt + \gamma(t, x).dW_t, \quad (6)$$

where, as in Kunita [25], the pair (β, γ) is called the *local characteristics* of U and are assumed to be progressively random fields with values in \mathbb{R} and \mathbb{R}^n respectively.

We are concerned with the properties of the utility of admissible wealth processes. Before that, we want to give precise definition of the progressive utility, its derivatives and their dynamic properties.

2.1 Regular stochastic flows and Itô-Ventzel's formula

Regular Stochastic flows There are several difficulties in the definition of semimartingales depending on a parameter, as explained in H. Kunita [25] and R.A. Carmona et al. [2], (see Appendix A).

First let us point out that in general equality (6) holds for any t except for a null set N_x . Then the semimartingale U is well defined for (t, x) if $\omega \in (\cup_{x \in \mathbb{R}_+} N_x)^c$. However the exceptional set $(\cup_{x \in \mathbb{R}_+} N_x)$ may not be a null set since it is an uncountable union of null sets. However if we suppose that local characteristic (β, γ) of U are δ -Hölder, for some $\delta > 0$ (see appendix A), then according to H. Kunita [25] (Theorems 3.1.2 p.75) using Kolmogorov's criterion, $U(t, x)$ has a continuous modification for which (6) holds almost surely.

A detailed discussion about these difficulties and their consequences in terms of dynamic representation and differential rules are provided in H. Kunita [25] and R.A. Carmona et al. [2]. The main results are also recalled in Appendix A. Here we only give a self-contained definition of the regularity in the sense of Kunita [25]. In particular, albeit the process U and its local characteristics (β, γ) are differentiable it is not enough as is showed in H. Kunita [25], to get that the dynamics of the derivative $\frac{\partial}{\partial x}U(t, x)$ is the derivative term by term of that of U . Let m be a non-negative integer, β be a real function in $\mathcal{C}^m([0, +\infty[\times [0, +\infty[)$ and γ be a $\mathcal{C}^m([0, +\infty[\times [0, +\infty[)$ vector. We define the following seminorms for any compact

K ,

$$\|\beta\|_{m:K}(t) = \sup_{x \in K} \frac{|\beta(t, x)|}{1 + |x|} + \sum_{1 \leq \alpha \leq m} \sup_{x \in K} |\partial_x^\alpha \beta(t, x)|.$$

$$\|\gamma\|_{\widetilde{m}:K}(t) = \sup_{x, y \in K} \frac{|\gamma^T(t, x) \cdot \gamma(t, y)|}{(1 + |x|)(1 + |y|)} + \sum_{1 \leq \alpha_1, \alpha_2 \leq m} \sup_{x, y \in K} |\partial_x^{\alpha_1} \partial_y^{\alpha_2} \gamma^T(t, x) \cdot \gamma(t, y)|$$

For simplicity if a random field $(G(t, x))_{t \geq 0, x \geq 0}$ is of class $\mathcal{C}^{0,2}([0, +\infty[\times [0, +\infty[)$ we use the notation G_x for $\frac{\partial}{\partial x} G$ and G_{xx} for $\frac{\partial^2}{\partial x^2} G$.

Definition 2.1. Let $m \geq 2$. A random field F is said to be $\mathcal{C}^{(m)}$ regular in the sense of Kunita if $F : [0, +\infty[\times [0, +\infty[\rightarrow \mathbb{R}$ is of class $\mathcal{C}^{0,m}([0, +\infty[\times [0, +\infty[)$ and satisfies

$$F(t, x) = F(0, x) + \int_0^t \beta(s, x) ds + \int_0^t \gamma(s, x) \cdot dW_s. \quad (7)$$

In the sequel, according to H. Kunita [25] the pair of adapted random fields (β, γ) is called the local characteristics of F and satisfy $\beta : [0, +\infty[\times [0, +\infty[\rightarrow \mathbb{R}$ and the N -dimensional vector $\gamma : [0, +\infty[\times [0, +\infty[\rightarrow \mathbb{R}^N$ are \mathcal{F} -adapted random field of class $\mathcal{C}^{0,m-1}([0, +\infty[\times [0, +\infty[)$ such that $\|\beta\|_{m-1:K}(t)$ and $\|\gamma\|_{\widetilde{m-1}:K}(t)$ are integrable with respect to t , for any compact $K \subset [0, +\infty[$.

Now we turn to the differential rules of semimartingales with spatial parameter. For this some other notations are needed. Let $0 < \delta \leq 1$ and K a compact of \mathbb{R}_+ . For some random fields $f(t, x)$ and $g(t, x, y)$ we set

$$\|f\|_{\delta:K} := \sup_{\substack{x, y \in K \\ x \neq y}} \frac{|\partial_x^\alpha f(x) - \partial_x^\alpha f(y)|}{|x - y|^\delta}, \quad \|g\|_{\delta:K} := \sup_{\substack{x, x', y, y' \in K \\ x \neq x', y \neq y'}} \frac{|g(x, y) - g(x', y) - g(x, y') + g(x', y')|}{|x - x'|^\delta |y - y'|^\delta}.$$

Using these notations and according to the results of H. Kunita [25], (Theorem 3.3.3 p.95, recalled in Appendix A), we have the following differential rule.

Theorem 2.1 (Differential Rules). Let F be a random field of class $\mathcal{C}^{0,1}([0, +\infty[\times [0, +\infty[)$ such that its local characteristics (β, γ) are of class $\mathcal{C}^{0,1}([0, +\infty[\times [0, +\infty[)$. Assume that the derivative β_x and γ_x are δ -Hölder, with $0 < \delta \leq 1$ such that for any compact K of \mathbb{R}_+ , $\|\beta\|_{\delta:K}(t)$ and $\|a^\gamma\|_{\delta:K}(t)$ are integrable with respect to t , with $a^\gamma(t, x, y) := \gamma(t, x)^T \cdot \gamma(t, y)$. Then the derivative F_x of F with respect to the spatial parameter x satisfies, almost surely,

$$F_x(t, x) = F_x(0, x) + \int_0^t \beta_x(s, x) ds + \int_0^t \gamma_x(s, x) \cdot dW_s \quad (8)$$

Furthermore, if F is of class $\mathcal{C}^{(m)}$, $m \geq 3$ then F_x is of class $\mathcal{C}^{(m-1)}$ with local characteristics (β_x, γ_x) which are of class $\mathcal{C}^{0,m-2}([0, +\infty[\times [0, +\infty[)$.

Itô-Ventzel's formula Now, we need to study the dynamics of $U(t, X_t^\kappa)$ (X^κ is a wealth process). Itô-Ventzel's formula is a generalization of classical Itô's formula where the deterministic function is replaced by a stochastic process depending on a real or multivariate parameter. This enables us to carry out computations in a stochastically modulated dynamic framework.

Theorem 2.2 (Itô-Ventzel's Formula). *Consider a random field $F : [0, +\infty[\times [0, +\infty[\rightarrow \mathbb{R}$ which is of class $\mathcal{C}^{(2)}$ in the sense of Kunita,*

$$F(t, x) = F(0, x) + \int_0^t \beta(s, x) ds + \int_0^t \gamma(s, x) \cdot dW_s, \text{ a.s.} \quad (9)$$

Furthermore, let X be a continuous semimartingale with decomposition

$$X_t = X_0 + \int_0^t \mu_s^X ds + \int_0^t \sigma_s^X \cdot dW_s$$

Then $(F(t, X_t))$ is also a continuous semimartingale with decomposition

$$\begin{aligned} F(t, X_t) &= F(0, X_0) + \int_0^t \beta(s, X_s) ds + \int_0^t \gamma(s, X_s) \cdot dW_s \\ &+ \int_0^t F_x(s, X_s) dX_s + \frac{1}{2} \int_0^t F_{xx}(s, X_s) \langle dX_s \rangle \\ &+ \int_0^t \gamma_x(s, X_s) \cdot \sigma_s^X ds. \end{aligned}$$

Let us now comment the dynamics of $F(t, X_t)$. The first line of the right hand side of this dynamic corresponds to the dynamics of the process $(F(t, x))_{t \geq 0}$ taken on $(X_t)_{t \geq 0}$, where the second one is none other than the classical Itô formula, and the last one represents a correction term which can be written $\gamma_x(s, X_s) \cdot \sigma_s^X = \langle dF_x(s, x), dX_s \rangle|_{x=X_s}$.

We refer to Kunita [25], (Theorem 3.3.1, p.92, Theorem 3.3.1, p.92) for more details and the proof of this result. We illustrate this formula from the classical Itô's formula.

Example: Itô's Formula Let $f(t, \theta, x)$ be a deterministic function $\mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ of class $\mathcal{C}^{1,2,2}$. Denote by ∇_θ the gradient with respect to θ and by $\Delta_{\theta\theta}$ the Hessian matrix with respect to θ where t and x are fixed.

Let $\Theta \in \mathbb{R}^n$ be a Itô's semimartingale $d\Theta_t = \mu_t^\Theta dt + \sigma_t^\Theta dW_t$, with the diffusion generator

$$L_t^\Theta = \mu_t^\Theta \nabla_\theta + \frac{1}{2} \text{trace}[\sigma_t^\Theta (\sigma_t^\Theta)^T \Delta_{\theta\theta}].$$

Denote by F the stochastic random field $F(t, x) \stackrel{\text{def}}{=} f(t, \Theta_t, x)$. By the classical Itô's formula

$$dF(t, x) = (f_t(t, x, \Theta_t) + L_t^\Theta f(t, x, \Theta_t)) dt + (\nabla_\theta f(t, \Theta_t, x) \cdot \sigma_t^\Theta dW_t)$$

such that F is a stochastic random field with local characteristics β^F and γ^F given by

$$\beta^F(t, x) = L_t^\Theta f(t, \Theta_t, x), \quad \gamma^F(t, x) = (\sigma_t^\Theta)^T \nabla_\theta f(t, x, \Theta_t).$$

Let now X be another real continuous semimartingale $dX_t = \mu_t^X dt + \sigma_t^X .dW_t$ and L^X its diffusion operator. We now compute the dynamics of $F(t, X_t) := f(t, Y_t, X_t)$ by the classical Itô's formula applied to the vector (Y_t, X_t) and compare it with the Itô-Ventzel's formula. We obtain

$$\begin{aligned} dF(t, X_t) &= L_t^{\Theta, X} f(t, \Theta_t, X_t) dt + \left((\sigma_t^\Theta)^T \nabla_\theta f(t, \Theta_t, X_t) + (\sigma_t^X)^T f'_x(t, \Theta_t, X_t) \right) .dW_t \\ &= \beta^F(t, X_t) dt + \gamma^F(t, X_t) .dW_t + \gamma_x^F(t, X_t) .\sigma_t^X dt + L_t^X f(t, \Theta_t, X_t) dt. \end{aligned}$$

Denoting $\Delta_{\theta, x} f := \frac{\partial}{\partial x} (\nabla_\theta f)$, $L_t^{\Theta, X} := \frac{\partial}{\partial t} + L_t^\Theta(t, \theta, x) + L_t^X(t, \theta, x) + \sigma_t^X .\sigma_t^\Theta \Delta_{\theta, x}(t, \theta, x)$, and then,

$$\gamma_x(t, X_t) .\sigma_t^X = \sigma_t^X .\sigma_t^\Theta \Delta_{\theta, x}(t, \Theta_t, X_t).$$

2.2 Stochastic PDE of \mathcal{X} -consistent Dynamic Utilities

Using the same ideas as in interest rate modeling when studying the dynamics of the forward rates, or in the stochastic volatility models to characterize the drift of the stochastic implied volatility, we show how the consistency property constraints the random fields $\beta(t, x)$ and $\gamma(t, x)$ in terms of the random field U , its derivatives and the market parameters (r_t, η_t^σ) .

Lemma 2.3 (Drift Constraint). *Let U be a progressive utility of class $\mathcal{C}^{(2)}$ in the sense of Kunita with local characteristics (β, γ) as in (6). Then, for any admissible portfolio X^κ ,*

$$\begin{aligned} dU(t, X_t^\kappa) &= \left(U_x(t, X_t^\kappa) X_t^\kappa \kappa_t + \gamma(t, X_t^\kappa) \right) .dW_t \\ &+ \left(\beta(t, X_t^\kappa) + U_x(t, X_t^\kappa) r_t X_t^\kappa + \frac{1}{2} U_{xx}(t, X_t^\kappa) \mathcal{Q}(t, X_t^\kappa, X_t^\kappa \kappa_t) \right) dt, \end{aligned}$$

where $\mathcal{Q}(t, x, \kappa) := \|x\kappa\|^2 + 2x\kappa . \left(\frac{U_x(t, x) \eta_t^\sigma + \gamma_x(t, x)}{U_{xx}(t, x)} \right)$.

Let γ_x^σ be the orthogonal projection of γ_x on \mathcal{R}^σ ; the previous expression is only depending on γ_x^σ since $\kappa \in \mathcal{R}^\sigma$. Let $\mathcal{Q}^*(t, x) = \inf_{\kappa \in \mathcal{R}^\sigma} \mathcal{Q}(t, x, \kappa)$; the minimum of this quadratic form is achieved at the optimal policy κ^* given by

$$\begin{cases} x\kappa_t^*(x) &= -\frac{1}{U_{xx}(t, x)} (U_x(t, x) \eta_t^\sigma + \gamma_x^\sigma(t, x)) \\ x^2 \mathcal{Q}^*(t, x) &= -\frac{1}{U_{xx}(t, x)^2} \|U_x(t, x) \eta_t^\sigma + \gamma_x^\sigma(t, x)\|^2 = -\|x\kappa_t^*(x)\|^2. \end{cases} \quad (10)$$

Proof. (i) The first assertion of the lemma is a direct consequence of Itô-Ventzel formula applied to the composite process $(U(t, X_t^\kappa))_{t \geq 0}$, where X_t^κ is an admissible wealth process with dynamics given by (4), $dX_t^\kappa = X_t^\kappa (r_t dt + \kappa_t . (dW_t + \eta_t^\sigma dt))$.

(ii) Let us now check the second assertion. In the minimization program, we can replace the vector $\gamma_x(t, x)$ by its orthogonal projection $\bar{\gamma}_x^\sigma(t, x)$ on \mathcal{R}_t^σ that yields to equation (10). Moreover, the minimum is given by $\mathcal{Q}^*(t, x) = -\|x\kappa_t^*(x)\|^2$. \square

This lemma suggests the constraint on the drift β implying the consistency condition. The idea of the next theorem is to reformulate this constraint as a natural candidate for β .

Theorem 2.4 (Utility-SPDE). *Let U be a progressive utility of class $\mathcal{C}^{(2)}$ in the sense of Kunita with local characteristics (β, γ) , and risk tolerance coefficient $\alpha_t^U(t, x) = -\frac{U_x(t, x)}{U_{xx}(t, x)}$. Given the volatility vector γ , we introduce the utility risk premium $\eta^U(t, x) = \frac{\gamma_x(t, x)}{U_x(t, x)}$, whose the orthogonal decomposition on \mathcal{R}_t^σ and $\mathcal{R}_t^{\sigma, \perp}$ is*

$$\eta^U(t, x) = \eta^{U, \sigma}(t, x) + \eta^{U, \perp}(t, x), \quad \eta^{U, \sigma}(t, x) \in \mathcal{R}_t^\sigma, \quad \eta^{U, \perp}(t, x) \in \mathcal{R}_t^{\sigma, \perp}.$$

With these notations, the quadratic form $x^2 \mathcal{Q}(t, x, \kappa) = \|x\kappa_t\|^2 - 2\alpha^U(t, x)(x\kappa_t) \cdot (\eta_t^\sigma + \eta^{U, \sigma}(t, x))$ achieves its minimum at

$$x\kappa_t^*(x) = -\frac{1}{U_{xx}(t, x)}(U_x(t, x)\eta_t^\sigma + \gamma_x^\sigma(t, x)) = \alpha^U(t, x)(\eta_t^\sigma + \eta^{U, \sigma}(t, x)) \quad (11)$$

a) Assume the drift constraint to be Hamilton-Jacobi-Bellman nonlinear type

$$\beta(t, x) = -U_x(t, x)r_t x + \frac{1}{2}U_{xx}(t, x)\|x\kappa_t^*(t, x)\|^2 \quad (12)$$

Then the progressive utility is solution of the following HJB-SPDE

$$dU(t, x) = -U_x(t, x) \left[xr_t + \frac{1}{2}\alpha^U(t, x)\|\eta_t^\sigma + \eta^{U, \sigma}(t, x)\|^2 \right] dt + \gamma(t, x) \cdot dW_t,$$

and for any admissible wealth X_t^κ , the process $U(t, X_t^\kappa)$ is a supermartingale.

b) Furthermore, if we assume that $\kappa^*(t, x)$ is sufficiently smooth so that for any initial wealth $x > 0$ the equation

$$dX_t^* = X_t^* [r_t dt + \kappa_t^*(X_t^*) \cdot (dW_t + \eta_t^\sigma dt)] \quad (13)$$

has at least one positive solution X^* , then $U(t, X_t^*)$ is a local martingale.

c) Moreover, if the local martingale $(U(t, X_t^*))_{t \geq 0}$ is a martingale then the progressive utility U is a \mathcal{X} -consistent stochastic utility with optimal wealth process X^* .

This theorem proves that the pair consisting on the investment universe and the derivative with respect to x of the volatility denoted by γ_x describes completely the evolution of the stochastic utility U . The drift $\beta(t, x)$ may be interpreted as the best compromise between the investment universe and volatility of the utility represented by the random field γ . Hence $\beta(t, x)$ can also be interpreted as the best combination between the market risk premium η^σ and utility risk premium represented by $\eta^{U, \sigma}(t, x) = \frac{\gamma_x^\sigma}{U_x}$.

The assumption (12) on the drift β is a sufficient condition under which the consistence with the investment universe of the second assertion of Definition 1.1 is satisfied. Nevertheless, additional assumptions are needed on the existence of the wealth process X^* for which $U(t, X_t^*)$ is a martingale. This explains the assumptions of the second part of the result.

The Utility-SPDE poses several challenges. It is a fully nonlinear and not elliptic SPDE; the latter is a direct consequence of the "forward in time" nature of the involved stochastic optimization problem, for which there is no maximum principle. Thus, existing results of existence, uniqueness and regularity of weak (viscosity) solutions are not directly applicable. An additional difficulty comes from the fact that the volatility coefficient may depend on higher order derivatives of U , in which case the SPDE cannot be turned into a regular PDE with random coefficients, using the method of stochastic characteristics. To overcome this difficulties we propose a new method based on stochastic change of variable; this method, that we call "stochastic flow method", allows us to construct explicit solutions of this Utility-SPDE. This will be the subject of Section 4.

Proof. All assertions are simple consequences of the previous lemma, since by the assumption on $\beta(t, x)$, $\beta(t, x) + x U_x(t, x) r_t + \frac{x^2 U_{xx}(t, x)}{2} \mathcal{Q}(t, x, \kappa) \leq 0$, *a.s.* $\forall \kappa \in \mathcal{R}^\sigma$, with equality for $\kappa^*(t, x)$. Therefore, $U(t, X_t^\kappa)$ is a positive supermartingale for any admissible strategy, moreover if equation (13) has a solution X^* , then the process $(U(t, X_t^*))_{t \geq 0}$ is a local martingale.

The additional assumption that $U(t, X_t^*)$ is a true martingale yields the characterization of the $U(t, x)$ as \mathcal{X} -consistent utility. \square

Example: Change of probability and numeraire in standard utility function

In this paragraph, we study the β -HJB constraint (12) of Theorem 2.4 in the case of a \mathcal{X} -consistent stochastic utilities obtained by combining a standard utility function v with some positive processes N and Z . The advantage here is that the drift β and the volatility γ of the utility are given explicitly from v , N and Z .

Let v an \mathcal{C}^2 utility function and let N and Z two positive processes satisfying

$$\frac{dN_t}{N_t} = \mu_t^N dt + \sigma_t^N .dW_t, \quad \frac{dZ_t}{Z_t} = \mu_t^Z dt + \sigma_t^Z .dW_t, \quad Z_0 = 1.$$

Define the strictly increasing and concave process (with respect to x) U by $U(t, x) = Z_t v(x/N_t)$. Applying Itô's Lemma and using identities $U_x(t, x) = \frac{Z_t}{N_t} v_x(\frac{x}{N_t})$, $U_{xx}(t, x) = \frac{Z_t}{N_t^2} v_{xx}(\frac{x}{N_t})$, it is straightforward to check that, for $x > 0$, the process $(U(t, x))_{t \geq 0}$ satisfies

$dU(t, x) = \beta(t, x)dt + \gamma(t, x).dW_t$ where the local characteristics β and γ are given by

$$\beta(t, x) = U(t, x)\mu_t^Z + xU_x(t, x)(-\mu_t^N + \|\sigma_t^N\|^2 - \sigma_t^N \cdot \sigma_t^Z) + \frac{1}{2}x^2U_{xx}(t, x)\|\sigma_t^N\|^2 \quad (14)$$

$$\gamma(t, x) = U(t, x)\sigma_t^Z - xU_x(t, x)\sigma_t^N. \quad (15)$$

Given that U is a progressive utility, we are interested in establishing conditions on the triplet (v, N, Z) for the drift β satisfies the HJB constraint (12).

Proposition 2.5. *Let v be an utility function.*

(i) *Except the case where v is a power or exponential utility, the process U defined by $U(t, x) = Z_t v(x/N_t)$ is a \mathcal{X} -Consistent stochastic utility iff Z is a martingale, ZX^κ/N , $\kappa \in \mathcal{R}^\sigma$ are positive local martingales and $\sigma \in \mathcal{R}^\sigma$. In this case the optimal policy is given by $\kappa_t^* = \sigma_t^N$.*

(ii) *If v is a power or exponential utility, then Condition : " Z is martingale, ZX^κ/N is a supermartingale for any $\kappa \in \mathcal{R}^\sigma$ " is not a necessary condition.*

– *If v is a power utility with risk aversion a , it suffices that the parameters of Z and N satisfied,*

$$\frac{1}{a}\mu_t^Z + r_t - \mu_t^N + \sigma_t^N \cdot \eta_t^\sigma - \sigma_t^{N,\perp} \cdot \sigma_t^{Z,\perp} + \frac{1}{2(1-a)}\|\eta_t^\sigma - \sigma_t^{N,\sigma} + \sigma_t^{Z,\sigma}\|^2 + \frac{1+a}{2}\|\sigma_t^{N,\perp}\|^2 = 0.$$

– *If v is an exponential utility it suffices to take Z and N satisfied*

$$\mu^N = r + \sigma^N \cdot \eta^\sigma, \quad \mu^Z = \frac{1}{2}\|\eta^\sigma - \sigma^{N,\sigma} + \sigma^{Z,\sigma}\|^2, \quad \sigma^N \in \mathcal{R}^\sigma$$

This result gives sufficient conditions under which U , defined above, is a \mathcal{X} -Consistent stochastic utilities. Note also that this example generalize that in [33] where the authors consider the case where u is an exponential utility and provides a similar sufficient condition.

Proof. To facilitate the exposition, let us denote by $\hat{r} = r - \mu^N + \sigma^N \cdot \eta^\sigma$, $\hat{\eta} = \eta^\sigma - \sigma^N$. The volatility vector γ being given by equation (15), Lemma 2.3 gives the optimal policy κ^*

$$\kappa_t^*(x) = -\frac{1}{xU_{xx}} \left(-xU_{xx}(t, x)\sigma_t^{N,\sigma} + U_x(t, x)(\hat{\eta}_t^\sigma + \sigma_t^{Z,\sigma}) \right). \quad (16)$$

Then, the drift of the utility process U satisfies the HJB constraint (12) if and only if,

$$U\mu_t^Z + U_x\hat{r}_t x - xU_x\sigma_t^{N,\perp} \cdot (\hat{\eta}_t^\sigma + \sigma_t^{Z,\perp}) - \frac{(U_x)^2}{2U_{xx}}\|\hat{\eta}_t^\sigma + \sigma_t^{Z,\sigma}\|^2 + \frac{x^2U_{xx}}{2}\|\sigma_t^{N,\perp}\|^2(t, x) = 0.$$

Using that $U(t, x) = Z_t v(x/N_t)$ and simplifying by $xv(x)Z$, it follows from the definition of $\hat{\eta}$ and \hat{r} that $\forall t \geq 0, x > 0$

$$\frac{v}{xv_x}\mu_t^Z + r_t - \mu_t^N + \sigma_t^N \cdot \eta_t^\sigma - \sigma_t^{N,\perp} \cdot \sigma_t^{Z,\perp} - \frac{v_x}{2xv_{xx}}\|\hat{\eta}_t^\sigma + \sigma_t^{Z,\sigma}\|^2 + \left(1 + \frac{xv_{xx}}{2v_x}\right)\|\sigma_t^{N,\perp}\|^2(t, x) = 0. \quad (17)$$

The case: v/xv_x and v_x/xv_{xx} are proportional, in turn v is a power or exponential utility.

• $v(x) = x^a/a$. Then equation (17) becomes, $\forall t \geq 0$

$$\frac{1}{a}\mu_t^Z + r_t - \mu_t^N + \sigma_t^N \cdot \eta_t^\sigma - \sigma_t^{N,\perp} \cdot \sigma_t^{Z,\perp} + \frac{1}{2(1-a)} \|\eta_t^\sigma - \sigma_t^{N,\sigma} + \sigma_t^{Z,\sigma}\|^2 + \frac{1+a}{2} \|\sigma_t^{N,\perp}\|^2 = 0.$$

• $v(x) = -\frac{1}{c}e^{-cx}$, $c > 0$. Then $\forall t \geq 0$, $x > 0$,

$$\mu_t^Z - \frac{1}{2} \|\eta_t^\sigma - \sigma_t^{N,\sigma} + \sigma_t^{Z,\sigma}\|^2 - cx \left(r_t - \mu_t^N + \sigma_t^N \cdot \eta_t^\sigma - \sigma_t^{N,\perp} \cdot \sigma_t^{Z,\perp} \right) + \left(\frac{cx^2}{2} - cx \right) \|\sigma_t^{N,\perp}\|^2 = 0.$$

Obviously, this is a second order polynomial identically null, consequently all coefficients are nulls, i.e., $\hat{r} = r - \mu^N + \sigma^N \cdot \eta^\sigma = 0$, $\mu^Z = \frac{1}{2} \|\eta^\sigma - \sigma^{N,\sigma} + \sigma^{Z,\sigma}\|^2$, $\sigma \in \mathcal{R}^\sigma$

Second case : v/xv' and v'/xv'' are not proportional, then it is immediate that all terms of (17) are equal to zero, in turn $\tilde{r} = 0$, $\mu^Z = 0$, $\sigma \in \mathcal{R}^\sigma$, $\eta^\sigma - \sigma^N + \sigma^Z \in \mathcal{R}^{\sigma,\perp}$ and hence the optimal strategy κ^* in (16) is simply σ^N .

To summarize the situation: Z is a martingale, X^κ/N is a martingale under the probability Q^Z defined by $dQ^Z/dP = Z$ and $\sigma^N \in \mathcal{R}^\sigma$.

□

As in the classical theory of optimal choice of portfolio in expected utility framework, the process $U_x(t, X_t^*)$ has nice properties and a central place in the dual problem we introduce in the next section.

Proposition 2.6. *Let U be a progressive utility of class $\mathcal{C}^{(3)}$ in the sense of Kunita, with local characteristics (β, γ) . Assume that all assumptions of Theorem 2.4 hold true, in particular that X^* is solution of $dX_t^* = X_t^* [r_t dt + \kappa_t^*(X_t^*) \cdot (dW_t + \eta_t^\sigma dt)]$.*

Let L^ be the diffusion operator of X^* , $L_{t,x}^* = \frac{1}{2} \|x\kappa_t^*(x)\|^2 \frac{\partial^2}{\partial x^2} + \{r_t x + (x\kappa_t^*(x)) \cdot \eta_t^\sigma\} \frac{\partial}{\partial x}$.*

i) Then, U_x is of class $\mathcal{C}^{(2)}$ in the sense of Kunita with local characteristics (β_x, γ_x) and

$$\begin{cases} \gamma_x(t, x) + U_{xx}(t, x)(x\kappa_t^*(x)) = -U_x(t, x)\eta_t^\sigma + \gamma_x^\perp(t, x) \\ \beta_x(t, x) = -U_x(t, x)r_t - L_{t,x}^* U_x(t, x) - (x\kappa_t^*(x)) \cdot \gamma_{xx}(t, x) \end{cases}$$

ii) The semimartingale $U_x(t, X_t^)$ has the following decomposition*

$$dU_x(t, X_t^*) = U_x(t, X_t^*) [-r_t dt + (\eta_t^{U,\perp}(t, X_t^*) - \eta_t^\sigma) \cdot dW_t] \quad (18)$$

In particular, for any admissible wealth process X^κ ($\kappa \in \mathcal{R}^\sigma$), $(X_t^\kappa U_x(t, X_t^))$ is a local martingale and a martingale if $X^\kappa = X^*$.*

This result shows that $U_x(t, X_t^*)$ plays the role of a state price density process.

Proof. Theorem 2.1 shows that U_x is of class $\mathcal{C}^{(2)}$ in the sense of Kunita, with local characteristics (β_x, γ_x) . On the other hand, by Theorem 2.4, we have the identities $\beta(t, x) = -xU_x(t, x)r_t + \frac{1}{2}x^2U_{xx}(t, x)\|\kappa_t^*(x)\|^2$ and $U_{xx}(t, x)(x\kappa_t^*(x)) = -(U_x(t, x)\eta_t^\sigma + \gamma_x^\sigma(t, x))$. This second identity is useful to calculate $\frac{1}{2}U_{xx}(t, x)\partial_x(\|x\kappa_t^*(x)\|^2) = U_{xx}(t, x)((x\kappa_t^*(x)) \cdot \partial_x(x\kappa_t^*(x)))$. Taking the derivative with respect to x in this second identity, it follows that

$$U_{xxx}(t, x)(x\kappa_t^*(x)) + U_{xx}(t, x)\partial_x(x\kappa_t^*(x)) = -(U_{xx}(t, x)\eta_t^\sigma + \gamma_{xx}^\sigma(t, x)).$$

In fact we are interested in the inner product with the vector $x\kappa_t^*(x)$ that yields to the following equality written in an appropriate form

$$\begin{aligned} & \frac{1}{2}U_{xxx}(t, x)\|x\kappa_t^*(x)\|^2 + U_{xx}(t, x)((x\kappa_t^*(x)) \cdot \partial_x(x\kappa_t^*(x))) \\ &= -\left\{\frac{1}{2}U_{xxx}(t, x)\|x\kappa_t^*(x)\|^2 + U_{xx}(t, x)\eta_t^\sigma \cdot (x\kappa_t^*(x))\right\} - \gamma_{xx}^\sigma(t, x) \cdot (x\kappa_t^*(x)). \end{aligned}$$

It is easy to recognize the first line as the derivative of $\frac{1}{2}x^2U_{xx}(t, x)\|\kappa_t^*(x)\|^2$ and the second line as related to the diffusion operator $L_{t,x}^*$. In this form the relation $\beta_x(t, x) = -U_x(t, x)r_t - L^*U_x(t, x) - (x\kappa_t^*(x)) \cdot \gamma_{xx}(t, x)$ is easy to establish.

Then we have all the elements to calculate the dynamics of $U_x(t, X_t^*)$ using Itô-Ventzel formula

$$\begin{aligned} dU_x(t, X_t^*) &= (\gamma_x(t, X_t^*) + U_{xx}(t, X_t^*)X_t^*\kappa_t^*(X_t^*)) \cdot dW_t \\ &+ \left\{\beta_x(t, X_t^*) + L^*U_x(t, X_t^*) + \gamma_{xx}(t, X_t^*) \cdot (X_t^*\kappa_t^*(X_t^*))\right\} dt. \end{aligned}$$

Note that in the last inner product, we can replace $\gamma_{xx}(t, X_t^*)$ by its orthogonal projection $\gamma_{xx}^\sigma(t, X_t^*)$ on the space \mathcal{R}^σ . Thanks to the previous calculation, the expression in the brackets of the second line is exactly $-r_t U_x(t, X_t^*)$. The diffusion coefficient may also be simplified in $-U_x(t, X_t^*)\eta_t^\sigma + \gamma_x^{\perp}(t, X_t^*)$. So that, we obtained the remarkably simple dynamics of $U_x(t, X_t^*)$

$$dU_x(t, X_t^*) = U_x(t, X_t^*) \left[-r_t dt + (\eta_t^{U,\perp}(t, X_t^*) - \eta_t^\sigma) \cdot dW_t \right].$$

We now consider the proof of the last sentence. Given an admissible wealth process, $dX_t^\kappa = X_t^\kappa(r_t dt + (\kappa_t(dW_t + \eta_t^\sigma dt)))$, standard Itô's calculus provides an explicit form for the dynamics of $Z_t^\kappa = X_t^\kappa U_x(t, X_t^*)$ as

$$\begin{aligned} \frac{dZ_t^\kappa}{Z_t^\kappa} &= \frac{dX_t^\kappa}{X_t^\kappa} + \frac{dU_x(t, X_t^*)}{U_x(t, X_t^*)} + \left\langle \frac{dX_t^\kappa}{X_t^\kappa}, \frac{dU_x(t, X_t^*)}{U_x(t, X_t^*)} \right\rangle \\ &= [\kappa_t - \eta_t^\sigma + \eta_t^{U,\perp}(t, X_t^*)] \cdot dW_t, \end{aligned}$$

which implies that $Z_t^\kappa = X_t^\kappa U_x(t, X_t^*)$ is a local martingale for any $\kappa \in \mathcal{R}^\sigma$. In particular the volatility coefficient of Z^* is $\sigma_t^{Z^*} = (\kappa^*(t, X_t^*) - \eta_t^\sigma + \eta_t^{U,\perp}(t, X_t^*))$. To show that the

positive local martingale Z^* is a martingale, we use the concavity of the utility U , with the fact that $U(0) = 0$. As consequence, $Z_t^* = X_t^* U_x(t, X_t^*) \leq U(t, X_t^*)$; since by assumption $U(\cdot, X_t^*)$ is a martingale, the same property is true for Z^* . This completes the proof. \square

3 Duality

After having introduced the consistent stochastic progressive utilities and established the associated SPDEs, several questions remain open at this stage. Indeed, we have shown that the volatility γ of these utilities plays a fundamental role since it completely describes the stochastic dynamics of utilities and the optimal policy. In particular, the projection of the derivative of the volatility γ_x^σ is proportional to the optimal policy. It now remains to give an interpretation of the orthogonal part γ_x^\perp . The concavity of $U(t, x)$ and the classical theory lead naturally to introduce the convex conjugate function $\tilde{U}(t, y)$ (also called the Legendre-Fenchel transform) of $U(t, x)$. We want to show that these conjugate random fields is solution of a dual Utility-SPDE whose the optimal risk premium is related to $\eta^{U, \perp} = \gamma_x^\perp / U_x$.

In the classical theory of concave function f and its conjugate \tilde{f} , the monotone functions f_x and $-\tilde{f}_y$ are inverse of each other, $-\tilde{f}_y(y) = f_x^{-1}(y)$; in the stochastic framework, monotone functions are replaced by stochastic monotone flows and there inverse flows whose dynamics are given by the Itô-Ventzel formula. For simplicity, we present these results separately and in an appropriate form.

3.1 Local characteristics of inverse flows

Let ϕ and ψ be two one-dimensional stochastic flows, with dynamics

$$\begin{aligned} d\phi(t, x) &= \mu^\phi(t, x)dt + \sigma^\phi(t, x).dW_t, \\ d\psi(t, x) &= \mu^\psi(t, x)dt + \sigma^\psi(t, x).dW_t. \end{aligned}$$

From Itô-Ventzel's formula, under regularity assumptions, the compound random field $\phi \circ \psi(t, x) = \phi(t, \psi(t, x))$ is a semimartingale whose the characteristics are given explicitly from those of ϕ and ψ and their derivatives.

Theorem 3.1. *Suppose that ϕ is a random field, regular in the sense of Kunita, and $\psi(t, x)$ is a continuous semimartingale. Then the random field $\phi \circ \psi(t, x) = \phi(t, \psi(t, x))$ is a continuous*

semimartingale with decomposition

$$\begin{aligned}
d(\phi \circ \psi)(t, x) &= \mu^\phi(t, \psi(t, x))dt + \sigma^\phi(t, \psi(t, x)).dW_t \\
&+ \phi_x(t, \psi(t, x))d\psi(t, x) + \frac{1}{2}\phi_{xx}(t, \psi(t, x))\|\sigma^\psi(t, x)\|^2 dt \\
&+ \sigma_x^\phi(t, \psi(t, x)).\sigma^\psi(t, x)dt.
\end{aligned} \tag{19}$$

The volatility of the compound process $\phi \circ \psi$ is given by

$$\sigma^{\phi \circ \psi}(t, x) = \sigma^\phi(t, \psi(t, x)) + \phi_x(t, \psi(t, x))\sigma^\psi(t, x).$$

The next proposition, used several times throughout this paper, gives the decomposition of the inverse of a strictly monotone stochastic flow.

Proposition 3.2 (Inverse flow dynamics). *Let ϕ be a strictly monotone flow, regular in the sense of Kunita, with characteristics $(\mu^\phi(t, x), \sigma^\phi(t, x))$. The inverse process ξ of ϕ is defined on the range of ϕ by $\phi(t, \xi(t, y)) = y$.*

i) *The inverse flow $\xi(t, y)$ has a dynamics given in terms of the old parameters by:*

$$d\xi(t, y) = -\xi_y(t, y)(\mu^\phi(t, \xi)dt + \sigma^\phi(t, \xi).dW_t) + \frac{1}{2}\partial_y(\xi_y(t, y)\|\sigma^\phi(t, \xi)\|^2)dt$$

ii) *With the new parameters, using that $\sigma^\xi(t, y) = -\xi_y(t, y)\sigma^\phi(t, \xi(t, y))$*

$$d\xi(t, y) = \sigma^\xi(t, y).dW_t + \left(\frac{1}{2}\partial_y\left(\frac{\|\sigma^\xi(t, y)\|^2}{\xi_y(t, y)}\right) - \mu^\phi(t, \xi(t, y))\xi_y(t, y)\right)dt$$

It is interesting to observe that the local characteristics of the inverse flow ξ can be easily interpreted as some derivatives. This point will play a crucial role in the sequel. The mathematical formulation of this remark is given in the following corollary, where the assumptions of Proposition 3.2 are made.

Corollary 3.3. *Let $(\Phi(t, x), M^\phi(t, x), \Sigma^\phi(t, x))$ be the primitives, null at $x = 0$, of $\phi(t, x)$, $(\mu^\phi(t, x), \sigma^\phi(t, x))$ respectively, so that the $\Phi(t, x)$ dynamics is $d\Phi(t, x) = M^\phi(t, x)dt + \Sigma^\phi(t, x).dW_t$.*

Then, the dynamics of the random field $\Xi(t, y) = \int_0^y -\xi(t, z)dz$ is

- *With old parameters,*

$$d\Xi(t, y) = \Sigma^\phi(t, \xi(t, y)).dW_t + M^\phi(t, \xi(t, y))dt + \frac{1}{2}\Xi_{yy}(t, y)\|\Sigma_x^\phi(t, \xi(t, y))\|^2 dt$$

- *In new variables, with the notations $M^\xi(t, y) = M^\phi(t, -\Xi_y(t, y))$ and $\Sigma^\xi(t, y) = -\Sigma^\phi(t, -\Xi_y(t, y))$,*

$$d\Xi(t, y) = \Sigma^\xi(t, y).dW_t + \left(M^\xi(t, y) + \frac{1}{2}\frac{\|\Sigma_y^\xi(t, y)\|^2}{\Xi_{yy}(t, y)}\right)dt.$$

Proof. of the proposition and the corollary. The proof is essentially based on the generalized Itô's formula established in the Appendix. For simplicity, we denote by (μ^ξ, σ^ξ) the local characteristic of ξ assumed to be regular. By Itô-Ventzel's formula, we have

$$\begin{aligned} d\phi(t, \xi(t, y)) &= 0 \\ &= \mu^\phi(t, \xi(t, y))dt + \sigma^\phi(t, \xi(t, y)).dW_t + \phi_x(t, \xi(t, y))d\xi(t, y) \\ &\quad + \frac{1}{2}\phi_{xx}(t, \xi(t, y)) \langle d\xi(t, y) \rangle + \sigma_x^\phi(t, \xi(t, y)).\sigma^\xi(t, y)dt \end{aligned}$$

Recalling the following identities

$$\phi_x(t, \xi(t, y)) = \frac{1}{\xi_y(t, y)}, \quad \phi_{xx}(t, \xi(t, y)) = -\frac{\xi_{yy}(t, y)}{(\xi_y(t, y))^3},$$

we can express the parameters of the decomposition in terms of $\xi, \xi_y,$ and ξ_{yy} and the diffusion coefficient $\sigma^\xi(t, y)$ of ξ , since $\sigma^\xi(t, y) = -\xi_y(t, y)\sigma^\phi(t, \xi(t, y))$. It is immediate that

$$\mu^\xi(t, y) = -\xi_y(t, y)\mu^\phi(t, \xi(t, y)) + \xi_y(t, y) \langle \sigma^\phi(t, \xi(t, y)) \rangle + \frac{1}{2}\xi_{yy}(t, y)\|\sigma^\phi(t, \xi(t, y))\|^2$$

In terms of the stochastic random fields μ^ϕ and σ^ϕ , this may be written as

$$\mu^\xi(t, y) = -\xi_y(t, y)\mu^\phi(t, \xi(t, y)) + \frac{1}{2}\partial_y [\xi_y(t, y)\|\sigma^\phi(t, \xi(t, y))\|^2].$$

In terms of their own parameters, it follows from the strict monotonicity of ξ that

$$\mu^\xi(t, y) = -\xi_y(t, y)\mu^\phi(t, \xi(t, y)) + \frac{1}{2}\partial_y [\|\sigma^\xi(t, y)\|^2/\xi_y(t, y)].$$

The proof of Proposition 3.2 is now complete.

The proof of Corollary 3.3 is achieved, first by reconciling the results of the previous proposition and the following identities,

$$(\Phi_x)^{-1}(t, y) = -\Xi_y(t, y), \quad \Phi_{xx}(t, -\Xi_y(t, y)) = -\frac{1}{\Xi_{yy}(t, y)}, \quad \text{and} \quad -C_x(t, -\Xi_y(t, y)) = \frac{D_y(t, y)}{\Xi_{yy}(t, y)}$$

and second by integrating with respect to y , using the initial condition $\Xi(t, 0) = 0$. \square

3.2 Convex conjugate of consistent stochastic utility and dual utility SPDE's

We now define the convex conjugate of a consistent stochastic utility.

Definition 3.1. Let \tilde{U} be the convex conjugate random field of the \mathcal{X} -consistent stochastic utility U , given by definition for $t \geq 0$,

$$\tilde{U}(t, y) \stackrel{def}{=} \inf_{x>0, x \in Q^+} (U(t, x) - xy)$$

$\tilde{U}(t, y)$ is a progressive decreasing convex random field, with first derivative $\tilde{U}_y(t, y) = -I(t, y)$, where $I(t, y) = U_x(t, \cdot)^{-1}(y)$ is the inverse flow of the decreasing flow $U_x(t, x)$.

The formula of inverse flows yields easily to the dynamics of $\tilde{U}(t, y)$ from the dynamics of $U(t, x)$. Based on this remark and Corollary 3.3, we derive a stochastic partial differential equation whose a solution is the convex conjugate processes of consistent stochastic utility.

Theorem 3.4. *Let U be a consistent progressive utility of class $\mathcal{C}^{(3)}$, in the sense of Kunita, satisfying the β constraint (12) with risk tolerance α^U and utility risk premium $\eta^U(t, x) = \frac{\gamma_x(t, x)}{U_x(t, x)}$. Let \tilde{U} be the dual convex conjugate of U , null if $y = 0$. Then*

(i) *In old variables, \tilde{U} satisfies $d\tilde{U}(t, y) = \beta^1(t, -\tilde{U}_y(t, y))dt + \gamma(t, -\tilde{U}_y(t, y)).dW_t$, where*

$$\begin{cases} \beta^1(t, x) = \beta(t, x) - \frac{1}{2U_{xx}(t, x)}\|\gamma_x(t, x)\|^2 \\ = U_x(t, x) \left[-xr_t - \frac{1}{2}\alpha^U(t, x) \left(\inf_{\theta \in \mathcal{R}^{\sigma, \perp}} \|\theta_t - (\eta_t^\sigma + \eta^U(t, x))\|^2 - \|\eta^U(t, x)\|^2 \right) \right]. \end{cases} \quad (20)$$

The optimization program is achieved on $\theta^(t, x) = \eta^{U, \perp}(t, x)$.*

(i) *In new variables,*

$$\begin{cases} d\tilde{U}(t, y) = \left[\frac{1}{2\tilde{U}_{yy}(t, y)} (\|\tilde{\gamma}_y(t, y)\|^2 - \|\tilde{\gamma}_y^\sigma(t, y) + y\tilde{U}_{yy}(t, y)\eta_t^\sigma\|^2) + y\tilde{U}_y(t, y)r_t \right] dt + \tilde{\gamma}(t, y).dW_t \\ \tilde{\gamma}(t, y) = \gamma(t, -\tilde{U}_y(t, y)). \end{cases}$$

Furthermore the drift $\tilde{\beta}(t, y) := \beta^1(t, -\tilde{U}_y(t, y))$ is the value of an optimization program achieved on the optimal policy $\nu^(t, y) = \theta^*(t, -\tilde{U}(t, y)) = -\tilde{\gamma}_y^\perp(t, y)/y\tilde{U}_{yy}(t, y)$.*

In particular, $\tilde{\beta}$ can be written us the solution of the following optimization program

$$\hat{\beta}(t, y) = y\tilde{U}_y(t, y)r_t - \frac{1}{2}y^2\tilde{U}_{yy}(t, y) \inf_{\nu_t \in \mathcal{R}^{\sigma, \perp}} \{ \|\nu_t - \eta_t^\sigma\|^2 + 2(\nu_t - \eta_t^\sigma) \cdot \left(\frac{\tilde{\gamma}_y(t, y)}{y\tilde{U}_{yy}(t, y)} \right) \} \quad (21)$$

with $-\tilde{\gamma}_y(t, y)/y\tilde{U}_{yy}(t, y) = \eta^U(t, -\tilde{U}(t, y)) = \gamma_x(t, -\tilde{U}(t, y))/y$.

First, observe that as $-\tilde{U}_y$ is the inverse flows of U_x , the dynamic of the convex conjugate \tilde{U} of U becomes a simple consequence of Corollary 3.3. Second, the orthogonal part of the utility prime risk $\eta^{U, \perp} := \gamma_x^\perp/U_x$ is the optimal policy of the dual problem in (i). Third, given that β is associated with an optimization program the dual drift $\tilde{\beta}$ is also constrained by HJB type relation in the new variables. Then, the convex conjugate is consistent with some given family of the state density processes.

Proof. By regularity assumptions, using Theorem 2.1, $U_x(t, x)$, $\beta_x(t, x)$ and $\gamma_x(t, x)$ are regular enough to apply Itô-Ventzel formula. The assumptions of Proposition 3.2 and Corollary 3.3 are satisfied and hence the dynamics of the convex conjugate is a direct consequence of Corollary 3.3. Let us now recall that the drift $\beta(t, x)$ of $U(t, x)$ is given in Theorem 2.4 by

$$\beta(t, x) = U_x(t, x) \left(-xr_t - \frac{\alpha^U(t, x)}{2} \|\eta^{U, \sigma}(t, x) + \eta_t^\sigma\|^2 \right).$$

To get the desired formula for $\beta^1(t, x)$ in (20), we use the following property of the orthogonal projection: the norm of the projection on \mathcal{R}^σ is the distance to the orthogonal vector space $\mathcal{R}^{\sigma, \perp}$. So, for any vector $a \in R^N$, $\|a^\sigma\|^2 = \inf_{\nu \in \mathcal{R}^{\sigma, \perp}} \|\nu - a\|^2$. By replacing a by $(\eta_t^\sigma + \eta^{U, \sigma}(t, x))$ yields the result.

Now, we focus on the drift $\tilde{\beta}$ of $\tilde{U}(t, y)$ in new variables, using essentially the following identities,

$$\frac{U_x^2(t, -\tilde{U}_y(t, y))}{2U_{xx}(t, -\tilde{U}_y(t, y))} = \frac{1}{2}y^2\tilde{U}_{yy}(t, y), \quad \frac{\tilde{\gamma}_y(t, y)}{\tilde{U}_{yy}(t, y)} = -\gamma_x(t, -\tilde{U}_y(t, y)), \quad \frac{\tilde{\gamma}_y(t, y)}{y\tilde{U}_{yy}(t, y)} = -\eta^U(t, -\tilde{U}_y(t, y))$$

We get the desired formula for $\tilde{\beta}$, i.e.

$$\tilde{\beta}(t, y) = y\tilde{U}_y(t, y)r_t - \frac{1}{2}y^2\tilde{U}_{yy}(t, y) \inf_{\nu_t \in \mathcal{R}^{\sigma, \perp}} \{ \|\nu_t + \eta_t^\sigma\|^2 + 2(\nu_t - \eta_t^\sigma) \cdot \left(\frac{\tilde{\gamma}_y(t, y)}{y\tilde{U}_{yy}(t, y)} \right) \}$$

On the other hand by orthogonal projection on $\mathcal{R}_t^{\sigma, \perp}$ and using the fact that $\eta_t^\sigma \in \mathcal{R}_t^\sigma$, there exists one and only one optimal process ν^* given by

$$\nu_t^*(y) = \frac{-\tilde{\gamma}_y^\perp(t, y)}{y\tilde{U}_{yy}(t, y)} = \eta^{U, \perp}(t, -\tilde{U}_y(t, y))$$

which achieves the proof. \square

Let us now focus on the dual optimization problem.

Definition 3.2 (State price density process). *A Itô semimartingale Y^ν is called a state price density process if for any wealth process X^κ , $\kappa \in \mathcal{R}^\sigma$, $Y^\nu X^\kappa$ is a local martingale. It follows that Y^ν satisfies,*

$$\frac{dY_t^\nu}{Y_t^\nu} = -r_t dt + (\nu_t - \eta_t^\sigma) \cdot dW_t, \quad \nu_t \in \mathcal{R}^{\sigma, \perp}. \quad (22)$$

\mathcal{Y} is the family of all state density processes $\mathcal{Y} := \{Y^\nu, \nu \in \mathcal{R}^{\sigma, \perp}, Y^\nu \text{ satisfies (22)}\}$

Obviously the class \mathcal{Y} is not empty, since taking $\nu \equiv 0$, Y^0 is the classical minimal density process where the pricing of future cash-flow at time T is obtained by first discounting between t and T the cash value at T with the short rate r_t , and then by taking the conditional expected value with respect to the minimal martingale measure. Moreover, any state density process Y^ν is the product of Y^0 by the density martingale $L_t^\nu = \exp\left(\int_0^t \nu_s \cdot dW_s - 1/2 \int_0^t |\nu_s|^2 ds\right)$.

We obtain an interesting interpretation of the volatility risk premium in terms of optimal density process.

Theorem 3.5. *Let U be a consistent progressive utility of class $\mathcal{C}^{(3)}$, in the sense of Kunita, satisfying the β HJB constraint. Then, its conjugate process $\tilde{U}(t, y)$ (convex decreasing stochastic flow) is consistent with the family of state density processes \mathcal{Y} , in the following sense:*

$\tilde{U}(t, Y^\nu)$ is a submartingale for any $Y^\nu \in \mathcal{Y}$, and a martingale for some process $Y^{\nu^} (:= Y^*)$. The optimal process can be chosen as $Y_t^* = U_x(t, X_t^*(-\tilde{u}_y(y)))$, whose the dynamics is*

$$\frac{dY_t^*}{Y_t^*} = -r_t dt + (\nu^*(t, Y_t^*) - \eta_t^\sigma) \cdot dW_t.$$

where the dual optimal parameter $\nu^*(t, y)$ is given by

$$\nu^*(t, y) = \frac{-\tilde{\gamma}_y^\perp(t, y)}{y\tilde{U}_{yy}(t, y)} = \frac{\gamma_x^\perp(t, -\tilde{U}_y(t, y))}{y} = \eta^{U, \perp}(t, -\tilde{U}_y(t, y)).$$

Remark . *Let $\mathcal{Y}(t, x) := U_x(t, X_t^*(x))$, if $X_t^*(x)$ is strictly monotone in x , by taking the inverse $\mathcal{X}(t, x)$, we can obtain $U_x(t, x)$ in terms of $\mathcal{Y}(t, x)$ and $\mathcal{X}(t, x)$.*

Proof. The first assertion of this result is essentially obtained by analogy with the primal problem. Indeed, using the $\tilde{\beta}$ expression's (21), which is

$$\hat{\beta}(t, y) = y\tilde{U}_y(t, y)r_t + \frac{1}{2}y^2\tilde{U}_{yy}(t, y) \sup_{\nu_t \in \mathcal{R}^{\sigma, \perp}} \{-\|\nu_t - \eta_t^\sigma\|^2 - 2(\nu_t - \eta_t^\sigma) \cdot (\frac{\tilde{\gamma}_y(t, y)}{y\tilde{U}_{yy}(t, y)})\}$$

One can easily remark, by analogy to expression of \mathcal{Q} in Lemma 2.3 and that of β , (equation (12), Theorem 2.4), that \tilde{U} is consistent with the family of processes \mathcal{Y} , that is $\tilde{U}(t, Y_t^\nu)$ is a submartingale for any $Y^\nu \in \mathcal{Y}$ and a local martingale for the optimal choice (Theorem 3.4) $\nu^*(t, y) = -\tilde{\gamma}_y^\perp(t, y)/y\tilde{U}_{yy}(t, y) = \gamma_x^\perp(t, -\tilde{U}(t, y))/y$, if there exists a solution to the SDE,

$$\frac{dY_t^{\nu^*}}{Y_t^{\nu^*}} = -r_t dt + (\eta^{U, \perp}(t, -\tilde{U}_y(t, Y_t^{\nu^*})) - \eta_t^\sigma) \cdot dW_t. \quad (23)$$

On the other hand we recall that according to Proposition 2.6 assertion *ii*) $U_x(t, X_t^*)$ satisfies

$$\frac{dU_x(t, X_t^*)}{U_x(t, X_t^*)} = -r_t dt + (\eta^{U, \perp}(t, X_t^*) - \eta_t^\sigma) \cdot dW_t,$$

Note that $Y_t^*(y) = \left(U_x(t, X_t^*(-\tilde{u}_y(y))) \right)_{t \geq 0}$ and that $-\tilde{U}_y(t, Y_t^*(y)) = X_t^*(-\tilde{u}_y(y))$ shows that that Y^* is a solution of (23) which in turn implies the optimality of Y^* .

To conclude, we have to show that $\tilde{U}(Y_t^*(y))$ is not only a local martingale but a "true" martingale, when $U(X_t^*(x))$ is a martingale. Put $x_y = -\tilde{u}_y(y)$, and use that the conjugacy relation implies that $\tilde{U}(Y_t^*(y)) = U(X_t^*(x_y)) - Y_t^*(y)X_t^*(x_y)$ with $Y_t^*(y) = (U_x(t, X_t^*(x_y)))$. Thanks to Proposition 2.6, $U(X_t^*(x_y))$ and $Y_t^*(y)X_t^*(x_y)$ and therefore $\tilde{U}(Y_t^*(y))$ are martingales. \square

Increasing Consistent Utilities An interesting class of consistent utilities is the class of increasing consistent utilities which was studied and fully characterized in the literature by Berrier & al. [12] and Musiela & al. [35]. These utilities have a volatility vector γ identically zero. It is an example where the dual SPDE is easier to study than the primal one. Indeed, taking $\gamma = 0$ it follows that U is a solution of the following PDE $dU(t, x) = \left[\frac{1}{2} \frac{U_x(t, x)^2}{U_{xx}(t, x)} \|\eta_t^\sigma\|^2 \right] dt$ where the convex conjugate \tilde{U} satisfies

$$d\tilde{U}(t, y) = \left[-\frac{1}{2} y^2 \tilde{U}_{yy}(t, y) \|\eta_t^\sigma\|^2 + r_t y \tilde{U}_y(t, y) \right] dt$$

Dividing by dt and writing that $\frac{\tilde{U}(t, y)}{dt} = U_t(t, y)$ yields

$$\tilde{U}_t(t, y)(\omega) = -\frac{1}{2} y^2 \tilde{U}_{yy}(t, y)(\omega) \|\eta_t^\sigma(\omega)\|^2 + r_t(\omega) y \tilde{U}_y(t, y)(\omega)$$

which implies, by convexity, that $t \mapsto \tilde{U}(t, y)$ is a decreasing function. Moreover, it is easy to recognize in this PDE that the right hand side of the equation is nothing other than the operator of diffusion of a geometrical Brownian motion with coefficients $\eta_t^\sigma(\omega)$ and $r_t(\omega)$ $L_{t, y}^{GB}(\omega)$ applied to \tilde{U} : $\tilde{U}_t(t, y)(\omega) = -L_{t, y}^{GB} \tilde{U}(t, y)(\omega)$. From this point, the idea is to look for positive solutions which are space-time harmonic functions of a geometric Brownian motion. Assume that the function $\tilde{U}(t, y)$ is of class \mathcal{C}^3 in y and such that $\frac{\partial^2}{\partial y \partial t} \tilde{U}(t, y) = \frac{\partial^2}{\partial t \partial y} \tilde{U}(t, y)$ is defined and continuous. First, put $\tilde{V}(t, y) = \tilde{U}(t, e^{-\int_0^t r_s ds} y)$ and $A(t) = \int_0^t \|\eta_s^\sigma\|^2 ds$; then \tilde{V} is a solution of the following PDE

$$\tilde{V}_t(t, y) = -\frac{1}{2} A_t(t) y^2 \tilde{V}_{yy}(t, y).$$

Second, define $H(\log(y) - \frac{1}{2} A_t, \frac{1}{2} A_t, \omega) = -\tilde{V}_{t, y}(t, y, \omega)$ and take the change of variable $\tau = \frac{1}{2} A_t$, it is straightforward to check that H solves the backward heat equation

$$H_\tau(\tau, z) + H_{zz}(\tau, z) = 0$$

The solutions of such equation are called space-time harmonic functions. Since the function H is strictly positive, using the result of Widder, D.V [42, 43], F. Berrier & al. [12] and Musiela & al. [35] show the following result which characterizes all decreasing consistent utilities

Theorem 3.6. *Let $U(t, x)$ be a regular random field of class \mathcal{C}^3 on x such that $\frac{\partial^2}{\partial y \partial t} \tilde{U}(t, y) = \frac{\partial^2}{\partial t \partial y} \tilde{U}(t, y)$ is defined and continuous. Assume U satisfies the utility SPDE with $\gamma = 0$ a.s.. Then U is a consistent stochastic utility if and only if there exists a constant $C \in \mathbb{R}$ and a finite Borel measure m , supported on the interval $(0, +\infty)$ with everywhere finite Laplace*

transform, such that

$$\begin{aligned}\tilde{U}(t, y) &= \int_{\mathbb{R}_+^*} \frac{1}{1 - \frac{1}{\alpha}} (1 - y^{1 - \frac{1}{\alpha}} e^{-\frac{1-\alpha}{2\alpha} \int_0^t \|\eta_s\|^2 ds}) dm(\alpha) + C. \\ \tilde{U}_y(0, y) &= - \int_{\mathbb{R}_+^*} y^{-\frac{1}{\alpha}} dm(\alpha)\end{aligned}$$

More over the optimal wealth process is strictly increasing and regular with respect to its initial condition x .

There is an interesting interpretation of these stochastic utilities: At date $t = 0$ the derivative $\tilde{U}_y(0, y)$ can be easily interpreted as the integral $-y^{-\frac{1}{\alpha}}$ weighted by the measure m , which is nothing than the derivative of the convex conjugate of power utility with risk aversion α . Hence, one can imagine that the investor starts from a power utility for which he pull at random the risk aversion α , for any realization α he associate the power utility u^α weighted by m . The derivative of the convex conjugate of his utility at any date t is then the integral of the derivatives of the convex conjugates of power utility where the deterministic measure m becomes stochastic $m_t(d\alpha) := e^{-\frac{1-\alpha}{2\alpha} \int_0^t \|\eta_s\|^2 ds} dm(\alpha)$. Moreover the stochastic measure $m_t(d\alpha)$ is the unique one which ensure that the process \tilde{U} constructed is the derivative of a consistent utility. This interpretation is the starting point of the work [22] where more general method to construct consistent utilities processes from a family of classical utilities functions is developed.

3.3 Change of numéraire

One of the our first interest in progressive utilities was the fact they are consistent with classical transformation in financial market in contrast to the classical utilities functions which are not stable by change of numéraire; so, the value function of the classical portfolio optimization problem is highly dependent on market parameters (r, η) . Moreover, it is easier to work with portfolios that are local martingales rather than semimartingales, that can be obtained using the market numéraire, we find that there is a genuine interest to provide details of this transformation on consistent stochastic utilities.

Proposition 3.7 (Stability by change of numéraire).

Let $U(t, x)$ be a \mathcal{X} consistent stochastic utility, N be a positive semimartingale and denote by \mathcal{X}^N the class of process defined by $\mathcal{X}^N = \{\hat{X} := X/N, X \in \mathcal{X}\}$, then the process V defined by

$$V(t, x) = U(t, xN_t)$$

is a \mathcal{X}^N -consistent stochastic utility in the market of numéraire N if and only if U is an \mathcal{X} -Consistent stochastic utility.

Roughly speaking the proposition says that the notion of \mathcal{X} -Consistent utilities is preserved by change of numeraire. An interpretation of this stability is that if the agent decides to invest in a second market (foreign market) his preferences (risk aversion) are still unchanged, given the uniqueness representation of his preferences.

Furthermore, there exists a market numeraire portfolio $M_t(z) = 1/Y_t^0(1/z) \in \mathcal{X}$, which transforms the classical wealth processes into positive local martingales, without changing the space of constraints \mathcal{R}^σ . When using any state price density process Y^ν and the associated numeraire $N_t^\nu(z) = 1/Y_t^\nu(1/z)$, the local martingale property holds true, but the space of constraints is modified. To show this result it is enough to verify the assertions of Definition 1.1 using identity $V(t, \hat{X}_t) = U(t, X_t)$.

Now, we turn to more quantitative aspects of the change of numeraire. The idea is to proof how, by change of numeraire techniques, we simplify the utility SPDE's of consistent stochastic utilities. To this end, some hypothesis and some regularity are needed.

Assumption 3.1. *The new market-numeraire N , is assumed to satisfy:*

$$\frac{dN_t}{N_t} = \mu_t^N dt + \delta_t^N \cdot dW_t, \quad N_0 = z.$$

The wealth process \hat{X} is defined by $\hat{X}_t := X_t/N_t$ where X denotes the wealth process in the initial market. By Itô's formula, we can easily write the dynamics of \hat{X} ,

$$\frac{d\hat{X}_t(\hat{x})}{\hat{X}_t(\hat{x})} = (r_t - \mu_t^N + \delta_t^N \cdot \eta_t^\sigma) dt + (\kappa_t - \delta_t^N) \cdot (dW_t + (\eta_t^\sigma - \delta_t^N) dt), \quad \hat{x} = \frac{x}{z}.$$

Denoting $\hat{r} = r - \mu^N - \delta^N \cdot \eta^\sigma$ the short interest rate in the new market and by $\hat{\eta} = \eta^\sigma - \delta^N$, the new market price of risk, we get

$$\frac{d\hat{X}_t(\hat{x})}{\hat{X}_t(\hat{x})} = \hat{r}_t dt + (\kappa_t - \delta_t^N) \cdot (dW_t + \hat{\eta}_t dt).$$

Let us now stress the fact that if $\delta^N \in \mathcal{R}^\sigma$ the volatility vector $\hat{\kappa} = \kappa - \delta^N$ of \hat{X} belongs to \mathcal{R}^σ if and only if $\kappa \in \mathcal{R}^\sigma$ since \mathcal{R}^σ is a vector space, and then the optimization problem is unchanged. Hence we get that a consistent utility V in this new market satisfies the same dynamics as U in the initial market only by replacing r , η by \hat{r} , $\hat{\eta}$.

Else if $\delta^{N,\perp} \neq 0$ the optimization problem is quite different and utility-SPDE is modified, as we will see in the next result.

Theorem 3.8. *Let $U(t, x)$ be a \mathcal{X} -consistent stochastic utility satisfying the assumptions of Theorem 2.4. The \mathcal{X}^N -consistent stochastic utility $V(t, x) := U(t, xN_t)$ is a solution of the following stochastic partial differential equation*

$$dV(t, x) = V_x(t, x) \left\{ \frac{1}{2\alpha^V(t, x)} \left(\|\hat{\eta}_t + \hat{\eta}_t^V\|^2 - \|\hat{\eta}_t^\sigma + \hat{\eta}_t^{V,\sigma} - x\alpha^V \delta_t^{N,\sigma}\|^2 \right) - x\hat{r}_t \right\} (t, x) dt + \gamma^V(t, x) \cdot dW_t$$

with $\alpha^V(t, x) := V_x(t, x)/V_{xx}(t, x)$ and $\eta^V(t, x) = \gamma^V(t, x)/V_x(t, x)$ denote the risk tolerance and the utility risk premium of V . The volatility of V is $\gamma^V(t, x) = \gamma^U(t, x N_t) + x V_x(t, x) \delta_t^N$ and the optimal policy \hat{k}^* is given by

$$x \hat{k}_t^*(x) = -\frac{1}{\alpha^V(t, x)} \left(\hat{\eta}_t^\sigma + \eta_t^{V, \sigma} \right) (t, x) - x \delta_t^{N, \sigma}. \quad (24)$$

Furthermore, taking N equal to the numeraire portfolio $1/Y^0$, the market has no risk premium and the ratio $\eta^{V, \sigma}$ has the same impact as a risk premium, but depending on the level of the wealth x at time t . In particular, the previous dynamic of V is simpler

$$dV(t, x) = \frac{(V_x)^2(t, x)}{2V_{xx}(t, x)} \left\| \frac{\gamma_x^{V, \sigma}(t, x)}{V_x(t, x)} \right\|^2 dt + \gamma^V(t, x) \cdot dW_t.$$

and the convex conjugate \tilde{V} of V satisfies

$$d\tilde{V}(t, y) = \frac{1}{2\tilde{U}_{yy}(t, y)} \left\| \tilde{\gamma}_y^\perp(t, y) \right\|^2 dt + \tilde{\gamma}(t, y) \cdot dW_t.$$

Let us comment on the content of this theorem and its relation to the previous results. The dynamic (24) of consistent utilities and the optimal policy (24) are more complicated than ones in the initial framework. We recognize in the optimal policy formula a first term (very similar to that of the initial market) which corresponds to an optimization program without δ^N added to a second one which correspond to a translation by δ^N . This is due to the fact that the dynamic of new wealth processes are a kind of combination of that of the initial market and the dynamic of the state price density processes in the dual problem studied in previous section. It suffices to take δ^N in the range of the matrix σ , to get the SPDE's of the old market and to take $\hat{\eta} = 0$ to get SPDE similar to the dual HJB-SPDE. Finally, in the second part of the result, taking a numeraire with good properties this HJB-SPDE is more simplified.

We close this section, by the the following corollary which is a consequence of the above theorem.

Corollary 3.9. *Under the assumptions of Theorem 3.8, taking $N = H^{r, \eta^\sigma}$, we have*

- $\gamma_x \in \mathcal{R}^\sigma$ implies \tilde{V} is a local martingale and the optimal dual process is constant: $Y^* \equiv 1$.
- $\gamma_x \in \mathcal{R}^{\sigma, \perp}$ implies V is a local martingale and the optimal wealth $X^*(x) \equiv x$.

The new market defined from the first one by change of numeraire $1/H^{r, \eta}$ (the market numeraire) is called a martingale market because new wealths are local martingales.

Proof. The proof of this result is based on the Itô- Ventzel's lemma applied to $U(t, x N_t)$ and the fact that the optimal wealth process $\hat{X}^* = X^*/N$ by definition of V , where X^* denote the optimal portfolio process associated to U . \square

4 Utility Characterization and Stochastic Flows

Method for Solving Utility-Stochastic PDE's

As we mentioned earlier, conventional methods of resolution of SPEs defined from their terminal conditions, such as the method of characteristics, cannot be used to solve utility SPDEs. To overcome this difficulty, we present here a new approach to these utility stochastic partial differential equations, based on a random change of variable. This new approach, that we call "stochastic flows method", is based on the properties of the optimal wealth X^* and the state price density process Y_t^* , supposed to either be a monotonic function of their initial condition. More precisely, we take the initial condition u and an admissible wealth process as given, and ask: what are the conditions on these data, to be an optimal wealth process of some consistent stochastic utility U , and also how to recover U from these information? In the classical expected utility framework, that is the question He and Huang [14] asked in (1992) in a complete market. From one point of view, our problem is easier to solve because we allow ourselves a larger class of utility functions. In particular, we establish in the following that the only restriction is the monotony of the wealth process with respect to the initial wealth, plus some integrability condition. Another difference between this work and that of He and Huang [14] is that we work directly on the path of wealth process while they work with the volatility of the wealth process $\kappa(t, x) = \kappa(t, S_t, x)$ in their setup.

Before presenting this new method, remember that the direct analysis gives us a natural way of finding U from the inputs (u, Y^*, X^*) : Let U be a consistent utility with optimal wealth X^* then, according to Theorem 3.5 and Proposition 2.6, the process Y^* defined by $Y_t^*(u_x(x)) = U_x(t, X_t^*(x))$ is optimal for the dual problem and such that Y^*X^* is a martingale. So, if X^* is strictly increasing with respect to the initial capital, with inverse flow \mathcal{X} , $U_x(t, x) = Y_t^*(u_x(\mathcal{X}(t, x)))$; integrating with respect to x we get U .

From this we assume for the rest of the paper the following main assumption.

Assumption 4.1. *The wealth process $X_t^*(x)$ is assumed to be continuous and increasing in x from 0 to $+\infty$ with $X_t^*(0) = 0$, $X_t^*(+\infty) = +\infty$ for any t and satisfies*

$$\frac{dX_t^*(x)}{X_t^*(x)} = r_t dt + \kappa^*(t, X_t^*(x)) \cdot (dW_t + \eta_t^\sigma dt), \quad \kappa^*(t, x) \in \mathcal{R}_t^\sigma, \quad \forall x > 0, \quad a.s.$$

Denote by $\mathcal{X}(t, z)$ the inverse flow such that $X_t^*(\mathcal{X}(t, z)) = z$.

Financially speaking this hypothesis which may be a consequence of no arbitrage opportunity says that: we do not invest more to earn less. On the other hand, this monotony assumption is true in many examples and, according to the classical results on the the

stochastic differential equations (SDE), is satisfied as soon as $x\kappa^*(t, x)$ is locally uniformly Lipschitz, (see Kunita [25]).

Note also, by conjugacy identity, that monotonicity of $X^*(x)$ implies that the dual process $Y^*(y)$ is, in turn, strictly increasing and therefore invertible with respect to its initial condition y for any date t . The converse property is also true. Consequently, we also do the following hypothesis.

Assumption 4.2. $Y_t^*(y)$ is continuous and increasing in y from $+\infty$ to 0 satisfying

$$\frac{dY_t^*(y)}{Y_t^*(y)} = -r_t dt + (\nu_t^*(Y_t^*(y)) - \eta_t^\sigma) \cdot dW_t, \quad \nu(t, y) \in \mathcal{R}^{\sigma, \perp}, \quad \forall y > 0, \quad a.s. \quad (25)$$

Starting from the idea above, Assumptions 4.1 and 4.2 allow us to compound X^* with the inverse of Y^* and Y^* with the inverse flow of X^* . Under some additional regularity assumptions, we establish one of our main contribution that involves the characterization of any consistent utilities generating X^* as an optimal portfolio. In particular, we give the decomposition of the derivative γ of the volatility vector as an operator of U_x and U_{xx} given κ^* . The second main result of this paper introduce a new method solve the utility stochastic PDE. The idea is to transform SPDE to a system of two stochastic differential equation (SDE). Herein, the method proposed can be used for a large class of SPDE with given optimal policy.

There are two different messages on our approach hence we decide to present the associated results separately. Note that the results of these section can be obtained first on the martingale market and, simply, by using results of Theorem 3.7 we get the similar ones on the initial market.

4.1 Utility Characterization from optimal wealth and state density processes

To fix the idea we consider a given wealth process X^* , a state density price process Y^* and an utility function $u(x)$. The objective is to construct a consistent utility U starting from the function $u(x)$ ($U(0, x) = u(x)$), generating X^* as optimal wealth and Y^* as optimal dual process. According to the necessary analysis above the constructed utility process may satisfies $U_x(\cdot, X^*(x)) = Y^*(u_x(x))$.

To illustrate our approach we first start by proving this result in a special case where we assume that the process $Y^*(y) = yH^{r, \eta^\sigma} = Y^0(y)$ since $\nu^* = 0$ a.s. The advantage of this case is that we can find all messages of our construction and a complete overview of main

calculations. This will better understand the calculations that are made in the general case, because they are more difficult to follow, even they are not really more complicated.

Assuming $Y^*(y) = yH^{r,\eta^\sigma}$, the identity (??) (Theorem 3.5) suggests a very simple way to associate a progressive utility $U(t, x)$ with the wealth process X^* . Indeed, if $\mathcal{X}(t, z)$ is the inverse flow of $X_t^*(x)$, then the increasing process $U(t, x)$ satisfying $U_x(t, x) = u_x(\mathcal{X}(t, x))H_t^{r,\eta^\sigma}$ is a good candidate to be a consistent stochastic utility. Another remarkable property of this random field is that $U_x(t, X_t^*(x)) = U_x(0, x)H_t^{r,\eta^\sigma}$, which is another way to express that the optimal dual process ν^* is null. We are then in measure to state one of the important results of this section.

Theorem 4.1. *In addition to the monotony assumption 4.1, assume that the given admissible portfolio $(X_t^*(x))$ has a volatility $\kappa^*(t, X_t^*)$, where the process $\kappa^*(t, x)$, is sufficiently regular to make the process $X_t^*(x)$ $\mathcal{C}^{(1)}$ regular in the sense of Kunita. In addition, we assume that $(H_t^{r,\eta^\sigma} X_t^*(x))$ is a martingale $\forall x > 0$. Recall that \mathcal{X} is the inverse flow of X . Let u be a utility function such that $x \mapsto u_{xx}(x)X_t^*(x)$ is integrable near to infinity. Then we define the processes U and \tilde{U} by,*

$$U(t, x) = H_t^{r,\eta^\sigma} \int_0^x u_x(\mathcal{X}(t, z))dz, \quad \tilde{U}(t, y) = \int_y^{+\infty} X_t^* \left(-\tilde{u}_y \left(\frac{z}{H_t^{r,\eta^\sigma}} \right) \right) dz. \quad (26)$$

U is a progressive utility, whose the convex conjugate is \tilde{U} , and the dynamics

$$\begin{cases} dU(t, x) &= \left(-U(t, x)r_t + \frac{1}{2U_{xx}(t, x)} \|\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma\|^2 \right) dt + \gamma(t, x).dW_t \\ \gamma_x(t, x) &= -U_{xx}(t, x)x\kappa^*(t, x) - U_x(t, x)\eta_t^\sigma \end{cases}$$

$\tilde{U}(t, yH_t^{r,\eta^\sigma})$ and $U(t, X_t^*)$ are martingale processes and U is a \mathcal{X} -consistent stochastic utility, with optimal wealth X^* .

Note that the fact that the state price density process Y^* is linear with respect to its initial condition greatly simplify this first result (true if ν^* is not depending on y) contrary to the next theorem where ν_t^* is a function of y .

Proof. First by definition U is strictly increasing concave random field and of class $\mathcal{C}^{(3)}$ in the sense of Kunita. Let us now focus on the dynamics of this progressive utility. Any things are simpler in a martingale market, when H_t^{r,η^σ} is equal to 1, but the PDEs is a little more complicated.

To get started, we introduce the intermediate process $\bar{U}(t, x) := \int_0^x u_x(\mathcal{X}(t, z))dz$ with a simpler convex conjugate $\tilde{\bar{U}}(t, y) = \int_y^{+\infty} X_t^* ((u_x)^{-1}(z))dz$. Denoting by $(\beta^{\bar{U}}, \gamma^{\bar{U}})$ and $(\tilde{\beta}^{\bar{U}}, \tilde{\gamma}^{\bar{U}})$ the local characteristic of \bar{U} and $\tilde{\bar{U}}$, it follows from the dynamics of X^* and the identity

$\tilde{U}_y(t, y) = -X_t^*(-\tilde{u}_y(y))$ that

$$\begin{cases} \tilde{\beta}_y^{\tilde{U}}(t, y) &= -r_t \tilde{U}_y(t, y) - \tilde{U}_y(t, y) \kappa^*(t, -\tilde{U}_y(t, y)) \cdot \eta_t^\sigma \\ \tilde{\gamma}^{\tilde{U}}(t, y) &= -\tilde{U}_y(t, y) \kappa^*(t, -\tilde{U}_y(t, y)) \end{cases} \quad (27)$$

On the other hand, using the correspondence between the diffusion parameters of \bar{U} and \tilde{U}_y given in Proposition 3.2 (or equivalently Corollary 3.3), we have

$$\begin{cases} \gamma_x^{\bar{U}}(t, x) &= -\bar{U}_{xx}(t, x) \tilde{\gamma}_y^{\tilde{U}}(t, \bar{U}_x(t, x)) \\ \beta^{\bar{U}}(t, x) &= \tilde{\beta}^{\tilde{U}}(t, U_x(t, x)) + \frac{1}{2\bar{U}_{xx}(t, x)} \|\gamma_x^{\bar{U}}(t, x)\|^2 \end{cases}$$

By this and (27), it is straightforward to check that

$$\begin{cases} \gamma_x^{\bar{U}}(t, x) &= -x \bar{U}_{xx}(t, x) \kappa^*(t, x) \\ \beta^{\bar{U}}(t, x) &= -x \bar{U}_{xx}(t, x) r_t + \gamma_x^{\bar{U}}(t, x) \cdot \eta_t^\sigma + \frac{\partial}{\partial x} \left(\frac{1}{2\bar{U}_{xx}(t, x)} \|\gamma_x^{\bar{U}}(t, x)\|^2 \right) \end{cases}$$

In turn, we get that $\bar{U}(t, x)$ satisfies $d\bar{U}(t, x) = \beta^{\bar{U}}(t, x) dt + \gamma^{\bar{U}}(t, x) \cdot dW_t$ with

$$\begin{cases} \gamma^{\bar{U}}(t, x) &= -\int_0^x \bar{U}_{xx}(t, z) z \kappa^*(t, z) dz \\ \beta^{\bar{U}}(t, x) &= -x \bar{U}_{xx}(t, x) r_t + \bar{U}(t, x) r_t + \gamma^{\bar{U}}(t, x) \cdot \eta_t^\sigma + \frac{1}{2\bar{U}_{xx}(t, x)} \|\gamma_x^{\bar{U}}(t, x)\|^2 \end{cases}$$

As $U(t, x) = H_t^{r, \eta^\sigma} \bar{U}(t, x)$, Itô's formula leads to

$$dU(t, x) = H_t^{r, \eta^\sigma} (\beta^{\bar{U}}(t, x) - \bar{U}(t, x) r_t - \gamma^{\bar{U}}(t, x) \cdot \eta_t^\sigma) dt + \left(H_t^{r, \eta^\sigma} \gamma^{\bar{U}}(t, x) - U(t, x) \eta_t^\sigma \right) \cdot dW_t.$$

Denote by $\gamma(t, x) := H_t^{r, \eta^\sigma} \gamma^{\bar{U}}(t, x) - U(t, x) \eta_t^\sigma = -\int_0^x \bar{U}_{xx}(t, z) z \kappa^*(t, z) dz - U(t, x) \eta_t^\sigma$, we obtain using $\beta^{\bar{U}}$ formula and identities $U = H^{r, \eta^\sigma} \bar{U}$, $U_x = H^{r, \eta^\sigma} \bar{U}_x$, $U_{xx} = H^{r, \eta^\sigma} \bar{U}_{xx}$ that U satisfies the desired dynamics given by

$$dU(t, x) = \left(-U(t, x) r_t + \frac{1}{2U_{xx}(t, x)} \|\gamma_x^\sigma(t, x) + U_x(t, x) \eta_t^\sigma\|^2 \right) dt + \gamma(t, x) \cdot dW_t$$

and $\gamma_x(t, x) = -U_{xx}(t, x) x \kappa^*(t, x) - U_x(t, x) \eta_t^\sigma$.

Then U is a progressive utility satisfying the utility non linear SPDE, with an optimal wealth satisfying equation (11) of Theorem 2.4. To conclude, it suffices to prove that $U(t, X_t^*)$ is a martingale. It is simpler to show this property on the conjugate dual process, since $\tilde{U}(y H_t^{r, \eta^\sigma}) = H_t^{r, \eta^\sigma} \int_y^{+\infty} X_t^*(-u(z)) dz$ is a martingale since $(H_t^{r, \eta^\sigma} X_t^*(z))$ is a martingale, by integrability assumption. By the conjugacy relation, the same property holds for $U(t, X_t^*)$. \square

The risk tolerance coefficient taken along the optimal wealth has nice properties easily proved. This martingale property has been established in He and Huang in [14]

Corollary 4.2. Let $\alpha^U(t, x) = -\frac{U_x(t, x)}{U_{xx}(t, x)}$ be the risk tolerance coefficient of U .

Then $\alpha^U(t, X_t^*(x)) = \alpha^u(x)X_x^*(t, x)$, where $X_x^*(t, x)$ is the derivative (assumed to exist) of $X_t^*(x)$ with respect to x . Moreover, $H_t^{r, \eta^\sigma} \alpha^U(t, X_t^*(x))$ is a local martingale, since $X_x^*(t, x)$ is also an admissible portfolio with initial wealth 1.

Proof. Observe that by definition $U_{xx}(t, x) = H_t^{r, \eta^\sigma} u_{xx}(\mathcal{X}(t, x))\mathcal{X}_x(t, x)$. Since $\mathcal{X}_x(t, x) = 1/X_x^*(t, \mathcal{X}(t, x))$, and $\mathcal{X}_x(t, X_x^*(t, x)) = x$, the formula $\alpha^U(t, X_t^*(x)) = \alpha^u(x)X_x^*(t, x)$ is a simple verification. Moreover, observe that the derivative $X_x^*(t, x)$ (assumed to exist) belongs to the same vector space of processes than $X_t^*(z)$, and $H_t^{r, \eta^\sigma} X_x^*(t, x)$ is a local martingale. \square

We have shown in Theorem 4.1 that for a monotone wealth process X^* , assumption (X^*H^{r, η^σ} is a martingale) is sufficient in order to construct at least a consistent utility whose the optimal wealth is X^* and the optimal dual process $Y^*(y) = yH^{r, \eta^\sigma}$. We ask now the question how to determine the progressive utilities associated with more general processes Y^* . As we saw in Theorem 3.5, the intuition is to look to U such that $U_x(t, x) = \mathcal{Y} \circ \mathcal{X}(t, x)$ where $\mathcal{Y}(t, x) = Y_t^*(u_x(x))$.

We now state the general consistent utility characterization theorem.

Theorem 4.3. Let $(X_t^*(x) \in \mathcal{X})$ be an admissible wealth process and $(Y_t^*(y)) \in \mathcal{Y}$ be an admissible state price density process, $\mathcal{C}^{(1)}$ regular in the sense of Kunita, such that in addition to Assumptions 4.1 and 4.2, $(X_x^*(t, x)Y_t^*(y))$ is a martingale, where by definition $X_x^*(t, x) = \partial_x X_t^*(t, x)$.

Let u be a utility function, and $\mathcal{Y}(t, x) = Y_t^*(u_x(x))$, $\mathcal{X}(t, z) = (X_t^*(\cdot))^{-1}$ two regular stochastic flows such that such $x \mapsto \mathcal{Y}(t, \mathcal{X}(t, z))$ is integrable near to zero.

Define the processes U and \tilde{U} by

$$U(t, x) = \int_0^x \mathcal{Y}(t, \mathcal{X}(t, z)) dz, \quad \tilde{U}(t, y) = \int_y^{+\infty} X_t^*((\mathcal{Y})^{-1}(t, z)) dz. \quad (28)$$

U is a progressive utility, whose the convex conjugate is \tilde{U} , and the dynamics

$$dU(t, x) = \left(-xU_x(t, x)r_t + \frac{1}{2U_{xx}(t, x)} \|\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma\|^2 \right) dt + \gamma(t, x) \cdot dW_t,$$

with volatility vector γ given by

$$\gamma(t, x) = -U(t, x)\eta_t^\sigma - \int_0^x \left(zU_{xx}(t, z)\kappa^*(t, z) - \nu_t^*(U_x(t, z)) \right) dz.$$

The associated optimal portfolio and the optimal dual process are X^* and Y^* . Moreover $U(t, X_t^*)$ is a martingale, so that U is a \mathcal{X} -consistent stochastic utility.

In the first theorem of this paragraph we build for a given initial utility function a consistent stochastic utility with given optimal wealth process. The extension which we give here characterizes all consistent stochastic utilities with the same optimal wealth process. This result expresses only how we must diffuse the function $U_x(0, x) = u_x(x)$ to stay within the framework of consistent stochastic utilities in incomplete market. The answer is intuitive because it expresses that it is enough to keep a monotone field Y^* which does not influence the reference market. On the other hand it is important to remark that the derivative with respect to x of the volatility vector γ is the sum of two orthogonal vectors and is given by

$$\begin{aligned}\gamma_x(t, x) &= \nu_t^*(U_x(t, x)) - U_{xx}(t, x)x\kappa^*(t, x) - U_x(t, x)\eta_t^\sigma \\ &= \nu_t^*(U_x(t, x)) - U_x(t, x)\left(\eta_t^\sigma + \frac{U_{xx}(t, x)x\kappa^*(t, x)}{U_x}\right),\end{aligned}$$

and consequently, given κ^* and ν^* , it is interpreted as an operator $\Upsilon(t, x, U_x, U_{xx})$ which is linear on U_{xx} , that depends on U_x through the volatility ν^* of Y^* and an affine term on η^σ , and depends on x only through the optimal policy κ^* . We also emphasize that the term U_x/U_{xx} in this formula is the risk tolerance of an investor with utility process U . In particular in the case of the market martingale ($\eta^\sigma = 0$), $\Upsilon(t, x, U_x, U_{xx})$ is linear on U_{xx} , depends on U_x only through the volatility ν^* of Y^* and on x only through the optimal policy κ^* .

Note that in the classical backward set-up of utility maximisation, similar idea is investigated by I. Karatzas & al [11]. The authors show also that the solution of backward SPDE can be represented as the composite of two invertible processes. But this differs from the approach proposed here because these processes are represented as an expectation of monotonic functions (characteristics method) where in this work are stochastic flows. Note that the authors also use the Itô-Ventzel formula to establish the backward SPDE.

Remark . *After giving the proof of this result, we want to draw the attention to the fact that this Theorem can be showed first in the case of the martingale market ($r = 0, \eta = 0$). This allows us to simplify calculation and we can always comeback to the initial market by a technique of change of numeraire.*

Proof. Under assumption 4.1 the inverse \mathcal{X} of X^* with respect to x satisfies by Proposition 3.2

$$d\mathcal{X}(t, x) = -x\mathcal{X}_x(t, x)\kappa^*(t, x).dW_t + \frac{1}{2}\partial_x(\mathcal{X}_x(t, x)\|x\kappa^*(t, x)\|^2)dt.$$

The hypothesis made on X^* and Y^* entail that we can apply the Itô-Ventzel formula to the compound flow $\mathcal{Y}\circ\mathcal{X}$. To study $U_x(t, x)$ we are first interested on the coefficient of dW_t of the dynamics of $\mathcal{Y}\circ\mathcal{X}$ because it represents the derivative of the volatility γ of the utility U .

As $(\mathcal{Y}_x \circ \mathcal{X})\mathcal{X}_x = U_{xx}$ and $U_x = \mathcal{Y} \circ \mathcal{X}$, formula (3.1) gives us that

$$\gamma_x(t, x) = \nu_t^*(U_x(t, x)) - xU_{xx}(t, x)\kappa^*(t, x) - U_x(t, x)\eta_t^\sigma.$$

This identity shows that the vector γ_x is the sum of two orthogonal vectors since the first term $\nu_t^*(U_x(t, x))$ belong by hypothesis to the orthogonal of the second which is $-U_x(t, x)\left(\eta_t^\sigma + (xU_{xx}/U_x)(t, x)\kappa^*(t, x)\right)$ that belongs by hypothesis to the space \mathcal{R}_t^σ . Throughout, the projection of γ_x on \mathcal{R}_t^σ is the vector $\gamma_x^\sigma(t, x) = -xU_{xx}(t, x)\kappa^*(t, x) - U_x(t, x)\eta_t^\sigma$.

As $U_x = \mathcal{Y} \circ \mathcal{X}$, γ_x is the volatility process of U_x , it is enough to integrate it with respect to x to obtain the result.

We now focus our interest on the drift μ^{U_x} of the derivative $\mathcal{Y} \circ \mathcal{X}$ of U . The idea and calculations are exactly identical to those of the proof of Corollary ?? . Indeed by the assumptions and equation (19) we have

$$\begin{aligned} \mu^{U_x}(t, x) &= -(x\mathcal{X}_x(t, x)\mathcal{Y}_x \circ \mathcal{X}(t, x) + \mathcal{Y} \circ \mathcal{X}(t, x))r_t \\ &+ \frac{1}{2}(\mathcal{Y}_x \circ \mathcal{X})(t, x)\partial_x(\mathcal{X}_x(t, x)\|x\kappa^*(t, x)\|^2) + \frac{1}{2}(\mathcal{Y}_x \circ \mathcal{X})(t, x)\|\mathcal{X}_x(t, x)x\kappa^*(t, x)\|^2 \\ &- x\mathcal{X}_x(t, x)\partial_x\left(\mathcal{Y} \circ \mathcal{X}(t, x)(\nu_t^*(\mathcal{Y} \circ \mathcal{X}(t, x)) - \eta_t^\sigma)\right).\kappa^*(t, x) - x\mathcal{X}_x(t, x)\mathcal{Y}_x \circ \mathcal{X}(t, x)\kappa^*(t, x).\eta_t^\sigma. \end{aligned}$$

Note that in the last line the term $-x\mathcal{X}_x(t, x)\partial_x\left(\mathcal{Y} \circ \mathcal{X}(t, x)(\nu_t^*(\mathcal{Y} \circ \mathcal{X}(t, x)) - \eta_t^\sigma)\right).\kappa^*(t, x)$ comes from Itô-Ventzel formula and corresponds to $\langle d\mathcal{Y}, d\mathcal{X} \rangle$.

To lead the proof we proceed by analyzing line by line the above equality. Using the identities $(\mathcal{Y}_x \circ \mathcal{X}(t, x))_x = \mathcal{Y}_{xx} \circ \mathcal{X}(t, x)\mathcal{X}_x(t, x)$ and $U_{xx}(t, x) = \mathcal{Y}_x \circ \mathcal{X}(t, x)\mathcal{X}_x(t, x)$, the first line becomes

$$-(x\mathcal{X}_x(t, x)\mathcal{Y}_x \circ \mathcal{X}(t, x) + \mathcal{Y} \circ \mathcal{X}(t, x))r_t = -(xU_{xx}(t, x) + U_x(t, x))r_t = -\partial_x(xU_x)(t, x)r_t. \quad (29)$$

Rewriting the second line, we obtain

$$\begin{aligned} &\frac{1}{2}\left[(\mathcal{Y}_x \circ \mathcal{X}(t, x)\mathcal{X}_x(t, x)(\mathcal{X}_x(t, x)\|x\kappa^*(t, x)\|^2) + (\mathcal{Y}_{xx} \circ \mathcal{X})(t, x)\mathcal{X}_x(t, x)\partial_x(\mathcal{X}_x(t, x)\|x\kappa^*(t, x)\|^2)\right] \\ &= \frac{1}{2}\partial_x[\mathcal{Y}_x \circ \mathcal{X}(t, x)\mathcal{X}_x(t, x)\|x\kappa^*(t, x)\|^2]. \end{aligned} \quad (30)$$

Finally, from the assumption that $\nu_t^*(\mathcal{Y}(t, x)).\kappa^*(t, X_t^*(x)) = 0$, we deduce $\nu_t^*(\mathcal{Y} \circ \mathcal{X}(t, x)).\kappa^*(t, x) = 0$ and $\partial_x(\nu_t^*(\mathcal{Y} \circ \mathcal{X}(t, x))).\kappa^*(t, x) = 0$. This yields in the last line

$$-x\mathcal{X}_x(t, x)\left[\partial_x\left(\mathcal{Y} \circ \mathcal{X}(t, x)(\nu_t^*(\mathcal{Y} \circ \mathcal{X}(t, x)) - \eta_t^\sigma)\right).\kappa^*(t, x) - \mathcal{Y}_x \circ \mathcal{X}(t, x)\kappa^*(t, x).\eta_t^\sigma\right] = 0. \quad (31)$$

Identities (29), (30) and (31) combined with the expression of μ^{U_x} and γ_x yield to

$$\mu^{U_x}(t, x) = \partial_x\left(-xU(t, x)r_t + \frac{1}{2U_{xx}(t, x)}\|x\kappa^*(t, x)\|^2\right).$$

As $U(t, 0) \equiv 0$ we get by integration that U satisfies

$$dU(t, x) = \{-xU_x(t, x)r_t + \frac{1}{2}U_{xx}(t, x)\|\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma\|^2\}dt + \gamma(t, x).dW_t.$$

Using $\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma = -xU_{xx}(t, x)\kappa^*(t, x)$ one easily sees that

$$dU(t, x) = \{-xU_x(t, x)r_t + \frac{1}{2}\left[\frac{\|\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma\|^2}{U_{xx}(t, x)}\right]\}dt + \gamma(t, x).dW_t.$$

It remains to show that $U(t, X_t^*)$ is a martingale, given that the positive process $Y_t^*(y)X_x^*(t, x)$ is a martingale by assumption. Then since $U(t, X_t^*) = \int_0^x \mathcal{Y}(t, z)X_x^*(t, z)dz$, $U(t, X_t^*)$ is also a martingale. The proof is complete. \square

Remark . *The risk tolerance coefficient along the optimal wealth $A(t, X_t^*(x))$ is no longer proportional to $X_x^*(t, x)$ since $A(t, X_t^*(x)) = \frac{Y_t^*(u_x(x))}{Y_y^*(t, u_x(x))u_{xx}(t, x)}X_x^*(t, x)$.*

Nevertheless, the both processes $\frac{Y_t^(u_x(x))}{Y_y^*(t, u_x(x))}$ and $Y_t^*(u_x(x))X_x^*(t, x)$ are local martingales, but their product is not a local martingale.*

4.2 Stochastic Flows Method for Solving Stochastic PDE's

In the previous paragraph using two invertible stochastic flows X^* and Y^* we construct a consistent utility with the desired dynamics. Naturally, the question of the converse point of view is required. Starting from a stochastic PDE that satisfy consistent utilities, the question is then under which assumptions we have existence and uniqueness of the solution? What can we deduce about the monotony and the concavity of a possible solution? Answering these questions is the purpose of this paragraph. In the following theorem we propose a new method that allows us to address the issue of such resolution of fully nonlinear second order stochastic PDEs .

Theorem 4.4. *Consider a utility stochastic PDE with initial condition $u(\cdot)$,*

$$dU(t, x) = \left(-xU_x(t, x)r_t + \frac{1}{2U_{xx}(t, x)}\|\gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma\|^2 \right)dt + \gamma(t, x).dW_t. \quad (32)$$

Put $-xU_{xx}(t, x)\kappa^*(t, x) = \gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma$ and $\nu_t^*(U_x(t, x)) = \gamma_x^\sigma(t, x)$.

Assume that the both equations

$$\frac{dX_t^*(x)}{X_t^*(x)} = r_t dt + \kappa^*(t, X_t^*(x)).(dW_t + \eta_t^\sigma dt), \quad \frac{dY_t^*(y)}{Y_t^*(y)} = -r_t dt + (\nu_t^*(Y_t^*(y)) - \eta_t^\sigma).dW_t \quad (33)$$

admit solutions which are increasing: $[0, +\infty) \rightarrow [0, +\infty)$ in their initial conditions and are a regular flow in the sense of Kunita. Let $\mathcal{Y}(t, x) = Y_t^*(u_x(x))$, $\mathcal{X}(t, z) = (X_t^*(\cdot))^{-1}$ and

assume $x \mapsto \mathcal{Y}(t, \mathcal{X}(t, z))$ is integrable near to zero. Then there exists an increasing and concave solution U of the SPDE (32) given by

$$U(t, x) = \int_0^x \mathcal{Y}(t, \mathcal{X}(t, z)) dz$$

Moreover, if X^* and Y^* are unique then U is the unique solution of (32).

Theorem 4.3 shows that for a given X^* and Y^* increasing solutions of SDEs (33) the random field $U(t, x) = \int_0^x \mathcal{Y}(t, \mathcal{X}(t, z)) dz$ is a consistent utility solution of the utility SPDE (32) with a volatility vector γ s.t. $\gamma_x(t, x) + U_x(t, x)\eta_t^\sigma = -xU_{xx}(t, x)\kappa^*(t, x) + \nu_t^*(U_x(t, x))$. In this result the converse point of view is investigated. Starting from the utility SPDE (32) with a given initial condition u , putting $-xU_{xx}(t, x)\kappa^*(t, x) = \gamma_x^\sigma(t, x) + U_x(t, x)\eta_t^\sigma$ and $\nu_t^*(U_x(t, x)) = \gamma_x^\sigma(t, x)$, the theorem shows, under the assumption that both two SDE's (33) admit a increasing solutions, that there exists an increasing concave solution U of SPDE (32) given by $U(t, x) = \int_0^x \mathcal{Y}(t, \mathcal{X}(t, z)) dz$. Moreover, the uniqueness of U is strongly related on the uniqueness of the solutions X^* and Y^* of SDE's (33). Finally, remark that the martingale property of the product X^*Y^* is not required to have a solution of the SPDE where it was necessary to conclude that U is a consistent utility in Theorem 4.3.

These is an interesting new approach in which the solution of the utility SPDE have a trajectory (path wise) representation contrary to the characteristics method where the solutions are represented as an expectation. In particular, note that there are several advantages of this connection between SPDE's and SDE's. For example, the existence of divers works in the domain of SDE's and seen in the multitude of results on the existence, uniqueness and on the integrability of solutions. The monotonicity of solutions X^* and Y^* gives several properties of the solution U of the SPDE which to the best of our knowledge there are no or few results that assert the monotonicity or the convexity of such solutions. Also, there may be other advantages in numerical methods and simulations of the SDE then of SPDE. We finish this section mentioning that the main assumption of Theorem 4.4 is to assume that the SDEs (33) admit a solutions which is a fairly strong assumption because κ^* and ν^* may depend on higher order derivatives of U . For example in the Markovian case where $U(t, x) = u(t, x, S_t)$, according to Example 2.1, the volatility vector γ of U is given by

$$\gamma(t, x) = (\sigma_t)^T \nabla_S u(t, x, S_t).$$

Hence, one can easily obtain that

$$-x\kappa^*(t, x) = \frac{(\sigma_t)^T \nabla_S u(t, x, S_t)}{u_{xx}(t, x, S_t)}$$

In other words, existence of a solution to the associated SDE in (33) is like an inverse problem.

Conclusion This paper investigates consistent stochastic utilities from the stochastic PDEs point of view. This leads therefore to make strong regularity assumptions: The market is a Brownian market and securities are modeled as continuous semimartingales. Utilities are at least of class $\mathcal{C}^{(2)}$ in the sense of Kunita in order to apply Itô-Ventzel Lemma and to deduce the SPDEs. Moreover, the method of stochastic utilities construction is based on the dynamics of stochastic flows and their inverses, and therefore additional regularity assumptions on X^* and Y^* are required. However, one can take a direct approach still based on monotony assumptions on optimal processes for the primal and dual problem, and on compound flows formula ; it is showed in [23], that these assumptions can be considerably weakened. Indeed, considering any financial market in which the securities are modeled as a bounded semimartingales, the stochastic utilities are of class \mathcal{C}^1 and wealth process are required to lie in a convex class $\mathcal{X} \subset \mathbb{X}^+$, the monotony assumption of X^* and Y^* is sufficient to show the validity of the construction proposed in this work, using analysis methods and optimality conditions.

Appendix

A Itô-Ventzel's formula

The Itô-Ventzel's formula is a generalization of classical Itô's formula where the deterministic function is replaced by a stochastic process depending on a real or multivariate parameter. There are several difficulties in the definition of semimartingale depending on a parameter, as explained in H. Kunita [25]. For instance, let us consider the Itô integral of a predictable process $f_t(x)$ with parameter x in some domain D of \mathbb{R}^+ with respect to some Brownian motion B . Suppose that $\int_0^T f_s(x)^2 ds < +\infty$ holds for each $x \in D$. Then the Itô integral $M_t(x) = \int_0^t f_s(x) dB_s$ is well defined for any t except for a null set N_x . It is a continuous local martingale with parameter $x \in D$. Then $M_t(x)$ is well defined for (t, x) if $\omega \in (\cup_{x \in \mathcal{I}} N_x)^c$. However the exceptional set $(\cup_{x \in \mathcal{I}} N_x)$ may not be a null set since it is an uncountable union of null sets. To overcome this technical problem we must take a good modification of the random field $M_t(x)$ so that it is well defined for all (t, x) a.s. and is continuous or continuously differentiable with respect to x for all t almost surely.

A.1 Notation and Definition

Functional spaces We shall first introduce some notations. Let D be a domain in \mathbb{R}^+ , m an non-negative integer and denote by $\mathcal{C}^m(D, \mathbb{R})$ the set of all functions $g : D \rightarrow \mathbb{R}$ which

are m -times continuously differentiable. Using the notation $g^{(m)}$ for the derivative of order m of some function g , we introduce the seminorms defined on some compact subset of D by

$$\|g\|_{m:K} = \sup_{x \in K} \frac{|g(x)|}{1 + |x|} + \sum_{1 \leq \alpha \leq m} \sup_{x \in K} |g^{(\alpha)}(x)|.$$

Equipped with these seminorms, $\mathcal{C}^m(D, \mathbb{R})$ is a Frechet space. When D itself is a compact space we drop out the reference to K .

We sometimes need to refer to more regular functions whose derivatives of order m are δ -Hölder continuous ($0 < \delta \leq 1$). Then we introduce a new family of seminorms,

$$\|g\|_{m+\delta:K} = \|g\|_{m:K} + \sup_{\substack{x, y \in K \\ x \neq y}} \frac{|g^{(m)}(x) - g^{(m)}(y)|}{|x - y|^\delta}.$$

on the set of $\mathcal{C}^m(D, \mathbb{R})$ whose last derivative is δ -Hölder continuous.

Definition A.1. A continuous function $g(t, x)$, $x \in \mathcal{I}$, $t \geq 0$ is said to belong to $\mathcal{C}^{m, \delta}$, $\delta \in [0, 1]$ if for every t , $f(t) = f(t, \cdot)$ belongs to $\mathcal{C}^{m, \delta}$ and $\|f(t)\|_{m+\delta:K}$ is integrable with respect to t for any compact subset K of \mathcal{I} . If the set K is \mathcal{I} , the function f is said to belong to the class $\mathcal{C}_b^{m, \delta}$. Furthermore, if $\|f(t)\|_{m+\delta}$ is bounded in t it is said to belong to $\mathcal{C}_{ub}^{m, \delta}$.

We also need to introduce the same kind of definition for functions depending on two parameters

$$\|g\|_{\tilde{m}+\delta:K} = \|g\|_{\tilde{m}:K} + \sum_{\alpha=m} \|\partial_x^\alpha \partial_y^\alpha g(x, y)\|_{\tilde{\delta}:K}$$

$$\|g\|_{\tilde{\delta}:K} = \sup_{\substack{x, x', y, y' \in K \\ x \neq x', y \neq y'}} \frac{|g(x, y) - g(x', y) - g(x, y') + g(x', y')|}{|x - x'|^\delta |y - y'|^\delta}.$$

Definition A.2. A continuous function $g(t, x, y)$, $x, y \in \mathcal{I}$, $t \in [0, T]$ is said to belong to $\tilde{\mathcal{C}}^{m, \delta}$, $\delta \in [0, 1]$ if for every t , $g(t) = g(t, \cdot, \cdot)$ belongs to $\tilde{\mathcal{C}}^{m, \delta}$ and $\|g(t)\|_{\tilde{m}+\delta:K}$ is integrable on $[0, T]$ with respect to t for any compact subset K of \mathcal{I} . If the set K is \mathcal{I} , the function g is said to belong to the class $\tilde{\mathcal{C}}_b^{m, \delta}$. Furthermore, if $\|g(t)\|_{\tilde{m}+\delta}$ is bounded in t it is said to belong to $\tilde{\mathcal{C}}_{ub}^{m, \delta}$.

$\mathcal{C}^{m, \delta}$ -process : Let $U(t, x)$ a family of real valued process with parameter $x \in \mathcal{I}$. We can regard it as a random field with double parameter t and x . If $U(t, x, \omega)$ is a continuous function of x for almost all ω for any t , we can regard $U(t, \cdot)$ as a stochastic process with values in $\mathcal{C} = \mathcal{C}(\mathcal{I}, \mathbb{R})$ or a \mathcal{C} -valued process. If $U(t, x, \omega)$ is m -times continuously differentiable with respect to x for almost all ω for any t , it can be regarded as a stochastic process with values in $\mathcal{C}^m = \mathcal{C}^m(\mathcal{I}, \mathbb{R})$ or a \mathcal{C}^m -valued process. If $U(t, x)$ is a continuous process with

value in \mathcal{C}^m , it is called a continuous \mathcal{C}^m -process. A $\mathcal{C}^{m,\delta}$ -valued process and continuous $\mathcal{C}^{m,\delta}$ -processes are defined similarly.

$\tilde{\mathcal{C}}^{m,\delta}$ -process : Let $G(t, x, y)$ be a stochastic process with parameter $x, y \in \mathcal{I}$. If it is m -times continuously differentiable with respect to each x and y a.s. for any t , it is called a stochastic process with values in $\tilde{\mathcal{C}}^m$ or a $\tilde{\mathcal{C}}^m$ -valued process. The $\tilde{\mathcal{C}}^{m,\delta}$ -valued process and continuous $\tilde{\mathcal{C}}^{m,\delta}$ -valued process are defined similarly.

Theorem A.1. *Let $M_t(x)$, $x \in \mathcal{I}$ be a family of continuous local martingales such that $M_0(x) \equiv 0$. Assume the joint quadratic variation $\langle M_t(x), M_t(y) \rangle$ has a modification $A(t, x, y)$ of a continuous $\tilde{\mathcal{C}}^{m,\delta}$ -process for some $m \geq 1$ and $\delta \in (0, 1]$. Then $M_t(x)$ has a modification of a continuous $\mathcal{C}^{m,\varepsilon}$ -process for any $\varepsilon < \delta$. Furthermore, for each $n \geq m$, $\partial_x^n M_t(x)$, $x \in \mathcal{I}$ is a family of continuous local martingales with joint quadratic variation $\partial_x^n \partial_y^n A(t, x, y)$.*

Definition A.3. *We shall call the random field $M_t(x)$ with the property of the previous Theorem a continuous local martingale with values in $\mathcal{C}^{m,\varepsilon}$ or a continuous $\mathcal{C}^{m,\varepsilon}$ -local martingales.*

Regular Itô's random fields $\mathcal{C}^{m,\delta}$ -semimartingale : Suppose $U(t, x)$, $x \in \mathcal{I}$ is a family of continuous semimartingale decomposed as $U(t, x) = B(t, x) + M(t, x)$, where $M(t, x)$ is a local martingale and $B(t, x)$ is a continuous process of bounded variation. $U(t, x)$, $x \in \mathcal{I}$ is said to belong to the class $\mathcal{C}^{m,\delta}$ or simply to be $\mathcal{C}^{m,\delta}$ -semimartingale if $M(t, x)$ is a continuous $\mathcal{C}^{m,\delta}$ -local martingale and $B(t, x)$ is a continuous $\mathcal{C}^{m,\delta}$ -process such that $D_x^\alpha B(t, x)$, $\alpha \leq m$ are all process of bounded variation. Further if $\delta = 0$ it is called a \mathcal{C}^m -semimartingale.

Let U be a semimartingale satisfying

$$dU(t, x) = \beta(t, x)dt + \gamma(t, x).dW_t, \quad U(r, x) = U(x),$$

where β and γ are predictable process.

Definition A.4 (Kunita).

- The pair (β, γ) is called the local characteristic of U .
- Let m be a non-negative integer and $a(t, x, y) := \gamma(t, x)^* \gamma(t, y)$. The local characteristic (β, γ) is said to be in the class $\mathcal{B}^{m,0}$ if both β and a are predictable process with value \mathcal{C}^m and if for any compact subset $K_1 \subset \mathbb{R}_+$ and $K_2 \subset \mathbb{R}_+ \times \mathbb{R}_+$ $\|\beta(t, \cdot)\|_{m;K_1}, \|a(t, \cdot, \cdot)\|_{m;K_2} \in L^1$.

Where

$$\|f\|_{m:K_1} = \sup_{x \in K_1} \frac{|f(x)|}{1+|x|} + \sum_{1 \leq |\alpha| \leq m} \sup_{x \in K_1} |D_x^\alpha f(x)|.$$

$$\|g\|_{\tilde{m}:K_2} = \sup_{x,y \in K_2} \frac{|g(x,y)|}{(1+|x|)(1+|y|)} + \sum_{1 \leq |\alpha| \leq m} \sup_{x,y \in K_2} |D_x^\alpha D_y^\alpha g(x,y)|$$

Definition A.5 (Itô-Ventzel Regularity). *A semimartingale random field U is said to be Itô-Ventzel regular if U is a continuous \mathcal{C}^2 -process and continuous \mathcal{C}^1 -semimartingale with local characteristic satisfying previous assumption .*

Theorem A.2 (Itô-Ventzel's Formula (Kunita)). *Let $(U(t,x))$ be an Itô-Ventzel regular semimartingale random field and let X_t be a continuous semimartingale with values in \mathcal{I} and volatility σ^X , then $U(t, X_t)$ is a continuous semimartingale and*

$$\begin{aligned} U(t, X_t) &= U(0, X_0) + \int_0^t \beta(s, X_s) ds + \int_0^t \gamma(s, X_s) \cdot dW_s \\ &+ \int_0^t \frac{\partial U}{\partial x}(s, X_s) dX_s + \int_0^t \frac{\partial^2 U}{\partial x^2}(s, X_s) \langle X \rangle_s ds \\ &+ \int_0^t \left\langle \frac{\partial \gamma}{\partial x}(s, X_s), \sigma_s^X \right\rangle ds. \end{aligned}$$

Furthermore, according to H. Kunita [25] Theorem 3.3.3 p.94 we have the following differential rules for stochastic integrals.

Theorem A.3 (Differential rules for stochastic integrals). *(i) Let $F(t,x)$ be a continuous $\mathcal{C}^{m,\delta}$ -semimartingale with local characteristic belonging to the class $\mathcal{B}^{m,\delta}$ where $\delta > 0$. Let $X(t,x)$, $x \in \Lambda$, $t \in [0, T]$ be a continuous predictable process with values in $\mathcal{C}^{k,\gamma}(\Lambda, \mathcal{I})$ where $\gamma > 0$ and $\Lambda \subset \mathbb{R}^e$. Set*

$$M(t,x) = \int_0^t F(X(s,x), ds).$$

Then $M(t,x)$ has a modification of continuous $\mathcal{C}^{m,\delta}$ -semimartingale with values in $\mathcal{C}^{m \wedge k, \varepsilon}(\Lambda, \mathbb{R})$ with local characteristic belonging to the class $\mathcal{B}^{m \wedge k, \gamma \delta}$ with $\varepsilon < \gamma \delta$.

Further if g_t is a continuous predictable process with values in Λ , then we have the equality:

$$\int_0^t M(ds, g_s) = \int_0^t F(X(s, g_s), ds). \quad (34)$$

(ii) If $m \geq 1$ and $k \geq 1$, then we have the equality:

$$\frac{\partial}{\partial x^i} M(t,x) = \sum_{i=1}^d \int_0^t \frac{\partial}{\partial x^j} X^i(s,x) \frac{\partial}{\partial x^i} F(X(s,x), ds). \quad (35)$$

References

- [1] A. Ben-Israel and T. N. E. Grenville. *Generalized Inverses: Theory and Applications*. Springer-Verlag, Berlin, 2002.
- [2] René A. Carmona and David Nualart. *Nonlinear stochastic integrators, equations and flows*, volume 6 of *Stochastics Monographs*. Gordon and Breach Science Publishers, New York, 1990.
- [3] Tahir Choulli, Christophe Stricker, and Jia Li. Minimal Hellinger martingale measures of order q . *Finance Stoch.*, 11(3):399–427, 2007.
- [4] M. G. Crandall, H. Ishii, and P. L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *ArXiv Mathematics e-prints*, June 1992.
- [5] J. Cvitanić and I. Karatzas. Convex duality in constrained portfolio optimization. *Ann. Appl. Probab.*, 2(4):767–818, 1992.
- [6] V. Henderson D. and Hobson. Horizon-unbiased utility functions. *Stochastic Process. Appl.*, 117(11):1621–1641, 2007.
- [7] F. Delbaen and W. Schachermayer. A general version of the fundamental theorem of asset pricing. *Math. Ann.*, 300(3):463–520, 1994.
- [8] F. Delbaen and W. Schachermayer. The no-arbitrage property under a change of numéraire. *Stochastics Stochastics Rep.*, 53(3-4):213–226, 1995.
- [9] F. Delbaen and W. Schachermayer. The fundamental theorem of asset pricing for unbounded stochastic processes. *Math. Ann.*, 312(2):215–250, 1998.
- [10] N. El Karoui. Les aspects probabilistes du contrôle stochastique. In *Ninth Saint Flour Probability Summer School—1979 (Saint Flour, 1979)*, volume 876 of *Lecture Notes in Math.*, pages 73–238. Springer, Berlin, 1981.
- [11] Nikolaos Englezos and Ioannis Karatzas. Utility maximization with habit formation: Dynamic programming and stochastic pdes. *SIAM J. Control Optim.*, 48(2):481–520, 2009.
- [12] M.R. Tehranchi F.P.Y.S. Berrier, L.C.G. Rogers. A characterization of forward utility functions. *Preprint*, 2009.
- [13] C. Groetsch. *Generalized inverses of linear operators*. Marcel Dekker. 1997.
- [14] Hua He and Chi fu Huang. Consumption-portfolio policies: An inverse optimal problem. *Journal of Economic Theory*, 62(2):257 – 293, 1994.

- [15] P. Huy en. *Optimisation et contr le stochastique appliqu s   la finance*, volume 61 of *Math matiques   Applications (Berlin) [Mathematics   Applications]*. Springer, Berlin, 2007.
- [16] John P. Lehoczky I. Karatzas, Steven E. Shreve, and Gan-Lin Xu. Martingale and duality methods for utility maximization in an incomplete market. *SIAM J. Control Optim.*, 29(3):702–730, 1991.
- [17] John M. Ingram and M. M. Marsh. Projections onto convex cones in hilbert space. *J. Approx. Theory*, 64(3):343–350, 1991.
- [18] R. J.Elliott and H. F llmer. Orthogonal martingale representation. In *Stochastic analysis*, pages 139–152. Academic Press, Boston, MA, 1991.
- [19] I. Karatzas and K. Constantinou. The num raire portfolio in semimartingale financial models. *Finance Stoch.*, 11(4):447–493, 2007.
- [20] I. Karatzas and Wang Hui. Utility maximization with discretionary stopping. *SIAM J. Control Optim.*, 39(1):306–329, 2000.
- [21] I. Karatzas and S. E. Shreve. *Methods of Mathematical Finance*. Springer, September 2001.
- [22] N. El Karoui and M. M’rad. Random risk aversion and consistent utilities. *preprint.*, 2010.
- [23] N. El Karoui and M. M’rad. Stochastic utilities with a given optimal portfolio : Approach by stochastic flows. *Preprint.*, 2010.
- [24] D. Kramkov and W. Schachermayer. The asymptotic elasticity of utility functions and optimal investment in incomplete markets. *Annals of Applied Probability*, 9:904–950, 1999.
- [25] H. Kunita. *Stochastic Flows and Stochastic Differential Equations*. Cambridges studies advanced mathematics, 1997.
- [26] M.Musiela and T. Zariphopoulou. A valuation algorithm for indifference prices in incomplete markets. *Finance and Stochastics*, 8(3):399–414, 08 2004.
- [27] M. Musiela and T. Zariphopoulou. Backward and forward utilities and the associated pricing systems: The case study of the binomial model. *Preprint*, 2003.
- [28] M. Musiela and T. Zariphopoulou. An example of indifference prices under exponential preferences. *Finance Stoch.*, 8(2):229–239, 2004.

- [29] M. Musiela and T. Zariphopoulou. Indifference prices of early exercise claims. In *Mathematics of finance*, volume 351 of *Contemp. Math.*, pages 259–271. Amer. Math. Soc., Providence, RI, 2004.
- [30] M. Musiela and T. Zariphopoulou. The single period binomial model (r. carmona, ed.). *Princeton University Press*, 2005.
- [31] M. Musiela and T. Zariphopoulou. Investment and valuation under backward and forward dynamic exponential utilities in a stochastic factor model. In *Advances in mathematical finance*, Appl. Numer. Harmon. Anal., pages 303–334. Birkhäuser Boston, Boston, MA, 2007.
- [32] M. Musiela and T. Zariphopoulou. Stochastic partial differential equations in portfolio choice. *Preliminary report*, 2007.
- [33] M. Musiela and T. Zariphopoulou. Optimal asset allocation under forward exponential performance criteria. volume 4 of *Contemp. Math.*, pages 285–300. Institute of Mathematical Statistics, 2008.
- [34] M. Musiela and T. Zariphopoulou. Portfolio choice under dynamic investment performance criteria. *Quantitative Finance*, 9(2):161–170, 2008.
- [35] M. Musiela and T. Zariphopoulou. Portfolio choice under space-time monotone performance criteria. *Preprint*, November-2008.
- [36] Eckhard Platen and David Heath. *A benchmark approach to quantitative finance*. Springer Finance. Springer-Verlag, Berlin, 2006.
- [37] R. Tyrrell Rockafellar. *Convex analysis*. Princeton Mathematical Series, No. 28. Princeton University Press, Princeton, N.J., 1970.
- [38] R. Rouge and N. El Karoui. Pricing via utility maximization and entropy. *Math. Finance*, 10(2):259–276, 2000. INFORMS Applied Probability Conference (Ulm, 1999).
- [39] W. Schachermayer. Optimal investment in incomplete financial markets. In *Mathematical Finance: Bachelier Congress 2000*, pages 427–462. Springer, 2001.
- [40] A.D. Ventzel. On equations of the theory of conditional Markov processes. *Theory of Probability and its Applications*, 10:357–361, 1965.
- [41] Z. H. Wei and S. G. Zhang. Relationship between optimal growth portfolio and martingale measure. *Chinese J. Appl. Probab. Statist.*, 19(1):14–18, 2003.
- [42] D. V. Widder. The role of the Appell transformation in the theory of heat conduction. *Trans. Amer. Math. Soc.*, 109:121–134, 1963.

- [43] D. V. Widder. *The heat equation*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975. Pure and Applied Mathematics, Vol. 67.
- [44] T. Zariphopoulou and G. Žitković. Maturity independent risk measures. *Preprint*, 2007.
- [45] G. Zitkovic. A dual characterization of self-generation and log-affine forward performances. *Preprint*, 2008.
- [46] L. Zongxia. Stochastic differential equations driven by spatial parameters semimartingale with non-lipschitz local characteristic. *Potential Analysis*, 26, 2007.