



HAL
open science

Energisation-dependent endogenous activation of proton conductance in skeletal muscle mitochondria

Nadeene Parker, Charles Affourtit, Antonio J Vidal-Puig, Martin D Brand

► **To cite this version:**

Nadeene Parker, Charles Affourtit, Antonio J Vidal-Puig, Martin D Brand. Energisation-dependent endogenous activation of proton conductance in skeletal muscle mitochondria. *Biochemical Journal*, 2008, 412 (1), pp.131-139. 10.1042/BJ20080006 . hal-00478942

HAL Id: hal-00478942

<https://hal.science/hal-00478942>

Submitted on 30 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Energisation-dependent endogenous activation of proton conductance in skeletal muscle mitochondria

Nadeene Parker^{*,†,‡}, Charles Affourtit^{*}, Antonio Vidal-Puig[†], Martin D. Brand^{*}

^{*}*MRC Dunn Human Nutrition Unit, Hills Road, Cambridge CB2 0XY, UK;* [†]*Metabolic Research Laboratories, Level 4, Institute of Metabolic Science, Box 289, Addenbrooke's Hospital, Cambridge CB2 0QQ, UK*

[‡]To whom correspondence should be addressed (email: np@mrc-dunn.cam.ac.uk)

Short title: Endogenous activation of proton conductance

Abbreviations used: ANT, adenine nucleotide translocase; BSA, bovine serum albumin; FCCP, carbonylcyanide *p*-trifluoromethoxyphenylhydrazone; TPMP, triphenylmethylphosphonium; UCP, uncoupling protein; KO, knockout; WT, wild type.

SYNOPSIS

Leak of protons into the mitochondrial matrix during substrate oxidation partially uncouples electron transport from phosphorylation of ADP, but the functions and source of basal and inducible proton leak *in vivo* remain controversial. We describe an endogenous activation of proton conductance in mitochondria isolated from rat and mouse skeletal muscle following addition of respiratory substrate. This endogenous activation increased with time, required a high membrane potential and was diminished by high concentrations of serum albumin. Inhibition of this endogenous activation by GDP (classically considered specific for uncoupling proteins, UCPs), carboxyatractylate and bongkrekate (considered specific for the adenine nucleotide translocase) was examined in skeletal muscle mitochondria from wild type and *UCP3*-knockout mice. Proton conductance through endogenously-activated UCP3 was calculated as the difference in leak between mitochondria from wild type and *UCP3*-knockout mice, and was found to be inhibited by carboxyatractylate and bongkrekate, but not GDP. Proton conductance in mitochondria from *Ucp3* knockout mice was inhibited strongly by carboxyatractylate, bongkrekate and partially by GDP. We conclude the following. (1) at high protonmotive force, an endogenously-generated activator stimulates proton conductance catalysed partly by UCP3 and partly by the adenine nucleotide translocase. (2) GDP is not a specific inhibitor of UCP3, but also inhibits proton translocation by the adenine nucleotide translocase. (3) The inhibition of UCP3 by carboxyatractylate and bongkrekate is likely to be indirect, acting through the adenine nucleotide translocase.

Key words: Uncoupling protein; adenine nucleotide translocase; UCP3; GDP; carboxyatractylate; proton leak

INTRODUCTION

During substrate oxidation, protons are pumped out of the mitochondrial matrix, creating an electrochemical proton gradient. This protonmotive force is used to drive ATP synthesis. However, substrate oxidation is not fully coupled to ATP synthesis; some of the electrochemical gradient is dissipated by protons leaking back into the matrix through natural proton conductance pathways across the mitochondrial inner membrane [1, 2]. Mitochondrial proton leak may cause up to 25% of resting metabolic rate [3, 4] and may still be significant when ATP turnover is high [5]. The prominence of mitochondrial proton leak suggests that it is physiologically important, and several roles have been suggested [4], including protection against mitochondrial free radical production [6] and regulation of insulin secretion in pancreatic beta cells [7].

Proton leak occurs through several pathways, some basal and some inducible. Membranes are themselves intrinsically leaky to protons [8] but proteins embedded in a bilayer can greatly increase permeability [4, 9, 10]. Half to two-thirds of the basal mitochondrial proton conductance is dependent upon the most abundant carrier protein in the mitochondrial inner membrane, the adenine nucleotide translocase (ANT) [11]. ANT-dependent basal proton conductance occurs even if the carrier is inhibited by carboxyatractylate, showing that this leak pathway does not require normal carrier function [11].

Unlike basal proton conductance, specific activators and inhibitors affect inducible proton conductance. Inducible conductance can be catalysed by ANT in the presence of fatty acids [12, 13], AMP [14] or the lipid peroxidation product hydroxynonenal [15]. This pathway is inhibited by carboxyatractylate, bongkrekate and by the ANT substrates, ATP or ADP. It is not known to be sensitive to other nucleotides. Other mitochondrial carriers, such as the glutamate [16] and phosphate carriers [17], will catalyse proton conductance in the presence of fatty acids. Specific uncoupling proteins (UCPs) also catalyse inducible proton conductance [18-21]. The best understood is UCP1, found primarily in brown adipose tissue and specialised for adaptive thermogenesis. Proton conductance through UCP1 is activated by fatty acids [22], superoxide [23], derivatives such as hydroxynonenal [15, 24], and hydroxynonenal analogues (such as retinoic acid [25]). UCP2, found in a variety of tissues [26], and UCP3, found primarily in skeletal muscle [27, 28], can be activated by fatty acids [29, 30], superoxide [23] and hydroxynonenal and its analogues [15].

A striking aspect of the UCPs is the inhibition of their proton conductance by a broad range of purine nucleotides. Unlike the proton conductance of the adenine nucleotide carrier, UCPs are strongly inhibited not only by ATP and ADP, but also by GTP and GDP [14]. This difference in specificity has led to widespread use of GDP as a diagnostic inhibitor of UCP function *in vitro* (e.g. [31-34]), particularly in systems that lack a UCP-ablated control [35-38].

Here we report an endogenous increase in the proton conductance of skeletal muscle mitochondria incubated at high membrane potential. It is catalysed by UCP3 and ANT. Although it is inhibited by GDP and carboxyatractylate, these inhibitors do not show the expected classical specificity.

EXPERIMENTAL

Animals

Female Wistar rats (Charles River) and male or female mice were housed at $21 \pm 2^\circ\text{C}$, $57 \pm 5\%$ humidity, 12/12 h light/dark, with standard laboratory chow and water *ad libitum*, following UK Home Office Guidelines for the Care and Use of Laboratory Animals. *Ucp3* knockout (*Ucp3KO*) [39] and wild type (WT) sibling paired mice were used at age 12 – 15 weeks. *Ucp3* ablation was confirmed by PCR analysis of genomic *Ucp3* and Western blot in skeletal muscle mitochondria.

Mitochondria

Animals (1 rat or 4 mice per preparation) were killed by stunning and cervical dislocation. Mitochondria were isolated from total hind limb skeletal muscle [15]. Mitochondria from WT and *Ucp3KO* mice were prepared and assayed in parallel.

Proton leak kinetics

Proton leak rates across the inner membrane of isolated mitochondria were measured as the oxygen consumption required to pump the protons back out of the mitochondrial matrix (in the presence of oligomycin to prevent ATP synthesis) [40]. Proton leak kinetics were measured as the dependence of proton leak rate on its driving force, protonmotive force (measured as membrane potential in the presence of nigericin to abolish pH gradients). Respiration rate and membrane potential were measured simultaneously using electrodes sensitive to oxygen or the membrane potential-dependent probe, triphenylmethylphosphonium (TPMP⁺). Assay medium contained 0.35 mg/ml mitochondrial protein, 120 mM KCl, 5 mM KH₂PO₄, 3 mM Hepes, 1 mM EGTA, 0.3 % w/v defatted bovine serum albumin (unless otherwise stated), 5 μM rotenone, 1 μg/ml oligomycin and 0.11 μM nigericin. The TPMP electrode was calibrated with sequential additions of 0.5 μM TPMP up to 2.5 μM and the mitochondria were energised with 4 mM succinate. Leak kinetics were measured by varying oxygen consumption and membrane potential by sequential additions of malonate (about 0.1 – 2.3 mM). After each run, 0.2 μM carbonylcyanide *p*-trifluoromethoxyphenylhydrazone (FCCP) was added to release matrix TPMP, allowing correction for any small baseline drift. The TPMP binding correction was taken to be $0.35 (\mu\text{l}/\text{mg protein})^{-1}$ [41].

Measurement of ANT content by carboxyatractylate titre

Respiration was measured without oligomycin or nigericin. 1 mM ADP was added to establish the maximum phosphorylating rate, and then respiration was successively inhibited by additions of carboxyatractylate (Calbiochem) until the resting non-phosphorylating rate was re-established. Carboxyatractylate binds ANT very tightly, so ANT content was interpolated as the minimum amount of carboxyatractylate required to abolish phosphorylating respiration [11, 42].

Statistics

To test for differences in proton conductance by ANOVA or paired Student's *t* tests, we calculated respiration rates (by linear interpolation between flanking points) at the highest

membrane potential common to all conditions on a given day. Means for each condition (with S.E.M.) are plotted in bar graphs, with the mean value of the highest common potential indicated.

RESULTS

Time-dependent endogenous activation of proton conductance in energised rat skeletal muscle mitochondria

Preliminary investigations of proton leak kinetics in skeletal muscle mitochondria suggested that proton conductance increased as the time of energisation with succinate was extended. Fig. 1 characterises this endogenous activation of proton conductance. Mitochondria were incubated in assay medium for a total of 8 min (Fig. 1A). They were energised with succinate for the last 0.75, 2.5 or 5.0 min, then proton leak kinetics were measured. Fig. 1B shows that respiration driving proton leak increased as incubation with succinate was extended. To allow comparison of proton leak rates at the same driving force, Fig. 1C shows rates interpolated at 175 mV, the highest common membrane potential in the experiment, indicated by the vertical line. The proton leak rate at 175 mV measured after 5 min with succinate was significantly higher than that measured after 0.75 min with succinate. The total time in the assay was always 8 min, therefore endogenous proton conductance increases as the time of incubation with succinate is extended.

Leak kinetics are measured by titrating membrane potential and respiration rate with malonate over three minutes. If time with succinate is important in inducing proton leak, endogenous activation should continue during the malonate titration, causing later points to be more uncoupled than earlier points. To test this, Fig. 1D shows a compilation of leak kinetics measured in five separate experiments, each with a different concentration of malonate added at $t = 0$. Time with succinate was the same for all titration points (dashed line). This is overlaid on the leak kinetics measured during one standard experiment using five sequential additions of malonate (as in Fig. 1B), where time with succinate differed between titration points (solid line). There was no difference in leak kinetics between the two protocols. Therefore energisation-dependent uncoupling does not increase over the three minutes taken to measure the proton leak kinetics suggesting no further increases after the first malonate addition.

Malonate attenuates the energisation-dependent increase in proton conductance by lowering membrane potential in rat skeletal muscle mitochondria.

To test the ability of malonate to blunt energisation-dependent, endogenous activation of proton leak, 430 μM malonate was added immediately before succinate. At $t = 0$, after incubation with succinate for 1.5 min, this concentration of malonate lowered the membrane potential by about 12 mV (X in Fig. 2). Succinate was added 1.5 min or 5.0 min before the leak titration, which was completed as normal with sequential additions of malonate (Fig. 2B). Fig. 2A shows the control without malonate pre-addition.

To allow comparison of proton leak rates at the same driving force, Fig. 2C shows the time-dependent increases in rates with succinate, interpolated at the highest common membrane potential, 169 mV. The leak induced between 1.5 and 5 min was significantly

lower when 430 μM malonate was present, therefore malonate does indeed attenuate the energisation-dependent increase in proton conductance.

Pre-addition of malonate could inhibit energisation-dependent induced leak directly, or by lowering protonmotive force, or by oxidising the electron transport chain. To test these possibilities, the experiment in Fig. 2C was repeated using the complex IV inhibitor, KCN, at a concentration that lowers the membrane potential to the same extent as 430 μM malonate. Leak kinetics were measured in the absence (Fig. 2D) or presence (Fig. 2E) of 43 μM KCN. At the highest common membrane potential, proton conductance induced by succinate between 1.5 and 5 min in the presence of 43 μM KCN (Fig. 2F) was significantly lower than in its absence. Unlike malonate, KCN reduces electron transport chain intermediates. The two inhibitors decreased energisation-dependent proton conductance to a similar extent, suggesting that it is high membrane potential, not a particular redox state, that is required for energisation-dependent induction of proton conductance.

The increase in proton conductance between 1.5 and 5 min incubation with succinate was also measured at intermediate concentrations of pre-added malonate or KCN. This produced intermediate drops in membrane potential at $t = 0$ following 1.5 min with succinate (equivalent to X in Fig. 2B and Fig. 2E). To characterise the ability of decreased membrane potential to blunt energisation-dependent induction of proton conductance, Fig. 2G shows the increase in leak rate (e.g. Fig. 2C and Fig. 2F) plotted against the drop (X) in membrane potential caused by different concentrations of pre-added malonate or KCN. There was a linear decrease in activation as membrane potential during the activatory period was decreased. The linear regressions for malonate and KCN overlap suggesting a conserved mechanism of inhibition through the lowering of membrane potential. Extrapolation to zero effect suggests that a decrease of 20 mV is sufficient to prevent activation of proton conductance.

Therefore, the endogenous energisation-dependent activation of proton conductance following addition of succinate requires a high membrane potential. Activation increases over 5 min with succinate but during subsequent titrations with malonate the membrane potential falls by more than 20 mV and further activation is prevented.

What is the activator of energisation-dependent increase in proton conductance in rat skeletal muscle mitochondria?

Bovine serum albumin (BSA) removes putative endogenous activators, including fatty acids [43] and reactive oxygen species [6] and their derivatives such as reactive alkenals [15]. To test whether these endogenous activators were involved in the observed energisation-dependent proton leak, the concentration of BSA was increased in the standard assay medium from 0.3% to 1% w/v (Fig. 3A-C). Proton leak kinetics were measured after energisation with succinate for 1.5 min or 5.0 min in the presence of 0.3% (Fig. 3A) or 1% (Fig. 3B) BSA. At 172 mV, leak induced between 1.5 and 5 min in assay medium containing 1% BSA was significantly lower than in 0.3% BSA (Fig. 3C), showing that BSA blunts energisation-dependent activation of proton conductance and suggesting that an endogenous activator that is quenched by BSA mediates the induced leak.

To test the possibility that the increase in energisation-dependent leak is due to lipid peroxidation products, phenylbutylnitron, a chain-breaking spin trap that quenches carbon-centred radical intermediates, was pre-added to the assay medium at concentrations known to inhibit lipid peroxidation [44]. Proton leak kinetics were measured after energisation with succinate for 1.5 or 5.0 min in the absence (Fig. 3D) or presence (Fig. 3E) of phenylbutylnitron. At 169 mV, there was no significant effect of phenylbutylnitron on leak induced between 1.5 and 5 min (Fig. 3F), suggesting that pathways that generate lipid peroxidation products through a carbon-centred radical sensitive to phenylbutylnitron are not involved in the energisation-dependent uncoupling. However, other reactive oxygen species-related activators remain candidates.

Energisation might either generate a pool of activator, or allow activation of uncoupling using a pre-existing activator pool (e.g. through an energisation-dependent change in conformation of an uncoupling protein). To distinguish between these alternatives, we tested the effect of pre-addition of fatty acids in medium containing 1% BSA. Proton leak kinetics were measured in 1% BSA in the absence (Fig. 3G) or presence (Fig. 3H) of 600 μ M palmitate after energisation with succinate for 1.5 or 5.0 min. Fig. 3I replicates the observation in Fig. 3C that 1% BSA alone prevented time-dependent activation. Pre-addition of palmitate did not overcome the inhibitory effect of 1% BSA, showing that time-dependent activation did not occur by potential-dependent changes in protein conformation that then allowed activation by a pre-existing pool of fatty acids. Fatty acids remain, however, one candidate for the endogenously-generated activator, since proton leak kinetics in the presence of palmitate (Fig. 3H) were the same after 1.5 min incubation with succinate as those seen in 0.3 % BSA after 5 min with succinate (Fig. 3A).

These experiments are consistent with at least three different interpretations. The endogenous activator may be (a) fatty acids, or (b) some other molecule, generated during 5 min incubation at high membrane potential, or (c) some molecule (e.g. hydroxynonenal) that is already present in the mitochondria before addition of succinate, but requires membrane potential to be high for several minutes to allow the interaction that causes uncoupling.

Which proteins catalyse the energisation-dependent increase in proton conductance in rat skeletal muscle mitochondria?

Uncoupling through the adenine nucleotide translocase is inhibited by carboxyatractylate [13], whereas uncoupling through UCPs is classically inhibited by GDP [14, 45]. Proton leak kinetics were measured after energisation with succinate for 2.5 min in the absence and presence of carboxyatractylate and GDP to determine whether ANT and UCP3 were involved (Fig. 4A). Fig. 4B shows that at the highest common membrane potential, 180 mV, GDP and carboxyatractylate significantly inhibited proton conductance separately and in combination. The extent of the carboxyatractylate inhibition of proton conductance suggests that the majority is attributable to ANT. As observed previously [15], inhibition of proton conductance by GDP and carboxyatractylate was non-additive; characterisation of the relative contributions of ANT and UCP3 therefore required further experiments.

UCP3 and ANT each catalyse part of the energisation-dependent increase in proton conductance in mouse skeletal muscle mitochondria

The contribution of UCP3 to energisation-dependent proton conductance was assessed using skeletal muscle mitochondria prepared from WT and *Ucp3*KO mice. Proton leak kinetics were measured after energisation with succinate for 1.5 or 5.0 min. Fig. 5A shows that skeletal muscle mitochondria from WT mice showed a similar time-dependent increase in proton leak after addition of succinate to mitochondria from rat (Fig. 1B). Fig. 5B shows the same experiment performed using skeletal muscle mitochondria from *Ucp3*KO mice. There was still a difference between the curves, but it was smaller than in WT. The difference is quantified in Fig. 5C, which shows proton leak through UCP3, expressed as the difference between rates in skeletal muscle mitochondria from WT and *Ucp3*KO mice at 169 mV. UCP3-associated proton conductance was significantly greater after incubation with succinate for 5.0 min than for 1.5 min. In the presence of UCP3, the increase in respiration rate between 1.5 and 5 min was 75 nmol/min/mg protein while in the absence of UCP3 it was 62 nmol/min/mg protein. Thus about 17% of the energisation-dependent proton leak induced between 1.5 and 5 min incubation with succinate requires the presence of UCP3.

The contribution of ANT to the proton conductance observed after 5 min incubation with succinate was assessed as the carboxyatractylate-sensitive effect in skeletal muscle mitochondria from *Ucp3*KO mice. Fig. 6B shows that carboxyatractylate substantially inhibited energisation-dependent proton conductance in the absence of UCP3. The effect is quantified in Fig. 6D, which shows UCP3-independent leak at 161 mV. Carboxyatractylate lowered the rate from 90 to 30 nmol O/min/mg protein, equivalent to about 50-55% of the total effect in mitochondria from WT mice in Fig. 6A. Together, these results show that the energisation-dependent proton conductance in mouse skeletal muscle mitochondria is catalysed about 15-20% by UCP3, 50-55% by ANT, and 25-30% by other pathways not sensitive to GDP or carboxyatractylate (e.g. the dicarboxylate carrier).

Specificity and lack of additivity of inhibition by GDP and carboxyatractylate in mouse skeletal muscle mitochondria

To investigate the lack of additivity between the inhibitors GDP and carboxyatractylate in rat skeletal muscle (Fig. 4), and also seen in other studies [15], inhibition was measured in skeletal muscle mitochondria from WT and *Ucp3*KO mice. The same lack of additivity was seen in skeletal muscle mitochondria from WT mice (Fig. 6A) as in rat: GDP partially inhibited proton conductance in the absence of carboxyatractylate but caused no further inhibition in the presence of carboxyatractylate. Fig. 6B shows the same effect in mitochondria from *Ucp3*KO mice.

Figs 6C and 6D dissect out the effects of the inhibitors. Fig. 6C represents the contribution of UCP3 (Fig. 6A minus 6B) at the highest common membrane potential, 161 mV, after 5 min incubation with succinate. Surprisingly (since GDP is the classical inhibitor of UCPs), the proton conductance through UCP3 under these conditions was insensitive to GDP. More surprisingly (since carboxyatractylate is thought to be a highly specific inhibitor of ANT), it was almost fully sensitive to carboxyatractylate. This sensitivity of UCP3 to carboxyatractylate partially explains the lack of additivity of the two inhibitors: once UCP3 is inhibited by carboxyatractylate, any inhibition by GDP would have no further effect.

Fig. 6D represents the contribution of ANT and other pathways (data from Fig. 6B). The energisation-dependent proton conductance of mitochondria from *Ucp3*KO mice was significantly lower in the presence of GDP, indicating that an endogenously activated pathway other than UCP3 was inhibited by GDP. Since this conductance was also strongly inhibited by carboxyatractylate, it was catalysed by ANT. This experiment shows that GDP is not specific for UCPs, but can also inhibit proton conductance through other pathways, specifically ANT. It completes the explanation of the lack of additivity of GDP and carboxyatractylate: once ANT is inhibited by carboxyatractylate, inhibition by GDP would have no further effect. Together, these results highlight potential promiscuity of the inhibitors GDP and carboxyatractylate: neither is specific for its classical target in our experiments.

A possible alternative explanation of the sensitivity of mitochondria from *Ucp3*KO mice to GDP is that the *Ucp3*KO targeting vector upregulates *Ucp2* expression, due to proximity of the *Ucp2* and *Ucp3* genes [46], and it is this UCP2 that is inhibited by GDP. We detected *Ucp2* transcript in total hind limb skeletal muscle using real-time PCR, but at levels 30 times lower than in spleen or lung, with no significant difference between WT and *Ucp3*KO (data not shown). UCP2 was undetectable by Western blot using skeletal muscle mitochondria (20 μ g protein) from either WT or *Ucp3*KO mice. The antibody can detect 125 pg affinity-purified recombinant UCP2, so UCP2 content is below 6 ng/mg mitochondrial protein, less than 5% of the UCP3 content of 140 ng/mg protein in mouse skeletal muscle mitochondria [47]. UCP1 is not expressed in skeletal muscle of WT or *Ucp3*KO mice [48]. Therefore, the putative presence of UCP1 or UCP2 does not explain the sensitivity of proton conductance in skeletal muscle mitochondria from *Ucp3*KO mice to GDP.

A possible alternative explanation of the decreased effect of carboxyatractylate in mitochondria from *Ucp3*KO mice is that ablation of UCP3 somehow decreased ANT content. We tested this possibility by titrating adenine nucleotide translocase activity with carboxyatractylate in skeletal muscle mitochondria from WT and *Ucp3*KO mice. UCP3 (4 pmol/mg mitochondrial protein [47]) is about 0.2% as abundant as ANT (2 nmol/mg [11]), so any sequestering of carboxyatractylate by UCP3 would be negligible. ANT content was not significantly different (WT, 1.52; *Ucp3*KO, 1.56 nmol/mg protein, Fig. 6E), eliminating this explanation.

Inhibition of UCP3 by carboxyatractylate could be direct or through ANT. To examine this, we tested the effects of an alternative ANT inhibitor, bongkrekate. Unlike carboxyatractylate, which binds to the C conformation of the carrier, bongkrekate binds to the M conformation [49]. BSA was omitted from the assay medium because it binds bongkrekate. Proton leak kinetics were measured in skeletal muscle mitochondria isolated from WT (Fig. 7A) and *Ucp3*KO (Fig. 7B) mice after energisation with succinate for 5.0 min in the absence or presence of bongkrekate or carboxyatractylate. Bongkrekate partially inhibited the energisation-dependent proton conductance in both cases. Fig. 7C shows the effects that were dependent on UCP3 (Fig. 7A minus Fig. 7B). At the highest common membrane potential, 158 mV, both carboxyatractylate and bongkrekate inhibited proton conductance through UCP3. It is unlikely that UCP3 and ANT share two different high-affinity inhibitor sites, therefore both carboxyatractylate and bongkrekate act on the UCP3-associated leak indirectly through primary effects on ANT.

DISCUSSION

We describe an endogenous activation of the proton conductance of skeletal muscle mitochondria isolated from both rats and mice that increases with time after energisation. This effect is not dependent upon redox state but requires a high membrane potential upon succinate addition. The effect of membrane potential may be to increase the production of endogenous fatty acids or other molecules that progressively activate uncoupling, or to sensitise target proteins to the effects of generated or pre-existing activators such as products of reactive oxygen species. The mechanism remains unclear. Membrane potential might change the conformation of membrane lipids to facilitate access by phospholipases, or might change the conformations of UCP3 and ANT to expose activator-binding sites. Mobilisation of fatty acids could explain the lack of GDP sensitivity of the UCP3-dependent proton conductance, by analogy with UCP1, where fatty acids functionally compete with inhibitory nucleotides [50]. Whatever the activator and mechanism, energisation-dependent uncoupling of skeletal muscle mitochondria may be important in limiting membrane potential during rest, to limit reactive oxygen species production [21] or prevent dielectric breakdown of the membrane [4].

Two catalysts of energisation-dependent proton conductance were found to be UCP3 and ANT. In WT mouse skeletal muscle mitochondria after 5 min incubation with succinate, the relative contributions of UCP3, ANT and other pathways to proton conductance were 15-20%, 50-55% and 25-30% respectively. Since UCP3 content is only about 0.2% of ANT content (see above), this result demonstrates that the specific proton flux through UCP3 is 100-200 times greater than that through ANT. The greater specific activity of UCP3 may allow finer control of high membrane potential in tissues that contain this protein in addition to ANT.

We found that GDP partially inhibited the contribution of ANT to energisation-dependent proton conductance. Khailova *et al* [51] also concluded that GDP inhibits proton translocation by ANT, although they did not use *Ucp* knockout controls, which provide our most compelling evidence that GDP can inhibit proton conductance in the absence of UCP3. GDP and GTP bind weakly to ANT [52], but compete poorly with ADP and ATP, so they are not regarded as inhibitors of adenine nucleotide exchange. However, adenine nucleotides are absent in assays of proton conductance, which may allow guanine nucleotides to bind and partially inhibit ANT-catalysed proton leak.

We found that carboxyatractylate inhibited the contribution of UCP3 to proton leak. Carboxyatractylate is a high-affinity inhibitor of ANT, and has not been shown previously to act on other proteins. However, most other studies have not looked for carboxyatractylate inhibition of UCP3 function. The observation that bongkrekate also inhibits UCP3-dependent proton conductance suggests that inhibition of UCP3 is indirect, via primary effects on ANT. It is unclear what these indirect effects might be, but they might involve protein-protein interactions between ANT and UCP3, or interactions through some other protein, lipid or small molecule. More exploration of the interaction is required to uncover the mechanism. However, the (indirect) inhibition of UCP3 by carboxyatractylate fully explains the hitherto puzzling lack of additivity between inhibition of proton conductance by GDP and carboxyatractylate.

Since inhibition by GDP and carboxyatractylate is also non-additive in mitochondria from kidney [15, 44], carboxyatractylate may also inhibit UCP2 by a

similar indirect mechanism. However, it appears that carboxyatractylate does not inhibit potato UCP [53] or mammalian UCP1, since guanine nucleotides are still inhibitory in the presence of carboxyatractylate in mitochondria from potato or mammalian brown adipose tissue. Thus caution should be shown when using sensitivity to GDP or carboxyatractylate as diagnostic of the involvement of a UCP or ANT in proton conductance. Observations of GDP- or carboxyatractylate-sensitivity should be verified by independent assays of the involvement of UCPs and ANT, such as sensitivity to genetic ablation or over-expression.

ACKNOWLEDGEMENTS

We thank Dr. Mary-Ellen Harper (Ottawa) for *Ucp3*KO mice and Julie Buckingham and Helen Boysen for technical assistance. This work was supported by the Medical Research Council and the Wellcome Trust (grants 065326/Z/01/Z and 066750/B/01/Z).

REFERENCES

- 1 Brand, M. D. (1990) The proton leak across the mitochondrial inner membrane. *Biochim. Biophys. Acta.* **1018**, 128-133
- 2 Brand, M. D., Chien, L. F., Ainscow, E. K., Rolfe, D. F. S. and Porter, R. K. (1994) The causes and functions of mitochondrial proton leak. *Biochim. Biophys. Acta.* **1187**, 132-139
- 3 Rolfe, D. F. S. and Brand, M. D. (1996) Contribution of mitochondrial proton leak to skeletal muscle respiration and to standard metabolic rate. *Am. J. Physiol.* **271**, C1380-1389
- 4 Rolfe, D. F. S. and Brand, M. D. (1997) The physiological significance of mitochondrial proton leak in animal cells and tissues. *Biosci. Rep.* **17**, 9-16
- 5 Rolfe, D. F. S., Newman, J. M., Buckingham, J. A., Clark, M. G. and Brand, M. D. (1999) Contribution of mitochondrial proton leak to respiration rate in working skeletal muscle and liver and to SMR. *Am. J. Physiol.* **276**, C692-699
- 6 Echtay, K. S., Murphy, M. P., Smith, R. A., Talbot, D. A. and Brand, M. D. (2002) Superoxide activates mitochondrial uncoupling protein 2 from the matrix side. Studies using targeted antioxidants. *J. Biol. Chem.* **277**, 47129-47135
- 7 Affourtit, C. and Brand, M. D. (2008) Uncoupling protein-2 contributes significantly to high mitochondrial proton leak in INS-1E insulinoma cells and attenuates glucose-stimulated insulin secretion. *Biochem. J.* **409**, 199-204
- 8 Dilger, J. P., McLaughlin, S. G., McIntosh, T. J. and Simon, S. A. (1979) The dielectric constant of phospholipid bilayers and the permeability of membranes to ions. *Science* **206**, 1196-1198
- 9 Brookes, P. S., Rolfe, D. F. S. and Brand, M. D. (1997) The proton permeability of liposomes made from mitochondrial inner membrane phospholipids: comparison with isolated mitochondria. *J. Membr. Biol.* **155**, 167-174
- 10 Brown, G. C. and Brand, M. D. (1991) On the nature of the mitochondrial proton leak. *Biochim. Biophys. Acta.* **1059**, 55-62

- 11 Brand, M. D., Pakay, J. L., Ocloo, A., Kokoszka, J., Wallace, D. C., Brookes, P. S. and Cornwall, E. J. (2005) The basal proton conductance of mitochondria depends on adenine nucleotide translocase content. *Biochem. J.* **392**, 353-362
- 12 Andreyev, A., Bondareva, T. O., Dedukhova, V. I., Mokhova, E. N., Skulachev, V. P., Tsofina, L. M., Volkov, N. I. and Vygodina, T. V. (1989) The ATP/ADP-antiporter is involved in the uncoupling effect of fatty acids on mitochondria. *Eur. J. Biochem.* **182**, 585-592
- 13 Andreyev, A., Bondareva, T. O., Dedukhova, V. I., Mokhova, E. N., Skulachev, V. P. and Volkov, N. I. (1988) Carboxyatractylate inhibits the uncoupling effect of free fatty acids. *FEBS Lett.* **226**, 265-269
- 14 Cadenas, S., Buckingham, J. A., St-Pierre, J., Dickinson, K., Jones, R. B. and Brand, M. D. (2000) AMP decreases the efficiency of skeletal-muscle mitochondria. *Biochem. J.* **351 Pt 2**, 307-311
- 15 Echtay, K. S., Esteves, T. C., Pakay, J. L., Jekabsons, M. B., Lambert, A. J., Portero-Otin, M., Pamplona, R., Vidal-Puig, A. J., Wang, S., Roebuck, S. J. and Brand, M. D. (2003) A signalling role for 4-hydroxy-2-nonenal in regulation of mitochondrial uncoupling. *EMBO J.* **22**, 4103-4110
- 16 Samartsev, V. N., Smirnov, A. V., Zeldi, I. P., Markova, O. V., Mokhova, E. N. and Skulachev, V. P. (1997) Involvement of aspartate/glutamate antiporter in fatty acid-induced uncoupling of liver mitochondria. *Biochim. Biophys. Acta.* **1319**, 251-257
- 17 Engstova, H., Zackova, M., Ruzicka, M., Meinhardt, A., Hanus, J., Kramer, R. and Jezek, P. (2001) Natural and azido fatty acids inhibit phosphate transport and activate fatty acid anion uniport mediated by the mitochondrial phosphate carrier. *J. Biol. Chem.* **276**, 4683-4691
- 18 Echtay, K. S., Pakay, J. L., Esteves, T. C. and Brand, M. D. (2005) Hydroxynonenal and uncoupling proteins: a model for protection against oxidative damage. *Biofactors* **24**, 119-130
- 19 Brand, M. D. and Esteves, T. C. (2005) Physiological functions of the mitochondrial uncoupling proteins UCP2 and UCP3. *Cell Metab.* **2**, 85-93
- 20 Esteves, T. C. and Brand, M. D. (2005) The reactions catalysed by the mitochondrial uncoupling proteins UCP2 and UCP3. *Biochim. Biophys. Acta.* **1709**, 35-44
- 21 Brand, M. D., Affourtit, C., Esteves, T. C., Green, K., Lambert, A. J., Miwa, S., Pakay, J. L. and Parker, N. (2004) Mitochondrial superoxide: production, biological effects, and activation of uncoupling proteins. *Free Radic. Biol. Med.* **37**, 755-767
- 22 Rafael, J., Ludolph, H. J. and Hohorst, H. J. (1969) [Mitochondria from brown adipose tissue: uncoupling of respiratory chain phosphorylation by long fatty acids and recoupling by guanosine triphosphate]. *Hoppe Seylers Z Physiol. Chem.* **350**, 1121-1131
- 23 Echtay, K. S., Roussel, D., St-Pierre, J., Jekabsons, M. B., Cadenas, S., Stuart, J. A., Harper, J. A., Roebuck, S. J., Morrison, A., Pickering, S., Clapham, J. C. and Brand, M. D. (2002) Superoxide activates mitochondrial uncoupling proteins. *Nature* **415**, 96-99

- 24 Esteves, T. C., Parker, N. and Brand, M. D. (2006) Synergy of fatty acid and reactive alkenal activation of proton conductance through uncoupling protein 1 in mitochondria. *Biochem. J.* **395**, 619-628
- 25 Rial, E., Gonzalez-Barroso, M., Fleury, C., Iturrizaga, S., Sanchis, D., Jimenez-Jimenez, J., Ricquier, D., Goubern, M. and Bouillaud, F. (1999) Retinoids activate proton transport by the uncoupling proteins UCP1 and UCP2. *EMBO J.* **18**, 5827-5833
- 26 Fleury, C., Neverova, M., Collins, S., Raimbault, S., Champigny, O., Levi-Meyrueis, C., Bouillaud, F., Seldin, M. F., Surwit, R. S., Ricquier, D. and Warden, C. H. (1997) Uncoupling protein-2: a novel gene linked to obesity and hyperinsulinemia. *Nat. Genet.* **15**, 269-272
- 27 Boss, O., Samec, S., Paoloni-Giacobino, A., Rossier, C., Dulloo, A., Seydoux, J., Muzzin, P. and Giacobino, J. P. (1997) Uncoupling protein-3: a new member of the mitochondrial carrier family with tissue-specific expression. *FEBS Lett.* **408**, 39-42
- 28 Vidal-Puig, A., Solanes, G., Grujic, D., Flier, J. S. and Lowell, B. B. (1997) UCP3: an uncoupling protein homologue expressed preferentially and abundantly in skeletal muscle and brown adipose tissue. *Biochem. Biophys. Res. Commun.* **235**, 79-82
- 29 Echtay, K. S., Winkler, E., Frischmuth, K. and Klingenberg, M. (2001) Uncoupling proteins 2 and 3 are highly active H(+) transporters and highly nucleotide sensitive when activated by coenzyme Q (ubiquinone). *Proc. Natl. Acad. Sci. U. S. A.* **98**, 1416-1421
- 30 Jaburek, M., Varecha, M., Gimeno, R. E., Dembski, M., Jezek, P., Zhang, M., Burn, P., Tartaglia, L. A. and Garlid, K. D. (1999) Transport function and regulation of mitochondrial uncoupling proteins 2 and 3. *J. Biol. Chem.* **274**, 26003-26007
- 31 Dejean, L., Camara, Y., Sibille, B., Solanes, G. and Villarroya, F. (2004) Uncoupling protein-3 sensitizes cells to mitochondrial-dependent stimulus of apoptosis. *J. Cell Physiol.* **201**, 294-304
- 32 Nadtochiy, S. M., Tompkins, A. J. and Brookes, P. S. (2006) Different mechanisms of mitochondrial proton leak in ischaemia/reperfusion injury and preconditioning: implications for pathology and cardioprotection. *Biochem. J.* **395**, 611-618
- 33 Carroll, A. M., Haines, L. R., Pearson, T. W., Fallon, P. G., Walsh, C. M., Brennan, C. M., Breen, E. P. and Porter, R. K. (2005) Identification of a functioning mitochondrial uncoupling protein 1 in thymus. *J. Biol. Chem.* **280**, 15534-15543
- 34 Ruzicka, M., Skobisova, E., Dlaskova, A., Santorova, J., Smolkova, K., Spacek, T., Zackova, M., Modriansky, M. and Jezek, P. (2005) Recruitment of mitochondrial uncoupling protein UCP2 after lipopolysaccharide induction. *Int J Biochem. Cell Biol.* **37**, 809-821
- 35 Fridell, Y. W., Sanchez-Blanco, A., Silvia, B. A. and Helfand, S. L. (2004) Functional characterization of a Drosophila mitochondrial uncoupling protein. *J Bioenerg. Biomembr.* **36**, 219-228

- 36 Czarna, M. and Jarmuszkiewicz, W. (2005) Activation of alternative oxidase and uncoupling protein lowers hydrogen peroxide formation in amoeba *Acanthamoeba castellanii* mitochondria. *FEBS Lett.* **579**, 3136-3140
- 37 Talbot, D. A., Duchamp, C., Rey, B., Hanuise, N., Rouanet, J. L., Sibille, B. and Brand, M. D. (2004) Uncoupling protein and ATP/ADP carrier increase mitochondrial proton conductance after cold adaptation of king penguins. *J. Physiol.* **558**, 123-135
- 38 Hourton-Cabassa, C., Mesneau, A., Miroux, B., Roussaux, J., Ricquier, D., Zachowski, A. and Moreau, F. (2002) Alteration of plant mitochondrial proton conductance by free fatty acids. Uncoupling protein involvement. *J. Biol. Chem.* **277**, 41533-41538
- 39 Gong, D. W., Monemdjou, S., Gavrilova, O., Leon, L. R., Marcus-Samuels, B., Chou, C. J., Everett, C., Kozak, L. P., Li, C., Deng, C., Harper, M. E. and Reitman, M. L. (2000) Lack of obesity and normal response to fasting and thyroid hormone in mice lacking uncoupling protein-3. *J. Biol. Chem.* **275**, 16251-16257
- 40 Rolfe, D. F. S., Hulbert, A. J. and Brand, M. D. (1994) Characteristics of mitochondrial proton leak and control of oxidative phosphorylation in the major oxygen-consuming tissues of the rat. *Biochim. Biophys. Acta.* **1188**, 405-416
- 41 Brand, M. D. (1995) Measurement of mitochondrial protonmotive force. In *Bioenergetics - A Practical Approach* (Brown, G. C. and Cooper, C. E., eds.), pp. 39-62, IRL Press, Oxford, UK
- 42 Streicher-Scott, J., Lapidus, R. and Sokolove, P. M. (1993) Use of carboxyatractylate and tight-binding inhibitor theory to determine the concentration of functional mitochondrial adenine nucleotide translocators in a reconstituted system. *Anal. Biochem.* **210**, 69-76
- 43 Mellors, A., Tappel, A. L., Sawant, P. L. and Desai, I. D. (1967) Mitochondrial swelling and uncoupling of oxidative phosphorylation by lysosomes. *Biochim. Biophys. Acta.* **143**, 299-309
- 44 Murphy, M. P., Echtay, K. S., Blaikie, F. H., Asin-Cayuela, J., Cocheme, H. M., Green, K., Buckingham, J. A., Taylor, E. R., Hurrell, F., Hughes, G., Miwa, S., Cooper, C. E., Svistunenko, D. A., Smith, R. A. and Brand, M. D. (2003) Superoxide activates uncoupling proteins by generating carbon-centered radicals and initiating lipid peroxidation: studies using a mitochondria-targeted spin trap derived from alpha-phenyl-N-tert-butylnitron. *J. Biol. Chem.* **278**, 48534-48545
- 45 Nicholls, D. G. (1976) Hamster brown-adipose-tissue mitochondria. Purine nucleotide control of the ion conductance of the inner membrane, the nature of the nucleotide binding site. *Eur. J. Biochem.* **62**, 223-228
- 46 Solanes, G., Vidal-Puig, A., Grujic, D., Flier, J. S. and Lowell, B. B. (1997) The human uncoupling protein-3 gene. Genomic structure, chromosomal localization, and genetic basis for short and long form transcripts. *J. Biol. Chem.* **272**, 25433-25436
- 47 Harper, J. A., Stuart, J. A., Jekabsons, M. B., Roussel, D., Brindle, K. M., Dickinson, K., Jones, R. B. and Brand, M. D. (2002) Artfactual uncoupling by uncoupling protein 3 in yeast mitochondria at the concentrations found in mouse and rat skeletal-muscle mitochondria. *Biochem. J.* **361**, 49-56

- 48 Vidal-Puig, A. J., Grujic, D., Zhang, C. Y., Hagen, T., Boss, O., Ido, Y.,
Szczepanik, A., Wade, J., Mootha, V., Cortright, R., Muoio, D. M. and Lowell, B.
B. (2000) Energy metabolism in uncoupling protein 3 gene knockout mice. *J.*
Biol. Chem. **275**, 16258-16266
- 49 Kramer, R. and Klingenberg, M. (1979) Reconstitution of adenine nucleotide
transport from beef heart mitochondria. *Biochemistry* **18**, 4209-4215
- 50 Shabalina, I. G., Jacobsson, A., Cannon, B. and Nedergaard, J. (2004) Native
UCP1 displays simple competitive kinetics between the regulators purine
nucleotides and fatty acids. *J. Biol. Chem.*
- 51 Khailova, L. S., Prikhodko, E. A., Dedukhova, V. I., Mokhova, E. N., Popov, V.
N. and Skulachev, V. P. (2006) Participation of ATP/ADP antiporter in oleate-
and oleate hydroperoxide-induced uncoupling suppressed by GDP and
carboxyatractylate. *Biochim. Biophys. Acta.* **1757**, 1324-1329
- 52 Huber, T., Klingenberg, M. and Beyer, K. (1999) Binding of nucleotides by the
mitochondrial ADP/ATP carrier as studied by ¹H nuclear magnetic resonance
spectroscopy. *Biochemistry* **38**, 762-769
- 53 Smith, A. M., Ratcliffe, R. G. and Sweetlove, L. J. (2004) Activation and function
of mitochondrial uncoupling protein in plants. *J. Biol. Chem.* **279**, 51944-51952

FIGURE LEGENDS

Figure 1 Energisation-dependent proton conductance of rat skeletal muscle mitochondria

Proton leak kinetics were measured in rat skeletal muscle mitochondria as described in the experimental section. (A) Timings of additions to the incubation chamber. Mitochondria plus inhibitors (M) were added at $t = -8$ min, 4 mM succinate (S) was added at the times indicated, then leak kinetics (LK) were measured between $t = 0$ and $t = 3$ min by titration of membrane potential and oxygen consumption rate with malonate from 0.14 to 2.3 mM. (B) Kinetic response of mitochondrial proton leak rate (measured as oxygen consumption rate) to membrane potential measured after energisation with 4 mM succinate for 0.75 (open squares), 2.5 (grey squares) or 5.0 min (black squares). (C) Proton leak rate at 175 mV (solid line (B)); the average highest protonmotive force common to all curves generated in the experiment. (D) Comparison of proton leak curves generated either in a single experiment using sequential malonate additions over 3 min (S at $t = -5$, leak kinetics measured between $t = 0$ and $t = 3$ min; squares) or in five separate experiments each with a different concentration of malonate added at $t = 0$ (S at $t = -5$; 0.14 to 2.3 mM malonate added at $t = 0$, leak kinetics measured between $t = 0$ and $t = 0.5$ min; triangles). Data are means \pm S.E.M. of duplicate experiments performed on four (B-C) or three (D) separate preparations. One-way ANOVA was performed for (C); $p = 0.0034$, with Dunnett's multiple comparison test used posthoc comparing other bars to 0.75 min: **significantly different: $p < 0.01$.

Figure 2 Sensitivity of energisation-dependent proton conductance in rat skeletal muscle mitochondria to membrane potential

(A-C), effect of malonate. (D-F), effect of KCN. (A,B,D,E), proton leak kinetics. 4 mM succinate was added at $t = -1.5$ min (open symbols) or $t = -5$ min (closed symbols) (see Fig. 1A). Proton leak kinetics were measured between $t = 0$ and $t = 3$ min after no treatment (A, D) or addition of 430 μ M malonate (B) or 43 μ M KCN (E) immediately prior to addition of succinate. (C, F), inhibition of energisation-dependent proton leak by malonate and KCN. The increase in proton conductance at $t = 0$ caused by incubation with succinate between -5.0 min and -1.5 min was interpolated at the highest common membrane potential, 169 mV (solid line (A,B)) in (C) and 170 mV (solid line (D,E)) in (F). 'X' indicates the drop in membrane potential measured at $t = 0$ caused by addition of malonate or KCN prior to addition of succinate at $t = -1.5$. (G), effect of this drop in membrane potential on energisation-dependent increase in proton conductance. The increases in proton leak rate from (C) and additional experiments (not shown) using 140 μ M and 290 μ M malonate (circles), and from (F) and additional experiments (not shown) using 14 μ M and 29 μ M KCN (triangles) are plotted against the drop ('X') in membrane potential at $t = 0$ after succinate addition at $t = -1.5$ min. Data are means \pm S.E.M. of 5 (A-C) or 6 (D-F) experiments each performed in duplicate. *Significantly different by paired Student's t test: $p < 0.03$. Lines in (G) are linear regressions.

Figure 3 Modulation of energisation-dependent leak in rat skeletal muscle mitochondria by BSA, phenylbutylnitron and palmitate

(A-C), effect of 1% BSA. (D-F), effect of phenylbutylnitron (PBN). (G-I), effect of palmitate. (A,B,D,E,G,H), proton leak kinetics. 4 mM succinate was added at $t = -1.5$ min (open symbols) or $t = -5$ min (closed symbols) (see Fig. 1A). Proton leak kinetics were measured between $t = 0$ and $t = 3$ min in medium containing 0.3% (A,D) or 1% w/v BSA (B,G,H), after no pre-treatment or pre-treatment with 0.1 mM (triangles) or 1 mM phenylbutylnitron (diamonds) (E) or 600 μ M palmitate (H). (C,F,I), effect on energisation-dependent proton leak of 1% BSA, phenylbutylnitron or palmitate. The increase in proton conductance at $t = 0$ caused by incubation with succinate between -5 and -1.5 min was interpolated at the highest common membrane potential, 172 mV (solid line (A,B,D,E)) in (C) and (I) and 167 mV (solid line (G,H)) in (F). Data are means \pm S.E.M. of 4 (A-C), 3 (D-F) or 3 (G-H) experiments, each performed in duplicate. *Significantly different by paired Student's t test: $p < 0.01$.

Figure 4 Inhibition of proton conductance in rat skeletal muscle mitochondria by GDP and carboxyatractylate

(A) Proton leak kinetics measured between $t = 0$ and $t = 3$ min following addition of inhibitors (where present) and mitochondria at $t = -5.5$ min. 4 mM succinate was added at $t = -2.5$ min (see Fig. 1A) in the presence of no additions (squares), 500 μ M GDP (triangles), 2.5 μ M carboxyatractylate (diamonds) or GDP and carboxyatractylate (circles). (B) Absolute respiration rate at 180 mV (solid line (A)) in each condition. Data are means \pm S.E.M. of 15 experiments, each performed in duplicate. One-way ANOVA was performed for (B); $p = 0.0001$, with Dunnett's multiple comparison test used posthoc comparing other bars to control: **significantly different: $p < 0.01$.

Figure 5 Contribution of UCP3 and ANT to energisation-dependent leak in mouse skeletal muscle mitochondria

(A, B) proton leak kinetics. (A) mitochondria from WT mice. (B), mitochondria from *UCP3*KO mice. 4 mM succinate was added at $t = -1.5$ min (open symbols) or $t = -5$ min (closed symbols) (see Fig. 1A). (C), proton leak due to UCP3 measured at 169 mV (solid line (A,B)) as (A) minus (B). Data are means \pm S.E.M. of 7 experiments, each performed in duplicate. **Significantly different by paired Student's t test: $p < 0.01$.

Figure 6 Inhibition of proton conductance in skeletal muscle mitochondria from *Ucp3*WT and *Ucp3*KO mice by GDP and carboxyatractylate

(A-B), proton leak kinetics. (A) mitochondria from WT mice. (B), mitochondria from *UCP3*KO mice. 4 mM succinate was added at $t = -5$ min (see Fig. 1A). Proton leak kinetics were measured after 5 min incubation with 4 mM succinate in the absence (closed symbols) or presence of 500 μ M GDP (dark grey symbols), 2.5 μ M carboxyatractylate (light grey symbols) or GDP and carboxyatractylate (open symbols). (C), proton leak due to UCP3 measured at 161 mV (solid line (A,B)) as (A) minus (B). (D), proton leak not due to UCP3 measured at 161 mV, data from (B). (E), carboxyatractylate titration of ANT content in mitochondria from WT mice (squares) and *Ucp3*KO mice (circles). Data are means \pm S.E.M. of 10 (A-D) and 4 (E) experiments, each performed in duplicate. One-way ANOVA was performed for (C) and (D); $p = 0.0001$, with Dunnett's multiple comparison test used posthoc comparing other bars to control: *significantly different: $p < 0.05$; **, $p < 0.01$.

Figure 7 Inhibition of proton conductance in *Ucp3*WT and *Ucp3*KO mouse skeletal muscle mitochondria by carboxyatractylate and bongkrekate
(A-B), proton leak kinetics. (A) mitochondria from WT mice. (B), mitochondria from *UCP3*KO mice. 4 mM succinate was added at $t = -5$ min (see Fig. 1A) in the absence of BSA in the absence (closed symbols) or presence of 2.0 μ M bongkrekate (dark grey) or 2.5 μ M carboxyatractylate (light grey). (C), proton leak due to UCP3 measured at 158 mV (solid line (A,B)) as (A) minus (B). Data are means \pm S.E.M. of 10 experiments, each performed in duplicate. One-way ANOVA was performed for (C); $p = 0.015$, with Dunnett's multiple comparison test used posthoc comparing other bars to control; *significantly different: $p < 0.05$.

Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

