



HAL
open science

Prohormone convertases 1/3 and 2 together orchestrate the site-specific cleavages of progastrin to release gastrin-34 and gastrin-17

Jens F. Rehfeld, Xiaorong Zhu, Christina Norrbom, Jens R Bundgaard, Anders H Johnsen, John E Nielsen, Jonas Vikesaa, Jeffrey Stein, Arunangsu Dey, Donald F Steiner, et al.

► To cite this version:

Jens F. Rehfeld, Xiaorong Zhu, Christina Norrbom, Jens R Bundgaard, Anders H Johnsen, et al.. Prohormone convertases 1/3 and 2 together orchestrate the site-specific cleavages of progastrin to release gastrin-34 and gastrin-17. *Biochemical Journal*, 2008, 415 (1), pp.35-43. 10.1042/BJ20080881 . hal-00479018

HAL Id: hal-00479018

<https://hal.science/hal-00479018>

Submitted on 30 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Prohormone convertases 1/3 and 2 together orchestrate the site-specific cleavages of progastrin to release gastrin-34 and gastrin-17

Jens F. REHFELD^{*1}, Xiaorong ZHU⁺⁺, Christina NORRBOM⁺⁺, Jens R. BUNDGAARD^{*}, Anders H. JOHNSEN^{*}, John E. NIELSEN⁺, Jonas VIKESAA^{*}, Jeffrey STEIN⁺⁺, Arunangsu DEY⁺⁺, Donald F. STEINER⁺⁺, Lennart FRIIS-HANSEN^{*}

^{*}Departments of Clinical Biochemistry and ⁺Growth & Reproduction, Rigshospitalet, University of Copenhagen, Denmark, and ⁺⁺The Department of Biochemistry & Molecular Biology, and Howard Hughes Medical Institute, University of Chicago, Illinois 60637, USA

Corresponding author:

Professor Jens F. Rehfeld
Copenhagen University Hospital
Department of Clinical Biochemistry, KB-3014
9 Blegdamsvej
DK-2100 Copenhagen
Tel: +45-3545 3018
Fax: +45-3545 4640
jens.f.rehfeld@rh.regionh.dk

Abbreviations used:

Author to whom correspondence should be addressed: jens.f.rehfeld@rh.regionh.dk

SYNOPSIS

Cellular synthesis of peptide hormones requires prohormone convertases (PCs) for the endoproteolysis of prohormones. Antral G-cells synthesize most gastrin and express PC1/3, 2 and 5/6 in rat and man. But the cleavage sites in progastrin for each PC have not been determined. Therefore, we measured the concentrations of progastrin, processing intermediates and α -amidated gastrins in antral extracts from PC1/3 null-mice and compared the results to those in mice lacking PC2 and wild-type controls. The expression of PCs was examined by immunocytochemistry and *in situ* hybridisation of mouse G-cells. Finally, the *in vitro* effect of recombinant PC5/6 on progastrin and progastrin fragments containing the relevant dibasic cleavage sites was also examined. The results showed that also mouse G-cells express PC1/3, 2 and 5/6. The concentration of progastrin in PC1/3 null-mice was threefold elevated. Chromatography showed that cleavage of the R₃₆R₃₇ and R₇₃R₇₄ sites were grossly decreased. Accordingly, the concentrations of progastrin products were markedly reduced, α -amidated gastrins (-34 and -17) being 25% of normal. Lack of PC1/3 was without effect on the third dibasic site (K₅₃K₅₄), which is the only processing site for PC2. Recombinant PC5/6 did not cleave any of the dibasic processing sites in progastrin and fragments containing the relevant dibasic processing sites. The complementary cleavages of PC1/3 and 2, however, suffice to explain most of the normal endoproteolysis of progastrin. Moreover, the results show that prohormone convertases react differently to the same dibasic sequences, suggesting that additional structural factors modulate the substrate specificity.

Short title: Prohormone convertases in gastrin biosynthesis

Key words: biosynthesis, gastrin, peptide hormones, posttranslational processing, progastrin, prohormone convertases

INTRODUCTION

Gastrin is a gastrointestinal hormone that regulates gastric acid secretion and mucosal cell growth [for review, see ref. 1]. Most gastrin is synthesized in antral G-cells where progastrin through multiple steps is processed to α -amidated gastrins of different lengths (Figure 1). Of these, gastrin-34 and gastrin-17 predominate in normal tissue and circulation. Their synthesis requires cleavage of three evolutionary conserved dibasic sequences in

progastrin: R₃₆R₃₇, K₅₃K₅₄ and R₇₃R₇₄ [2-6]. Outside antrum the gastrin gene is expressed at low levels in the small intestinal mucosa, in the fetal pancreas, in pituitary corticotrophs and melanotrophs, in hypothalamo-pituitary neurons and in spermatozoa [7-11] as well as in a number of common cancers [for review, see ref. 12]. The extra-antral processing of progastrin is often cell-type specific. For instance, there is no cleavage of the K₅₃K₅₄ site in neither pituitary corticotrophs nor ileal TG-cells [13,14].

The G-cell maturation of progastrin requires several processing enzymes (Figure 1). While the involvement of tyrosyl protein sulfotransferases, carboxypeptidase E and the amidating enzyme complex (peptidylglycyl α -amidating monooxygenase (PAM)) appears well-established [15-18], the role of the individual prohormone convertase (PC) remains to be settled. Mammals express seven basic amino acid specific PCs of which PC1/3 and 2 are assumed to be involved primarily in endoproteolytic processing of neuroendocrine propeptides [for reviews, see refs. 19-21]. Both PC1/3 and PC2 have been demonstrated by immunocytochemistry in human and rat antral G-cells [22,23] and, at mRNA level, in rat antral extracts [24]. PC5/6, however, also appears to be neuroendocrine as it is expressed in gastrointestinal endocrine cells and cerebral neurons [23,25,26]. Accordingly, small amounts of PC5/6 mRNA were also detected in extracts of the rat antrum, and immunoreactive PC5/6 demonstrated in rat antral G-cells [23,24].

Recently, we showed in PC2 and 7B2 knockout mice that PC2 plays a distinct role limited to partial cleavage of the K₅₃K₅₄ site of progastrin [27]. Hence, PC2 determines the concentrations and ratio between the two primary gastrins, gastrin-34 and gastrin-17. But PC2 was without effect on any of the two di-arginyl sites of which processing of the latter, R₇₃R₇₄, is decisive for production of bioactive gastrins (Figure 1). In the present study we have examined the roles of PC1/3 and PC5/6, which became possible with the development of mice lacking PC1/3 [28], and the availability of active recombinant PC5/6. Taken together, the results should reflect the normal endoproteolytic profile of progastrin processing in antral G-cells since no other PCs are expressed in endocrine G-cells.

EXPERIMENTAL

Mice

The PC1/3 null mutant mice were generated as described earlier [28]. For the present study, tissue was obtained from 12-week-old PC1/3 null mice and wild-type (+/+) littermates of the same age. The care and treatment of all animals were in accordance with NIH and institutional guidelines.

Tissue isolation and extraction

Mice in groups of four were sacrificed by CO₂ inhalation, and stomach tissue was rapidly dissected and frozen on dry ice. The tissue was gently rinsed in phosphate-buffered saline on ice before freezing. Tissue extracts for radioimmunoassay (RIA) were prepared as previously described [29]. Briefly, frozen tissues were boiled in water (1 ml/mg) for 20 min, homogenized (polytron) and centrifuged for 30 min at 10,000 rpm. The supernatants were withdrawn and the pellets re-extracted in 0.5 M CH₃COOH (1 ml/mg), rehomogenized, incubated at room temperature for 30 min and centrifuged. The neutral water and acid supernatants were stored at -20°C until RIA analysis. The acid extracts, however, turned out to contain only negligible amounts of progastrin and its products. Consequently they were not included in further examinations.

Chromatography

One or two ml of extracts were applied to Sephadex G-50 superfine columns (10 × 1,000 mm) which were eluted at 4°C with 0.02 M barbital buffer, pH 8.4, containing 0.1% bovine serum albumin. Fractions of 1.0 ml were collected at a rate of 4.0 ml/h. The columns were calibrated with [¹²⁵I]albumin (void volume), gastrin-34 and gastrin-17, and with ²²NaCl (total volume). The eluted fractions were assayed with a library of sequence-specific radioimmunoassays against different progastrin epitopes and fragments using antibodies nos. 2604, 2605, 8017, 2145, 5284, and 3208, as detailed below. The percentages of carboxyamidated gastrin were calculated by planimetry. Four different extracts from each of the two groups of mice (PC1/3 knockout, and corresponding wild-type controls) were subjected to gel chromatography.

Radioimmunoassays

A library of six sequence-specific antisera against progastrin and its fragments was used to measure the different forms of gastrin and progastrin processing intermediates (Figure 1). The sum of carboxyamidated gastrins

was measured using the gastrin-specific antiserum no. 2604, [¹²⁵I]gastrin-17 as tracer and gastrin-17 as standard. Antibody no. 2604 binds the carboxyamidated gastrin-34 and gastrin-17 with equimolar affinity, irrespective of size and degree of tyrosyl O-sulfation [30]. The cross-reactivity with homologous cholecystokinin peptides is negligible [30,31]. Non-sulfated carboxyamidated gastrin-34 and -17 were measured in parallel using antibody no. 2605 [32]. Glycine-extended processing intermediates of progastrin were measured using antisera nos. 5284 and 3208 with [¹²⁵I]glycine-extended gastrin-17 as tracer and glycine-extended gastrin-17 as standard [33]. Glycine-arginine-extended intermediate precursors were measured using antiserum nos. 5284 and 3208 following enzymatic pretreatment with only carboxypeptidase B [33]. To measure all precursor forms of gastrin, samples were pretreated with both trypsin and carboxypeptidase B followed by radioimmunoassay measurement using antiserum no. 5284, as previously described [33]. Carboxypeptidase B mimics the effect of carboxypeptidase E, while trypsin mimics the effects of PCs. Antiserum no. 2145 was used to measure fragment 38-52 of human progastrin, corresponding to the pyroglutaminated N-terminus of gastrin-34 using [¹²⁵I]gastrin-34 as tracer and gastrin-34 as standard [34]. Correspondingly, antiserum no. 8017 was used to measure the pyroglutaminated N-terminus of human gastrin-17 using [¹²⁵I]gastrin-17 as tracer and gastrin-17 as standard [35]. Due to species differences in the N-terminal sequences of mammalian gastrin-34 and -17, it was necessary to use antisera nos. 2145 and 8017 for characterization of possible progastrin cleavage products after incubation of human progastrin with recombinant PC5/6. To monitor the purification of recombinant human progastrin, we also used a radioimmunoassay based on antiserum no. 94023 directed against the very N-terminus of human progastrin together with the corresponding N-terminal decapeptide fragment tyrosylated at its C-terminus as tracer and standard [36].

In vitro cleavage of progastrin and progastrin fragments with recombinant PC5/6A

To produce recombinant human progastrin, an expression vector of human preprogastrin, as previously described [37], was transiently transfected into CHO cells using a standard Fugene6 transfection protocol (Roche Applied Science). After 48 h, culture media was recovered, separated by gel filtration on a 1000x 50 mm Superdex G50 Superfine column and eluted, as previously described [37]. Recovered fractions were analysed for progastrin using the radioimmunoassay specific to the N-terminus of human progastrin [36]. Fractions eluting at the position of intact progastrin were pooled and subjected to an

in vitro PC5/6A cleavage assay. Half of the peptide pool was added 10 units of human PC5/6 (Phenoswitch Bioscience Inc., Sherbrooke, Canada) and reaction buffer to a concentration as suggested by Phenoswitch Bioscience (vide *intra*) and incubated for 4 hours at 37°C. A control pool was only added reaction buffer and incubated like the reaction pool. Both pools were subjected to gel filtration and analysed using specific radioimmunoassays towards the N-terminals of progastrin, gastrin-34 and gastrin-17 as well as C-terminal assay against glycine-extended gastrin [30,33,34,36].

Gastrin-34 from Sigma and the peptides GPASHHRRQLGPOGP (progastrin cleavage site for gastrin-34 N-terminus) and PEELVHRVKRRADPDKM (PC substrate [38]) were custom synthesized by Schafer-N (Copenhagen, Denmark). Incubation of the peptides were performed as suggested by PhenoSwitch in a total volume of 100 µl 20 mM Bis-Tris buffer pH 6.5, containing 1 mM CaCl₂, BSA (2 g/l), 100 µM peptide (substrate) and 10 U of PC5/6 for 2 h at 37°C. The reaction was terminated by addition of 10 µl 10% trifluoroacetic acid. The reaction mixture was separated by HPLC using a 2.1 × 150 mm C18 column (Vydac, Hisperia, USA) eluted with a gradient from 10-40% acetonitrile over 30 min, the elution buffers contained 0.1% trifluoroacetic acid. The chromatograms were monitored at 214 nm and peaks were collected manually. The peptides collected were analysed by MALDI-TOF mass spectrometry (AutoFlex II, Bruker, Bremen, Germany) using α-cyano-4 hydroxycinnamic acid as matrix.

Immunocytochemistry

To examine the cellular co-localization between gastrin and PC1/3 or PC2, the stomach was removed from three wild-type mice aged 3-4 months, cut open, washed, and immersion fixed overnight in 4% paraformaldehyde. Six µm traverse paraffin sections were cut, mounted on superfrosted glass slides and baked overnight at 60°C. Antigen retrieval was performed by boiling the slides in a microwave oven for 3 minutes at 800W, followed by 3 times 5 minutes at 400W in a 10mM Tris-buffered EGTA solution at pH 9.0. Preblocking was performed in 10% ovine serum in TBS for 20 minutes. The sections were incubated with PC1/3 or PC2 antisera diluted 1:10,000 overnight at 4°C [39,40]. The primary antiserum was detected with Envision+ (Dako, Glostrup, Denmark) for 1 hr followed by incubation with a Cy3 conjugated anti-horseradish peroxidase antibody (Jackson, West Grove, Pennsylvania, USA) diluted 1:400 for 1 hr. Gastrin was detected with overnight incubation with antiserum no.

8007 (dilution 1:1000) at 4°C and subsequently reacted to Alexa 488 conjugated donkey-anti rabbit F(ab)-fragments (dilution 1:1000) (Molecular Probes, Inc., Eugene, Oregon, USA). Antiserum no. 8007 is specific to the carboxyamidated C-terminus of the gastrins [31]. Cover slips were mounted with Aquamount (Dako). Analysis of the sections was performed with Zeiss LSM510 confocal microscope (Zeiss, Oberkochen, Germany).

In situ hybridisation

Serial sections were probed to the PC5/6 antisense probe, gastrin antiserum no. 8007 [31] and PC5/6 sense probe. *In situ* hybridisation was performed as previously described [41]. In brief, mouse stomach was fixed in 4% paraformaldehyde, 8 µm sections were placed on SuperFrostRPlus slides (Menzel GmbH, Braunschweig Germany), deparaffinised and PFA (4%) fixed, and treated with proteinase K (P2308; Sigma, St. Louis, Missouri, USA), postfixed (4% PFA), prehybridised for 1h at 50°C, and incubated over night with the labelled PC5/6 probe (50°C), specific to nucleotide. Visualization was made with anti-digoxigenin-AP (Roche Diagnostics, Mannheim, Germany) and chromogens BCIP (Sigma B-8503) and NBT (Sigma N-6876).

RESULTS AND DISCUSSION

Progastrin processing in PC1/3 and PC2 knockout mice

As shown in Table 1 and Figure 2, the antral concentrations of progastrin and its products are grossly altered in PC1/3 knockout mice. The concentration of progastrin itself is increased nearly three-fold, whereas concentrations of the processing intermediates are reduced to one third, and the bioactive, carboxyamidated gastrins are reduced to one fourth of the concentrations in wild-type controls. The reductions contrast the earlier findings in PC2 knockout mice and the carboxypeptidase E-deficient (CPE^{fat}) mice, in which the antral concentrations of the bioactive, carboxyamidated gastrins are maintained at a normal level in spite of major disturbances at earlier steps in the processing pathway and in the ratio between gastrin-34 and gastrin-17 (Table 1, Figure 2 and [18,27]).

Gel chromatography of neutral antral extracts revealed an unchanged molecular pattern of carboxyamidated gastrins in the PC1/3 knockout mice, despite the marked difference in tissue concentrations (Figure 3). Hence, the ratios between gastrin-34 and gastrin-17, and between tyrosyl-sulfated and non-sulfated gastrins were as in the wild-type controls (Figure 3). The low concentrations ex-

cluded chromatographic examination of the processing intermediates, but it was possible to study the progastrin immunoreactivity, which showed a significant difference between the null mice and their wild-type littermates (Figure 4). Thus, the progastrin molecule is only partly cleaved at the dibasic R₃₆R₃₇ site and even less at the K₅₃K₅₄ site in the PC1/3 null mice (Figure 4 upper panel), while progastrin in the controls is almost completely processed to shorter fragments (Figure 4 lower panel). The modest differences in antral concentrations of progastrin and its processing products between the three groups of wild-type controls (Table 1) supposedly reflect differences in mouse strains and age. Hence, the mice used in the present study are younger (3 months of age) than those used in the studies of carboxypeptidase E-deficiency and PC2 knockout that were up to 6 months of age [18,27]. As shown before, the antral concentration of progastrin declines with age in rodents [42].

The processing observed in PC1/3 knockout mice indicates that PC1/3 initiates cleavage at the N-terminal di-arginyl site (R₃₆R₃₇) at an early stage in the processing, since more progastrin and a large fragment corresponding to sequence 38-83 of progastrin were found in the antral extract of PC1/3 null mice (Figure 4). Subsequently, PC1/3 continues to cleave the C-terminal di-arginyl site R₇₃R₇₄. PC1/3, however, appears to be without effect on the di-lysyl site, K₅₃K₅₄ (Figure 3). This endoproteolytic cleavage pattern of progastrin is opposite to that of PC2 (Table 1, Figure 2 and [27]). Thus, PC2 only cleaves the single di-lysyl site of progastrin, K₅₃K₅₄, the cleavage for half of which PC2 appears to be responsible [27]. The results hence support the contention that PC1/3 generally prefers arginine containing cleavage sites, while PC2 is activated later in the processing pathway to cleave mainly lysine containing dibasic sites at the lower pH characteristic of mature secretory granules [20].

The present data support our earlier observation that pituitary corticotrophs which contain only PC1/3 and no PC2 cannot process the K₅₃K₅₄ site of progastrin. Therefore, corticotrophs synthesize gastrin-34, but no gastrin-17 [9,10,13]. This notion is also supported by Sawada et al. who found that PC2 transfected into AtT-20 cells but not PC1/3 can accomplish cleavage of the di-lysyl site in progastrin [43]. Also proopiomelanocortin (POMC) contains a KK site within β -endorphin which remains unprocessed in AtT-20 cells but can be processed upon cotransfection with PC2 [44,45]. However, cleavage of the KK site within proenkephalin is apparently largely mediated by PC1/3 since it is efficiently cleaved in AtT-20 cells [46]. The primary structure of the three precursors (progastrin, POMC and proenkephalin) re-

veals extensive differences in the vicinity of the KK sites, which could account for the differential ability of PC1/3 to recognize this type of processing site within different contexts.

A surprising finding in the present study was the increased concentration of progastrin in the antrum of PC1/3 null mice, while the concentrations of the processing products were drastically reduced (Table 1 and Figure 2). It is obvious that the lack of PC1/3 reduces the processing of progastrin severely. But it is an unsolved question why the total translational product in PC1/3 null mice is reduced, whereas it is increased in other processing enzyme deficiencies [18]. The C-terminal dibasic cleavage sites in progastrin are, however, part of the motif for sorting to the regulated secretory pathway [47]. Moreover, PC1/3 has been shown to enhance the post-Golgi sorting of some prohormones [48]. It is therefore possible that the lack of PC1/3 per se reduces the storage of progastrin and its products in the secretory granules of G-cells.

Progastrin cleavage by PC5/6

Since half of the KK site within progastrin is cleaved in the absence of PC2 [27], and since 30% of the two di-arginyl sites are cleaved in the absence of PC1/3 (Table 1), other convertases might participate in the *in vivo* endoproteolytic processing of progastrin in normal G-cells. A candidate for this activity is PC5/6, which has been shown to be expressed in endocrine cells of the gastrointestinal tract [25,49]. As shown here and in the rat, PC5/6 gene expression can be demonstrated in antral G-cells (Figure 7 and [23]). PC5/6 is known to cleave basic processing sites [19,20]. In order to examine possible effects of PC5/6, we first looked at the processing of progastrin in SK-N-MC cell line, since PC5/6 knockout mice were not available, and since SK-N-MC cells have been claimed to express the PC5A isoform but neither PC1/3 nor PC2 [50]. Resent re-examinations, however, have shown that SK-N-MC cells may indeed express PC1/3 at protein level [Bundgaard J, unpublished results] and also PC2, although without its chaperone 7B2 [Lindberg I, personal communication]. Consequently, we decided to exclude the results obtained of progastrin processing in this cell-line.

Instead we examined directly *in vitro* the substrate activity of PC5/6 against intact progastrin. Human progastrin was expressed in CHO cells, which do not cleave progastrin. Progastrin was subsequently purified by size exclusion gel chromatography, and recovered peptide was incubated *in vitro* with recombinant PC5/6A. Prod-

ucts were subjected to repeated chromatography and analysed using radioimmunoassays that are specific to the dibasic cleavage sites of human progastrin [34,35]. Enzyme treated progastrin eluted identically to untreated control fractions (Figure 5), and the site-specific radioimmunoassays did not detect processing at any of the cleavage sites. Moreover, digestion with recombinant PC5/6 of gastrin-34 and the synthetic peptide spanning the N-terminus of gastrin-34 within progastrin resulted in no cleavage, while the general PC substrate [38] used as control was cleaved as expected. Consequently, we conclude that PC5/6 is without effect on progastrin, and that the processing of progastrin originally observed in S-K-N-MC cells is due to other endoproteases than PC5/6.

Prohormone convertase expression in G-cells

As shown in Figure 6, both PC1/3 and PC2 are expressed in normal antral G-cells in mice. The immunocytochemical results are in accordance with earlier results in human and rat antrum [22,23]. In addition, as shown here by *in situ* hybridisation (Figure 7), also PC5/6 is expressed in murine antral G-cells.

The interplay of PC1/3 and PC2 and the role of PC5/6 in G-cells

Since no cleavage was found in the *in vitro* studies using PC5/6 against intact progastrin and progastrin fragments containing the dibasic cleavage sites, whereas the same PC5/6 was fully active in cleaving a control peptide. PC5/6 hardly plays a significant role in the G-cell maturation of progastrin. It is possible that PC5/6 is instead involved in the processing of other neuroendocrine G-cell products such as chromogranin A and neuropeptide W [51,53]. Thus, the endoproteolytic maturation of progastrin as seen in normal G-cells appears to require an interplay, primarily between PC1/3 and PC2. We believe that the processing begins with PC1/3, which is solely responsible for the cleavage of R₃₆R₃₇. But, subsequently, PC1/3 cleaves the crucial R₇₃R₇₄. Later, in secretory granules, PC2 performs the partial cleavage of K₅₃K₅₄ to ensure the production of gastrin-17.

ACKNOWLEDGEMENTS

The skilful technical and secretarial assistance of Alice von der Lieth, Rikke Kröncke, Allan Kastrup, and Diana Skovgaard is gratefully acknowledged. The study was supported by grants from the Danish Biotechnology Program for Peptide Research; NIH grants DK-13914, DK-20595, and the Howard Hughes Medical Institute.

REFERENCES

- 1 Rehfeld, J.F. (1998) The new biology of gastrointestinal hormones. *Physiol. Rev.* **78**, 1087-1108
- 2 Yoo, O.J., Powell, C.T., and Agarwal, K.L. (1982) Molecular cloning and nucleotide sequence of full-length of cDNA coding for porcine gastrin. *Proc. Natl. Acad. Sci. USA* **79**, 1049-1053
- 3 Boel, E., Vuust, J., Norris, F., Norris, K., Wind, A., Rehfeld, J.F., and Marcker, K.A. (1983) Molecular cloning of human gastrin cDNA: evidence for evolution of gastrin by gene duplication. *Proc. Natl. Acad. Sci. USA* **80**, 2866-2869
- 4 Gregory, R.A., and Tracy, H.J. (1964) The constitution and properties of two gastrins extracted from hog antral mucosa. *Gut* **5**, 103-117
- 5 Gregory, R.A., and Tracy, H.J. (1972) Isolation of two "big gastrins" from Zollinger-Ellison tumour tissue. *Lancet* **2**, 797-799
- 6 Rehfeld, J.F., and Johnsen, A.H. (1994) Identification of gastrin component I as gastrin-71. The largest possible bioactive progastrin product. *Eur. J. Biochem.* **223**, 765-773
- 7 Larsson, L.I., and Rehfeld, J.F. (1979) A peptide resembling the COOH-terminal tetrapeptide amide of gastrin from a new gastrointestinal endocrine cell type. *Nature* **277**, 575-578
- 8 Larsson, L.I., Rehfeld, J.F., Sundler, F., and Håkanson, R. (1976) Pancreatic gastrin in foetal and neonatal rats. *Nature* **262**, 609-610
- 9 Rehfeld, J.F. (1978) Localisation of gastrins to neuro- and adenohypophysis. *Nature* **271**, 771-773
- 10 Rehfeld, J.F., and Larsson, L.I. (1981) Pituitary gastrin: Different processing in corticotrophs and melanotrophs. *J. Biol. Chem.* **256**, 10426-10429
- 11 Schalling, M., Persson, H., Pelto-Huikko, M., Ødum, L., Ekman, P., Gottlieb, C., Hökfelt, T., and Rehfeld, J.F. (1990) Expression and localization of gastrin messenger RNA and peptide in spermatogenic cells. *J. Clin. Invest.* **86**, 660-669
- 12 Rehfeld, J.F., and van Solinge, W.W. (1994) The tumor biology of gastrin and cholecystokinin. *Adv. Cancer Res.* **63**, 295-347
- 13 Rehfeld, J.F. (1986) Accumulation of nonamidated preprogastrin and preprocholecystokinin products in porcine pituitary corticotrophs: Evidence of post-translational control of cell differentiation. *J. Biol. Chem.* **261**, 5841-5847
- 14 Friis-Hansen, L., and Rehfeld, J.F. (1994) Ileal expression of gastrin and cholecystokinin: In search of a related hormone. *FEBS Lett.* **343**, 115-119
- 15 Hilsted, L., and Rehfeld, J.F. (1987) Alpha-carboxyamidation of antral progastrin: Relation to other post-translational modifications. *J. Biol. Chem.* **262**, 16953-16957
- 16 Eipper, B.A., Milgram, S.L., Husten, E.J., Yun, H.Y., and Mains, R.E. (1993) Peptidylglycine alpha-amidation monooxygenase: a multifunctional protein

- with catalytic, processing, and routing domains. *Protein Sci.* **2**, 489-497
- 17 Bundgaard, J.R., Vuust, J., and Rehfeld, J.F. (1995) Tyrosine O-sulfation promotes proteolytic processing of progastrin. *EMBO J.* **14**, 3073-3079
 - 18 Lacourse, K.A., Friis-Hansen, L., Rehfeld, J.F., and Samuelson, L.C. (1997) Disturbed progastrin processing in carboxypeptidase E-deficient fat mice. *FEBS Lett.* **416**, 45-50
 - 19 Zhou, A., Webb, G., Zhu, X., and Steiner, D.F. (1999) Proteolytic processing in the secretory pathway. *J. Biol. Chem.* **274**, 20745-20748
 - 20 Müller, L., and Lindberg, I. (2000) The cell biology of the prohormone convertases PC1 and PC2. *Prog. Nucleic. Acid. Res. Mol. Biol.* **63**, 69-108
 - 21 Scamuffa, N., Calvo, F., Chretien, M., Seidah, N.G. and Khatib, A.-M. (2006) Proprotein convertases: Lessons from knockouts. *FASEB J.* **20**, 1954-1963
 - 22 Scopsi, L., Gullo, M., Rilke, F., Martin, S., and Steiner, D.F. (1995) Proprotein convertases (PC1/PC3 and PC2) in normal and neoplastic human tissues: Their use as markers of neuroendocrine differentiation. *J. Clin. Endocrinol. Metab.* **80**, 294-301
 - 23 Macro, J.A., Bate, G.W., Varro, A., Vaillant, C., Seidah, N.G., Dimaline, R., and Dockray, G.J. (1997) Regulation by gastric acid of the processing of progastrin-derived peptides in rat antral mucosa. *J. Physiol.* **502**, 409-419
 - 24 Macro, J.A., Dimaline, R., and Dockray, G.J. (1996) Identification and expression of prohormone-converting enzymes in the rat stomach. *Am. J. Physiol.* **270**, G87-G93
 - 25 Lusson, J., Vieau, D., Hamelin, J., Day, R., Chretien, M., and Seidah, N.G. (1993) cDNA structure of the mouse and rat subtilisin/kexin-like PC5: a candidate proprotein convertase expressed in endocrine and nonendocrine cells. *Proc. Natl. Acad. Sci. USA* **90**, 6691-6695
 - 26 Villeneuve, P., Seidah, N.G., and Beaudet, A. J. (1999) Immunohistochemical distribution of the prohormone convertase PC5-A in rat brain. *Neurosci.* **92**, 641-654
 - 27 Rehfeld, J.F., Lindberg, I., and Friis-Hansen, L. (2002) Progastrin processing differs in 7B2 and PC2 knockout animals : a role for 7B2 independent of action on PC2. *FEBS Lett.* **510**, 89-93
 - 28 Zhu, X., Zhou, A., Dey, A., Norrbom, C., Caroll, R., Zhang, C., Laurent, V., Lindberg, I., Ugleholdt, R., Holst, J.J., and Steiner, D.F. (2002) Disruption of PC1/3 expression in mice causes dwarfism and multiple neuroendocrine peptide processing defects. *Proc. Natl. Acad. Sci. USA* **99**, 10293-10298
 - 29 Rehfeld, J.F., Bardram, L., and Hilsted, L. (1992) Ontogeny of procholecystokinin maturation in rat duodenum, jejunum, and ileum. *Gastroenterology* **103**, 424-430
 - 30 Rehfeld, J.F., Stadil, F., and Rubin, B. (1972) Production and evaluation of antibodies for the radioimmunoassay of gastrin. *Scand. J. Clin. Lab. Invest.* **30**, 221-232
 - 31 Rehfeld, J.F. (1998) Accurate measurement of cholecystokinin in plasma. *Clin. Chem.* **44**, 991-1001
 - 32 Rehfeld, J.F., de Magistris, L., and Andersen, B.N. (1981) Sulfation of gastrin : effect on immunoreactivity. *Regul. Pept.* **2**, 333-342
 - 33 Hilsted, L., and Rehfeld, J.F. (1986) Measurement of precursors for alpha-amidated hormones by radioimmunoassay of glycine-extended peptides after trypsin-carboxypeptidase B cleavage. *Anal. Biochem.* **152**, 119-126
 - 34 van Solinge, W.W., and Rehfeld, J.F. (1990) Radioimmunoassay for sequence 38-54 of human progastrin: increased diagnostic specificity of gastrin-cell diseases. *Clin. Chim. Acta* **192**, 35-46
 - 35 Bardram, L., and Rehfeld, J.F. (1988) Processing-independent radioimmunoanalysis: a general analytical principle applied to progastrin and its products. *Anal. Biochem.* **175**, 537-543
 - 36 Goetze, J.P., Hansen, C.P., Rehfeld, J.F. (2006) Antral content, secretion and metabolism of N-terminal progastrin fragments. *Regul. Peptides.* **133**, 47-53
 - 37 Bundgaard, J.R., Cowland, J.B., Vuust, J., Rehfeld, J.F. (1996) An efficient cellular system for mutational analysis of prohormone processing. *DNA Cell. Biol.* **15**, 147-57
 - 38 Spour, N., Lebel, A., McMahon, S., Fournier, I., Fugère, M., Day, R., and Dubois, C.M. (2003) TACE/ADAM-17 maturation and activation of sheddase activity require proprotein convertase activity. *FEBS Lett.* **555**, 275-283
 - 39 Shen, F.S., Seidah, N.G., and Lindberg, I. (1993) Biosynthesis of the prohormone convertase PC2 in Chinese hamster ovary cells and in rat insulinoma cells. *J. Biol. Chem.* **268**, 24910-24915
 - 40 Hornby, P.J., Rosenthal, S.D., Mathis, J.P., Vindrola, O., and Lindberg, I. (1993) Immunocytochemical localization of the neuropeptide-synthesizing enzyme PC1 in AtT-20 cells. *Neuroendocrinol.* **58**, 555-563
 - 41 Nielsen, J.E., Hansen, M.A., Jørgensen, M., Tarnaka, M., Almstrup, K., Skakkebaek, N.E., and Leffers, H. (2003) Germ cell differentiation-dependent and stage-specific expression of LANCL1 in rodent testis. *Eur. J. Histochem.* **47**, 215-222
 - 42 Hilsted, L., Bardram, L., and Rehfeld, J.F. (1988) Progastrin maturation during ontogenesis. Accumulation of glycine-extended gastrins in rat antrum at weaning. *Biochem. J.* **255**, 397-402
 - 43 Sawada, M., Finniss, S., and Dickinson, C.J. (2000) Diminished prohormone convertase 3 expression (PC1/PC3) inhibits progastrin post-translational processing. *Regul. Pept.* **19**, 19-28
 - 44 Zhou, A., Bloomquist, B.T., and Mains, R.E. (1993) The prohormone convertases PC1 and PC2 mediate distinct endoproteolytic cleavages in a strict temporal order during proopiomelanocortin biosynthetic processing. *J. Biol. Chem.* **268**, 1763-1769

- 45 Zhou, A., and Mains, R.E. (1994) Endoproteolytic processing of proopiomelanocortin and prohormone convertases 1 and 2 in neuroendocrine cells overexpressing prohormone convertases 1 or 2. *J. Biol. Chem.* **269**, 17440-17447
- 46 Mathis, J.P., and Lindberg, I. (1992) Posttranslational processing of proenkephalin in AtT-20 cells: evidence for cleavage at a Lys-Lys site. *Endocrinology* **131**, 2287-2296
- 47 Bundgaard, J.R., Birkedal, H., and Rehfeld, J.F. (2004) Progastrin is directed to the regulated secretory pathway by synergistically acting basic and acidic motifs. *J. Biol. Chem.* **279**, 5488-5493
- 48 Mulcahy, L.R., Vaslet, C.A., and Nilni, E.A. (2005) Prohormone-convertase 1 processing enhances post-Golgi sorting of prothyrotropin-releasing hormone-derived peptides. *J. Biol. Chem.* **280**, 39818-39826
- 49 Nakagawa, T., Hosaka, M., Torii, S., Watanabe, T., Murakami, K., and Nakayama, K. (1993) Identification and functional expression of a new member of the mammalian Kex2-like processing endoprotease family: its striking structural similarity to PACE4. *J. Biochem. Tokyo* **113**, 132-135
- 50 Cain, B.M., Vishnuvardhan, D., and Beinfeld, M.C. (2001) Neuronal cell lines expressing PC5, but not PC1 or PC2, process Pro-CCK into glycine-extended CCK 12 and 22. *Peptides* **22**, 1271-1277
- 51 Helle, K.B. (2004) The granin family of uniquely acidic proteins of the diffuse neuroendocrine system: Comparative and functional aspects. *Biol. Rev.* **79**, 769-794
- 52 Tanaka, H., Yoshida, T., Mayamoto, N., Motoike, T., Kurosu, H., Shibata, K., Yamanaka, A., Williams, S.C., Richardson, J.A., Tsujino, N., Garry, M.G., Lerner, M.R., King, D.S., O'Dowd, B.F., Sakurai, T., Yanagisawa, M. (2003) Characterization of a family of endogenous neuropeptide ligands for the G protein-coupled receptors, GPR7 and GPR8. *Proc. Natl. Acad. Sci. USA.* **100**, 6251-6256
- 53 Mondal, M.S., Yamaguchi, H., Date, Y., Toshinai, K., Kawagoe, T., Tsuruta, T., Kageyama, H., Kawamura, Y., Shioda, S., Shimomura, Y., Mori, M., Nakazato, M. (2006) Neuropeptide W is present in antral G cells of rat, mouse and human stomach. *J. Endocrinol.* **29**, 61-72

Table 1 Progastrin and its products in antral mucosa of PC1/3 and PC2* null mice, carboxypeptidase E deficient mice** and corresponding wild-type mice (pmol/g tissue; mean \pm SEM)

	PC1/3 knockout (n = 4)	Wild-type (n = 4)
Progastrin	98.0 \pm 49.6	35.8 \pm 7.3
Gastrin-Gly-Arg	0.5 \pm 0.3	1.0 \pm 0.6
Gastrin-Gly	3.5 \pm 1.8	10.3 \pm 1.8
Gastrin-amide	145.5 \pm 65.0	568.3 \pm 135.6
	PC2 knockout (n = 8)	Wild-type (n = 8)
Progastrin	21.0 \pm 6.5	14.0 \pm 3.5
Gastrin-Gly-Arg	0.0 \pm 0.0	0.5 \pm 0.2
Gastrin-Gly	6.5 \pm 1.5	5.5 \pm 1.5
Gastrin-amide	750.0 \pm 140.5	650.5 \pm 131.0
	CPE-deficiency (n = 8)	Wild-type (n = 8)
Progastrin	0.0 \pm 0.0	11.6 \pm 5.2
Gastrin-Gly-Arg	1054.0 \pm 122.0	12.2 \pm 6.3
Gastrin-Gly	34.7 \pm 7.1	4.8 \pm 0.5
Gastrin-amide	601.0 \pm 28.0	715.0 \pm 43.0

* data from ref. 27; ** data from ref. 18

Table 2 The site-specific cleavages at dibasic sequences in progastrin by prohormone convertases (PCs) expressed in antral G-cells

Prohormone Convertase	R ₃₆ R ₃₇	K ₅₃ K ₅₄	R ₇₃ R ₇₄
PC1/3	++	-	++
PC2	-	+	-
PC5/6	-	-	-

LEGENDS TO FIGURES

Figure 1 (A) The mouse progastrin sequence

is given in single letter code with the three dibasic cleavage sites printed in bold and asterisk indicating the single tyrosyl-sulfation site. The bioactive tetrapeptide sequence in the gastrins is boxed.

Figure 1 (B) Diagram of the co- and posttranslational modification of progastrin in antral G-cells

Activation of the gastrin amidation site (-Phe71-Gly72-Arg73-Arg74-) occurs via a series of carboxyterminal cleavages and modifications. Endoproteolytic cleavage by prohormone convertases (PCs) produces the carboxypeptidase E substrate (-Phe-Gly-Arg). Carboxypeptidase E then acts in secretory granules to remove the C-terminal arginyl residue yielding glycine-extended gastrin (-Phe-Gly). Carboxyamidation by PAM subsequently results in the production of bioactive gastrin (-Phe-NH₂). Concomitant N-terminal cleavage by PCs produces bioactive gastrins of varying size (e.g. gastrin-34 and gastrin-17). The indications of PC1/3, PC2 and PC5/6 in the diagram is based on the results of the present and an earlier study [27]. A library of sequence-specific antibodies was used in combination with *in vitro* protease treatment to measure bioactive gastrin and various precursor peptides, as described in the text.

Figure 2 Average percentages of the processing parameters measured in antral extracts of PC1/3 knockout mice

The parameters (progastrin, gastrin-Gly-Arg, gastrin-Gly and bioactive gastrin-amide) are compared with those measured in PC2 knockout mice and in carboxypeptidase E deficient (CPE) mice [19,28], as well as with their respective wild-type controls (broken line at 100%).

Figure 3. Gel chromatography on Sephadex G-50 superfine columns of antral extracts from PC1/3 knockout mice (left panels) and from corresponding wild-type control mice (WT, right panels)

The chromatographic elutions were monitored using antibody 2604 that binds carboxyamidated gastrins (the bioactive gastrins), irrespective of the degree of tyrosyl-sulfation (upper panels) and antibody 2605 that binds only non-sulfated (ns) gastrin (lower panels). Antral extracts from four different mice in each of the two groups

(PC1/3 and controls) were subjected to chromatography. Extracts containing similar amounts of gastrin were applied to the columns. Those shown above are characteristic of each group.

Figure 4. Gel chromatography on Sephadex G-50 superfine columns of antral extracts from PC1/3 knockout mice (KO, upper panel) and littermate wild-type controls (WT, lower panel)

The chromatographic elutions were monitored using antibody 5284 that binds glycine-extended gastrins after trypsin and carboxypeptidase B treatment of each fraction. Antral extracts from four different mice in each of the two groups (PC1/3 and controls) were subjected to chromatography. Those shown are characteristic of each group.

Figure 5 Gel chromatography on Sephadex G-50 superfine columns of human progastrin before and after incubation with recombinant PC5/6

The chromatographic elution was monitored by a radioimmunoassay specific to the intact N-terminus of human progastrin using antiserum No. 94023 (ref. 36). The upper panel shows the elution without prior treatment with PC5/6. The lower panel shows the elution after treatment with PC5/6, which has been shown to cleave a relevant control peptide.


Figure 6 Gastrin and PC1/3 co-localization

Gastrin (green) co-localizes with PC1/3 (red) (Panel A). Gastrin (green) also co-localizes with PC2 (red) (Panel B). In contrast to PC1/3, which is mainly found in the antral G-cells, many other antral endocrine cells also express PC2 (Panel B). (Spacebar 50 μ m). Gastrin was identified with antibody no. 8007.

Figure 7 Gastrin and PC5/6 co-localization

Serial sections were examined for PC5/6 expression by *in situ* hybridisation (Panel A) and gastrin expression by immunohistochemistry (Panel B). Panel C shows the PC5/6 *in situ* using the sense probe. The black arrows point to cells that express both PC5/6 and gastrin (Panel A and B). The white arrow identifies a cell expressing PC5/6 but not gastrin (Panel A and B). (Spacebar 50 μ m). Gastrin was identified using antibody no. 8007

Figure 1



THIS IS NOT THE FINAL VERSION - see doi:10.1042/BJ20080881

Figure 2






Figure 3



THIS IS NOT THE FINAL VERSION - see doi:10.1042/BJ20080881


Figure 4

PC 1/3



THIS IS NOT THE FINAL VERSION - see doi:10.1042/BJ20080881

Figure 5



THIS IS NOT THE FINAL VERSION - see doi:10.1042/BJ20080881

Figure 6
G-CELLS





Figure 7
G-CELLS



THIS IS NOT THE FINAL VERSION - see doi:10.1042/BJ20080881