



HAL
open science

Active site substitutions delineate distinct classes of eubacterial flap endonuclease

Lee M Allen, Michael R G Hodkinson, Jon R Sayers

► **To cite this version:**

Lee M Allen, Michael R G Hodkinson, Jon R Sayers. Active site substitutions delineate distinct classes of eubacterial flap endonuclease. *Biochemical Journal*, 2009, 418 (2), pp.285-292. 10.1042/BJ20081637 . hal-00479083

HAL Id: hal-00479083

<https://hal.science/hal-00479083>

Submitted on 30 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Active site substitutions delineate distinct classes of eubacterial flap endonuclease

Lee M. Allen¹, Michael R. G. Hodkinson¹ and Jon R. Sayers*

The University of Sheffield School of Medicine and Biomedical Sciences, Henry Wellcome Laboratories for Medical Research, Section of Infection, Inflammation and Immunity, Sheffield, UK S10 2RX

*Corresponding Author: Phone: +44 (0)114 271 2327 Fax: +44 (0)114 273 9926
Email: j.r.sayers@shef.ac.uk

¹ These authors contributed equally to this work.

ABSTRACT

Flap endonucleases (FENs) play essential roles in DNA replication, pivotally in the resolution of Okazaki fragments. In eubacteria, DNA Polymerase I (Poll) contains a flap processing domain, the N-terminal 5' → 3' exonuclease. We present evidence of paralogous FEN-encoding genes present in many eubacteria. Two distinct classes of these independent FEN-encoding genes exist with four groups of eubacteria being identified based on the number and type of FEN gene encoded. The respective proteins possess distinct motifs hallmarking their differentiation. Crucially, based on primary sequence and predicted secondary structural motifs, we reveal key differences at their active site. These data are supported by biochemical characterization of two family members - exonuclease IX (ExoIX) from *Escherichia coli* and *Staphylococcus aureus* flap endonuclease (SaFEN). These proteins displayed marked differences in their ability to process a range of branched and linear DNA structures. On bifurcated substrates, SaFEN exhibited similar substrate specificity to previously characterised FENs. In quantitative exonuclease assays, SaFEN maintained a comparable activity to that reported for Poll. However, ExoIX showed no observable enzymatic activity. A threaded model is presented for SaFEN, demonstrating the probable interaction of this newly identified class of flap endonuclease with divalent metal ions and a branched DNA substrate. These data provide an intriguing model for the cellular role of these FEN sub-classes and illustrate the evolutionary importance of processing aberrant DNA, which has led to their maintenance alongside DNA Poll in many eubacteria.

SHORT (PAGE HEADING) TITLE: Eubacterial flap endonucleases

KEYWORDS

Exonuclease IX, FEN, DNA polymerase I, Replication, metallonuclease.

ABBREVIATIONS

Exonuclease IX, ExoIX; Flap endonuclease, FEN; DNA polymerase I, Poll.

ACKNOWLEDGEMENTS

We acknowledge the BBSRC for funding to JRS (ref. no. B19466) and studentships for LMA (ref. no. 99B1B05418) and MRGH (ref. no. 02B1B08371). We thank Prof. Peter Artymiuk (University of Sheffield) for helpful advice on model building.

INTRODUCTION

Classical biochemistry conducted by Kornberg and co-workers revealed that *Escherichia coli* DNA Polymerase I (Poll) contained two readily separable enzymatic activities within the same polypeptide [1, 2]. The C-terminal 'Klenow' fragment exhibited DNA polymerase and 3' → 5' 'proofreading' activity, while the N-terminal domain accounted for 5' → 3' exonuclease activity. The 5' → 3' exonuclease domain of Poll falls within of a wider class of essential metallonucleases, the flap endonuclease (FEN) family [3]. FENs demonstrate both 5' → 3' exonuclease and structure-specific endonuclease activity, playing important roles in DNA replication, repair and recombination [4-6]. These enzymes are ubiquitous and highly conserved in all three kingdoms of life (Eubacteria, Archaea and Eukaryota), presumably due to their role in these crucial biological processes.

The importance and biological similarity of FENs has been demonstrated in several cross-species studies. In bacteriophage T4, disruption of the *rnh* gene (which encodes a FEN, known historically as T4 RNase H) resulted in slower, less accurate DNA replication. Furthermore, no measurable progeny were recovered from a host carrying a deletion in the Poll 5' → 3' exonuclease domain [7], clearly implicating the essential requirement of an active FEN for viral replication. The lethal phenotype of $\Delta rad27$ (encoding another FEN homologue) in *Saccharomyces cerevisiae* was suppressed by the Poll 5' → 3' exonuclease domain of *E. coli* [8]. In higher organisms, heterozygous (*Fen1*^{+/-}) mice rapidly developed lymphoma and thymic abnormalities, while homozygous knockouts (*Fen1*^{-/-}) displayed the embryonic lethal phenotype [9]. Moreover, mice with a single point mutation (E160D) in one of the two divalent metal ion binding sites present in the FEN enzyme were reported to be predisposed to autoimmunity, chronic inflammation and cancers [10].

FENs recognise substrates with a DNA duplex and a single-stranded 5' overhang such as might arise during strand displacement synthesis. The principle site of hydrolytic phosphodiester backbone cleavage is one nucleotide into the DNA duplex, at the bifurcation of single and double stranded DNA [11]. In addition, FENs that play a role in nucleotide excision repair (e.g. XPG in humans) can cleave 'bubble' substrates that have no free 5' end at the bifurcation of single and double-stranded DNA. As exonucleases, FENs cleave free 5' termini and nicked DNA duplexes liberating mono-, di- and short poly-nucleotides [12, 13]. Both endo- and exonucleolytic activities are generally accepted to play a role in Okazaki fragment processing during DNA replication [6].

A number of FEN structures have been elucidated using X-ray crystallography. The structures show striking similarities, with a core β -sheet adorned with a number of surrounding α -helices. The FEN domain of *Thermus aquaticus* DNA Poll, T4RNaseH and T5FEN all reveal the presence of at least two divalent metal ion binding sites [14-16]. Experimental evidence based on isothermal titration calorimetry and mutagenesis studies support the presence of two metal binding sites [17, 18]. Metal site I is required for generation of an attacking hydroxyl nucleophile during phosphodiester backbone hydrolysis [19] and metal site II is required for substrate binding, particularly during exonucleolytic cleavage [17]. In T5FEN a key structural motif is formed from α -helices bridging above the active site and has been described as a 'helical arch' [15] or 'helical clamp' [20, 21]. This feature has been observed in a number of enzymes, and based on biochemical studies, has been proposed to play a role in substrate binding [20-23]. Most recently, a 3 Å X-ray diffraction structure of T4 RNase H in complex with a forked DNA substrate was solved [24]. These data were

in close agreement with models of FEN-substrate interaction previously proposed from site directed mutagenesis and substrate binding studies [25, 26].

An early interrogation of the incomplete *E. coli* genome sequence identified a putative gene (later termed *xni* [27].), whose product shared significant homology to Poll 5'→3' exonuclease domain and other known flap endonucleases [4, 28]. Shafritz and coworkers reported the purification and characterization of the product of the *xni* gene [27]. Surprisingly, given the significant homology to a family of enzymes demonstrating 5'→3' exonuclease activity the authors suggested that the *xni* product was a 3'→5' exonuclease and designated the *xni* gene product exonuclease IX (ExoIX) [27]. Recently we have shown that exonuclease III (a 3'→5' exonuclease) co-purifies with ExoIX over a multiple-stage purification process [29]. Therefore, we suggested that the 3'→5' exonuclease activity described by Shafritz *et al* in ExoIX was due to co-purifying exonuclease III.

Here we show that a number of prokaryotes encode a second FEN paralogue, in addition to that present as an N-terminal domain of Poll. Based on amino acid sequence analysis, these single-domain eubacterial FEN homologues can be delineated into two distinct groups. One group appears to possess binding sites for two divalent metals ions, as observed in all FENs characterized to date. The second group, exemplified by *E. coli* ExoIX, seems to lack a number of normally conserved amino acid residues that act as metal-binding ligands. We demonstrate that, in contrast to ExoIX which apparently lacks activities expected of a FEN when assayed in isolation, a discrete FEN homologue encoded by *S. aureus* has both flap endonuclease and 5'→3' exonuclease activities. We compare and contrast these two representatives of the cryptic single-domain eubacterial FEN homologues and speculate as to their biological roles.

EXPERIMENTAL

Alignments and phylogenetics

DNA Poll paralogues were identified by TBLASTN searches [30] and multiple sequence alignments prepared with CLUSTALW [31], edited in Jalview [32] and annotated by hand. Identified primary motifs were analysed with Pattrinprot [33]. A phylogenetic tree was constructed from the alignment using Phylip algorithms [34]. These were defined using default parameters in Protdist and Neighbor-joining clustering [35], followed by radial plots, generated using the Phylemon server (<http://phylemon.bioinfo.cipf.es>) [36].

Secondary and tertiary structural predictions

Structural predictions for SaFEN were generated using FUGUE ([http://www-cryst.bioc.cam.ac.uk/~fugue/prfsearch.html](http://www.cryst.bioc.cam.ac.uk/~fugue/prfsearch.html)) [37] and Phyre (<http://www.sbg.bio.ic.ac.uk/~phyre/>) [38]. The quality of the derived models was checked by eye in the first instance, for the local geometry of strictly conserved residues. A model of SaFEN, docked with a branched substrate was prepared by superimposition of the Phyre Apo SaFEN model with bacteriophage T4 RNase H in complex with DNA (pdb code 2IHN), using an all atom best-fit algorithm in Deepview (<http://www.expasy.org/spdbv/>) [39], followed by manual rigid body docking of DNA within a 1.5 Å constraint and final remodeling of ssDNA regions to facilitate best fit. Final models were submitted to PROCHECK [40], hosted on the JCSB server (http://www.icsg.org/scripts/prod/validation/sv_final.cgi) for assessment of their

stereo-chemical quality. MolProbity (<http://molprobity.biochem.duke.edu/>) was used for additional checks on the validity of ssDNA remodeling, in comparison with the original DNA template (chain CD, of pdb 2IHN [24]).

Bacterial strains and plasmids

Escherichia coli strain M72 pJONEX4ch was kindly provided by Dr. C. E. Meadows (University of Sheffield) and purified genomic DNA from *Staphylococcus aureus* strain 8325-4 was a generous gift from Prof. Simon Foster (University of Sheffield). Recombinant bacteriophage T5 FEN and *E. coli* exonuclease IX (ExoIX) were over-expressed and purified, as described previously [12, 29].

Cloning recombinant *S. aureus fen*.

Polymerase chain reaction (PCR) was performed using *S. aureus* strain 8325-4 genomic DNA template with primers for MW1329 (forward, 5'-TAT TGA ATT CAA AGA AGG AAC GTT TAA ATG CCT-3'; reverse, 5'-TTT ATA CGG ATC CAA AAA TGG GAT GAA ATA TAT TTT CC-3'), using the Expand High Fidelity PCR System, according to the manufacturer's instructions (Roche Diagnostics GmbH). Gel-purified product was digested with BamHI and EcoRI, ligated into pJONEX4ch (a generous gift from Dr. C. Meadows, University of Sheffield) and transformed into electrocompetent *E. coli* M72. Positive clones were confirmed by capillary sequencing on an ABI 3730 instrument (Genetics Core Facility, University of Sheffield). The resulting expression plasmid was designated pJONEX4/fench. This plasmid encoded the *S. aureus* FEN-like protein as a C-terminal polyhistidine fusion linked via an enterokinase cleavage motif resulting in the addition of the sequence DDDDKHHHHH to the C-terminus of the protein.

Preliminary inductions

Culture medium (30 ml 2xYT with 100 µg ml⁻¹ carbenicillin [2xYT/C]) was inoculated with a 1/100 volume (300 µl) of an overnight culture of M72 pJONEX4/fench. The culture was grown to an A_{595nm} of 0.5 and induced at 42°C for 0.5, 1, 1.5 or 2 h duration, followed by incubation at 30°C, for up to a total of 2 h post induction (*i.e.* the 2 h, 42°C culture received no further incubation). A control strain containing the empty vector was grown with continuous heat-shock or without temperature-induction. Cells were harvested and analysed by denaturing 10 % acrylamide SDS-PAGE using either Coomassie stain or DNase substrate gel assays [41].

Over-expression and purification of SaFEN

A fermenter vessel containing approx. 5 L 2xYT/C with 0.01% v/v antifoam (Sigma) was seeded with a 1/15 inoculum of an overnight culture of M72 pJONEX4/fench. The culture was grown to an absorbance of 0.6 (A_{595 nm}) at ≤ 30°C, followed by heat-shock induction at 42°C for 2 h. The culture was cooled to 25°C and incubated for a further 4.5 h, prior to harvesting. Cell pellet was recovered by centrifugation at 3,000 x g for 30 min at 4°C, washed with 50 mM Tris·HCl, pH 8.0 containing 0.9% w/v NaCl and stored in aliquots at -80°C. Cell pellet (7.5 g) was resuspended in 40 ml of 25 mM potassium phosphate buffer, pH 7.4, containing 20% w/v glycerol, 500 mM NaCl and 30 mM imidazole (EQ buffer). Lysozyme (200 µg ml⁻¹) and sodium deoxycholate (500 µg ml⁻¹) were added and the reaction incubated on ice for up to 1 h, followed by

sonication at 20% max. amplitude in 10 s bursts. Insoluble material was removed via centrifugation at 19,000 x *g* for 30 min and the supernatant cleared by filtration (Whatman No. 1) to remove residual precipitate, before loading onto a 1 ml HisTrap column (GE Healthcare), equilibrated with EQ buffer, at $\leq 0.5 \text{ ml min}^{-1}$. The column was washed in excess of 30 ml EQ buffer (returning $A_{280 \text{ nm}}$ to baseline), followed by elution over a 15 ml 40 - 500 mM imidazole gradient, collecting 0.5 ml fractions. Fractions were analysed by SDS-PAGE using Coomassie and zymogram assays for exonuclease activity [41]. Fractions of greatest purity were equilibrated against a 100-fold excess of LSQ buffer (25 mM Tris•HCl, pH 8.0 containing 5% w/v glycerol, 1 mM EDTA and 5 mM DTT), adjusted to 30% w/v ammonium sulphate and incubated for 15 min at RT. Precipitated material was collected via centrifugation at 4,000 x *g* for 10 min and the buffer adjusted to LSQ Buffer, containing 50% w/v glycerol, for storage at -20°C.

Nuclease assays

Structure-specific nuclease assays were performed using oligonucleotides synthesized by MWG Biotech AG (table 1) and purified by butanol precipitation [42]. Oligonucleotides were radiolabelled in a reaction (30 μl) containing 5 pmol DNA, [1x] kinase buffer (70 mM Tris•HCl, pH 7.6, 10 mM MgCl_2 , 5 mM DTT), 6 pmol [γ - ^{32}P] ATP and 30 units polynucleotide kinase at 37°C for 1 h, quenched by the addition of an equal volume of stop-mix (95 % deionised formamide, 10 mM EDTA and 0.05 % w/v bromophenol blue). Labelled oligonucleotides were purified using a QIAquick Nucleotide Removal Kit (Qiagen Ltd) and quantified by comparison with a serial dilution of [γ - ^{32}P] ATP using a BioRad Personal FX phosphorimager. Substrates for structure-specific assays were formed in annealing reactions (200 μl) containing 450 pM [γ - ^{32}P]-labelled DNA and 13.5 nM cold complementary oligonucleotide(s), in a buffer comprising 25 mM Tris•HCl, pH 8.0 and 500 mM KCl, incubated at 80°C for 5 min and passively cooled to ambient temperature. Standard nuclease reactions (10 μl) were performed with 3 μl annealed labelled oligonucleotide, 5 μl of 2 X reaction buffer (50 mM Tris•HCl, pH 8.0, 200 mM KCl, 20 mM MgCl_2 and 0.6 mg ml^{-1} acetylated BSA), and 2 μl enzyme prepared in enzyme dilution buffer (25 mM Tris•HCl, pH 8.0, containing 10 % w/v glycerol, 1 mM EDTA, 1 mM DTT and 1 mg ml^{-1} acetylated BSA). The final mixture, which contained 100 nM enzyme and 135 pM radiolabelled oligonucleotide was incubated at 37°C for 30 min before the reaction was quenched by the addition of an equal volume of stop-mix. Reaction products were analysed by denaturing PAGE (15% 19:1 acrylamide: bisacrylamide) [42]. In addition to the structure-specific nuclease assays described above, an exonuclease assay was deployed based on that described by Fraser [43]. This assay used high molecular weight DNA as a substrate. Exposure to exonuclease releases acid soluble mononucleotides and polynucleotides from acid precipitable substrate. Release of acid soluble products was monitored by UV spectroscopy after incubation with perchloric acid and centrifugation as described [29].

RESULTS AND DISCUSSION

Delineation of eubacterial FENs

We originally identified the gene now known as *xni* by homology searches of the sequence databases in 1994 [4, 28]. The *xni* gene encodes ExoIX, a paralogue of the N-terminal FEN domain of DNA polymerase I. The defining feature of all the eubacterial FEN family is a highly conserved cluster of carboxylate ligands at the

putative active site (Supplementary figure S1). In *E. coli* ExoIX (*EcExoIX*), these are Asp10, Asp51, Glu103, Asp105 and Asp128. A key feature of the *EcExoIX* polypeptide sequence is the absence of certain aspartic acid residues, conserved in previously characterized FENs (Figure 1A). These conserved residues form the ligands that coordinate one of two divalent metal ions, directly observed in several FEN crystal structures [14-16]. *EcExoIX* possesses the conserved aspartates which make up the high affinity, catalytic metal binding site, known as site I [17]. However, the multiple sequence alignment (Figure 1) reveals that key aspartate residues, normally involved in binding the second divalent metal ion in classical FENs have been replaced by small side chains Gly130, Ser175 and the aliphatic residue, Ile178 in *EcExoIX* (boxed residues in Figure 1A). These residues are unable to chelate a divalent metal ion. Indeed, when only two of the analogous mutations (D201I and D204S) were introduced into T5 FEN, the resulting mutant lost its ability to bind metal at site II [17] as well as much of its potency as an exonuclease. However, the structure-specific endonuclease activity of T5FEN D201I-D204S mutant remained intact. The metal-site II substitutions form a striking motif that delineates the ExoIX subset from the wider FEN family of polypeptides (Figures 1B and S1).

Our sequence analysis revealed a further class of FEN paralogues. We observed a subset encoding short FEN-like proteins that retain all the conserved aspartic acid residues associated with both metal sites I and II. This subset encoded FEN-like proteins that were independent of the N-terminal FEN domains found in DNA Poll orthologues. Figure 1A shows exemplars of the two subsets of autonomous FEN-like proteins. The *S. aureus* (*SaFEN*) represents those containing all the conserved features of the FENs while *E. coli* ExoIX (*EcExoIX*), represents those lacking the aspartates necessary for binding metal site II. Phylogenetic analysis reveals that ExoIX and FEN fall broadly, though not exclusively in classification based on Gram type (Figure 1B). Here we analysed nineteen previously identified Poll FEN paralogues [44], performing ClustalW multiple alignments (Figure S1) to generate a phylogenetic tree (Figure 1B).

In our analysis we also observed another, intriguing class of Poll FEN paralogues in the marine bacteria, *Shewanella* sp. and *Colwellia* sp. These organisms appear to fall at the divergent branch point between autonomous eubacterial FEN and ExoIX subsets (Figure 1B). They retain more conserved residues than *EcExoIX* (*Colwellia* sp. Gly150, Asp195 and N198) in place of typical FEN site II aspartate ligands. Based on site directed mutagenesis in T4 RNase H and other FENs, we suggest these gene products would be likely to exhibit diminished enzymatic activity[22].

We tentatively suggest that prokaryotes can be classified into four groups based on the number and type of FEN homologue that they encode: (i) a FEN only as an N-terminal domain of DNA Poll, as exemplified in *Streptococcus pneumoniae* (ii) N-terminal DNA Poll with a separate FEN paralogue with one divalent metal binding site e.g. ExoIX in *E. coli*, (iii) N-terminal DNA Poll with a separate FEN paralogue with two divalent metal binding sites e.g. *S. aureus* and (iv) a Klenow-like Poll domain (with no attached FEN domain) and a separate FEN domain with two divalent metal ion binding sites as exists in *Mycoplasma genitalium* (NP_072928). Accordingly, we therefore propose a more precise nomenclature for prokaryotic FEN-like genes, to allow for this simple but important distinction. The gene name *xni* should be used to specify the subset containing only the site I carboxylate ligands (*i.e.* those genes encoding ExoIX orthologues). The gene *fen* should specify all other homologues, whether carried alongside *polA* or as the sole chromosomal copy. We support this proposition with the following biochemical analysis of the *S. aureus fen* gene product.

***S. aureus fen* encodes an active nuclease**

Based on the retention of a full complement of metal-binding aspartyl ligands, we reasoned that the newly identified staphylococcal enzyme should display similar structure-specific endo- and exo-nucleolytic activities to previously identified flap endonucleases. We cloned and expressed *S. aureus fen*, under the control of the temperature-inducible bacteriophage λ P_L promoter in the pJONEXCH expression vector [45]. The recombinant protein was expressed with a C-terminal polyhistidine tag. Figure 2A shows the properties of the recombinant SaFEN protein in preliminary inductions. A distinct band of the anticipated molecular mass (35 kDa) was induced upon heat-shock (0.5 - 2 h) but was absent in control lysates (-/+ , 0 h). Furthermore, using an *in situ* zymogram assay we observed a nuclease activity co-induced, under conditions of heat shock (Figure 2B, panel ii) with some evidence of proteolysis. The protein was purified to > 95% homogeneity (Figure 2B, panel i). Zymogram assays confirmed that the DNase activity shared chromatographic and physicochemical properties with the induced SaFEN protein (Figure 2B, panel ii). These data show that the novel *S. aureus fen* gene encodes a *bona fide* nuclease. The nuclease activity required divalent metal ions as expected and was not adversely affected by the C-terminal fusion (data not shown). We also obtained very similar zymogram data for an untagged version of the SaFEN protein but it proved difficult to purify without the affinity tag unlike the untagged T5FEN and ExoIX (data not shown).

Comparison of SaFEN with other flap endonuclease family members

Using a standard spectrophotometric assay, we quantified the exonucleolytic activity of purified SaFEN against the well-characterized control flap endonuclease T5FEN (Figure 3). This has proven to be a simple, highly reproducible and robust assay, in the characterisation of other flap endonucleases [12, 23, 29]. Under similar experimental conditions (pH 8.0 at 37°C), the specific activity for T5FEN, calculated in a previous study (610 units μg^{-1}) [29] was similar to that reported here (420 \pm 10 units μg^{-1}). The specific activity of SaFEN (90 \pm 5 units μg^{-1}) was approximately five-fold lower than that of T5FEN, equivalent to a turnover rate of 106 min⁻¹. T5FEN is a highly processive exonuclease, which serves as a stringent control. By comparison, the reported turnover of *E. coli* Pol I FEN, under similar assay conditions is 37 min⁻¹ (37°C, pH 7.4) [2]. Thus, the activity of SaFEN is approximately three-fold greater than *E. coli* Pol I FEN activity under similar conditions, suggesting the activity of the novel polypeptide is biologically relevant. As reported previously ExoIX, showed no detectable activity under any assay conditions tested [29].

Structure-specific activity of SaFEN and EcExoIX

We hypothesised that the alterations in predicted active site of the two classes of divergent FENs, would lead to pronouncedly different catalytic activity. To test this we employed a substrate screen formed from partial or completely complementary oligonucleotides (Table 1). The catalytic activity of the proteins was examined for exo- and endonucleolytic cleavage under denaturing PAGE (Figure 4).

Single-stranded and fully duplexed DNA substrates were cleaved exonucleolytically by SaFEN, liberating products of the same electrophoretic mobility as those released from T5FEN, albeit to a lesser extent (Figure 4, panel A and B). The enzyme also liberated exo-product (\leq 4mer) from the 5' terminus of the downstream strand of a nicked DNA strand. Under the conditions tested, all substrate was degraded (Figure 4, panel C). The nicked substrate (36mer) was larger than the duplex DNA (24mer), though this difference is unlikely to account for the heightened activity on the former substrate. This is supported by the similar substrate turnover of the strand upstream of a nick, compared with purely duplex DNA (*cf.* Panels D and B). These data

suggest SaFEN favours a free 3' terminus, upstream of the free 5' end, which perhaps facilitates loading of the enzyme onto the DNA substrate. In this assay the activity of SaFEN was apparently equivalent to T5FEN. Such activity correlates with the involvement of this class of proteins in nick translation [46].

While SaFEN did not endonucleolytically hydrolyse a 5' DNA overhang, 5' pseudo-Y (PsY) and 3' invader (Inv) bifurcations were cleaved by the staphylococcal enzyme (Panels E-G). On PsY and Inv substrates, SaFEN exhibited both endo- and exo-nucleolytic cleavage, the products for which corresponded to decamers (closed triangle) and tetramers (open triangle). Interestingly, a free 5' flap of a partially formed junction (panel H) was not hydrolysed by SaFEN. This structure resembled the double flap substrate used in the early characterization of human FEN [47]. In those assays, human FEN cleaved the 5' flap of double flap structures efficiently, when 3' overlaps of up to 10 nucleotides were tested. T5FEN cut the large double flap substrate at a number of sites, the most prominent product of which was a 20mer, formed from hydrolysis at the bifurcation junction. The complex cleavage pattern with this substrate is probably due to transient base pairing, causing the formation of substrates with different free 5' arms.

The length of duplex DNA also appears to play a role in cleavage efficiency. This hypothesis is supported by the cleavage properties on the dsDNA region of larger (20 bp) Holliday junction intermediates (Panels I and J). Duplex DNA regions of partially formed junctions were hydrolyzed more efficiently than shorter substrates (3' PsY and 5' Inv substrates, data not shown), with comparable activity to all duplex DNA (panel B). SaFEN activity was limited to exonucleolytic degradation of duplex DNA in Holliday junction intermediates.

In contrast to SaFEN, ExoIX did not demonstrate flap endonuclease activity when incubated with a range of substrates (Figure 4). Therefore, ExoIX (when assayed in isolation) did not possess either 5-3' exonuclease or 5' flap endonuclease activity that would be expected of a FEN enzyme. We hypothesize that the lack of observable characteristic FEN activity is due to the absence of divalent metal ion binding site II in this protein.

Modelling cofactor and substrate interactions of SaFEN

We used a structural bioinformatics approach to gain insight into the catalytic differences observed in the different FEN homologues characterised above (Figure 5). Using a coordinated approach of different structural prediction algorithms we were able to ascertain with confidence the likely structure of the newly identified staphylococcal enzyme (FUGUE z-score 43.6, where ≥ 6 represents 99% confidence [37]). A Phyre-generated model was used for analysis based on its favourable structural constraints [38]. The model bore striking semblance to many experimentally derived FEN structures, most notably at the DNA binding helix-three-turn helix (H3TH) motif and the putative active site (panel A and B). The model reveals that both divalent metal binding sites are present in SaFEN, whereas, metal site II ligands are absent in ExoIX. We postulate this accounts for the different catalytic activities observed in our screen when the staphylococcal enzyme was compared to ExoIX. Comparison of this threaded model with other FEN family members revealed a smaller arch domain than that found in T5FEN. The first co-crystal structure of a FEN in complex with a bifurcated substrate has recently been solved [24]. We sought to use this as a model to understand the behaviour of SaFEN in its activity on a similar Ps-Y substrate assayed above (Figure 5, panel C *cf.* Figure 4, panel F). The restricted arch of the SaFEN model could still allow DNA to thread

though upon interaction with a branched substrate, positioning the scissile phosphate diester close to the site I metal binding site which is known to be essential for catalysis in the FENs [17,18]. The more restricted arch present in SaFEN may account for the slightly different cleavage pattern it exhibits compared to the T5FEN (Figure 4).

Biological role of the cryptic FEN-like proteins

The gene now known as *xni* was initially designated *exo* by this laboratory. It was speculated at the time that the gene might encode a backup function for the FEN-domain of PolI [28]. This inference was initially undermined by the observation that *Streptococcus pneumoniae polA* null mutants are non-viable [48]. However, this organism does not encode an additional FEN-like protein. Recently, Fukushima and co-workers demonstrated that the FEN-domain encoded by either *polA* or *xni* was required for the viability of *E. coli* [44]. Similarly, they also showed that either *polA* or *ypcP* is essential for *B. subtilis*. The *ypcP* gene encodes a two-metal-site FEN that shares 48% identity and 64% similarity to SaFEN (Figure S1). Thus, it would appear that either subclass of the single domain FEN-like proteins discussed here acts in redundancy with DNA polymerase I. The actual biochemical activity of the *xni*-encoded proteins remains elusive, as we were unable to identify any nucleolytic activity when tested in isolation in biochemical assays, despite testing a wide range of substrates and cofactors. Interestingly, amino acid sequence alignments also suggest that the helical arch motif identified as being important in FEN/substrate interactions is truncated in *EcExoIX* which may be an additional factor in the lack of expected nuclease activity. Perhaps the *ExoIX*-type proteins need to be part of a complex of protein(s) to display nuclease activity or require some as yet unidentified cofactor. Alternatively, it could be that they process some form of unusual nucleic acid structure.

However, it is clear from genetic experiments and their widespread occurrence that these small FEN-like proteins are capable of fulfilling a critical biological role. The reasons why each group of organisms has retained FEN-like proteins with differing properties is unclear and will require further investigation in to how these proteins participate in biological processes.

Accepted Manuscript

Sequence		Used in Panel
5'-d(<i>G</i> ATCCTAACTGGGGTTCAGAATCC)	This study	A-G
5'-d(GGATTCTGAACCCCAGTTAGGATC)	"	B
5'-d(GGATTCTGAACCCCAGTTAGGATCTCGGATCCTGTA)	"	C
5'-d(TCGGATCCTGTAGGATTCTGAACCCCAGTTAGGATC)	"	D
5'-d(TACAGGATCCGA)	"	C, D
5'-d(GGATTCTGAACCCC)	"	E
5'-d(GGATTCTGAACCCCACGGTAGTCAACGTG)	"	F, G
5'-d(CACGTTGACTACCGTC)	"	G
5'-d(<i>A</i> GTCTAGACTGCAGTTGAGTCCTTGCTAGGACGGATCCCT)	Ref. [49]	H, I, J
5'-d(AGGAATTCAACCACCGCTCAACTCAACTGCAGTCTAGACT)	Ref. [49]	I, J
5'-d(AGGGATCCGTCTAGCAAGGGGCTGCTACCGGAAGCTTCT)	Ref. [49]	H, J
5'-d(AGAAGCTCCGGTAGCAGCCTGAGCGGTGGTTGAATTCCT)	Ref. [49]	H, I

Table 1. Oligonucleotides used in structure-specific nuclease assays, presented in Figure 4.

Substrates were prepared as described in the Materials and Methods, in brief, cold oligonucleotides were present at a 30-fold excess (13.5 nM) when annealed to the complementary radiolabelled strand (450 pM). Radiolabelled oligonucleotides are *italicised* and the labelled 5' terminal nucleotide indicated in bold typeface.

Accepted Manuscript

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20081637

Figure Legends

Figure 1. Evaluation of eubacterial DNA Polymerase I 5' - 3' exonuclease paralogues.

A. ClustalW multiple sequence alignment of 5'-3'-like nucleases with their related Pol I paralogue. Both classes of eubacteria contain two putative paralogous FEN-like proteins, one associated with the Klenow fragment in the form of DNA Pol I (*polA*) and the other encoded autonomously. The autonomous FEN-like proteins can be divided into those which retain all the putative ligands to bind two metal ions (e.g. *Staphylococcus aureus* FEN, SaFEN) and those which are retain the ligands to bind a single divalent metal ion (*Escherichia coli* Exo IX). Perfectly conserved residues are boxed in grey. The amino acid substitutions affecting binding at the second metal site in Exo IX are shown in clear boxes. **B.** Phylip radial phylogenetic tree, derived from ClustalW alignment of nineteen secondary FENs from eubacteria (supplementary Figure S1). A clear differentiation of Exo IX and FEN classes. Eubacterial paralogous FENs fall into two discrete classes with predicted affinity for one (hatched area) or two (grey area) metal ions. This broadly, but not exclusively correlates with Gram positive and negative classification (bold and underlined type, respectively).

Figure 2. SDS-PAGE (10%) analysis of recombinant *S. aureus* FEN.

A. Heat-shock induction (0 - 2 h at 42°C) of recombinant SaFEN demonstrates protein of the anticipated molecular mass (35 kDa; black arrow); **i)** stained with Commassie or **ii)** zymogram for DNase activity. **B.** Fractions taken during the purification of SaFEN were analysed by SDS-PAGE (10% acrylamide) and stained with **i)** Coomassie blue or **ii)** a zymogram assay of DNase activity. A replicate gel was prepared with 40 µg ml⁻¹ Type XIV herring sperm DNA, and assayed for *in situ* DNase activity, furnished with 10 mM MgCl₂ as cofactor [41]. Abbreviations; M, Marker (BioRad precision unstained), molecular mass of standards is given in kDa; TL, total lysate; SE, soluble extract; Ch, pooled Ni-chelate eluate; AS; ammonium sulphate precipitated fraction (final 30% saturated).

Figure 3. Comparison of exonucleolytic properties of prokaryotic FEN homologues.

Release of acid soluble nucleotides (nmol) in a standard reaction (1 ml) at pH 8.0, containing an excess of type XIV herring sperm DNA substrate, measured spectrophotometrically, adjusted to 1 µg protein to aid comparison: ExoIX (▲), SaFEN (□), T5FEN (◇). Mean data from three independent readings were plotted, shown with standard error (error bars). Nucleotide release was determined as described elsewhere [12].

Figure 4. Comparison of enzymes' structure-specific nucleolytic cleavage on a selection of exo- and endo- nuclease substrates.

An excess of enzyme (100 nM) was incubated with preformed 5'-labelled substrate (135 pM) in the presence of 10 mM MgCl₂, incubated at 37 °C for 30 min. and the reactions analysed on denaturing 15% PAGE. Filled triangles (▶) denote structure-specific endonucleolytic cleavage products and open triangles (▷) exonucleolytic cleavage products. Abbreviations: -ve, negative control (BSA carrier, no enzyme); T5, bacteriophage T5 FEN; Sa, *S. aureus* FEN; IX; *E. coli* ExoIX.

Figure 5. Homology modelling SaFEN in complex with M^{2+} cofactor and a branched Pseudo-Y DNA substrate.

Bacteriophage T5FEN (pdb 1UT8) served as a template for the Phyre-threaded model of SaFEN [38], confirmed by two independent algorithms and structural validation (refer to experimental section). **A.** Cartoon representation of SaFEN. α -helices are blue and β -sheets yellow. The DNA binding H3TH motif is illustrated in white and the predicted active site region is hatched, shown in detail in **B.** Perfectly conserved eubacterial FEN residues include Tyr and Lys, in addition to a cluster of carboxyl ligands at the active site, here modelled with two Mn^{2+} ions (cyan spheres). The ligand substitutions affecting site II in the ExoIX sub-class are in brackets. **C.** A branched DNA substrate was modelled by rigid body docking, based on structural superimposition of the SaFEN model with 2IHN, and steric clashes reduced to within 1.5 Å. The free 5' end (in gold) fits best with a threading model, as originally proposed for T5FEN [15], although the arch is disordered and more spatially constricted in SaFEN.

Accepted Manuscript

References

- 1 Setlow, P., Brutlag, D. and Kornberg, A. (1972) Deoxyribonucleic acid polymerase: two distinct enzymes in one polypeptide. I. A proteolytic fragment containing the polymerase and 3' leads to 5' exonuclease functions. *J. Biol. Chem.* **247**, 224-231
- 2 Setlow, P. and Kornberg, A. (1972) Deoxyribonucleic acid polymerase: two distinct enzymes in one polypeptide. II. A proteolytic fragment containing the 5' leads to 3' exonuclease function. Restoration of intact enzyme functions from the two proteolytic fragments. *J. Biol. Chem.* **247**, 232-240
- 3 Robins, P., Pappin, D. J., Wood, R. D. and Lindahl, T. (1994) Structural and functional homology between mammalian DNase IV and the 5'-nuclease domain of *Escherichia coli* DNA polymerase I. *J. Biol. Chem.* **269**, 28535-28538
- 4 Lieber, M. R. (1997) The FEN-1 family of structure-specific nucleases in eukaryotic DNA replication, recombination and repair. *Bioessays* **19**, 233-240
- 5 Ceska, T. A. and Sayers, J. R. (1998) Structure-specific DNA cleavage by 5' nucleases. *Trends Biochem. Sci.* **23**, 331-336
- 6 Liu, Y., Kao, H. I. and Bambara, R. A. (2004) Flap endonuclease 1: a central component of DNA metabolism. *Annu. Rev. Biochem.* **73**, 589-615
- 7 Hobbs, L. J. and Nossal, N. G. (1996) Either bacteriophage T4 RNase H or *Escherichia coli* DNA polymerase I is essential for phage replication. *J. Bacteriol.* **178**, 6772-6777
- 8 Sun, X., Wu, P., Zheng, L., Thrower, D., Partikian, A., Qiu, J. and Shen, B. (2002) Suppression of *Saccharomyces cerevisiae* rad27 null mutant phenotypes by the 5' nuclease domain of *Escherichia coli* DNA polymerase I. *Curr. Genet.* **41**, 379-388
- 9 Kucherlapati, M., Yang, K., Kuraguchi, M., Zhao, J., Lia, M., Heyer, J., Kane, M. F., Fan, K., Russell, R., Brown, A. M., Kneitz, B., Edelmann, W., Kolodner, R. D., Lipkin, M. and Kucherlapati, R. (2002) Haploinsufficiency of Flap endonuclease (Fen1) leads to rapid tumor progression. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 9924-9929
- 10 Zheng, L., Dai, H., Zhou, M., Li, M., Singh, P., Qiu, J., Tsark, W., Huang, Q., Kernstine, K., Zhang, X., Lin, D. and Shen, B. (2007) Fen1 mutations result in autoimmunity, chronic inflammation and cancers. *Nat. Med.* **13**, 812-819
- 11 Rossi, M. L. and Bambara, R. A. (2006) Reconstituted Okazaki fragment processing indicates two pathways of primer removal. *J. Biol. Chem.* **281**, 26051-26061
- 12 Sayers, J. R. and Eckstein, F. (1990) Properties of Overexpressed Phage T5 D15 Exonuclease. *J. Biol. Chem.* **265**, 18311-18317
- 13 Lundquist, R. C. and Olivera, B. M. (1982) Transient generation of displaced single-stranded DNA during nick translation. *Cell* **31**, 53-60
- 14 Kim, Y., Eom, S. H., Wang, J. M., Lee, D. S., Suh, S. W. and Steitz, T. A. (1995) Crystal structure of *Thermus aquaticus* DNA polymerase. *Nature* **376**, 612-616
- 15 Ceska, T. A., Sayers, J. R., Stier, G. and Suck, D. (1996) A helical arch allowing single-stranded DNA to thread through T5 5'- exonuclease. *Nature* **382**, 90-93
- 16 Mueser, T. C., Nossal, N. G. and Hyde, C. C. (1996) Structure of bacteriophage-T4 RNASE-H, a 5' to 3' RNA-DNA and DNA-DNA exonuclease with sequence similarity to the RAD2 family of eukaryotic proteins. *Cell* **85**, 1101-1112
- 17 Feng, M., Patel, D., Dervan, J. J., Ceska, T., Suck, D., Haq, I. and Sayers, J. R. (2004) Roles of divalent metal ions in flap endonuclease-substrate interactions. *Nat. Struct. Mol. Biol.* **11**, 450-456
- 18 Zheng, L., Li, M., Shan, J., Krishnamoorthi, R. and Shen, B. (2002) Distinct roles of two Mg²⁺ binding sites in regulation of murine flap endonuclease-1 activities. *Biochemistry* **41**, 10323-10331
- 19 Tock, M. R., Frary, E., Sayers, J. R. and Grasby, J. A. (2003) Dynamic evidence for metal ion catalysis in the reaction mediated by a flap endonuclease. *EMBO J.* **22**, 995-1004
- 20 Hosfield, D. J., Mol, C. D., Shen, B. and Tainer, J. A. (1998) Structure of the DNA repair and replication endonuclease and exonuclease FEN-1: coupling DNA and PCNA binding to FEN-1 activity. *Cell* **95**, 135-146
- 21 Hwang, K. Y., Baek, K., Kim, H. Y. and Cho, Y. (1998) The crystal structure of flap endonuclease-1 from *Methanococcus jannaschii*. *Nat. Struct. Biol.* **5**, 707-713

- 22 Bhagwat, M., Meara, D. and Nossal, N. G. (1997) Identification of residues of T4
RNase H required for catalysis and DNA binding. *J. Biol. Chem.* **272**, 28531-28538
- 23 Garforth, S. J., Ceska, T. A., Suck, D. and Sayers, J. R. (1999) Mutagenesis of
conserved lysine residues in bacteriophage T5 5'-3' exonuclease suggests separate
mechanisms of endo- and exonucleolytic cleavage. *Proc. Natl. Acad. Sci. U.S.A.* **96**,
38-43
- 24 Devos, J. M., Tomanicek, S. J., Jones, C. E., Nossal, N. G. and Mueser, T. C. (2007)
Crystal structure of bacteriophage T4 5' nuclease in complex with a branched DNA
reveals how flap endonuclease-1 family nucleases bind their substrates. *J. Biol.*
Chem. **282**, 31713-31724
- 25 Dervan, J. J., Feng, M., Patel, D., Grasby, J. A., Artymiuk, P. J., Ceska, T. A. and
Sayers, J. R. (2002) Interactions of mutant and wild-type flap endonucleases with
oligonucleotide substrates suggest an alternative model of DNA binding. *Proc. Natl.*
Acad. Sci. U.S.A. **99**, 8542-8547
- 26 Allawi, H. T., Kaiser, M. W., Onufriev, A. V., Ma, W. P., Brogaard, A. E., Case, D. A.,
Neri, B. P. and Lyamichev, V. I. (2003) Modeling of flap endonuclease interactions
with DNA substrate. *J. Mol. Biol.* **328**, 537-554
- 27 Shafritz, K. M., Sandigursky, M. and Franklin, W. A. (1998) Exonuclease IX of
Escherichia coli. *Nucleic Acids Res.* **26**, 2593-2597
- 28 Sayers, J. R. (1994) Computer aided identification of a potential 5'-3' exonuclease
gene encoded by *Escherichia coli*. *J. Theor. Biol.* **170**, 415-421
- 29 Hodkinson, M. R., Allen, L. M., Thomson, D. P. and Sayers, J. R. (2007) Molecular
interactions of *Escherichia coli* ExoIX and identification of its associated 3'-5'
exonuclease activity. *Nucleic Acids Res.* **35**, 4094-4102
- 30 Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. and
Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Res.* **25**, 3389-3402
- 31 Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A.,
McWilliam, H., Valentin, F., Wallace, I. M., Wilm, A., Lopez, R., Thompson, J. D.,
Gibson, T. J. and Higgins, D. G. (2007) Clustal W and Clustal X version 2.0.
Bioinformatics **23**, 2947-2948
- 32 Clamp, M., Cuff, J., Searle, S. M. and Barton, G. J. (2004) The Jalview Java
alignment editor. *Bioinformatics* **20**, 426-427
- 33 Combet, C., Blanchet, C., Geourjon, C. and Deleage, G. (2000) NPS@: network
protein sequence analysis. *Trends Biochem. Sci.* **25**, 147-150
- 34 Retief, J. D. (2000) Phylogenetic analysis using PHYLIP. *Methods Mol. Biol.* **132**,
243-258
- 35 Felsenstein, J. (1996) Inferring phylogenies from protein sequences by parsimony,
distance, and likelihood methods. *Methods Enzymol.* **266**, 418-427
- 36 Tarraga, J., Medina, I., Arbiza, L., Huerta-Cepas, J., Gabaldon, T., Dopazo, J. and
Dopazo, H. (2007) Phylemon: a suite of web tools for molecular evolution,
phylogenetics and phylogenomics. *Nucleic Acids Res.* **35**, W38-42
- 37 Shi, J., Blundell, T. L. and Mizuguchi, K. (2001) FUGUE: sequence-structure
homology recognition using environment-specific substitution tables and structure-
dependent gap penalties. *J. Mol. Biol.* **310**, 243-257
- 38 Bennett-Lovsey, R. M., Herbert, A. D., Sternberg, M. J. and Kelley, L. A. (2008)
Exploring the extremes of sequence/structure space with ensemble fold recognition in
the program Phyre. *Proteins* **70**, 611-625
- 39 Guex, N. and Peitsch, M. C. (1997) SWISS-MODEL and the Swiss-PdbViewer: an
environment for comparative protein modeling. *Electrophoresis* **18**, 2714-2723
- 40 Laskowski, R. A., Moss, D. S. and Thornton, J. M. (1993) Main-chain bond lengths
and bond angles in protein structures. *J. Mol. Biol.* **231**, 1049-1067
- 41 Rosenthal, A. and Lacks, S. A. (1977) Nuclease detection in SDS-polyacrylamide gel
electrophoresis. *Analytical Biochemistry* **80**, 76-90
- 42 Sambrook, J., Fritsch, E. F. and Manniatis, T. (1989) *Cloning: A Laboratory Manual*.
Cold Spring Harbor Laboratory Press, New York
- 43 Fraser, M. J. (1980) Purification and properties of *Neurospora crassa* endo-
exonuclease, an enzyme which can be converted to a single-strand specific
endonuclease. *Methods Enzymol.* **65**, 255-263


- 44 Fukushima, S., Itaya, M., Kato, H., Ogasawara, N. and Yoshikawa, H. (2007) Reassessment of the in vivo functions of DNA polymerase I and RNase H in bacterial cell growth. *J. Bacteriol.* **189**, 8575-8583
- 45 Sayers, J. R. and Eckstein, F. (1991) A single-strand specific endonuclease activity copurifies with overexpressed T5 D15 exonuclease. *Nucleic Acids Res.* **19**, 4127-4132
- 46 Kornberg, A. and Baker, T. A. (1992) *DNA Replication*. W.H. Freeman, New York
- 47 Harrington, J. J. and Lieber, M. R. (1995) DNA structural elements required for FEN-1 binding. *J. Biol. Chem.* **270**, 4503-4508
- 48 Díaz, A., Lacks, S. A. and López, P. (1992) The 5' to 3' exonuclease activity of DNA polymerase I is essential for *Streptococcus pneumoniae*. *Molecular Microbiology* **6**, 3009-3019
- 49 White, M. F. and Lilley, D. M. (1997) The resolving enzyme CCE1 of yeast opens the structure of the four-way DNA junction. *J. Mol. Biol.* **266**, 122-134


Accepted Manuscript

A


<i>Sa_Poll</i>	1	-----MNKLV	I	DGNSLS	S	FRAFYAL	PLLS	----	N	
<i>Sa_FEN</i>	1	-----MPNKI	L	LDGMALL	FRHFYAT	SLHKQF	MYN			
<i>Ec_Poll</i>	1	-----MVQI	PQNPLI	LVDGSSY	LYRAYHAF	PPLT	----	N		
<i>Ec_ExoIX</i>	1	MRGLFP	I	SHPAVAC	SGIECY	PYRLIF	KGVIVAVHLL	IVDALN	LIRRI	
<i>Sa_Poll</i>	26	KAGIHTNAVY	GFAMLLEK	I	LKEEK	FNHFLVAF	DAGKTT	--	FRHEKYS	
<i>Sa_FEN</i>	31	SQGEPTNG	IQGFVRH	I	FSAIHE	IRPTHVAVCW	DMGQST	--	FRNDMFD	
<i>Ec_Poll</i>	31	SAGEPTGAMY	GVNLMLRSL	I	MQYKPT	HAAVVFD	DAKGKT	--	FRDELFE	
<i>Ec_ExoIX</i>	48	HAVQGS	PCVETCQHALDQL	I	MHSQ	PTHAVAVF	DENR	SSGWR	HQRLP	
<i>Sa_Poll</i>	71	EYKGG	RQKTPPEL	SEQFPY	IRQLLDAYH	IKRYEL	LDNYEADD	I	IGTLS	
<i>Sa_FEN</i>	78	GYKQNR	SAPPEEL	I	PQFDYVKE	I	SEQFGFVN	I	GVKNEYADDV	IGTLA
<i>Ec_Poll</i>	78	HYKSHR	PPMPDDLRAQ	I	EPLHAM	VKAMGLP	LLAVSGVE	ADDV	IGTLA	
<i>Ec_ExoIX</i>	95	DYKAGR	PPMPEEL	HDEMPAL	LRAAFE	QRGVPCW	STSGNE	ADD	LAATLA	
<i>Sa_Poll</i>	118	KEADKAG	FQTI	I	TGDRDLT	DLATDNVT	I	YYTKK	GVTDVDHYTPDF	I
<i>Sa_FEN</i>	123	QQYSTDN	-DVYI	I	TGDKDL	LOCINDNVE	VWL	I	KKGFNI	YNRYTLHRF
<i>Ec_Poll</i>	123	REAEKAGR	PVLI	STGDK	DMAQLVTPN	ITL	INTMTNT	I	LGPEEV	VNKY
<i>Ec_ExoIX</i>	142	VKVTQAGH	QATIV	STK	GYCQLLSPTLR	IRDYF	QKRWLDAPF	I	DKEF	
<i>Sa_Poll</i>	165	AEKYNG	LTPNQI	I	DMKGLM	GDTSDNI	PCVAG	VG	GEKTAIKLLN	QFDTV
<i>Sa_FEN</i>	169	NEEY-	ALEPQQL	I	IKAFM	EDTADGYAG	VKGI	GEKTAIKLI	QQYQSV	
<i>Ec_Poll</i>	170	----	GVPELI	I	DFLALM	GSSDNIP	GV	PGV	GEKTAQALLQGLGGL	
<i>Ec_ExoIX</i>	189	----	GVQPQQL	P	YWG	LPISS	SKV	PGVAG	IGPKSATQLLVEFQSL	
<i>Sa_Poll</i>	212	EGVYEH	-----	LDEI	SGKLLKEK	LQNSKEDALMSKEL	AT	INVDSP	I	
<i>Sa_FEN</i>	215	ENVVEN	-----	TDALSAG	-QRNK	INDNLD	EYLSKRLAE	IHTQVP	I	
<i>Ec_Poll</i>	212	DTLYAE	PEKI	AGLSFR	GAKTMAAK	LEQNKEVAYLSYQL	AT	I	KTDVEL	
<i>Ec_ExoIX</i>	231	EGIYEN	-----	LDAVAEK	-WRKK	LETHKEMAF	L	CRDI	ARLQTDLHI	


B






THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20081637







THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20081637