



HAL
open science

Loop quantum black hole

Leonardo Modesto

► **To cite this version:**

| Leonardo Modesto. Loop quantum black hole. 2005. hal-00479682

HAL Id: hal-00479682

<https://hal.science/hal-00479682>

Preprint submitted on 12 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Loop quantum black hole

Leonardo Modesto

Department of Physics, Bologna University V. Iriero 46, I-40126 Bologna & INFN Bologna, EU

&

Centre de Physique Théorique de Luminy,

Université de la Méditerranée, F-13288 Marseille, EU

Abstract

In this paper we consider the Kantowski-Sachs space-time in Ashtekar variables and the quantization of this space-time starting from the complete loop quantum gravity theory. The Kantowski-Sachs space-time coincides with the Schwarzschild black hole solution inside the horizon. By studying this model we can obtain information about the black hole singularity and about the dynamics across the point $r = 0$. We studied this space-time in ADM variables in two previous papers where we showed that the classical black hole singularity disappears in quantum theory. In this work we study the same model in Ashtekar variables and we obtain a regular space-time inside the horizon region and that the dynamics can be extended further the classical singularity.

Introduction

In this work we study the space-time inside the horizon of a Schwarzschild black hole using the connection variables introduced by Ashtekar [1]. We start the quantization program from the complete theory of “loop quantum gravity” [2] and we reduce the theory to consider an homogeneous but anisotropic minisuperspace model.

We studied the singularity problem in two previous papers [3] and [4]. There we used the same non Schrödinger quantization procedure as in the work of V. Husain and O. Winkler on quantum cosmology. This formalism was introduced by Halvorson [13] and also by A. Ashtekar, S. Fairhurst and J. Willis [10]. In the previous papers we used ADM formulation of general relativity and we obtained that the quantum minisuperspace model for the black hole is singularity free; in fact in the first very simple model [3] we showed that the operator $1/r$ and so the curvature invariant $\mathcal{R}_{\mu\nu\rho\sigma}\mathcal{R}^{\mu\nu\rho\sigma} = 48M^2G_N^2/r^6$ are not divergent in the quantum theory. In the second paper [4] we considered a general two dimensional minisuperspace with space section of topology $\mathbf{R} \times \mathbf{S}^2$ (the Kantowski-Sachs space-time [6]) and we obtained that the inverse volume operator and the curvature invariant are singularity free in $r = 0$ in quantum gravity. In both models we have that at the quantum level the Hamiltonian constraint acts like a difference operator. A quantum extension on the other side of the classical singularity is straightforward.

We acknowledge Ioannis Raptis’ work on a mathematical approach to the singularity problem in general relativity [7].

The paper is organized as follows : in the first section we introduce the classical theory, in particular we recall the invariant connection 1-form for the Kantowski-Sachs space-time. We introduce the holonomies and the form of the Hamiltonian constraint and the inverse volume operator in terms of such holonomies. We also express the quantity $1/r$ in terms of holonomies. This quantity is

interesting because it is connected with the Schwarzschild curvature invariant $\mathcal{R}_{\mu\nu\rho\sigma} \mathcal{R}^{\mu\nu\rho\sigma}$ in the metric formulation of general relativity. In the second section we quantize the system. In particular we calculate explicitly the spectrum of the inverse volume operator, the spectrum of the operator $1/r$ and the solution of the Hamiltonian constraint in the dual to the kinematical Hilbert space.

1 Classical theory

In this section first of all we summarize the fundamental Hamiltonian formulation of “loop quantum gravity” [2] then we recall the formulation of the Kantowski-Sachs space-time in Ashtekar variables [9]. At this point we define the holonomies for the connection and we rewrite the classical Hamiltonian constraint and the inverse volume operator in terms of holonomies.

1.1 Loop quantum gravity preliminary

The classical starting point of loop quantum gravity (LQG) is the Hamiltonian formulation of general relativity. Historically in the ADM Hamiltonian formulation of the Einstein theory the fundamental variables are the three-metric q_{ab} of the spatial section Σ of a foliation of the $4-d$ manifold $\mathcal{M} \cong \mathbb{R} \times \Sigma$ and the extrinsic curvature K_{ab} . In LQG the fundamental variables are the Ashtekar variables: they consist of an $SU(2)$ connection A_a^i and the electric field E_i^a , where $a, b, c, \dots = 1, 2, 3$ are tensorial indexes on the spatial section and $i, j, k, \dots = 1, 2, 3$ are indexes in the $su(2)$ algebra. The density weighted triad E_i^a is define in terms of the triad e_a^i by the relation $E_i^a = \frac{1}{2} \epsilon^{abc} \epsilon_{ijk} e_b^j e_c^k$. And the metric can be expressed in terms of the triad by $q_{ab} = e_a^i e_b^j \delta_{ij}$.

We can express also the metric in terms of the weighted triad as

$$q q^{ab} = E_i^a E_j^b \delta^{ij}, \quad q = \sqrt{\det(q_{ab})}. \quad (1)$$

The relation between the variables (A_a^i, E_i^a) and the ADM variables (q_{ab}, K_{ab}) is

$$A_a^i = \Gamma_a^i + \gamma K_{ab} E_j^b \delta^{ij} \quad (2)$$

where γ is the Immirzi parameter and Γ_a^i is the spin connection, being the solution of Cartan’s equation: $\partial_{[a} e_{b]}^i + \epsilon_{jk}^i \Gamma_{[a}^j e_{b]}^k = 0$.

The action in the new variables is

$$S = \frac{1}{k} \int dt \int_{\Sigma} d^3x \left[-2Tr(E^a A_a) - NH - N^a H_a - N^i G_i \right], \quad (3)$$

where N^a is the shift vector, N is the lapse function and N^i is the Lagrange multiplying for the Gauss constraint G_i . The functions H , H_a and G_i are respectively the Hamiltonian, diffeomorphism and Gauss constraints .

The constrains are given by

$$\begin{aligned} H(E_i^a, A_a^i) &= -\frac{4}{\sqrt{|\det(E)|}} Tr [F_{ab} E^a E^b] - 2(1 + \gamma^2) \frac{E_i^a E_j^b K_{[a}^i K_{b]}^j}{\sqrt{|\det(E)|}} \\ H_b(E_i^a, A_a^i) &= E_j^a F_{ab}^j - (1 + \gamma^2) K_a^i G_i \\ G_i(E_i^a, A_a^i) &= \partial_a E_i^a + \epsilon_{ij}^k A_a^j E_k^a, \end{aligned} \quad (4)$$

and the curvature field straight is $F_{ab} = \partial_a A_b - \partial_b A_a + [A_a, A_b]$.

After this brief introduction to ”loop quantum gravity” we can study the ”Kantowski-Sachs” symmetric reduced model.

1.2 Kantowski-Sachs space-time in Ashtekar variables

At this point after we recalled the basic ingredients of LQG we can get back to study the Kantowski-Sachs space-time. This is a simplified model of a generical homogeneous but anisotropic space-time of coordinates (t, r, θ, ϕ) .

In general a homogeneous but anisotropic space-time of spatial section Σ of topology $\Sigma \cong \mathbb{R} \times S^2$ is characterized by a invariant 1-form connection $A_{[1]}$ of the form [8]

$$A_{[1]} = A_r(t) \tau_3 dr + (A_1(t) \tau_1 + A_2(t) \tau_2) d\theta + (A_1(t) \tau_2 - A_2(t) \tau_1) \sin \theta d\phi + \tau_3 \cos \theta d\phi. \quad (5)$$

The τ_i generators of the $SU(2)$ fundamental representation are relate to the Pauli σ_i matrix by $\tau_i = -\frac{i}{2}\sigma_i$.

On the other side the dual invariant densitized triad is

$$E_{[1]} = E^r(t) \tau_3 \sin \theta \frac{\partial}{\partial r} + (E^1(t) \tau_1 + E^2(t) \tau_2) \sin \theta \frac{\partial}{\partial \theta} + (E^1(t) \tau_2 - E^2(t) \tau_1) \frac{\partial}{\partial \phi}. \quad (6)$$

In this case the space-time is homogeneous and the diffeomorphism constraint is automatically satisfied; instead the Gauss constraint and the Euclidean part of the Hamiltonian constrains are similar to

$$\begin{aligned} G &\sim A_1 E^2 - A_2 E^1 \\ H_E &= \frac{2 \sin \theta \operatorname{sgn}[\det(E_{[1]})]}{\sqrt{|E^r|[(E^1)^2 + (E^2)^2]}} \left[2A_r E^r (A_1 E^1 + A_2 E^2) + \left((A_1)^2 + (A_2)^2 - 1 \right) [(E^1)^2 + (E^2)^2] \right] \end{aligned} \quad (7)$$

In this paper we want to study the Kantowski-Sachs space-time with space section of topology $R \times S^2$; the connection $A_{[1]}$ is more simple than in (5) with $A_2 = A_1$, and in the triad (6) we can choose the gauge $E^2 = E^1$ [9]. There is another residual gauge freedom on the pair (A_1, E^1) . This is a discrete transformation $P : (A_1, E^1) \rightarrow (-A_1, -E^1)$; we have to fix this symmetry on the Hilbert space.

The Gauss constraint in (7) is automatically satisfied and the Euclidean part of the Hamiltonian constraint becomes

$$H_E = \frac{2\sqrt{2} \sin \theta \operatorname{sgn}(E^r)}{\sqrt{|E^r||E^1|}} \left[2A_r E^r A_1 E^1 + (2(A_1)^2 - 1)(E^1)^2 \right] \quad (8)$$

from now on we redefine $E^r \equiv E$ and $A_r \equiv A$.

In the metric formalism the kantowski-Sachs space-time assume the form

$$ds^2 = -dt^2 + a^2(t)dr^2 + b^2(t)(\sin^2 \theta d\phi^2 + d\theta^2), \quad (9)$$

where $a(t)$ and $b(t)$ are two independent functions of the time coordinate "t".

The connection between the metric and density triad formulation is

$$q_{ab} = \begin{pmatrix} \frac{2(E^1)^2}{|E|} & 0 & 0 \\ 0 & |E| & 0 \\ 0 & 0 & |E| \sin^2 \theta \end{pmatrix}. \quad (10)$$

Using (10) we can calculate the volume of the spatial section Σ

$$V = \int dr d\phi d\theta \sqrt{q} = 4\pi\sqrt{2}R\sqrt{|E||E^1|}, \quad (11)$$

where R is a cut-off on the space radial coordinate. We can use also radial densities because the model is homogeneous and all the following results remain identical. In another way, the spatial homogeneity enable us to fix a linear radial cell \mathcal{L}_r and restrict all integrations to this cell [5].

To obtain the classical symplectic structure of the phase space we can calculate the first term of the action (3) introducing in it our variables.

This very simple calculation produces

$$S = \frac{1}{\kappa} \int dt \int dr d\phi d\theta \sin \theta [Tr(-2E^a A_a) + \dots] = \frac{R}{\kappa} \int dt \int d\phi d\theta \sin \theta [EA + 4E^1 A_1 + \dots]. \quad (12)$$

Concerning what we said before we can restrict the linear radial cell to the Planck length l_P and so we can take $R \equiv l_P$. Observing the reduced action (12) we obtain the symplectic structure of the classical phase space. The phase space consists of two canonical pairs A, E and A_1, E^1 and the symplectic structure is given by the poisson brackets, $\{A, E\} = \frac{\kappa}{l_P}$ and $\{A_1, E^1\} = \frac{\kappa}{4l_P}$. The coordinates and the momentums have dimensions: $[A] = L^{-1}$, $[A_1] = M^0$, $[E] = L^2$ and $[E^1] = L$.

1.3 Holonomies, curvature and Hamiltonian constraint

In this section we introduce the fundamental classical ingredients necessary to quantize the theory. To define the holonomies we don't consider the field independent term in the connection; this means that we define the model on particular spin networks which are based on graph made only of radial edges, or of edges along circles in the θ -direction or at $\theta = \pi/2$. Considering this graph type we can omit the term " $\tau_3 \cos \theta d\phi$ " in the holonomy in the " ϕ " direction.

We introduce the background triad ${}^o e_I^a = \text{diag}(1, 1, \sin^{-1} \theta)$ and co-triad ${}^o \omega_a^I = \text{diag}(1, 1, \sin \theta)$, and we define the holonomy

$$h = e^{\int A_{[1]}} = e^{\int A_a dx^a} = e^{\int A_a^i \frac{dx^a}{d\lambda} \tau_i d\lambda} = e^{\int A_I^i {}^o \omega_a^I {}^o e_J^a u^J \tau_i d\lambda} = e^{\int A_I^i u^I \tau_i d\lambda}, \quad (13)$$

where $u^a = \frac{dx^a}{d\lambda} = (R \frac{dr'}{d\lambda}, \frac{d\theta}{d\lambda}, \frac{d\phi}{d\lambda})$ and $u^I = {}^o \omega_a^I u^a$, r' is a dimensionless radial coordinate and R is the scale introduced in (11). In the rest of the paper as we wrote in the previous section we will take $R = l_P$ to simplify the formulas.

We can define now the holonomy along the direction "I"

$$\begin{aligned} h_I &= \exp \int A_I^i \tau_i d\lambda, \\ h_1 &= \exp \int A_1^i \tau_i d\lambda = \exp[A \mu_0 l_P \tau_3], \\ h_2 &= \exp \int A_2^i \tau_i d\lambda = \exp[A_1 \mu_0 (\tau_2 + \tau_1)], \\ h_3 &= \exp \int A_3^i \tau_i d\lambda = \exp[A_1 \mu_0 (\tau_2 - \tau_1)], \end{aligned} \quad (14)$$

and introducing the normalized vectors $n_1^i = (0, 0, 1)$, $n_2^i = \frac{1}{\sqrt{2}}(1, 1, 0)$, $n_3^i = \frac{1}{\sqrt{2}}(-1, 1, 0)$ we can rewrite h_I as

$$\begin{aligned} h_I &= \exp(\bar{A}_I n_I^i \tau_i) = \cos\left(\frac{\bar{A}_I}{2}\right) + 2n_I^i \tau_i \sin\left(\frac{\bar{A}_I}{2}\right), \\ h_1 &= \cos\left(\frac{A \mu_0 l_P}{2}\right) + 2\tau_3 \sin\left(\frac{A \mu_0 l_P}{2}\right), \\ h_2 &= \cos\left(\frac{\sqrt{2} A_1 \mu_0}{2}\right) + \sqrt{2}(\tau_2 + \tau_1) \sin\left(\frac{\sqrt{2} A_1 \mu_0}{2}\right), \\ h_3 &= \cos\left(\frac{\sqrt{2} A_1 \mu_0}{2}\right) + \sqrt{2}(\tau_2 - \tau_1) \sin\left(\frac{\sqrt{2} A_1 \mu_0}{2}\right) \end{aligned} \quad (15)$$

We remember that the Hamiltonian constraint can be written in terms of the curvature F_{ab} and the Poisson bracket between A_a and the volume V . The Euclidean part of the Hamiltonian constraint becomes

$$H_E = -2 \int d^3x \epsilon^{abc} \text{Tr} \left[F_{ab} \{A_c, V\} \right]. \quad (16)$$

We now rewrite the hamiltonian constraint (16) in terms of holonomies. We start to define the curvature F_{ab} using the holonomies (14)

$$F_{ab}^i \tau_i = {}^o\omega_a^I {}^o\omega_b^J \left[\frac{h_I h_J h_I^{-1} h_J^{-1} h_{[IJ]} - 1}{\epsilon(I)\epsilon(J)} \right], \quad (17)$$

where $\epsilon(1) = R\mu_0$, $\epsilon(2) = \epsilon(3) = \mu_0$. We have introduced also the field independent holonomy $h_{[IJ]}$ to obtain the correct classical curvature in the limit $\epsilon(I) \rightarrow 0$. This term is defined by

$$h_{[IJ]} = \exp(-\mu_0^2 C_{IJ} \tau_3) = \cos\left(\frac{\mu_0^2 C_{IJ}}{2}\right) - 2\tau_3 \sin\left(\frac{\mu_0^2 C_{IJ}}{2}\right), \quad (18)$$

where C_{IJ} is a constant antisymmetric matrix, $C_{IJ} = \delta_{2I}\delta_{3J} - \delta_{3I}\delta_{2J}$.

The other term we need is “ $\tau_i \{A_a^i, V\}$ ” and in terms of holonomies becomes

$$\tau_i \{A_a^i, V\} = {}^o\omega_a^I \frac{h_I^{-1} \{h_I, V\}}{\epsilon(I)}. \quad (19)$$

At this point we have all the elements to define the Euclidean part of the Hamiltonian constraint

$$\begin{aligned} H_E &= -2 \int d^3x \epsilon^{abc} {}^o\omega_a^I {}^o\omega_b^J {}^o\omega_c^K \text{Tr} \left[\frac{(h_I h_J h_I^{-1} h_J^{-1} h_{[IJ]} - 1) h_K^{-1} \{h_K, V\}}{\epsilon(I)\epsilon(J)\epsilon(K)} \right] \\ &= -2 \int d^3x \sqrt{\sigma q} \epsilon^{IJK} \text{Tr} \left[\frac{(h_I h_J h_I^{-1} h_J^{-1} h_{[IJ]} - 1) h_K^{-1} \{h_K, V\}}{\epsilon(I)\epsilon(J)\epsilon(K)} \right]. \end{aligned} \quad (20)$$

Recalling the $\epsilon(I)$'s definitions and that $\int d^3x \sqrt{\sigma q} = 4\pi l_P$ we can simplify the expression (20)

$$H_E = -\frac{8\pi}{\mu_0^3} \sum_{IJK} \epsilon^{IJK} \text{Tr} [h_I h_J h_I^{-1} h_J^{-1} h_{[IJ]} h_K^{-1} \{h_K, V\}], \quad (21)$$

where we have also eliminated the identity term because it doesn't contribute when traced.

1.4 Inverse volume & Schwarzschild curvature invariant in terms of holonomies

We can define the inverse volume operator following Thiemann. We remember that the volume operator is $V = 4\pi\sqrt{2}l_P |E|^{1/2} |E^1|$ and we define $\bar{V} = \frac{\sqrt{\det(q)}}{\sin\theta} = |E|^{1/2} |E^1|$.

A useful property to define the inverse volume operator is

$$\frac{\det(e_a^i)}{(\det(q))^{\frac{3}{4}}} = \frac{\sqrt{\det(q)}}{(\det(q))^{\frac{3}{4}}} = \frac{1}{\sqrt{\bar{V}}}, \quad (22)$$

using this relation and the identity $e_a^i = -\frac{(16)^{\frac{1}{2}} l_P}{\kappa} \{A_a^i, V\}$ we obtain

$$\frac{\text{sgn}(E)}{\sqrt{\bar{V}}} = -\frac{64}{3} \left(\frac{l_P}{\kappa}\right)^3 \epsilon_{ijk} \epsilon^{abc} \{A_a^i, \bar{V}^{\frac{1}{2}}\} \{A_b^j, \bar{V}^{\frac{1}{2}}\} \{A_c^k, \bar{V}^{\frac{1}{2}}\}. \quad (23)$$

To rewrite the volume in terms of holonomies we use extensively the holonomy version of the triad

$$e_a^i = \frac{2(16)^{\frac{1}{3}} l_P}{\kappa} o_{\omega_a^I} \frac{\text{Tr}[\tau^i h_I^{-1} \{h_I, \bar{V}\}]}{\epsilon(I)}. \quad (24)$$

At this point using the definition (24) in (23) we obtain the final version of the inverse volume operator in terms of holonomies

$$\frac{\text{sgn}(E)}{\sqrt{V}} = \left(\frac{512 l_P^3}{3 \kappa^3} \right) \epsilon_{ijk} \sum_{IJK} \frac{\epsilon^{IJK} \text{Tr}[\tau^i h_I^{-1} \{h_I, \bar{V}^{\frac{1}{2}}\}] \text{Tr}[\tau^j h_J^{-1} \{h_J, \bar{V}^{\frac{1}{2}}\}] \text{Tr}[\tau^k h_K^{-1} \{h_K, \bar{V}^{\frac{1}{2}}\}]}{\epsilon(I) \epsilon(J) \epsilon(K)}. \quad (25)$$

We consider now another operator that is useful to understand the black hole singularity. In the metric formulation of the Schwarzschild black hole the invariant quantity $\mathcal{R}_{\mu\nu\rho\sigma} \mathcal{R}^{\mu\nu\rho\sigma} \sim 1/r^6$ diverges in $r = 0$ where the singularity is localized. The relevant quantity $1/r^2 = 1/b^2(t)$ in metric formulation is given by $1/|E|$ in Ashtekar variables.

Now we express $1/\sqrt{|E|}$ in terms of holonomies

$$\frac{\text{sgn}(E) \text{sgn}(E^1)}{\sqrt{|E|}} = \frac{64 l_P}{\mu_0^2 \kappa^2} \text{Tr} \left[\tau_3 h_1^{-1} \{h_1, \bar{V}^{\frac{1}{2}}\} \right] \text{Tr} \left[(\tau_2 + \tau_1) h_2^{-1} \{h_2, \bar{V}^{\frac{1}{2}}\} \right] \quad (26)$$

We will use the quantities (25) and (26) in the next section where we study the black hole singularity for $E = 0$ in the quantum theory.

2 Quantum Theory

We want quantize the symmetry reduced theory using techniques from the full theory (loop quantum gravity). The Hilbert space for the Kantowski-Sachs minisuperspace model in Ashtekar variables is the tensor product of two Hilbert spaces, one for the anisotropy operator \hat{E}^1 and the other for the operator $\hat{E} : \mathcal{H}_E \otimes \mathcal{H}_{E^1}$; this is the kinematical Hilbert space (\mathcal{H}_{kin}) for the reduced model obtained representing the almost periodic function algebra (holonomy algebra for the homogeneous space-time) using the Bohr compactification of the real line.

A basis on this space is given by

$$|\mu_E, \mu_{E^1}\rangle \equiv |\mu_E\rangle \otimes |\mu_{E^1}\rangle \rightarrow \langle A | \mu_E \rangle \otimes \langle A_1 | \mu_{E^1} \rangle = e^{\frac{i\mu_E l_P A}{2}} \otimes e^{\frac{i\mu_{E^1} A_1}{\sqrt{2}}}. \quad (27)$$

The basis states (27) are normalizable in contrast to the standard quantum mechanical representation and they satisfy

$$\langle \mu_E, \mu_{E^1} | \nu_E, \nu_{E^1} \rangle = \delta_{\mu_E, \nu_E} \delta_{\mu_{E^1}, \nu_{E^1}} \quad (28)$$

We can quantize the theory using the standard quantization procedure $\{, \} \rightarrow -\frac{i}{\hbar} [,]$. We recall the fundamental difference from the standard Schrödinger quantization program. In the ordinary Schrödinger representation of the Weyl-Heisenberg algebra the classical fields A and A_1 translate in operators. In loop quantum gravity (or polymer representation of the Weyl algebra) on the contrary the operators A and A_1 don't exist. We can't promote the Poisson brackets to commutators $[\hat{A}, \hat{E}] = i\sqrt{l_P}$, $[\hat{A}_1, \hat{E}^1] = i\sqrt{l_P}/4$, but instead the quantum fundamental operators are $\hat{E}, \hat{E}^1, \hat{h}_I$.

The momentum operators can be represented on the Hilbert space by

$$\begin{aligned} \hat{E} &\rightarrow -i l_P \frac{d}{dA}, \\ \hat{E}^1 &\rightarrow -i \frac{l_P}{4} \frac{d}{dA_1}, \end{aligned} \quad (29)$$

It is very simply to calculate the spectrum of these two momentum operators on the Hilbert space basis, and so we obtain

$$\begin{aligned}\hat{E}|\mu_E, \mu_{E^1}\rangle &= \frac{\mu_E l_P^2}{2}|\mu_E, \mu_{E^1}\rangle, \\ \hat{E}^1|\mu_E, \mu_{E^1}\rangle &= \frac{\mu_{E^1} l_P}{4\sqrt{2}}|\mu_E, \mu_{E^1}\rangle.\end{aligned}\tag{30}$$

We have to fix the residual gauge freedom on the Hilbert space. We consider the operator $\hat{P} : |\mu_E, \mu_{E^1}\rangle \rightarrow |\mu_E, -\mu_{E^1}\rangle$ and we impose that only the invariant states (under \hat{P}) are in the kinematical Hilbert space. The states in the Hilbert space are of the form $\frac{1}{2} [|\mu_E, \mu_{E^1}\rangle + |\mu_E, -\mu_{E^1}\rangle]$.

In the following, we calculate the action of the volume, of the inverse volume and of the Schwarzschild curvature invariant operators on the kinematical Hilbert space, obtaining a bounded spectrum for these operators.

2.1 Inverse volume & $1/\sqrt{|E|}$ operators

In this section we study the black hole singularity problem in loop quantum gravity . We recall that the particular form of the volume for the space-time inside the horizon is $V = 4\sqrt{2} \pi l_P \sqrt{|E||E^1|}$.

The action of the volume operator on the basis states is

$$\hat{V}|\mu_E, \mu_{E^1}\rangle = \frac{4\pi R l_P^2}{\sqrt{2}} \sqrt{|\mu_E||\mu_{E^1}|} |\mu_E, \mu_{E^1}\rangle.\tag{31}$$

Now we will show that the operator $1/\sqrt{\hat{V}}$ doesn't diverge in quantum theory.

The operator version of the quantity $\text{sgn}(E)/\sqrt{\hat{V}}$ defined in the formula (25) is

$$\frac{\widehat{\text{sgn}(E)}}{\sqrt{\hat{V}}} = \frac{512 i}{3 l_P^4 \mu_0^3} \epsilon_{ijk} \sum_{IJK} \epsilon^{IJK} \text{Tr} \left[\tau^i \hat{h}_I^{-1} [\hat{h}_I, \hat{V}^{\frac{1}{2}}] \right] \text{Tr} \left[\tau^j \hat{h}_J^{-1} [\hat{h}_J, \hat{V}^{\frac{1}{2}}] \right] \text{Tr} \left[\tau^k \hat{h}_K^{-1} [\hat{h}_K, \hat{V}^{\frac{1}{2}}] \right].\tag{32}$$

At this point to obtain the spectrum of the inverse volume operator we express (32) using the $SU(2)$ identity in (15) and we arrive to the following form

$$\begin{aligned}\frac{\widehat{\text{sgn}(E)}}{\sqrt{\hat{V}}} &= -\frac{512 i}{3 l_P^4 \mu_0^3} \epsilon_{ijk} \sum_{IJK} \epsilon^{IJK} n_I^i n_J^j n_K^k \\ &\quad \left[\cos\left(\frac{A_I}{2}\right) \hat{V}^{\frac{1}{2}} \sin\left(\frac{A_I}{2}\right) - \sin\left(\frac{A_I}{2}\right) \hat{V}^{\frac{1}{2}} \cos\left(\frac{A_I}{2}\right) \right] \\ &\quad \left[\cos\left(\frac{A_J}{2}\right) \hat{V}^{\frac{1}{2}} \sin\left(\frac{A_J}{2}\right) - \sin\left(\frac{A_J}{2}\right) \hat{V}^{\frac{1}{2}} \cos\left(\frac{A_J}{2}\right) \right] \\ &\quad \left[\cos\left(\frac{A_K}{2}\right) \hat{V}^{\frac{1}{2}} \sin\left(\frac{A_K}{2}\right) - \sin\left(\frac{A_K}{2}\right) \hat{V}^{\frac{1}{2}} \cos\left(\frac{A_K}{2}\right) \right].\end{aligned}\tag{33}$$

All the operators in brackets commute with the volume operator and so the operator (33) is diagonal on the basis (27). Using the relation $\epsilon_{ijk} n_I^i n_J^j n_K^k = \epsilon_{IJK}$ it is straightforward to obtain the spectrum

on the Hilbert space basis

$$\begin{aligned} \frac{\widehat{\text{sgn}}(E)}{\sqrt{\widehat{V}}} |\mu_E, \mu_{E^1}\rangle &= \frac{8}{\sqrt{2} l_P \mu_0^3} |\mu_E|^{\frac{1}{2}} |\mu_{E^1}|^{\frac{1}{2}} \\ &\quad \left(|\mu_E + \mu_0|^{\frac{1}{4}} - |\mu_E - \mu_0|^{\frac{1}{4}} \right) \left(|\mu_{E^1} + \mu_0|^{\frac{1}{2}} - |\mu_{E^1} - \mu_0|^{\frac{1}{2}} \right)^2 |\mu_E, \mu_{E^1}\rangle. \end{aligned} \quad (34)$$

The spectrum of the inverse volume operator is upper bounded near the classical singularity which is in $E = 0$ or $\mu_E = 0$ and reproduce the correct classical spectrum of $1/\bar{V}$ for large volume eigenvalues ($\mu_E \sim \mu_{E^1} \gg 0$),

$$\frac{\widehat{\text{sgn}}(E)}{\sqrt{\widehat{V}}} |\mu_E, \mu_{E^1}\rangle \rightarrow \frac{2\sqrt{2}}{l_P} \frac{1}{|\mu_E|^{\frac{1}{4}} |\mu_{E^1}|^{\frac{1}{2}}} |\mu_E, \mu_{E^1}\rangle. \quad (35)$$

We see that the spectrum of the inverse volume operator is discrete. It is not divergent differing from the classical operator which has a singularity in $E = 0$ and reproduce the correct classical behavior for large values of the volume eigenvalues.

Now we pass to study the operator $1/\sqrt{|E|}$. The classical quantity defined in (26) becomes the operator

$$\begin{aligned} \frac{\widehat{\text{sgn}}(E) \widehat{\text{sgn}}(E^1)}{\sqrt{|E|}} &= -\frac{64}{\mu_0^2 l_P^3} \text{Tr} \left[\tau_3 \hat{h}_1^{-1} [\hat{h}_1, \hat{V}^{\frac{1}{2}}] \right] \text{Tr} \left[(\tau_2 + \tau_1) \hat{h}_2^{-1} [\hat{h}_2, \hat{V}^{\frac{1}{2}}] \right] \\ &= -\frac{64\sqrt{2}}{\mu_0^2 l_P^3} \left[\cos\left(\frac{A\mu_0 l_P}{2}\right) \hat{V}^{\frac{1}{2}} \sin\left(\frac{A\mu_0 l_P}{2}\right) \hat{V}^{\frac{1}{2}} - \sin\left(\frac{A\mu_0 l_P}{2}\right) \hat{V}^{\frac{1}{2}} \cos\left(\frac{A\mu_0 l_P}{2}\right) \hat{V}^{\frac{1}{2}} \right] \\ &\quad \left[\cos\left(\frac{A_1 \mu_0 l_P}{\sqrt{2}}\right) \hat{V}^{\frac{1}{2}} \sin\left(\frac{A_1 \mu_0 l_P}{\sqrt{2}}\right) \hat{V}^{\frac{1}{2}} - \sin\left(\frac{A_1 \mu_0 l_P}{\sqrt{2}}\right) \hat{V}^{\frac{1}{2}} \cos\left(\frac{A_1 \mu_0 l_P}{\sqrt{2}}\right) \hat{V}^{\frac{1}{2}} \right]. \end{aligned} \quad (36)$$

We consider the action of the operator (36) on the basis states of the Hilbert space and we obtain the spectrum of such operator

$$\begin{aligned} \frac{\widehat{\text{sgn}}(E) \widehat{\text{sgn}}(E^1)}{\sqrt{|E|}} |\mu_E, \mu_{E^1}\rangle &= \frac{\sqrt{2}}{\mu_0^2 l_P} |\mu_E|^{\frac{1}{4}} |\mu_{E^1}|^{\frac{1}{2}} \\ &\quad \left[|\mu_{E^1} + \mu_0|^{\frac{1}{2}} - |\mu_{E^1} - \mu_0|^{\frac{1}{2}} \right] \left[|\mu_E + \mu_0|^{\frac{1}{4}} - |\mu_E - \mu_0|^{\frac{1}{4}} \right] |\mu_E, \mu_{E^1}\rangle. \end{aligned} \quad (37)$$

The spectrum of $1/\sqrt{|E|}$ is well defines in $E = 0$, where the classical Schwarzschild singularity is localized.

If we consider the limite of large eigenvalues $\mu_E \rightarrow \infty$ and $\mu_{E^1} \rightarrow \infty$ we obtain

$$\frac{\widehat{\text{sgn}}(E) \widehat{\text{sgn}}(E^1)}{\sqrt{|E|}} |\mu_E, \mu_{E^1}\rangle \rightarrow \frac{\sqrt{2}}{l_P |\mu_E|^{\frac{1}{2}}} |\mu_E, \mu_{E^1}\rangle. \quad (38)$$

This is the correct classical value of $1/\sqrt{|E|}$.

In this section we have studied the operators relevant for the singularity analysis in quantum gravity and we have found that the singularity disappears and that the space-time can be extend beyond the classical singular point localized in $\mu_E = 0$.

2.2 Hamiltonian constraint and quantum dynamics

In this section we study the dynamics of the model. We want resolve the Hamiltonian constraint in the dual space of the Hilbert space.

The quantum version of the Hamiltonian constraint defined in (21) can be obtained promoting the classical holonomies to operators and the poisson bracket to the commutator

$$\hat{H}_E = \frac{8\pi i}{\mu_0^3 \hbar} \sum_{IJK} \epsilon^{IJK} \text{Tr} \left[\hat{h}_I \hat{h}_J \hat{h}_I^{-1} \hat{h}_J^{-1} \hat{h}_{[IJ]} \hat{h}_K^{-1} [\hat{h}_K, \hat{V}] \right]. \quad (39)$$

Using the relations in (15) we can express the Hamiltonian operator as

$$\begin{aligned} \hat{H}_E = & -\frac{8\pi i}{\mu_0^3 \hbar} \left[4 \sin(2x) \sin(2y) \left(\sin(y) \hat{V} \cos(y) - \cos(y) \hat{V} \sin(y) \right) + \right. \\ & \left. + 2 \left(\cos(\beta) \sin^2(2y) - \sin(\beta) \left(\sin^2(2y) + \cos(2y) \right) \right) \left(\sin(x) \hat{V} \cos(x) - \cos(x) \hat{V} \sin(x) \right) \right], \end{aligned} \quad (40)$$

where we have introduced the following notations: $x = A\mu_0 l_P/2$, $y = \sqrt{2}A_1\mu_0/2$ and $\beta = \mu_0^2/2$.

Before calculating the solution of the Hamiltonian constraint we must consider the action of the operator $\sin(\dots)$ and $\cos(\dots)$ on the Hilbert space for all possible trigonometric arguments which appear in the Hamiltonian constraint (40),

$$\begin{aligned} \cos(x)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2} \left(|\mu_E + \mu_0, \mu_{E^1}\rangle + |\mu_E - \mu_0, \mu_{E^1}\rangle \right) \\ \sin(x)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2i} \left(|\mu_E + \mu_0, \mu_{E^1}\rangle - |\mu_E - \mu_0, \mu_{E^1}\rangle \right) \\ \cos(y)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2} \left(|\mu_E, \mu_{E^1} + \mu_0\rangle + |\mu_E, \mu_{E^1} - \mu_0\rangle \right) \\ \sin(y)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2i} \left(|\mu_E, \mu_{E^1} + \mu_0\rangle - |\mu_E, \mu_{E^1} - \mu_0\rangle \right) \\ \sin(2x)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2i} \left(|\mu_E + 2\mu_0, \mu_{E^1}\rangle - |\mu_E - 2\mu_0, \mu_{E^1}\rangle \right) \\ \sin(2y)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2i} \left(|\mu_E, \mu_{E^1} + 2\mu_0\rangle - |\mu_E, \mu_{E^1} - 2\mu_0\rangle \right) \\ \cos(2y)|\mu_E, \mu_{E^1}\rangle &= \frac{1}{2} \left(|\mu_E, \mu_{E^1} + 2\mu_0\rangle + |\mu_E, \mu_{E^1} - 2\mu_0\rangle \right) \\ \sin^2(2y)|\mu_E, \mu_{E^1}\rangle &= -\frac{1}{4} \left(|\mu_E, \mu_{E^1} + 4\mu_0\rangle - 2|\mu_E, \mu_{E^1}\rangle + |\mu_E, \mu_{E^1} - 4\mu_0\rangle \right). \end{aligned} \quad (41)$$

Using all these ingredients we can calculate the action of the Hamiltonian constraint on the Hilbert space basis

$$\begin{aligned} \hat{H}_E|\mu_E, \mu_{E^1}\rangle &= \frac{8\pi^2 l_P^3}{\sqrt{2} \mu_0^3 \hbar} \left[-\alpha(\mu_E, \mu_{E^1}) \left(|\mu_E + 2\mu_0, \mu_{E^1} + 2\mu_0\rangle - |\mu_E - 2\mu_0, \mu_{E^1} + 2\mu_0\rangle \right) \right. \\ &\quad \left. - |\mu_E + 2\mu_0, \mu_{E^1} - 2\mu_0\rangle + |\mu_E - 2\mu_0, \mu_{E^1} - 2\mu_0\rangle \right) \\ &+ \beta(\mu_E, \mu_{E^1}) \left(\frac{\sin(\mu_0^2/2) - \cos(\mu_0^2/2)}{2} \right) \left(|\mu_E, \mu_{E^1} + 4\mu_0\rangle - 2|\mu_E, \mu_{E^1}\rangle + |\mu_E, \mu_{E^1} - 4\mu_0\rangle \right) \\ &\quad \left. - \beta(\mu_E, \mu_{E^1}) \sin(\mu_0^2/2) \left(|\mu_E, \mu_{E^1} + 2\mu_0\rangle + |\mu_E, \mu_{E^1} - 2\mu_0\rangle \right) \right] \end{aligned} \quad (42)$$

At this point we can resolve the Hamiltonian constraint. The solutions of the Hamiltonian constraint are in the \mathcal{C}^* space that is the dual of the dense subspace \mathcal{C} of the kinematical space. A generic element of this space is

$$\langle \psi | = \sum_{\mu_E, \mu_{E^1}} \psi(\mu_E, \mu_{E^1}) \langle \mu_E, \mu_{E^1} |. \quad (43)$$

The constraint equation $\hat{H}|\psi\rangle = 0$ is now interpreted as an equation in the dual space $\langle \psi | \hat{H}^\dagger$; from this equation we can derive a relation for the coefficients $\psi(\mu_E, \mu_{E^1})$

$$\begin{aligned} & -\alpha(\mu_E - 2\mu_0, \mu_{E^1} - 2\mu_0) \psi(\mu_E - 2\mu_0, \mu_{E^1} - 2\mu_0) + \alpha(\mu_E + 2\mu_0, \mu_{E^1} - 2\mu_0) \psi(\mu_E + 2\mu_0, \mu_{E^1} - 2\mu_0) \\ & + \alpha(\mu_E - 2\mu_0, \mu_{E^1} + 2\mu_0) \psi(\mu_E - 2\mu_0, \mu_{E^1} + 2\mu_0) - \alpha(\mu_E + 2\mu_0, \mu_{E^1} + 2\mu_0) \psi(\mu_E + 2\mu_0, \mu_{E^1} + 2\mu_0) \\ & + \left(\frac{\sin(\mu_0^2/2) - \cos(\mu_0^2/2)}{2} \right) \left(\beta(\mu_E, \mu_{E^1} - 4\mu_0) \psi(\mu_E, \mu_{E^1} - 4\mu_0) - \beta(\mu_E, \mu_{E^1}) \psi(\mu_E, \mu_{E^1}) \right. \\ & \quad \left. + \beta(\mu_E, \mu_{E^1} + 4\mu_0) \psi(\mu_E, \mu_{E^1} + 4\mu_0) \right) \\ & - \sin(\mu_0^2/2) \left(\beta(\mu_E, \mu_{E^1} - 2\mu_0) \psi(\mu_E, \mu_{E^1} - 2\mu_0) + \beta(\mu_E, \mu_{E^1} + 2\mu_0) \psi(\mu_E, \mu_{E^1} + 2\mu_0) \right) = 0, \quad (44) \end{aligned}$$

where the functions $\alpha(\mu_E, \mu_{E^1})$ and $\beta(\mu_E, \mu_{E^1})$ are define by

$$\begin{aligned} \alpha(\mu_E, \mu_{E^1}) & \equiv |\mu_E|^{\frac{1}{2}} (|\mu_{E^1} + \mu_0| - |\mu_{E^1} - \mu_0|) \\ \beta(\mu_E, \mu_{E^1}) & \equiv |\mu_{E^1}| \left(|\mu_E + \mu_0|^{\frac{1}{2}} - |\mu_E - \mu_0|^{\frac{1}{2}} \right) \end{aligned} \quad (45)$$

How can be seen from equation (44) the quantization program produce a difference equation and imposing a boundary condition we can obtain the wave function $\psi(\mu_E, \mu_{E^1})$ for the black hole. We can interpret $\psi(\mu_E, \mu_{E^1})$ as the wave function of the anisotropy “ E^1 ” at the time “ E ”. As in loop quantum cosmology also in this case the state $\psi(0, 0)$ decouples from the dynamics.

Conclusions

In this paper we studied the Kantowski-Sachs space time in Ashtekar variables. We started from the complete “loop quantum gravity” and we reduced to the space-time of spatial topology $\mathbf{R} \times \mathbf{S}^2$. This space-time is very important if we want to understand what happens very closed to the black hole singularity where the quantum gravity effects are dominant and the classical Schwarzschild solution it isn’t correct. In particular we concentrated on the Schwarzschild black hole solution inside the horizon where the time exchanges with space and the model reduces to a cosmological type space-time: a general space-time with this topology is called a Kantowski-Sachs space-time.

The quantization procedure is induced by the full “loop quantum gravity”. We have introduced holonomies for the homogeneous space-time and we have expressed the inverse volume operator, the classically singular quantity $1/\sqrt{|E|}$ and the Euclidean part of the Hamiltonian constraint in terms of holonomies. All these operators are well defined on the kinematical Hilbert space.

The main results are :

1. the inverse volume operator has a finite spectrum in all the region inside the horizon and we can conclude that the classical singularity disappears in “loop quantum gravity”; for large eigenvalues of the volume operator on the other side we find the classical inverse volume behaviour,
2. the quantity $1/\sqrt{|E|}$, connected to the Schwarzschild curvature invariant $R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \sim 1/b(t)^6$ by $b(t)^2 = |E|$, is singularity free in quantum gravity and reproduces the classical behaviour for large eigenvalues,

3. the solution of the Hamiltonian constraint gives a discrete difference equation for the coefficients of the physical states which are defined in the dual space of the dense subspace of the kinematical Hilbert space.

An important consequence of the quantization is that, unlike the classical evolution, the quantum evolution doesn't stop at the classical singularity and the "other side" of the singularity corresponds to a new domain where the triad reverses its orientation. This work is useful if we want understand what is the mechanism to resolve the problem of the "information loss" in the process of black hole formation [16].

Acknowledgements

We are grateful to Carlo Rovelli, Eugenio Bianchi and Guido Cossu for many important and clarifying discussion. This work is supported in part by a grant from the Fondazione Angelo Della Riccia.

Appendix

We report some technical properties extensively used in this paper and the independent components of the curvature F_{ab} for the Kantowski-Sachs space time.

The $SU(2)$ generators used in the paper are $\tau^i = -\frac{i}{2}\sigma^i$, where σ^i are the Pauli matrices

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (46)$$

Other important properties of the Pauli matrix and of the $SU(2)$ generators τ^i are

$$\begin{aligned} [\sigma^i, \sigma^j] &= 2i\epsilon^{ijk}\sigma^k, \\ \sigma^i\sigma^j &= \delta^{ij} + i\epsilon^{ijk}\sigma^k, \\ [\tau^i, \tau^j] &= 2i\epsilon^{ijk}\tau^k, \\ \text{Tr}[\tau^i\tau^j] &= -\frac{1}{2}\delta^{ij}, \\ \text{Tr}[\tau^i\tau^j\tau^k] &= -\frac{1}{4}\epsilon^{ijk}. \end{aligned} \quad (47)$$

The field strenght F_{ab} independent components are

$$\begin{aligned} F_{12} &= AA_1(\tau_2 - \tau_1), \\ F_{13} &= -AA_1(\tau_2 + \tau_1)\sin\theta, \\ F_{23} &= (2A_1^2 - 1)\tau_3\sin\theta. \end{aligned} \quad (48)$$

References

- [1] Abhay Ashtekar, *New Hamiltonian formulation of general relativity*, Phys. Rev. D **36** 1587-1602
- [2] Carlo Rovelli, *Quantum Gravity*, (Cambridge University Press, Cambridge, 2004)
- [3] Leonardo Modesto, *Disappearance of the black hole singularity in loop quantum gravity*, Phys. Rev. D **70** (2004) 124009, gr-qc/0407097
- [4] Leonardo Modesto, *The kantowski-Sachs space-time in loop quantum gravity*, gr-qc/0411032
- [5] M. Bojowald, *Inverse scale factor in isotropic quantum geometry*, Phys. Rev. **D64** 084018 (2001); M. Bojowald, *Loop Quantum Cosmology IV: discrete time evolution*, Class. Quant. Grav. **18**, 1071 (2001); A. Ashtekar, M. Bojowald and J. Lewandowski, *Mathematica structure of loop quantum cosmology*, Adv. Theor. Math. Phys. **7** (2003) 233-268, gr-qc/0304074
- [6] R. Kantowski and R. K. Sachs, J. Math. Phys. **7** (3) (1966)
- [7] Ioannis Raptis, *Finitary-Algebraic ‘Resolution’ of the Inner Schwarzschild Singularity*, gr-qc/0408045; Anastasios Mallios and Ioannis Raptis, *Smooth Singularities Exposed: Chimeras of the Differential Spacetime Manifold*, gr-qc/041112; Ioannis Raptis, *‘Iconoclastic’, Categorical Quantum Gravity*, gr-qc/0509089
- [8] Igemar Bengtsson, *Note on Ashtekar’s variables in the spherically symmetric case*, Class. Quant. Grav. **5** (1988) L139-L142; Igemar Bengtsson, *A new phase for general relativity?*, Class. Quant. Grav. **7** (1990) 27-39; H.A. Kastrup & T. Thiemann, *Spherically symmetric gravity as a complete integrable system*, Nucl. Phys. B **425** (1994) 665-686, gr-qc/9401032; M. Bojowald & H.A. Kastrup, *Quantum symmetry reduction of Diffeomorphism invariant theories of connections*, JHEP **0002** (2000) 030, hep-th/9907041; M. Bojowald, *Spherically symmetric quantum geometry: states and bases operators*, Class. Quant. Grav. **21** (2004) 3733-3753, gr-qc/0407017;
- [9] Luca Bombelli & R. J. Torrence *Perfect fluids and Ashtekar variables, with application to Kantowski-Sachs models* Class. Quant. Grav. **7** (1990) 1747-1745
- [10] A. Ashtekar, S. Fairhurst and J. Willis, *Quantum gravity, shadow states, and quantum mechanics*, Class. Quant. Grav. **20** 1031-1062 (2003)
- [11] C Rovelli, “Partial observables”, Phy Rev D **65** (2002) 124013; gr-qc/0110035
- [12] Viqar Husain and Oliver Winkler, *On singularity resolution in quantum gravity* ; gr-qc/0312094
- [13] H. Halvorson, *Complementary of representations in quantum mechanics*, Studies in History and Philosophy of Modern Physics **35**, 45-56 (2004), quant - ph/0110102
- [14] T. Thiemann, *Quantum Spin Dynamics*, Class. Quant. Grav. **15**, 839 (1998)
- [15] T. Thiemann, *Introduction to Modern Canonical Quantum General Relativity* ; gr -qc/0110034; *Lectures on Loop Quantum Gravity*, gr-qc/0210094
- [16] Abhay Ashtekar & Martin Bojowald, *Black hole evaporation : A paradigm* Class. Quant. Grav. **22** (2005) 3349-3362, gr-qc/0504029