



**HAL**  
open science

## **Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils**

Camille Dumat, K. Quenea, A. Bermond, S. Toinen, M.F. Benedetti

### ► **To cite this version:**

Camille Dumat, K. Quenea, A. Bermond, S. Toinen, M.F. Benedetti. Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils. *Environmental Pollution*, 2006, 142 (3), pp.521-529. <10.1016/j.envpol.2005.10.027>. <hal-00481120>

**HAL Id: hal-00481120**

**<https://hal.science/hal-00481120v1>**

Submitted on 2 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils

C. Dumat<sup>a,b,\*</sup>, K. Quenea<sup>c</sup>, A. Bermond<sup>c</sup>, S. Toinen<sup>a</sup>, M.F. Benedetti<sup>d</sup>

<sup>a</sup> Laboratoire de Biogéochimie des Milieux Continentaux, UMR CNRS 7618, UPMC, 75252 Paris cedex 05 France

<sup>b</sup> Laboratoire Environnement Ecotoxicologie, EA 3715 INP-ENSAT, Paris, France

<sup>c</sup> INA-PG, 16 rue Claude Bernard, 75231 Paris Cedex 05, France

<sup>d</sup> Lab. de Géochimie des Eaux, Univ. Denis Diderot, UMR CNRS 7047, Paris, France

*Measure of  $\delta^{13}C$  and total metal concentrations in size fractions of contaminated soils suggests an influence of metals on the soil organic matter dynamic.*

## Abstract

The role of metals in the behaviour of soil organic matter (SOM) is not well documented. Therefore, we investigated the influence of metals (Pb, Zn, Cu and Cd) on the dynamic of SOM in contaminated soils where maize (C<sub>4</sub> plant) replaced C<sub>3</sub> cultures. Three pseudogley brown leached soil profiles under maize with a decreasing gradient in metals concentrations were sampled. On size fractions, stable carbon isotopic ratio ( $\delta^{13}C$ ), metals, organic carbon and nitrogen concentrations were measured in function of depth. The determined sequence for the amount of C<sub>4</sub> organic matter in the bulk fractions: M<sub>3</sub> (0.9) > M<sub>2</sub> (0.4) > M<sub>1</sub> (0.3) is in agreement with a significant influence of metals on the SOM turnover. New C<sub>4</sub> SOM, mainly present in the labile coarser fractions and less contaminated by metals than the stabilised C<sub>3</sub> SOM of the clay fraction, is more easily degraded by microorganisms.

*Keywords:* Soil organic matter dynamic; Trace metal; Size fractionation; Stable isotopic carbon ratio

## 1. Introduction

With respect to the important environmental functions of soil organic matter (SOM): potential source or sink for greenhouse gases, contribution to soil aggregation, influence on soil fertility and bioactivity (Tiessen et al., 1994; Vanlauwe et al., 1999), the impact on its dynamics of some parameters like soil characteristics (Krull and Skjemstad, 2003), climate (Hevia et al., 2003) and agricultural practices (Pinheiro et al., 2004; Beare et al., 1994) are studied in detail.

Due to various anthropogenic activities, potentially toxic metals are accumulated in soils, with a risk of biota and

groundwater contamination. Among the various reactive soil constituents, SOM has a large sorption capacity towards metal ions (Benedetti et al., 1996; Sauve et al., 2000; Yin et al., 2002) and therefore metals often accumulate in the organic topsoils. Metals influence the soil biota in its diversity, abundance and activity (Mhatre and Pankhurst, 1997; Barajas-Aceves et al., 1999). But nevertheless, their impact on the SOM turnover is not well documented.

SOM is composed of various organic constituents chemically heterogeneous (materials in various stages of decomposition for instance) with a large range of reactivity and turnover time in soil (Rethemeyer et al., 2004). In the literature, the affinity of metal ions towards humic substances is well described, but the reactivity with plant debris or bio-polymers is less documented (Spark et al., 1997). Moreover, the reactive components of the soil are in interaction (Franzluebbers and

\* Corresponding author. Tel.: +33 5 59 77 30 53; fax: +33 5 62 19 39 01.  
E-mail address: dumat@ccr.jussieu.fr (C. Dumat).

Arshad, 1997; Dumat et al., 1997). When plant litter compounds enter the soil, soil-inherent stabilisation mechanisms such as inclusion into aggregates or protection due to the interaction with soil minerals are operative (Sollins et al., 1996; Kaiser and Guggenberger, 2000). The interaction of SOM with the mineral surfaces may be responsible for the occurrence of a stable carbon pool in the fine clay fraction (Baldock and Skjemstad, 2000). Therefore the studies concerning the affinity of one extracted soil component for metals are open to criticism. Conversely, methods of soil particle-size fractionation can be used to assess the distribution of trace elements in soils (Ducaroir and Lamy, 1995). These size fractions can inform on the soil biogeochemistry and the evolution with time of soil components. Changes in the natural abundance of  $^{13}\text{C}$  in SOM over time can be used to identify sources of the organic C in soil (Schwartz et al., 1996) and to determine the turnover rate of SOM (Balesdent and Mariotti, 1996).

Three pseudogley brown leached soils under maize (the primary  $\text{C}_4$  plants cultivated in temperate countries), with a decreasing gradient in total metal concentrations were collected near a smelter plant. On samples collected at various depths, size fractions were obtained and the following measurements were done: metal ions, organic carbon and nitrogen concentrations and  $\delta^{13}\text{C}$ . The aim of the present work was to determine the effect of metals on the dynamics of SOM by comparing the patterns of the  $\delta^{13}\text{C}$  signals for similar soils where maize replaced the previous  $\text{C}_3$  vegetation and differing by their metal concentration levels.

## 2. Materials and methods

### 2.1. Soils characteristics and management

The 100 year old smelter (Metaleurop) site is located near Evin-Malmaison (North of France). Dust emissions with high contents of Pb, Zn, Cd and Cu have contaminated the nearby agricultural field over tens of  $\text{km}^2$ . Dust particles were characterised by Sobanska et al. (1999) and consist mainly of sulphide and oxide metal ions. The soils classified as pseudogley brown leached soils (Morin et al., 1999) develop over quaternary clays (0.6–1.5 m) covering the Ostrincourt sand. Three soils were collected:  $\text{M}_1$ , continuous maize (*Zea mays* L.) since 10 years and 1 km away from the smelter;  $\text{M}_2$ , continuous maize since 14 years and 3 km away from the smelter and  $\text{M}_3$ , continuous maize since 13 years and 10 km away from the smelter. The field sampling has been oriented to collect representative soil samples weighing 2–40 kg in closed bags. After mixing, all the samples were air dried and sieved at 2 mm.

For these three NPK fertilised soils, conventional tillage (ploughing to 30 cm, followed by superficial tillage and sowing) was performed and liming was done every few years. In preliminary experiments we checked that soil characteristics were homogeneous in the first 30 cm. Tillage favours the SOM mineralisation, because it breaks soil aggregates, increases aeration and thereby exposes SOM, which was previously protected within the aggregate structure (Pinheiro et al., 2004; Beare et al., 1994; Chenu, 1995). Arrouays et al. (1995) observed an increase of the ratio between labile and recalcitrant SOM when a soil under forest was cultivated.

For around 80 000 plants  $\text{ha}^{-1}$  cultivated on the three soils ( $\text{M}_1$ ,  $\text{M}_2$  and  $\text{M}_3$ ), the maize yield is 75 000–80 000  $\text{kg ha}^{-1}$ . Before maize, previous cultivated  $\text{C}_3$  plants were potatoes (50 000  $\text{kg ha}^{-1}$ ), peas (1200  $\text{kg ha}^{-1}$ ) and beets (70 000  $\text{kg ha}^{-1}$ ). The type of land use is an important factor controlling organic matter storage in soils since it affects the amount and quality of litter input, the litter decomposition rates and the processes of organic matter stabilisation in soils (Shepherd et al., 2001; John et al., 2005).

One objective of the present work was to study soils contaminated by a real old industrial activity. Such long term continuous  $\text{C}_3$  and then  $\text{C}_4$  monoculture soils with similar characteristics (texture, land-use history, etc.) and differing only by metals concentrations are difficult to find. Therefore, it should be noted that soil  $\text{M}_1$  is under maize monoculture only since 10 years. Moreover, for the fodder maize cultivated soil  $\text{M}_1$ , only maize roots were incorporated in the soil, whereas in  $\text{M}_2$  and  $\text{M}_3$ , additionally the maize straw was incorporated in order to improve the soil characteristics.

### 2.2. Physical fractionation of the soil samples

Before particle size fractionation, an initial physical dispersion of the soil samples was performed by end-to-end shaking overnight in pure de-ionised water. The suspensions were then sieved through a series of grids (AFNOR French reference) with decreasing mesh size (2000, 1000, 500, 200 and 50  $\mu\text{m}$ ). Then the fractions (20  $\mu\text{m}$  < size < 50  $\mu\text{m}$ ), (2  $\mu\text{m}$  < size < 20  $\mu\text{m}$ ) and <2  $\mu\text{m}$  were obtained by sedimentation. All the recovered particles were air dried at 60 °C, weighed, ground (200  $\mu\text{m}$ ) and homogenised. Particle size fractionation of soil samples resulted in the recovery of  $98 \pm 0.5\%$  of the initial sample mass.

### 2.3. Measure of total metal concentrations

The total concentrations of Pb, Cd, Zn and Cu in the size fractions of the soil samples were analysed by inductively coupled plasma atomic emission spectrometry (ICP-AES) on a Jobin Yvon JYESS 4.03 apparatus, after a tri-acid (HF, HCl and  $\text{HNO}_3$ ) digestion of the solid samples. A standard reference soil (GRX2) was used to control the digestion protocol and the measurements accuracies. All the results correspond to the mean of two replicates.

### 2.4. Organic carbon amounts and stable carbon isotopic signals ( $\delta^{13}\text{C}$ )

All terrestrial plants discriminate against  $^{13}\text{C}$  (in favour of  $^{12}\text{C}$ ) during photosynthesis, inducing differences between atmospheric  $\text{CO}_2$  and the carbon isotopic composition of the plant tissues (Smith and Epstein, 1971). The degree of discrimination is dependant on the photosynthetic pathway of the plant (Ehleringer et al., 1997). It increases from  $\text{C}_4$  plants like maize (the metabolic pathway forms initial molecules with four carbon atoms) to  $\text{C}_3$  plants. In natural ecosystems, the fractionation of the stable carbon during decomposition of the organic matter is small relative to the original fractionation during C fixation by plants (Peterson and Fry, 1987). Therefore in soils where the vegetation changes from  $\text{C}_3$  to  $\text{C}_4$  (or vice versa), the changes in the natural abundance of  $^{13}\text{C}$  in SOM over time can be used to identify sources of organic C in the soil (Schwartz et al., 1996) or to determine the turnover rate of SOM (Balesdent and Mariotti, 1996).

When  $\text{C}_4$  plants grow on a soil that had always received  $\text{C}_3$  plants, the  $\delta^{13}\text{C}$  value of the SOM can be relevant to the proportion of the carbon from  $\text{C}_3$  and  $\text{C}_4$  plants to study the dynamic of the SOM (Balesdent and Mariotti, 1996). The effect of such a vegetation change on SOM isotope composition can be described as follows, with the indices 4 and 3 corresponding to the different photosynthetic pathway types. At the time of vegetation change ( $t_0 = 10, 13$  and 14 years ago for  $\text{M}_1, \text{M}_3$  and  $\text{M}_2$ , respectively), SOM have an isotopic composition  $\delta_{3-0}$ , which is close to the composition of the original vegetation  $\text{C}_3$ . SOM progressively decays and is partially replaced by SOM derived from the new vegetation  $\text{C}_4$ . At a given time  $t$ , the total carbon content ( $C$ ) can be expressed as:  $C = C_3 + C_4$ . The isotopic composition of SOM ( $\delta$ ) is given by the following mixing equation:  $\delta(C_3 + C_4) = \delta_3 C_3 + \delta_4 C_4$  (with the approximation that the  $\delta$  unit is related linearly to  $^{13}\text{C}/^{12}\text{C}$ ). Where  $C_3$  stands for the amount of carbon derived from the  $\text{C}_3$  old vegetation,  $\delta_3$  for the isotopic composition of that carbon,  $C_4$  for the amount of carbon derived from the new vegetation  $\text{C}_4$  (maize), and  $\delta_4$  for its isotopic composition. The fraction of new carbon ( $C_4$ ) in the sample ( $F$ ) =  $C_4/C = (\delta - \delta_3)/(\delta_4 - \delta_3)$ . The function  $F(t)$  can be obtained from successive samples taken on the same site. The kinetics of decay of the initial organic carbon ( $C_3$ ) is equivalent to  $C(1 - F)$ , whereas the kinetics of accumulation of the new carbon is equivalent to  $CF$ . Because the  $\delta$  of the soil sample is linearly related to  $F$ , its change with time gives a direct expression of the turnover of soil carbon.  $\delta_3$  and  $\delta_4$  cannot be measured directly and must be estimated.

As there is evidence that the isotopic composition of SOM is only slightly different from that of the plant material from which it derives, most investigators using the natural  $^{13}\text{C}$ -labelling technique assume  $\delta_4$  is equivalent to the isotopic composition of the new vegetation  $C_4$  or its litter ( $\delta_{\text{veg}C_4}$ ), and  $\delta_3$  is equivalent to the initial  $\delta^{13}\text{C}$  of the soil (at  $t_0$ ) or to the isotopic composition of soil at a reference site kept under the initial vegetation ( $\delta_{\text{ref}C_3}$ ). Under these assumptions:  $F = (\delta - \delta_{\text{ref}C_3})/(\delta_{\text{veg}C_4} - \delta_{\text{ref}C_3}) = C_4/C$ .

Organic C and N contents of soil samples were determined by dry combustion in a Carlo Erba NA 1500 CHN elemental analyser. Stable carbon isotope ratios of SOM and plants were measured on a Fisons SIRA 10 Isotope Ratio Mass Spectrometer.  $^{13}\text{C}$  natural abundance was expressed in  $\delta$  units, by reference to the international standard PDB (marine carbonate: belemnite from the PEE Dee formation), according to the following equation:

$$\delta^{13}\text{C}_{\text{‰}} = \frac{[^{13}\text{C}/^{12}\text{C}]_{\text{sample}} - [^{13}\text{C}/^{12}\text{C}]_{\text{PDB}}}{[^{13}\text{C}/^{12}\text{C}]_{\text{PDB}}} \times 1000$$

All samples were sieved to 1 mm, ground to 100  $\mu\text{m}$  and homogenised in order to minimise the effects of sample heterogeneity. Organic carbon, nitrogen and  $\delta^{13}\text{C}$  of the soil samples were measured in duplicate, with previous elimination of carbonates, by acidification. The  $^{13}\text{C}$  natural labelling technique requires information on  $^{13}\text{C}$  abundance of all plant materials entering the soil. Plants samples associated with the soils were collected, crushed, homogenised and analysed to obtain reference values for the stable ( $^{13}\text{C}/^{12}\text{C}$ ) carbon isotopic analysis.

### 3. Results and discussion

#### 3.1. General characteristics of the soil samples

Trace metals and organic carbon concentrations of the bulk samples (<2 mm) for each soil profile are reported in Table 1. All the results are given as the mean of the two replicates for each sample. The relative standard deviation of all the measured values never exceeded 7%. The organic carbon is mainly located in the topsoils (plough horizons), where most of the metals occur. The values for Pb, Zn, Cd and Cu concentrations are sometimes higher than the legal French limit NFU 744-041 (100, 300, 2 and 100 ppm for Pb, Zn, Cd and Cu, respectively). The depth versus concentration profiles depend on the metal. For Pb and Cu, below 25 cm, the metal concentrations are close to the geological background concentrations of the area estimated (Sterckeman et al., 2000). At the opposite,

Table 1  
Trace metals concentrations ( $\text{mg kg}^{-1}$ ) and organic carbon ( $\text{g kg}^{-1}$ ) of the bulk samples (fraction <2 mm) for each soil ( $M_1$ ,  $M_2$  and  $M_3$ )

Depth (cm)	[Pb] ( $\text{mg kg}^{-1}$ )	[Zn] ( $\text{mg kg}^{-1}$ )	[Cu] ( $\text{mg kg}^{-1}$ )	[Cd] ( $\text{mg kg}^{-1}$ )	C ( $\text{g kg}^{-1}$ )
Soil $M_1$					
0–30	503	852	34	16.8	15
30–35	216	280	18	9.8	9
35–40	118	175	13	6	5
55–65	71	41	4.6	2.1	2.4
170–180	33	39	4.15	1.8	1.15
Soil $M_2$					
0–30	187	172.5	17.9	12.05	27
30–40	145	133	13.1	9.3	19
40–50	62.3	39	9.8	5.1	11
50–65	38.3	29	7.9	3.3	8.5
Soil $M_3$					
0–30	54.5	64.2	13.2	3.5	25

for the more mobile metal ions (Zn and Cd), the enhanced concentrations are found down to 50 cm.

As reported in Table 2, the  $\text{pH}_{\text{water}}$  (between 6.4 for  $M_3$  and 7.5 for  $M_1$ ) and the CEC (between  $13.9 \text{ C mol}^+ \text{ kg}^{-1}$  for  $M_1$  and  $14.7$  for  $M_3$ ) are quite similar for  $M_1$ ,  $M_2$  and  $M_3$ , whereas the concentrations of the metals are significantly different and follow the sequence:  $M_1 > M_2 > M_3$ . Between 1.5% (for  $M_1$ ) and 2.7 (for  $M_2$ ), accordingly with John et al., 2005, the organic matter amount in the plough horizons is lower for the soil  $M_1$  where the straws are exported.

The size repartition as a function of depth for the three soils is reported in Table 3 (only the 0–30 cm depth was the studied for the soil  $M_3$ ). Soil texture is quite similar for  $M_1$ ,  $M_2$  and  $M_3$ : clay (<2  $\mu\text{m}$ ) and silt particles (2  $\mu\text{m}$  < fine silt <20  $\mu\text{m}$  and 20  $\mu\text{m}$  < coarse silt < 50  $\mu\text{m}$ ) dominate and that texture is relatively constant with the depth. But the soil  $M_2$  is significantly less clayed (9.1%) than the two others soils  $M_1$  (25.2%) and  $M_3$  (28.2%). According to Oades (1995), a high amount of clay decreases the biodegradation of SOM and therefore increases the stock of carbon. Muller and Hoper (2004) concluded that the capacity of soil to protect organic matter against microbial decomposition depends on its clay content. Soil clay content is an abiotic factor defining the size of protected pools of organic matter in models of SOM turnover (Parton, 1996). Therefore, the protection of SOM could be lower for  $M_2$  in comparison with  $M_1$  and  $M_3$ , and induces a higher turnover of the SOM in that soil.

#### 3.2. The $\delta^{13}\text{C}$ constraints

Fig. 1 gives the  $\delta^{13}\text{C}$  of the topsoils (0–30 cm)  $M_1$ ,  $M_2$  and  $M_3$  as a function of the size fraction. For the cultivated soils under maize  $M_1$  (10 years),  $M_2$  (14 years) and  $M_3$  (13 years), the  $\delta^{13}\text{C}$  of the top bulk samples are respectively:  $-22.8\text{‰}$ ;  $-22.3\text{‰}$  and  $-15.3\text{‰}$ . These data were compared with the  $\delta^{13}\text{C}$  signals measured for plants sampled in the field and reference  $\delta^{13}\text{C}$  values (Balesdent and Mariotti, 1996; Boutton, 1996) of  $C_3$  and  $C_4$  plant tissues and actual  $\text{CO}_2$  reported in Table 4. First, the measure of  $\delta^{13}\text{C}$  informs on the origin of the organic matter: a variable part of the organic matter of these soils comes from the incorporation of maize residues, and old  $C_3$  organic matter is still remains.

In agreement with previous studies, a decrease of the  $\delta^{13}\text{C}$  with size fraction is observed in relation to an increase of old SOM from  $C_3$  origin. For 13 years of maize cultivation on an uncontaminated soil, Balesdent and Mariotti (1996) calculated turnover times of  $4 \pm 1.5$  years and  $13 \pm 2$  years, respectively, for the 2000–200  $\mu\text{m}$  and the 200–50  $\mu\text{m}$  fractions. In contrast, all fractions smaller than 10  $\mu\text{m}$  exhibited an annual

Table 2  
Cation exchange capacity (CEC) and  $\text{pH}_{\text{water}}$  of the topsoil bulk fractions

	$M_1$	$M_2$	$M_3$
CEC <sup>a</sup> ( $\text{C mol}^+ \text{ kg}^{-1}$ )	13.9	14.1	14.7
$\text{pH}_{\text{water}}$	7.5	6.8	6.4

<sup>a</sup> The cation exchange capacity was determined by the ammonium acetate method at pH 7 (Metson, 1956).

Table 3  
Size repartitions (%) for each soil ( $M_1$ ,  $M_2$  and  $M_3$ ; sizes are expressed in  $\mu\text{m}$ )

Depth (cm)	2000 > x > 1000	1000 > x > 500	500 > x > 200	200 > x > 50	50 > x > 20	20 > x > 2	2 $\mu\text{m}$ > x
$M_1$							
0–30	0.7	1.2	5.4	8.0	17.1	41.0	25.2
30–35	1.0	1.1	25.7	4.2	19.4	26.3	22.1
35–40	0.7	2.1	7.0	15.2	12.7	32.9	29.3
55–65	1.8	4.4	4.2	23.8	21.2	22.8	21.7
170–180	0.5	1.9	14.9	30.2	30.3	3.4	18.6
$M_2$							
0–30	2.30	3.70	10.00	23.30	16.40	32.30	9.10
30–40	0.70	1.60	17.50	26.00	19.50	25.60	8.80
40–50	0.40	0.60	3.00	18.60	28.10	39.70	8.30
50–62	1.20	3.70	1.30	22.60	26.70	29.90	13.00
$M_3$							
0–30	0.90	2.00	4.60	14.80	30.40	19.20	28.20

rate of new organic matter input equal to  $1.3 \pm 0.4\%$  of the carbon amount in the fraction. Monreal et al. (1997) reported in a long-term experiment with continuous maize cultivation that the mean turnover time of soil organic carbon was 14 years in macroaggregates, 61 years in microaggregates and 275 years in the fraction  $<50 \mu\text{m}$ .

The  $C_4/C$  ratios were calculated from the  $\delta^{13}\text{C}$  values measured (as described in Section 2):  $C_4/C = (\delta - \delta_{\text{ref}C_3}) / (\delta_{\text{veg}C_4} - \delta_{\text{ref}C_3})$  with  $\delta_{\text{ref}C_3}$  fixed to  $-27\text{‰}$  and  $\delta_{\text{veg}C_4}$  fixed to  $-14\text{‰}$ . The results are reported in Fig. 2:  $C_4/C$  as a function of the size fraction for the topsoils  $M_1$ ,  $M_2$  and  $M_3$ . From that curve we can conclude that maize replaced  $C_3$  vegetation in various proportions on these three sites. The amount of  $C_4$  organic matter in the bulk fractions of the soils under maize follows the sequence  $M_3$  ( $C_4/C = 0.9$ )  $>$   $M_2$  ( $0.4$ )  $>$   $M_1$  ( $0.3$ ). A low  $C_4/C$  ratio (in comparison with the value 1) corresponds to a situation where a high amount of old ( $C_3$ ) soil organic matter is always present. The lowest value of the  $C_4/C$  ratio for the contaminated soils ( $M_1$  and  $M_2$ ) in comparison with the uncontaminated soil  $M_3$  could be due to the influence of metals on the SOM turnover.

### 3.3. Organic carbon amounts for the soils as a function of depth and size fraction

Values of organic carbon of particle-size fractions as a function of depth are reported in Table 5 for  $M_1$ ,  $M_2$  and  $M_3$  soil. The organic carbon concentration decreases quickly with the depth: SOM is mainly concentrated in the first centimetres (0–30 cm). For the soil  $M_1$ , between 0 and 180 cm depth, the clay fraction presents the highest organic carbon concentration: the ratio between clay and bulk fractions concentrations is between 1.1 and 2.5. The C/N ratio measured for  $M_1$  decreases with depth (between 12.5 at 0–30 cm and 7.7 at 170–180 cm for the bulk fraction) and size fraction; this trend is in agreement with a change in chemical nature of SOM (Besnard et al., 2001). For the soil  $M_2$ , between 0 and 60 cm depth, the 1000–500  $\mu\text{m}$  and the clay fractions have similar organic carbon concentrations, around two or three times more than in the bulk fraction. For the topsoil  $M_3$ , the clay fraction presents the highest organic carbon concentration (around 2.5 times in comparison with the bulk) and then the concentration in the 1000–200  $\mu\text{m}$  fraction is around two times higher in comparison with the bulk concentration.

### 3.4. Trace metal amounts for the soils as a function of depth and size fraction

The total metal concentrations as a function of depth and size fractions are given in Table 6 for the soils  $M_1$ ,  $M_2$  and  $M_3$ . For the soil  $M_1$ , whatever the depth and the nature of




Fig. 1.  $\delta^{13}\text{C}$  of topsoil samples ( $M_1$ ,  $M_2$  and  $M_3$ ) as a function of size fractions.

Table 4  
 $\delta^{13}\text{C}$  values of actual plants collected on the various soils and reference values

Sample	$\delta^{13}\text{C}$ (‰)
Wastes of maize	$-14.8 \pm 0.5$
$C_4$ plants: maize, tropical grasses and gramineous	Mean of $-14$ ( $-17$ to $-10$ )
$C_3$ plants: most other plants (trees, rice, wheat, etc.) and certain grass species	Mean of $-27$ ( $-32$ to $-22$ )
Present $\delta^{13}\text{C}$ value of the atmospheric $\text{CO}_2$	$-8$




Fig. 2.  $C_4/C$  ratios of the topsoils samples ( $M_1$ ,  $M_2$  and  $M_3$ ) as a function of size fractions.

the metal, the clay fraction is always the more enriched (between two and five times more in comparison with the bulk). The soil  $M_2$  is less contaminated in comparison with  $M_1$  and whatever the depth and the nature of the metal, the repartition among the size fractions is less heterogeneous. For lead and zinc, between 0 and 65 cm depth, the clay fraction is the more enriched and then the 1000–200  $\mu\text{m}$  fraction. For Cu and Cd, the clay fraction is the only one significantly enriched in comparison with the bulk fraction. For the reference uncontaminated topsoil  $M_3$ , the clay fraction presents the highest metal concentrations. Except for cadmium, the coarse 1000–200  $\mu\text{m}$  fraction of each soil presents a slight enrichment in comparison with the bulk for Pb, Zn and Cu, but always remains much less contaminated than the clay fraction.

From the results presented in Table 6 (total metal concentrations) and in Table 3 (size repartition), we calculated the stocks of the various metals as a function of the size fraction for the topsoils (0–30 cm)  $M_1$ ,  $M_2$  and  $M_3$ . These data are given in Fig. 3 for lead (a), zinc (b), copper (c) and cadmium (d). For these cultivated soils under maize, at each depth the stock of metals in the fine fraction is always higher than in the coarse fraction.

### 3.5. Influence of metals on the organic matter dynamics

Several mechanisms have been proposed to contribute to SOM stabilisation in soils (Sollins et al., 1996): SOM can be (1) biochemically stabilised through the formation of recalcitrant SOM compounds; (2) protected by intimate association with silt and clay particles; and (3) physically stabilised through aggregate formation. The calculated amount of  $C_4$  organic matter in the bulk fractions of the soils under maize follows the sequence  $M_3$  ( $C_4/C = 0.9$ ) >  $M_2$  (0.4) >  $M_1$  (0.3). These differences observed between the three soils in term of SOM turnover could perhaps result from the following parameters: (i) clay amount, (ii) age of the  $C_4$  culture and (iii) soil management (incorporation or not of the straws). Considering the individual influence of each parameter on the SOM turnover, the expected  $C_4/C$  sequence is: (i)  $M_2$  (9% of clay) >  $M_1$  (25%) >  $M_3$  (28%), (ii)  $M_2$  (14 years) >  $M_3$  (13 years) >  $M_1$  (10 years) and (iii)  $M_3 = M_2 > M_1$ . Therefore, if the clay amount was the main factor controlling the SOM of the studied soils, the  $C_4/C$  ratio of  $M_2$  will be the highest. On the other hand, if the age of the  $C_4$  culture or soil management were the main factors controlling the SOM of the studied soils, the  $C_4/C$  ratios of  $M_2$  and  $M_3$  will be quite similar and much higher than the  $C_4/C$  ratio of  $M_1$ . One hypothesis to explain the  $C_4/C$  sequence observed for the three soils is that the SOM turnover was influenced by the metal contamination (with an intensity depending on the metal concentrations).

For the three soils studied, mainly the clay fraction, and to a lesser degree the particulate (1000–200  $\mu\text{m}$ ) SOM fraction, are both enriched with metals and organic carbon. Chemical extractions (with  $\text{CaCl}_2$ ,  $\text{Na}_4\text{P}_2\text{O}_7$ , dithionite and  $\text{HNO}_3$ ) performed for a cultivated soil similar to  $M_1$  (Ponthieu, 2003), indicated that globally, SOM and metal oxides (Fe and Mn) control the speciation of metals. For Pb, EXAFS data confirmed that trend (Morin et al., 1999). SOM displays a high reactivity for metals, lead and copper in particular (Lamy et al., 1994; Harter and Naidu, 1995). From all the results obtained for the atmospheric contaminated soils  $M_1$ ,  $M_2$  and  $M_3$ , we can hypothesis that the metal–organic matter interaction

Table 5  
Organic C (%) of the samples (various depths and size fractions) for each soil

Depth (cm)	1000–500 $\mu\text{m}$	500–200 $\mu\text{m}$	200–50 $\mu\text{m}$	50–20 $\mu\text{m}$	20–2 $\mu\text{m}$	<2 $\mu\text{m}$
<i>(a) Soil profile <math>M_1</math></i>						
0–30		1.22 ± 0.065	1.26 ± 0.05		0.48 ± 0.055	1.75 ± 0.1
30–35		0.55 ± 0.032	0.18 ± 0.04		0.24 ± 0.05	1.36 ± 0.07
35–40		0.69 ± 0.04	0.21 ± 0.03		0.11 ± 0.02	0.93 ± 0.1
55–65		0.26 ± 0.01	0.15 ± 0.02		0.09 ± 0.005	0.5 ± 0.004
170–180		0.19 ± 0.012	0.11 ± 0.05		0.08 ± 0.005	0.27 ± 0.1
<i>(b) Soil profile <math>M_2</math></i>						
0–30	7 ± 0.42	3.85 ± 0.1	2.7 ± 0.1	1.5 ± 0.1	1.2 ± 0.02	8.3 ± 0.3
30–40	5.1 ± 0.3	2.7 ± 0.1	1.9 ± 0.12	1.8 ± 0.11	1 ± 0.02	4.5 ± 0.16
40–50	2.9 ± 0.17	1.6 ± 0.03	1 ± 0.02	1 ± 0.02	0.85 ± 0.01	3.2 ± 0.13
50–65	1.3 ± 0.06	2 ± 0.06	1.1 ± 0.04	1 ± 0.05	0.7 ± 0.03	1.6 ± 0.06
<i>(c) Topsoil <math>M_3</math></i>						
	4.5 ± 0.4	4.7 ± 0.42	3.1 ± 0.27	1.1 ± 0.09	2.35 ± 0.24	6.1 ± 0.53

Table 6  
Metals concentrations (mg kg<sup>-1</sup>) for the samples of each soil at various depths and size fractions

Depth (cm)	1000–500 μm	500–200 μm	200–50 μm	50–20 μm	20–2 μm	<2 μm
<i>(a) Soil M<sub>1</sub></i>						
[Pb] (mg kg <sup>-1</sup> )						
0–30	514.5 ± 15.4	450 ± 13.5	139.3 ± 4.1	115.5 ± 6.8	298.5 ± 11.8	1327.3 ± 39.5
30–35	83 ± 2.5	67.7 ± 2	32 ± 1	29 ± 1.65	83 ± 3.3	742 ± 2.5
35–40	77 ± 2.5	62.1 ± 1.8	24.3 ± 0.7	18.7 ± 1	67 ± 2.6	280 ± 8.4
55–65	68 ± 2.1	62.3 ± 1.9	24 ± 0.7	17 ± 1	45 ± 1.65	140 ± 7.3
170–180	24 ± 0.7	25.5 ± 0.65	21.3 ± 0.6	16.5 ± 0.95	25 ± 1	112 ± 3.4
[Zn] (mg kg <sup>-1</sup> )						
0–30	394 ± 11.8	321 ± 9.5	203 ± 6.1	112.5 ± 6.7	274 ± 10.8	3286 ± 98.7
30–35	72 ± 2.15	49.1 ± 1.5	50.4 ± 1.45	32 ± 1.8	102.3 ± 4.1	1081 ± 45.6
35–40	47 ± 4	37.2 ± 1.1	22 ± 0.65	17 ± 1.05	98 ± 3.75	603 ± 33
55–65	38 ± 1.1	39.7 ± 1.2	20.3 ± 0.65	19 ± 1.1	58 ± 2.3	154 ± 11.2
170–180	26 ± 0.75	20.8 ± 0.65	16.2 ± 0.55	13 ± 0.75	16 ± 0.66	174 ± 5.7
[Cu] (mg kg <sup>-1</sup> )						
0–30	17.5 ± 0.55	14.05 ± 0.4	4.3 ± 0.42	10 ± 0.55	28.2 ± 1.2	93.7 ± 3
30–35	11 ± 0.3	8.7 ± 0.25	10.7 ± 0.3	7 ± 0.4	15.3 ± 0.65	51 ± 1.7
35–40	10 ± 0.3	8.6 ± 0.25	3.7 ± 0.1	2.3 ± 0.15	17 ± 0.7	55 ± 1.9
55–65	7 ± 0.2	7.3 ± 0.2	3.2 ± 0.1	2.5 ± 0.15	7 ± 0.25	22 ± 0.05
170–180	3.5 ± 0.1	2.4 ± 0.07	1.7 ± 0.05	0.8 ± 0.05	2 ± 0.08	21 ± 0.7
[Cd] (mg kg <sup>-1</sup> )						
0–30	11.7 ± 0.35	10 ± 0.3	5.8 ± 0.17	4.5 ± 0.15	14.4 ± 0.55	53.5 ± 1.6
30–35	8.3 ± 0.25	6 ± 0.18	4.1 ± 0.12	3 ± 0.18	8.1 ± 0.35	30 ± 0.9
35–40	6.4 ± 0.18	5.8 ± 0.17	3.6 ± 0.1	1.5 ± 0.09	5 ± 0.2	13 ± 0.4
55–65	5.1 ± 0.15	5.3 ± 0.15	1.9 ± 0.05	1 ± 0.06	3.8 ± 0.15	8 ± 0.3
170–180	0.8 ± 0.02	0.5 ± 0.015	0.4 ± 0.012	0.2 ± 0.012	2 ± 0.08	7 ± 0.3
<i>(b) Soil M<sub>2</sub></i>						
[Pb] (mg kg <sup>-1</sup> )						
0–30	273 ± 13.7	224.5 ± 18	207.5 ± 10.4	155 ± 5.4	163 ± 7.5	318 ± 15.9
30–40	165 ± 8.5	178 ± 14.2	142 ± 7	131 ± 4.7	142 ± 6.4	187 ± 9.5
40–50	86 ± 4.3	84 ± 6.7	75 ± 3.8	62 ± 2.25	65 ± 3.1	86 ± 4.5
50–65	89 ± 4.5	81 ± 6.5	49 ± 2.3	33 ± 1.2	25 ± 1.2	91 ± 4.7
[Zn] (mg kg <sup>-1</sup> )						
0–30	235.5 ± 11.9	200 ± 16	184 ± 9	161 ± 5.8	162.5 ± 7.6	258 ± 12.8
30–40	153 ± 7.7	169 ± 13.7	125.5 ± 6.3	122 ± 4.7	127 ± 5.9	164 ± 8.5
40–50	38.5 ± 1.9	42 ± 3.3	36.1 ± 1.75	27 ± 0.9	28 ± 1.3	67 ± 3.7
50–65	34 ± 1.7	34 ± 2.5	22.4 ± 1.3	23 ± 0.65	32 ± 1.45	51 ± 2.8
[Cu] (mg kg <sup>-1</sup> )						
0–30	23.8 ± 1.2	17.15 ± 1.35	17.4 ± 0.9	14.95 ± 0.55	14.25 ± 0.65	55 ± 2.65
30–40	15.3 ± 0.8	11.4 ± 0.9	10.4 ± 0.55	9.7 ± 0.35	8.5 ± 0.4	44 ± 2.5
40–50	9.5 ± 0.45	9.2 ± 0.7	6.2 ± 0.4	8 ± 0.3	9 ± 0.4	39 ± 1.65
50–65	6.9 ± 0.3	7.4 ± 0.5	4.3 ± 0.2	3.5 ± 0.12	2.8 ± 0.1	41 ± 2.1
[Cd] (mg kg <sup>-1</sup> )						
0–30	16.25 ± 0.8	12.95 ± 1	10.05 ± 0.5	11.9 ± 0.4	9.7 ± 0.4	30.4 ± 1.5
30–40	12.5 ± 0.65	8.8 ± 0.7	7.7 ± 0.4	6.3 ± 0.22	6.8 ± 0.305	33 ± 1.65
40–50	7.3 ± 0.365	5.9 ± 0.45	3.8 ± 0.2	4.8 ± 0.165	5.1 ± 0.225	17 ± 0.85
50–65	6.7 ± 0.335	7.1 ± 0.565	6.2 ± 0.35	2.7 ± 0.095	1.9 ± 0.08	13.2 ± 0.66
<i>(c) Topsoil M<sub>3</sub></i>						
[Pb] (ppm)	57 ± 3.8	66 ± 3.3	49 ± 1.5	41 ± 0.8	39 ± 1.5	84 ± 2.4
[Zn] (ppm)	91 ± 6.2	86 ± 4.2	48 ± 1.4	40 ± 0.75	35 ± 1.35	121 ± 3.5
[Cu] (ppm)	16 ± 1.1	14 ± 0.7	11 ± 0.35	7.5 ± 0.15	8.2 ± 0.3	21 ± 0.65
[Cd] (ppm)	3.6 ± 0.25	2.4 ± 0.12	2.5 ± 0.065	2.7 ± 0.4	2.1 ± 0.08	5.3 ± 0.2

depends in two different ways on two size fractions: fraction <2 μm and fraction >50 μm. Plants are contaminated (soil–plant transfer or direct absorption by leaves) and after their death, a selective degradation takes place depending in particular on their metal content and the soil characteristics. Even in metal contaminated soils, biological activity occurred (Aoyama and Nagumo, 1997), but an alteration of the

decomposition of organic matter was reported (Berg et al., 1991; Balabane et al., 1999; Cotrufo et al., 1995; Valsecchi et al., 1995), generally explained by an inhibition of microbial activity by bio-available toxic metals (Brookes, 1995; Mhatre and Pankhurst, 1997). Therefore an accumulation of highly contaminated coarse plant debris can result (Hughes et al., 1980; Harter and Naidu, 1995). However, the particulate




Fig. 3. Stocks of metals for the topsoil samples as a function of size fractionations: (a) lead, (b) zinc, (c) copper and (d) cadmium.

organic matter fraction, the coarse sized SOM fraction composed by recently accumulated organic matter, is poorly associated with the soil mineral matrix and has a relatively rapid turnover in comparison with other SOM (Cambaradella and Elliott, 1992; Balesdent, 1996). Metals of the coarse fractions are progressively released during the decay of particulate organic matter. A part of these metals is then sorbed by the colloid organic matter of the clay fraction (high CEC and specific surface) (Besnard et al., 2001). According to Wattel-Koekkoek et al. (2003) and Gregorich et al. (2003), the interaction between SOM and polyvalent cations can induce SOM stabilisation.

Comparison of the  $C_4/C$  ratios for the 500–200 μm fraction ( $M_1$ :0.3,  $M_2$  0.45,  $M_3$  0.78) and the clay fraction ( $M_1$  0.08,  $M_2$  0.15,  $M_3$  0.3) of the three topsoils suggest that the transformation of coarse organic matter (from  $C_4$  origin) into humic substances present in the clay fraction (Baldock et al., 1992) could be reduced by metals. Moreover, the distribution of recently incorporated organic carbon is not homogeneous in soil aggregates. For instance, Puget et al. (2000), found that maize residue carbon was enriched in particular in the particulate organic matter located in macro-aggregates. In the studied soils, metals are mainly concentrated in the clay fraction enriched by the old  $C_3$  SOM. In contrast, the particulate organic matter is less contaminated. We can therefore conclude that the stabilisation effect of metals is stronger for the old  $C_3$

SOM in comparison to the  $C_4$  young SOM (mainly present in the labile coarser fractions). Consecutively to metal contamination, if the  $C_4$  SOM is more degraded by the microorganisms than the  $C_3$  SOM, then the  $C_4/C$  ratio decreases.

#### 4. Conclusions

The determined sequence for the amount of  $C_4$  organic matter in the bulk fractions:  $M_3$  (0.9) >  $M_2$  (0.4) >  $M_1$  (0.3), cannot result only from the differences observed between the three soils in terms of clay amount, age of the  $C_4$  culture and soil management. On the contrary, the lowest value of the  $C_4/C$  ratio for the contaminated soils ( $M_1$  and  $M_2$ ) in comparison with the uncontaminated soil  $M_3$  could be due to the influence of metals on the SOM turnover.

The organic carbon is mainly located in the topsoils (plough horizons), where most of the metals occur. For the three studied soils, the clay fraction, and to a lesser degree (except for cadmium) the particulate (1000–200 μm) SOM fraction, are both enriched with metals and organic carbon. Chemical extractions performed for a cultivated soil similar to  $M_1$  (Ponthieu, 2003), indicated that SOM and metal oxides control the speciation of metals.

The  $\delta^{13}C$  results demonstrated differences in the nature of the SOM as a function of the size fraction and depth. The clay fraction is strongly enriched with old  $C_3$  SOM in

comparison with the coarse fractions. The comparison of the  $C_4/C$  ratios for the 500–200  $\mu\text{m}$  and the clay fractions of the three topsoils suggest that the transformation of coarse organic matters (from  $C_4$  origin) into humic substances present in the clay fraction could be reduced by metals.

The less contaminated (in comparison with the clay fraction) particulate organic matter fraction composed by recently accumulated organic residue has a relatively rapid turnover in comparison with the clay SOM (Cambaradella and Elliott, 1992; Balesdent, 1996). Metals of the coarse fractions are progressively released during the decay of particulate organic matter. A part of these metals is then sorbed by the colloid organic matter of the clay fraction, therefore the clay fraction (enriched with old  $C_3$  SOM) is progressively more contaminated and more stabilised (Wattel-Koekkoek et al., 2003; Gregorich et al., 2003). Consecutively to metal contamination, the proportion of the young  $C_4$  organic matter relative to old  $C_3$  (or  $C_4/C$  ratio) decreases because the young  $C_4$  SOM (mainly present in the labile coarser fractions) is more easily degraded by the microorganisms than the  $C_3$  SOM.

Understanding of mechanisms involved in SOM dynamic is an important challenge in regard to the global influence of that natural compartment. The size fractionation and profile sampling permit access to various pools of SOM with different stability and reactivity for pollutants. In order to better understand the mechanisms, we could test in detail, on various pieces of the same experimental long term site: the influence of the nature of the metal on the turnover of SOM; the effect of increasing metal concentrations; and investigate various metallic sources (influence of initial metal speciation).

## Acknowledgements

The authors thank the French National Program Environment (PNSE) for financial support.

## References

- Aoyama, M., Nagumo, T., 1997. Comparison of the effects of Cu, Pb, As on plant residue decomposition, microbial biomass and soil respiration. *Soil Science and Plant Nutrition* 43, 613–622.
- Arrouays, D., Balesdent, J., Mariotti, A., Girardin, C., 1995. Modelling organic carbon turnover in cleared temperate forest soils converted to maize cropping by using  $^{13}\text{C}$  natural abundance measurements. *Plant and Soil* 173, 191–196.
- Balabane, M., Faivre, D., van Oort, F., Dahmani-Muller, H., 1999. Mutual effects of soil organic matter dynamics and heavy metals fate in a metallophyte grassland. *Environmental Pollution* 105, 45–54.
- Baldock, J.A., Oades, J.M., Waters, A.G., Peng, X., Vassallo, A.M., Wilson, M.A., 1992. Aspects of the chemical structure of soil organic materials as revealed by solid-state  $^{13}\text{C}$  NMR spectroscopy. *Biochemistry* 16, 1–42.
- Baldock, J.A., Skjemstad, J.O., 2000. Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Organic Geochemistry* 31, 697–710.
- Balesdent, J., 1996. The significance of organic separates to carbon dynamics and its modelling in some cultivated soils. *European Journal of Soil Science* 47, 485–493.
- Balesdent, J., Mariotti, A., 1996. Measurement of soil organic matter turnover using  $^{13}\text{C}$  natural abundance. In: Boutton, T.W., Yamasaki, S. (Eds.), *Mass Spectrometry of Soil*. Marcel Dekker, New York, pp. 83–111.
- Barajas-Aceves, M., Grace, C., Ansorena, J., Dendooven, L., Brookes, P.C., 1999. Soil microbial biomass and organic C in a gradient of zinc concentrations in soils around a mine spoil tip. *Soil Biology and Biochemistry* 31, 867–876.
- Beare, M.H., Cabrera, M.L., Hendrix, P.F., Coleman, D.C., 1994. Aggregate-protected and unprotected organic matter pools in conventional-tillage and no-tillage soils. *Soil Science Society of America Journal* 58, 787–795.
- Benedetti, M.F., Van Riemsdijk, W.H., Koopal, L.K., Kinniburgh, D.G., Goody, D.C., Milne, C.J., 1996. Metal binding by natural organic matter: from the model to the field. *Geochimica Cosmochimica Acta* 60 (14), 2503–2513.
- Berg, B., Ekbohm, G., Soderstrom, B., Staaf, H., 1991. Reduction in the decomposition rates of Scots Pine needle litter due to heavy-metal pollution. *Water Air and Soil Pollution* 59, 165–177.
- Besnard, E., Chenu, C., Robert, M., 2001. Influence of organic amendments on copper distribution among particle-size and density fractions in Champagne vineyard soils. *Environmental Pollution* 112, 329–337.
- Boutton, T.W., 1996. Stable carbon isotope ratio of soil organic matter and their use as indicators of vegetation and climate change. In: Boutton, T.W., Yamasaki, S. (Eds.), *Mass Spectrometry of Soil*. Marcel Dekker, New York, pp. 47–82.
- Brookes, P.C., 1995. The use of microbial parameters in monitoring soil pollution by heavy metals. *Biology and Fertility of Soils* 19, 269–279.
- Cambaradella, C.A., Elliott, E.T., 1992. Particulate soil organic matter changes across a grassland cultivation sequence. *Soil Science Society of America Journal* 56, 777–783.
- Chenu, C., 1995. Extracellular polysaccharides: An interface between microorganisms and soil constituents. In: Huang, P.M., Berthelin, J., Bollag, J.M., McGill, W.B., Page, A.L. (Eds.), *Environmental Impact of Soil Component Interactions*. Lewis, Boca Raton, pp. 217–233.
- Cotrufo, M.F., De Santo, A.V., Alfani, A., Bartoli, G., De Cristofaro, A., 1995. Effects of urban heavy metal pollution on organic matter decomposition in *Quercus ilex* L. woods. *Environmental Pollution* 89, 81–87.
- Ducaroir, J., Lamy, I., 1995. Evidence of trace metal association with soil organic matter using particle size fractionation after physical dispersion treatment. *Analyst* 120, 741–745.
- Dumat, C., Cheshire, M.V., Fraser, A., Shand, C., Staunton, S., 1997. The effect of removal of soil organic matter and iron on the adsorption of Cs. *European Journal of Soil Science* 48 (4).
- Ehleringer, J.R., Cerling, T.E., Heliker, B.R., 1997.  $C_4$  photosynthesis, atmospheric  $\text{CO}_2$  and climate. *Oecologia* 112, 285–299.
- Franzluebbers, A.J., Arshad, M.A., 1997. Particulate organic carbon content and potential mineralization as affected by tillage and texture. *Soil Science Society of America J* 61, 1382–1386.
- Gregorich, E.G., Beare, M.H., Stocklas, U., St-Georges, P., 2003. Bio degradability of soluble organic matter in maize cropped soils. *Geoderma* 113, 237–252.
- Harter, R.D., Naidu, R., 1995. Role of the metal-organic complexation in metal sorption by soils. *Advances in Agronomy* 55, 219–263.
- Hevia, G.G., Buschiazzo, D.E., Hepper, E.N., Urioste, A.M., Anton, E.L., 2003. Organic matter in size fractions of soils of the semiarid Argentina. Effects of climate, soil texture and management. *Geoderma* 116, 265–277.
- Hughes, M.K., Lepp, N.W., Phipps, D.A., 1980. Aerial heavy metal pollution and terrestrial ecosystems. *Advances in Ecology Research* 11, 217–227.
- John, B., Yamashita, T., Ludwig, B., Flessa, H., 2005. Storage of organic carbon in aggregate and density fractions of silty soils under different types of land use. *Geoderma* 128, 63–79.
- Kaiser, K., Guggenberger, G., 2000. The role of DOM sorption to mineral surfaces in the preservation of organic matter in soils. *Organic Geochemistry* 31, 711–725.
- Krull, E.S., Skjemstad, J.O., 2003.  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  profiles in  $^{14}\text{C}$ -dated Oxisol and Vertisols as a function of soil chemistry and mineralogy. *Geoderma* 112, 1–29.
- Lamy, I., Cambier, P., Bourgeois, S., 1994. Pb and Cd complexation. *Environmental Geochemistry and Health* 16, 1–16.

- Metson, A.J., 1956. Methods of chemical analysis for soil survey samples. NZ Department of Scientific and Industrial Research Soil Bureau Bulletin 12, pp. 208.
- Mhatre, G.N., Pankhurst, C.E., 1997. Bioindicators to detect contamination of soils with special reference to heavy metals. In: Pankhurst, C.E., Doube, B.M., Gupta, V.V. (Eds.), *Biological Indicators of Soil Health*. CAB International, Wallingford, pp. 349–369.
- Monreal, C.M., Schulten, H.R., Kodama, H., 1997. Age, turnover and molecular diversity of soil organic matter in aggregates of a Gleysol. *Canadian Journal of Soil Science* 77, 379–388.
- Morin, G., Juillot, F., Ostergren, J., Rytuba, J.J., Ildefonse, P., Calas, G., Brown, G.E., 1999. XAFS determination of the chemical form of lead in smelter-contaminated soils and mine tailings: importance of sorption processes. *American Mineralogy* 84, 420–434.
- Muller, T., Hoper, H., 2004. Soil organic matter turnover as a function of the soil clay content: consequences for model applications. *Soil Biology and Biochemistry* 36, 877–888.
- Oades, J.M., 1995. An overview of processes affecting the cycling of organic carbon in soils. In: Zepp, R.G., Sonnatg, C. (Eds.), *Role of Nonliving Organic Matter in the Earth's Carbon Cycle*. John Wiley and Sons, New York, pp. 293–303.
- Parton, W.J., 1996. The CENTURY model. In: Powlson, D.S., Smith, P., Smith, J.U. (Eds.), *Evaluation of Soil Organic Matter Models*. Springer, Berlin, pp. 283–291.
- Peterson, B.J., Fry, B., 1987. Stable isotopes in ecosystem studies. *Annual Review of Ecology and Systematics* 18, 293–320.
- Pinheiro, E.F.M., Pereira, M.G., Anjos, L.H.C., 2004. Aggregate distribution and soil organic matter under different tillage systems for vegetable crops in a Red Latosol from Brazil. *Soil & Tillage Research* 77, 79–84.
- Ponthieu, M., 2003. Spéciation des éléments traces métalliques dans les sols et les solutions des sols: du modèle au terrain. Thèse de l'Univ. d'Aix Marseille III.
- Puget, P., Chenu, C., Balesdent, J., 2000. Dynamics of soil organic matter associated with particle-size fractions of water-stable aggregates. *European Journal of Soil Science* 51, 595–605.
- Rethemeyer, J., Grootes, P.M., Bruhn, F., Andersen, N., Nadeau, M.J., Kramer, C., Gleixner, G., 2004. Age heterogeneity of soil organic matter. *Nuclear Instruments and Methods in Physics Research Section B Beam Interactions with Materials and Atoms* 24, 521–527.
- Sauvé, S., Martinez, C.E., McBride, M., Hendershot, W., 2000. Adsorption of free lead by pedogenic oxides, ferrihydrite and leaf compost. *Soil Science Society of America Journal* 64, 595–599.
- Schwartz, D., de Foresta, H., Mariotti, A., Balesdent, J., Massimba, J.P., Girardin, C., 1996. Present dynamics of the savanna-forest boundary in the Congolese Mayombe: a pedological, botanical and isotopic ( $^{13}\text{C}$  and  $^{14}\text{C}$ ) study. *Oecologia* 106, 516–524.
- Shepherd, T.G., Saggat, S., Newman, R.H., Ross, C.W., Dando, J.L., 2001. Tillage-induced changes to soil structure and organic carbon fractions in New Zealand soils. *Australian Journal of Soil Research* 39, 465–489.
- Smith, B.N., Epstein, S., 1971. Two categories of  $^{13}\text{C}/^{12}\text{C}$  ratios for higher plants. *Plant Physiology* 47, 380–384.
- Sobanska, S., Ricq, N., Laboudigues, A., Guillermo, R., Brémard, C., Laureyns, J., Merlin, J.C., Wignacourt, J.P., 1999. Microchemical investigations of dust emitted by lead smelter. *Environmental Science and Technology* 33, 1334–1339.
- Sollins, P., Homann, P., Caldwell, B.A., 1996. Stabilization and destabilization of soil organic matter: mechanisms and controls. *Geoderma* 74, 65–105.
- Spark, K.M., Wells, J.D., Johnson, B.B., 1997. The interaction of a humic acid with heavy metals. *Australian Journal of Soil Research* 35, 89–101.
- Sterckeman, T., Douay, F., Proix, N., Fourrier, H., 2000. Vertical distribution of Cd, Pb and Zn in soils near smelters in the north of France. *Environmental Pollution* 107, 377–389.
- Tiessen, H., Cuevas, E., Chacon, O., 1994. The role of soil organic matter in sustaining soil fertility. *Nature* 371, 783–785.
- Valsecchi, G., Gigliotti, C., Farini, A., 1995. Microbial biomass, activity and organic matter accumulation in soils contaminated with heavy metals. *Biology and Fertility of Soils* 20, 253–259.
- Vanlauwe, B., Nwoke, O.C., Sanginga, N., Merckx, R., 1999. Evaluation of methods for measuring microbial biomass C and N and relationships between microbial biomass and soil organic matter particle size classes in West-African soils. *Soil Biology and Biochemistry* 31, 1017–1082.
- Wattel-Koekkoek, E.J.W., Buurman, P., van der Plicht, J., Wattel, E., van Breemen, N., 2003. Mean residence time of soil organic matter associated with kaolinite and smectite. *European Journal of Soil Science* 54, 269–278.
- Yin, Y., Impellitteri, C.A., You, S.J., Allen, H.E., 2002. The importance of organic matter distribution and exact soil: solution ratio on the desorption of heavy metals from soils. *The Science of the Total Environment* 287, 107–119.