

Population dynamics of swine influenza virus in farrow-to-finish and specialised finishing herds in The Netherlands

W.L.A. Loeffen, W.A. Hunneman, J. Quak, J.H.M. Verheijden, J.A. Stegeman

▶ To cite this version:

W.L.A. Loeffen, W.A. Hunneman, J. Quak, J.H.M. Verheijden, J.A. Stegeman. Population dynamics of swine influenza virus in farrow-to-finish and specialised finishing herds in The Netherlands. Veterinary Microbiology, 2009, 137 (1-2), pp.45. 10.1016/j.vetmic.2009.01.004 . hal-00485528

HAL Id: hal-00485528 https://hal.science/hal-00485528

Submitted on 21 May 2010 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Population dynamics of swine influenza virus in farrow-to-finish and specialised finishing herds in The Netherlands

Authors: W.L.A. Loeffen, W.A. Hunneman, J. Quak, J.H.M. Verheijden, J.A. Stegeman

PII:	S0378-1135(09)00003-0
DOI:	doi:10.1016/j.vetmic.2009.01.004
Reference:	VETMIC 4316
To appear in:	VETMIC
Received date:	24-6-2008
Revised date:	19-12-2008
Accepted date:	2-1-2009

Please cite this article as: Loeffen, W.L.A., Hunneman, W.A., Quak, J., Verheijden, J.H.M., Stegeman, J.A., Population dynamics of swine influenza virus in farrow-tofinish and specialised finishing herds in The Netherlands, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2009.01.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1	POPULATION DYNAMICS OF SWINE INFLUENZA VIRUS IN FARROW-TO-FINISH AND										
2	SPECIALISED FINISHING HERDS IN THE NETHERLANDS										
3											
4	W.L.A. Loeffen, W.A. Hunneman, J. Quak, J.H.M. Verheijden, J.A. Stegeman										
5											
6	W.L.A. Loeffen, DVM, Department of Swine Health, Animal Health Service, PO Box 9,										
7	7400AA Deventer, The Netherlands (present address: Central Veterinary Institute of										
8	Wageningen UR (CVI), P.O. Box 65, 8200AB Lelystad, The Netherlands)										
9	W.A. Hunneman, DVM, PhD, Department of Swine Health, Animal Health Service, PO Box										
10	9, 7400AA Deventer, The Netherlands										
11	J. Quak, Central Veterinary Institute of Wageningen UR (CVI), P.O. Box 65, 8200AB										
12	Lelystad, The Netherlands										
13	J.H.M. Verheijden, DVM, PhD, Faculty of Veterinary Medicine, University of Utrecht, PO										
14	Box 80151, 3508 TD Utrecht, The Netherlands										
15	J.A. Stegeman, DVM, PhD, Faculty of Veterinary Medicine, University of Utrecht, PO Box										
16	80151, 3508 TD Utrecht, The Netherlands										
17											
18											
19 20 21 22 23 24	Corresponding author: W.L.A. Loeffen Central Veterinary Institute of Wageningen UR (CVI) P.O. Box 65 8200AB Lelystad										
25 26 27 28 29 30	The Netherlands Telephone number : +31 320 238696 Fax number : +31 320 238668 Email : willie.loeffen@wur.nl										

31 Influenza virus infections with subtypes H1N1, H3N2 and H1N2 are very common in 32 domestic pigs in Europe. Data on possible differences of population dynamics in 33 finishing pigs in farrow-to-finish herds and in specialised finishing herds are, 34 however, scarce. The presence of sows and weaned piglets on the same premises may, however, affect the exposure of finishing pigs to influenza viruses. In a 35 36 longitudinal study on 14 farrow-to-finish herds and 15 finishing herds, groups of pigs 37 were followed by repeatedly testing the same animals for antibodies against all three 38 influenza virus subtypes (H1N1, H3N2 and H1N2). At the end of the finishing period, 39 the seroprevalences in farrow-to-finish and specialised finishing herds were 44.3% and 62.0% respectively for H1N1, 6.6% and 19.3% respectively for H3N2, and 57.2% 40 and 25.6% respectively for H1N2. For all three subtypes, the incidence of influenza 41 42 virus infections was highest at the beginning of the finishing period in farrow-to-finish 43 herds, while the incidence of influenza virus infections was highest at the end of the finishing period in finishing herds. Respiratory disease, probably related to the 44 45 influenza infections, was observed in five of these herds only, but also occurred at the beginning of the finishing period in farrow-to-finish herds and at the end of the 46 47 finishing period in finishing herds. The observed differences of population dynamics of influenza virus may affect choice and timing of intervention measures. 48

49

50 Keywords: swine influenza; H1N1; H3N2; H1N2; incidence; herd type; population 51 dynamics

52 **1. Introduction**

Influenza virus infections in swine are very common. In several studies in European
countries, seroprevalences for the swine influenza strains H1N1 and H3N2 were found to be
in the range of 20-80% in finishing pigs at the end of the finishing period and in sows
(Masurel et al., 1983; Haesebrouck and Pensaert, 1986b; Yus et al., 1989; Elbers et al.,

57 1990; Teuffert et al., 1991; Elbers et al., 1992; Groschup et al., 1993; Ewald et al., 1994; 58 Brown et al., 1995b; Maes et al., 1999; Maldonado et al., 2006). A more recent subtype, 59 H1N2, seems to originate from the UK where it was found for the first time in 1994 (Brown et 60 al., 1995a). H1N2 was subsequently reported from Belgium (Van Reeth et al., 2000), Italy and France (Marozin et al., 2002), Germany (Schrader and Suss, 2003), and Spain 61 62 (Maldonado et al., 2006). Many, if not all of these strains are related to the original H1N2 63 strain from the UK (Marozin et al., 2002). Seroprevalence studies in Belgium (Van Reeth et 64 al., 2000) and Spain (Maldonado et al., 2006) resulted in high seroprevalences of 65 approximately 70% and 50% respectively for H1N2, but both studies were carried out in 66 SOWS. 67 Studies in finishing pigs so far give overall estimates of the seroprevalence at the end of the finishing period. However, it is not clear at what moment during the finishing period these 68

infections take place, nor do they differentiate between finishing pigs in specialised finishing
herds and finishing pigs in farrow-to-finish herds. The presence of sows and piglets on the
same premises as the finishing pigs may, however, affect the population dynamics of swine
influenza in these herd types, and thus also affect the choice and timing of intervention
measures, like vaccination or zoosanitary measures.

Longitudinal studies were therefore carried out to be able to compare the seroprevalences and incidences of swine influenza virus infections at different ages in finishing pigs in both herd types. This allowed us also to determine whether there are differences in the population dynamics of swine influenza virus infections in finishing pigs in farrow-to-finish herds versus finishing pigs in specialised finishing herds.

79 **2. Materials and methods**

80 2.1. Selection of herds

81 In the Netherlands, there are three regions with a high pig density (figure 1). In these 82 regions the average pig density is more than 600 pigs per square kilometre. In contrast, the

average pig density in the remaining part of the Netherlands is below 100 pigs per square
kilometre. From the three pig dense regions all herds with >400 finishing pigs were selected
for possible participation in the study. This selection included 27% of the Dutch swine herds,
housing 69% of the Dutch finishing pig population.

87 In each category (farrow-to-finish and finishing herds) 64 herds were randomly selected. 88 A written request for participation in a longitudinal serological survey was sent to these farmers. Vaccination against swine influenza was not allowed. The first 15 in each category 89 90 returning a positive response were included in the study. One farrow-to-finish herd that 91 applied for the survey withdrew shortly before the survey actually started and could not be 92 replaced on such short notice. Thus, 29 herds finally participated. In eleven of the finishing 93 herds the investigated pigs originated from only one breeding herd, in three finishing herds the pigs originated from two breeding herds and in one finishing herd the pigs originated 94 95 from three breeding herds.

96 2.2. Sampling

All farms were visited three (finishing herds) or four (farrow-to-finish herds) times during
the months of January to May 1999. In each herd one compartment was followed in a
longitudinal study and blood samples were collected during each visit.

100 In finishing herds one compartment where the piglets were 12 weeks old was sampled 101 during the first visit. On average, these piglets arrived 2-3 weeks before on the farm. Pigs 102 were tagged individually during the first sampling to allow for resampling of the same pigs 103 during subsequent visits. The second blood samples were taken four weeks later (age of 16 104 weeks) and a final sample was taken within one week before the first pigs from that 105 compartment were delivered to the slaughter house (on average at the age of 22 weeks). 106 Piglets within the sampled compartment always originated from one breeding herd. During 107 the finishing period no other pigs were added to that compartment. 108 In farrow-to-finish herds one compartment where the piglets were 8 weeks old was

sampled during the first visit. Pigs were tagged individually during the first sampling to allow

110 for resampling of the same pigs during subsequent visits. The second series of blood 111 samples was taken four weeks later (age of 12 weeks), the third another four weeks later 112 (age of 16 weeks) and the final series of samples was taken within one week before the first 113 pigs from that compartment were delivered to the slaughter house (on average at the age of 114 22 weeks). Tagged piglets were kept together in the same compartment until slaughter, but 115 while being transferred to the finishing facilities, piglets from two compartments were 116 sometimes mixed. However, during the finishing period no other pigs were added to that 117 compartment anymore.

The sample size for each compartment was calculated so that with an estimated seroprevalence of 50% and a confidence of 0.95 the margin of error was less than 20%. This resulted in sample sizes of 16-24 pigs per compartment using the formula for simple random sampling (Thrusfield, 1995). Within a compartment an equal number of pigs were sampled per pen, as far as the total number of samples allowed for this. Within each pen, pigs were selected at random (haphazardly).

All farmers were asked to record all clinical signs and medications in the compartmentunder study.

126 2.3. Serological examination

All sera were tested in a hemagglutination inhibition (HI) test (Kendal et al., 1982) for antibodies against influenza virus strains A/swine/Neth/Best/96 (H1N1), A/swine/Neth/St Oedenrode/96 (H3N2) and A/swine/Gent/7625/99 (H1N2). Sera were pre-treated with chicken erythrocytes and cholera filtrate to remove non-specific hemagglutinating factors and non-specific inhibitors of hemagglutination. Four hemagglutinating units (HAU) were used in each test.

A/swine/Neth/Best/96 (H1N1) and A/swine/Neth/St Oedenrode/96 were isolated in the
Netherlands and are representative for influenza strains circulating in the Netherlands
(Loeffen et al., 1999). A/swine/Gent/7625/99 (H1N2) is a Belgian strain, representative for
H1N2 strains circulating among pigs in Belgium, which in turn is expected to be

representative for H1N2 strains all over Western Europe (Marozin et al., 2002). Although
H1N1 and H1N2 strains both possess a hemagglutinin H1, these are of different origin
(Brown et al., 1998) and little or no cross reactivity was found in serological tests after
experimental infections (Van Reeth et al., 2000; Van Reeth et al., 2004; Van Reeth et al.,
2006).

142 Sera were tested in serial twofold dilutions, starting at 1:9. Titres ≥18 were considered 143 positive. A fourfold rise in titres in consecutive samples was considered evidence of an 144 influenza virus infection somewhere during that period (taking into account approximately 145 one week for titres to develop after an influenza virus infection). In the presence of maternal 146 antibodies, titres of the second sera were not compared to the titres of the first sera, but to 147 expected titres of the second sera if these were due to maternal antibodies only. To do this, 148 the decay of maternal antibodies, having a half-life of approximately 12 days, was taken into 149 consideration, extrapolating titres from the first sera (Loeffen et al., 2003).

150 2.4. Statistical analysis

151 Statistical analyses for the differences between farrow-to-finish herds (FFH) and 152 specialised finishing herds (FH) were performed per strain and per time interval. The 153 incidence, cumulative incidence and prevalence data were expressed as fractions per farm. 154 The incidence being the number of seroconversions since the previous sampling, the 155 cumulative incidence the total number of seroconversions until the moment of sampling, and 156 the prevalence the fraction of seropositive pigs at each sampling. The test statistic (t) for the 157 null hypothesis of no difference between the herd types was:

$$158 t = a - b .$$

wherein a and b are the fractions averaged over FFH and FH farms respectively. The null-distribution was obtained by random permutation of the labels FFH and FH over the farms.

161 **3. Results**

175

162 3.1. Seroprevalence

163 For H1N2, at the end of the finishing period the seroprevalences in farrow-to-finish and 164 specialised finishing herds were 57.2% and 25.6% respectively (table 1). The 165 seroprevalence was in all age groups higher in farrow-to-finish herds than in specialised 166 finishing herds (P=0.01, P<0.01 and P=0.03 for 12, 16 and 22 weeks of age respectively). 167 For H1N1 and H3N2, at the end of the finishing period the seroprevalences in farrow-to-168 finish and specialised finishing herds were 44.3% and 62.0% respectively for H1N1 and 169 6.6% and 19.3% respectively for H3N2. The differences of the seroprevalences in farrow-to-170 finish herds and finishing herds were not statistically significant. 171 3.2. Incidence For H1N1 the differences between both herd types are statistically significant between 16 172 and 22 weeks of age (P<0.01) and for H1N2 between 0 and 12 (P=0.05) and 12 and 16 173 174 (P=0.01) weeks of age (table 2). No significant differences (P<0.05) were seen for H3N2.

seroprevalence, however, corrected for antibodies due to maternal immunity or antibodies
from an infection that declined to a level below the detection limit of the test.

The cumulative incidences are shown in table 3. They are essentially comparable to the

178 3.3. Clinical symptoms of respiratory disease

Twenty-four farmers (from 11 farrow-to-finish herds and 13 finishing herds) kept a record of clinical signs and medications. In eleven of them, an episode of respiratory disease was recorded: five in farrow-to-finish herds and six in finishing herds. The severity of the symptoms was variable, ranging from only coughing for a few days up to a combination of coughing, laboured breathing and decrease of feed intake. Five of these episodes coincided with an incidence of 80-95% of the pigs for one of the influenza subtypes in the same period. Two of these were in farrow-to-finish herds and both occurred between the age of 8 and 12

weeks (1x H1N2 and 1x H3N2). The other three occurred in finishing herds, all of them
between the age of 16 and 22 weeks (1x H1N1 and 2x H1N2).

188 4. Discussion

In this study we found that the timing of influenza infections in finishing pigs is quite different in finishing pigs from farrow-to-finish herds compared to finishing pigs from specialised finishing herds. In farrow-to-finish herds the incidence of influenza virus infections was highest at the beginning of the finishing period, while in finishing herds the incidence of influenza virus infections was highest at the end of the finishing period. This pattern was observed for all three influenza virus subtypes. It can be speculated that the exposure of finishing pigs to influenza virus is different in both herd types.

196 Sows and especially weaned piglets may be a reservoir for continuous circulation of 197 influenza viruses (Loeffen et al., 2003). Indirect exposure of finishing pigs to influenza virus 198 from sows or weaned piglets will be more easily achieved in farrow-to-finish herds than in 199 specialised finishing herds. Multiple virus introductions into a compartment of finishing pigs in 200 a farrow-to-finish herd could thus result in a high incidence at the beginning of the finishing period. Large outbreaks are apparently rare at that time, possibly due to the combination of 201 202 decreasing numbers of piglets with maternal antibodies and increasing numbers of piglets 203 with antibodies due to an infection. Introductions later during the finishing period may then 204 die out guickly due to the already high seroprevalence. A compartment of pigs in a finishing 205 herd on the other hand may be subject to only a few virus introductions. On average this 206 may result in populations with more susceptible animals in the second half of the finishing 207 period, with any introduction at that time resulting in a large outbreak within that 208 compartment.

The presence of finishing pigs in a herd might also affect influenza virus infections in sows and weaned piglets. Especially with subtype H1N2, at least 8.6% percent of the weaned piglets seroconverted before they were 8 weeks old. This could also explain the higher seroprevalence at the age of 12 weeks in farrow-to-finish herds, compared to

213 finishing herds. It may be that in the breeding herds that supplied the piglets to the 214 specialised finishing herds, virus circulation in weaned piglets was lower than in weaned 215 piglets on farrow-to-finish herds. This would suggest that the presence of finishing pigs may 216 result in an additional exposure of weaned piglets to the virus and thus may affect virus 217 circulation in weaned piglets. It's also possible that the relatively recent introduction of the 218 H1N2 subtype in the population somehow causes this effect. This would, however, indicate 219 that farrow-to-finish herds and specialised breeding herds are at a different risk for virus 220 introduction. Given that the two herd types were not located in exactly the same place, these 221 results have to interpreted with some care anyway.

222 Even though serological differentiation between H1N1 and H1N2 is possible in 223 experimental sera (Van Reeth et al., 2004; Van Reeth et al., 2006), nothing is known about 224 differentiation in field sera. In the field multiple infections can be expected, and variations 225 occur in haemagglutinin, even within one subtype (De Jong et al., 2001). Seroprevalence 226 and incidence of H1N1 and H1N2 might as a result be overestimated. However, if the 227 serological results of these two subtypes are compared at the age of 22 weeks, the 228 agreement beyond chance is very low (Kappa value 0.078). Under the assumption that both 229 subtypes occur independently, this suggests a very low level of cross-reactivity.

230 The seroprevalence of H1N1 is comparable to the levels that were found in finishing pigs 231 in 1980 (44%) (Masurel et al., 1983) and 1987 (62%) (Elbers et al., 1990) in the Netherlands. 232 The seroprevalence found for H3N2 is, however, much lower than the ones found in those 233 same studies (68% in 1980; 33% in 1987). All other seroprevalence studies carried out in 234 Europe in the past two decades also showed much higher seroprevalences (Haesebrouck 235 and Pensaert, 1986b; Yus et al., 1989; Teuffert et al., 1991; Groschup et al., 1993; Ewald et 236 al., 1994). The reason for the low seroprevalence therefore remains unknown and it remains 237 to be seen whether this is a structural decrease or just an uncommonly low seroprevalence 238 during a single season. The seroprevalence of H1N2 is not comparable to published results 239 from other countries, because they were carried out in sows (Van Reeth et al., 2000;

240 Maldonado et al., 2006).

Page 9 of 20

241 While the population dynamics of influenza virus infections is different between farrow-to-242 finish and finishing herds, it would be equally important to determine whether the occurrence 243 of clinical symptoms, and subsequent economic losses, are also different between both herd 244 types. Unfortunately, also with respect to this, no information can be found in existing 245 literature. Some studies that tried to determine the importance of influenza virus infections as 246 a cause for respiratory disease, found that approximately 50% of the respiratory outbreaks 247 were caused by influenza virus infections (Loeffen et al., 1999; Barigazzi and Donatelli, 2003; Choi et al., 2003). There was however no differentiation in herd type, or time of 248 249 infection. In this study, eleven episodes of respiratory disease were noticed, almost equally 250 distributed over both herd types. Only five of them coincided with a seroconversion against 251 one of the influenza subtypes of at least 80% of the pigs in the same period, also equally 252 distributed over both herd types. These five episodes are the ones most likely related to the 253 influenza virus infections, which is also consistent with the 50% of respiratory disease caused by influenza virus infections (Loeffen et al., 1999). Although this is not enough to 254 255 allow for any statistical analysis, it is noticeable that the two episodes of respiratory disease 256 in farrow-to-finish herds occurred between the age of 8 and 12 weeks, while the three 257 episodes in finishing herds occurred between the age of 16 and 22 weeks. Finishing pigs in 258 both herd types may be equally subject to respiratory disease, but at a different moment 259 during the finishing period. All three subtypes may have been involved in the occurrence of 260 clinical symptoms.

261 Differences in time of influenza virus infections in both herd types may ask for a different 262 approach in preventive measures, whether these are zoosanitary measures or vaccination. If 263 infections, and subsequent economic losses, in specialised finishing herds occur mainly at 264 the end of the finishing period, it would be possible to wait with vaccination, until maternal 265 antibodies have mostly disappeared. Interference of vaccination with maternal antibodies 266 (Wesley and Lager, 2006) would thus not be a real problem and full and equal protection 267 may be reached. It might even be considered to vaccinate only once at such a time. In 268 general, single vaccinations are considered to be insufficient for full protection. However,

partial protection was shown after single vaccination, that may be enough to sufficiently
reduce damages due to influenza virus infections under the right circumstances
(Haesebrouck and Pensaert, 1986a).

In farrow-to-finish herds on the other hand, there seems to be no optimum with respect to the time of vaccination. Before all maternal antibodies disappear in a population, most of the influenza virus infections, and possibly also the economic losses due to the infection, already occur. Vaccination at a young age, and probably multiple vaccinations to circumvent the negative effect of maternal antibodies, may be indicated.

With respect to zoosanitary intervention measures, the possibilities for influenza seem 277 278 somewhat limited. Zoosanitary measures may focus on reduction of damages (either due to 279 the primary infection or due to secondary infections). Mainly climate control and general 280 hygiene seem to be relevant zoosanitary measures for this purpose. Zoosanitary measures 281 may also focus on preventing infection by keeping the virus out. Given the ease with which 282 influenza viruses may spread, hygienic measures for this purpose probably have to be rather 283 strict, involving air filtration and strict hygiene protocols for anything and anyone entering the 284 premises. In farrow-to-finish herds this may be more difficult to achieve, also being more 285 expensive because it involved a broader range of age groups, than in specialized finishing 286 herds, given that most infections seem to occur at a later stage in the latter herd type.

287 While these conclusions and recommendations may apply to swine herds in general, 288 variations in individual herds may occur. It is therefore advisable that, where possible, results 289 from investigations in an individual herd are also taken into account before putting together a 290 strategy of vaccination or any other kind of intervention.

291

292 Acknowledgements

The authors thank the farmers for their willing cooperation and hospitality in this study.
The people at the lab are thanked for their dedication in testing all these samples. Bas Engel

and Willem Buist are thanked for the statistical analysis. This study was partly financed by
the Animal Health Service and Fort Dodge Animal Health.

297 **References**

298

- Barigazzi, G., Donatelli, I., 2003. Swine influenza in Italy. Vet. Res. Commun. 27 Suppl. 1,
 300 93-99.
- Brown, I.H., Chakraverty, P., Harris, P.A., Alexander, D.J., 1995a. Disease outbreaks in pigs
 in Great Britain due to an influenza A virus of H1N2 subtype. Vet. Rec. 136, 328-329.
- Brown, I.H., Harris, P.A., Alexander, D.J., 1995b. Serological studies of influenza viruses in
 pigs in Great Britain 1991- 2. Epidemiol. Infect. 114, 511-520.
- Brown, I.H., Harris, P.A., McCauley, J.W., Alexander, D.J., 1998. Multiple genetic
 reassortment of avian and human influenza A viruses in European pigs, resulting in the
 emergence of an H1N2 virus of novel genotype. J. Gen. Virol. 79 (Pt. 12), 2947-2955.
- 308 Choi, Y.K., Goyal, S.M., Joo, H.S., 2003. Retrospective analysis of etiologic agents 309 associated with respiratory diseases in pigs. Can. Vet. J. 44, 735-737.
- 310 De Jong, J.C., Heinen, P.P., Loeffen, W.L., Van Nieuwstadt, A.P., Claas, E.C., Bestebroer,
- 311 T.M., Bijlsma, K., Verweij, C., Osterhaus, A.D., Rimmelzwaan, G.F., Fouchier, R.A.,
- 312 Kimman, T.G., 2001. Antigenic and molecular heterogeneity in recent swine influenza
- 313 A(H1N1) virus isolates with possible implications for vaccination policy. Vaccine 19,
- 314 4452-4464.
- 315 Elbers, A.R.W., Tielen, M.J.M., Cromwijk, W.A.J., Hunneman, W.A., 1990. Sero-

316 epidemiological screening of pig sera collected at the slaughterhouse to detect herds

- 317 infected with Aujeszky's disease virus, porcine influenza virus and Actinobacillus
- 318 (Haemophilus) pleuropneumoniae in the framework of an integrated quality control (IQC)
- 319 system. Vet. Q. 12, 221-230.
- Elbers, A.R.W., Tielen, M.J.M., Cromwijk, W.A.J., Hunneman, W.A., 1992. Variation in
 seropositivity for some respiratory disease agents in finishing pigs: epidemiological

322	studies on some health parameters and farm and management conditions in the herds.
323	Vet. Q. 14, 8-13.
324	Ewald, C., Heer, A., Havenith, U., 1994. Factors Associated with Influenza-A Virus-Infections
325	in Finishing Pigs. Berliner und Münchener Tierärztliche Wochenschrift 107, 256-262.
326	Groschup, M.H., Brun, A., Haas, B., 1993. Serological studies on the potential synergism of
327	porcine reproductive and respiratory syndrome virus and influenza-, corona- and
328	paramyxoviruses in the induction of respiratory symptoms in swine. Zentralbl.
329	Veterinarmed. [B] 40, 681-689.
330	Haesebrouck, F., Pensaert, M.B., 1986a. Effect of intratracheal challenge of fattening pigs
331	previously immunised with an inactivated influenza H1N1 vaccine. Vet. Microbiol. 11,
332	239-49.
333	Haesebrouck, F., Pensaert, M.B., 1986b. Prevalence of H1N1- and H3N2-influenza A
334	viruses in fatteners [swine] in Belgium (1984). Vlaams Diergeneeskundig Tijdschrift 55,
335	12-16.
336	Kendal, A.P., Skehel, J.J., Pereira, M.S., 1982. Concepts and procedures for laboratory-
337	based influenza surveillance. US Department of Health and Human Services, Public
338	Health Service, Centers for Disease Control, Atlanta.
339	Loeffen, W.L.A., Kamp, E.M., Stockhofe-Zurwieden, N., Van Nieuwstadt, A.P.K.M.I.,
340	Bongers, J.H., Hunneman, W.A., Elbers, A.R.W., Baars, J., Nell, T., Van Zijderveld, F.G.,
341	1999. Survey of infectious agents involved in acute respiratory disease in finishing pigs.
342	Vet. Rec. 145, 123-129.
343	Loeffen, W.L.A., Nodelijk, G., Heinen, P.P., Van Leengoed, L.A., Hunneman, W.A.,
344	Verheijden, J.H., 2003. Estimating the incidence of influenza-virus infections in Dutch
345	weaned piglets using blood samples from a cross-sectional study. Vet. Microbiol. 91,
346	295-308.
347	Maes, D., Deluyker, H., Verdonck, M., Castryck, F., Miry, C., Vrijens, B., De Kruif, A., 1999.
348	Risk indicators for the seroprevalence of Mycoplasma hyopneumoniae, porcine influenza

349	viruses and Aujeszky's disease virus in slaughter pigs from fattening pig herds. Zentralbl.
350	Veterinarmed. [B] 46, 341-352.
351	Maldonado, J., Van Reeth, K., Riera, P., Sitja, M., Saubi, N., Espuna, E., Artigas, C., 2006.
352	Evidence of the concurrent circulation of H1N2, H1N1 and H3N2 influenza A viruses in
353	densely populated pig areas in Spain. Vet. J. 172, 377-381.
354	Marozin, S., Gregory, V., Cameron, K., Bennett, M., Valette, M., Aymard, M., Foni, E.,
355	Barigazzi, G., Lin, Y., Hay, A., 2002. Antigenic and genetic diversity among swine
356	influenza A H1N1 and H1N2 viruses in Europe. J. Gen. Virol. 83, 735-745.
357	Masurel, N., De Boer, G.F., Anker, W.J., Huffels, A.D., 1983. Prevalence of influenza viruses
358	A-H1N1 and A-H3N2 in swine in the Netherlands. Comp. Immunol. Microbiol. Infect. Dis.
359	6, 141-149.
360	Schrader, C., Suss, J., 2003. Genetic characterization of a porcine H1N2 influenza virus
361	strain isolated in Germany. Intervirology 46, 66-70.
362	Teuffert, J., Sinnecker, R., Karge, E., 1991. Seroepidemiologische Untersuchungen mit dem
363	HAHT zum Vorkommen porciner und humaner Infl-A-Viren bei Haus- und Wildschweinen
364	in der ehemaligen DDR (in German). Monatsh. Veterinarmed. 46, 171-174.
365	Thrusfield, M., 1995. Veterinary epidemiology, second edition. Blackwell Science Ltd,
366	Cambridge, 479 pp.
367	Van Reeth, K., Brown, I., Essen, S., Pensaert, M., 2004. Genetic relationships, serological
368	cross-reaction and cross-protection between H1N2 and other influenza A virus subtypes
369	endemic in European pigs. Virus Res. 103, 115-124.
370	Van Reeth, K., Brown, I.H., Pensaert, M., 2000. Isolations of H1N2 influenza A virus from
371	pigs in Belgium. Vet. Rec. 146, 588-589.
372	Van Reeth, K., Labarque, G., Pensaert, M., 2006. Serological profiles after consecutive
373	experimental infections of pigs with European H1N1, H3N2, and H1N2 swine influenza
374	viruses. Viral Immunol. 19, 373-382.

- Wesley, R.D., Lager, K.M., 2006. Overcoming maternal antibody interference by vaccination
 with human adenovirus 5 recombinant viruses expressing the hemagglutinin and the
- 377 nucleoprotein of swine influenza virus. Vet. Microbiol. 118, 67-75.

- Yus, E., Laviada, M.D., Moreno, L., Castro, J.M., Escribano, J.M., Simarro, I., 1989. The
- 379 prevalence of antibodies to influenza virus and respiratory coronavirus in fattening pigs in
- 380 Spain. Zentralbl. Veterinarmed. [B] 36, 551-556.
- 381
- 382
- 383

384	Table 1: Seroprevalences for swine influenza subtypes H1N1, H3N2, and H1N2, in
385	farrow-to-finish herds and finishing herds. Any significant (P<0.05) differences
386	between farrow-to-finish and finishing herds (pair wise comparisons for the same
387	influenza virus subtype over the same period) are indicated with a shaded
388	background. Any trends towards significance (P<0.1) are indicated by a square.
389	
390	Table 2: Incidences for swine influenza subtypes H1N1, H3N2, and H1N2, in farrow-to-
391	finish herds and finishing herds. Incidences were calculated based on the number of
392	seroconversions per period; in farrow-to-finish herds the periods of 0-8 and 8-12
393	weeks are added to each other to be able to compare them to the period of 0-12 weeks
394	in finishing pigs. Legend as in table 1.
395	
396	Table 3: Cumulative incidences for swine influenza subtypes H1N1, H3N2, and H1N2,
397	in farrow-to-finish herds and finishing herds. Incidences were calculated based on the
398	number of seroconversions per period. Legend as in table 1.
399	
400	Figure 1: Location of the three most pig-dense regions in the Netherlands and the
401	location of the herds participating in the longitudinal study (farrow-to-finish herds ($ abla$)
402	and finishing herds (Δ))
403	

ACCEPTED MANUSCRIPT

 \blacklozenge

								7				
			H1N1				H3N2		H1N2			
Herd type	Age (weeks)	Number of herds	Number of samples	Prevalence (%)	95%CI	Number of samples	Prevalence (%)	95%CI	Number of samples	Prevalence (%)	95%CI	
Farrow-	8	14	290	21.7	11.1-32.3	290	21.7	12.9-30.6	290	53.8	38.4-69.2	
to-finish	12	14	287	20.9	5.9-35.9	287	15.7	2.4-29.0	287	48.8	28.9-68.7	
	16	14	286	44.4	22.3-66.5	286	8.7	0.0-20.1	286	60.8	42.7-79.0	
	22	14	287	44.3	23.1-65.4	287	6.6	0.0-15.7	283	57.2	38.2-76.3	
Finishing	12	15	322	22.0	8.1-36.0	322	35.7	19.7-51.8	322	18.6	6.6-30.6	
	16	15	321	22.1	8.7-35.6	321	20.2	7.0-33.5	320	10.3	0.0-21.1	
	22	15	321	62.0	44.6-79.4	321	19.3	3.7-34.9	320	25.6	7.6-43.7	
		8										

ACCEPTED MANUSCRIP

			H1N1				H3N2		H1N2		
Herd type	Period (weeks of age)	Number of herds	Number of samples	Incidence (%)	95%CI	Number of samples	Incidence (%)	95%CI	Number of samples	Incidence (%)	95%CI
Farrow-	0-8	14	290	-	-	290	-	-	290	8.6	0.0-19.0
to-finish	8-12	14	287	15.7	2.7-28.6	287	10.1	0.0-23.3	287	24.4	8.5-40.3
	12-16	14	286	31.1	8.1-54.1	286	1.4	0.0-2.9	286	25.9	7.9-43.9
	16-22	14	285	2.5	0.0-7.1	285	1.1	0.0-2.6	281	1.4	0.3-2.6
Finishing	0-12	15	320	6.9	0.0-15.7	320	4.1	0.0-8.5	320	5.0	0.0-14.5
	12-16	15	320	13.8	4.8-22.7	320	10.3	0.9-19.7	319	2.8	0.0-5.9
	16-22	15	308	41.2	22.1-60.4	308	8.8	0.0-21.1	307	18.6	1.4-35.7

ACCEPTED MANUSCRIPT

			H1N1				H3N2		H1N2		
Herd type	Period (weeks of age)	Number of herds	Number of samples	Incidence (%)	95%CI	Number of samples	Incidence (%)	95%CI	Number of samples	Incidence (%)	95%CI
Farrow-	0-8	14	290	-	-	290	-	-	290	8.6	0.00-19.0
to-finish	0-12	14	287	15.7	2.7-28.6	287	10.1	0.0-23.3	287	33.1	12.7-53.4
	0-16	14	286	46.9	24.2-69.5	286	11.5	0.0-24.6	286	59.1	40.0-78.2
	0-22	14	285	49.5	26.6-72.3	285	12.6	0.0-25.8	281	60.5	41.4-79.6
Finishing	0-12	15	320	6.9	0.0-15.7	320	4.1	0.0-8.5	320	5.0	0.0-14.5
	0-16	15	320	20.6	8.6-32.6	320	14.4	3.3-25.4	319	7.8	0.0-17.5
	0-22	15	308	62.7	45.4-79.9	308	23.4	7.2-39.5	307	26.7	8.6-44.9
		~	C	X							

ACCEPTED MANUSCRIPT

