



HAL
open science

Berry-Esseen's central limit theorem for non-causal linear processes in Hilbert space

Mohamed El Machkouri

► **To cite this version:**

Mohamed El Machkouri. Berry-Esseen's central limit theorem for non-causal linear processes in Hilbert space. 2010. hal-00488676

HAL Id: hal-00488676

<https://hal.science/hal-00488676>

Preprint submitted on 3 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Berry-Esseen's central limit theorem for non-causal linear processes in Hilbert space

Mohamed EL MACHKOURI

June 2, 2010

Abstract

Let H be a real separable Hilbert space and $(a_k)_{k \in \mathbb{Z}}$ a sequence of bounded linear operators from H to H . We consider the linear process X defined for any k in \mathbb{Z} by $X_k = \sum_{j \in \mathbb{Z}} a_j(\varepsilon_{k-j})$ where $(\varepsilon_k)_{k \in \mathbb{Z}}$ is a sequence of i.i.d. centered H -valued random variables. We investigate the rate of convergence in the CLT for X and in particular we obtain the usual Berry-Esseen's bound provided that $\sum_{j \in \mathbb{Z}} |j| \|a_j\|_{\mathcal{L}(H)} < +\infty$ and ε_0 belongs to L_H^∞ .

Short title: Berry-Esseen's CLT for Hilbertian linear processes.

Key words: Central limit theorem, Berry-Esseen bound, linear process, Hilbert space.

1 Introduction and notations

Let $(H, \|\cdot\|_H)$ be a separable real Hilbert space and $(\mathcal{L}, \|\cdot\|_{\mathcal{L}(H)})$ be the class of bounded linear operators from H to H with its usual uniform norm. Consider a sequence $(\varepsilon_k)_{k \in \mathbb{Z}}$ of i.i.d. centered random variables, defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$, with values in H . If $(a_k)_{k \in \mathbb{Z}}$ is a sequence in \mathcal{L} , we define the (non-causal) linear process $X = (X_k)_{k \in \mathbb{Z}}$ in H by

$$X_k = \sum_{j \in \mathbb{Z}} a_j(\varepsilon_{k-j}), \quad k \in \mathbb{Z}. \quad (1)$$

If $\sum_{j \in \mathbb{Z}} \|a_j\|_{\mathcal{L}(H)} < \infty$ and $E\|\varepsilon_0\|_H < +\infty$ then the series in (1) converges almost surely and in $L_H^1(\Omega, \mathcal{A}, \mathbb{P})$ (see Bosq [2]). The condition $\sum_{j \in \mathbb{Z}} \|a_j\|_{\mathcal{L}(H)} < \infty$ is known to be sharp

for the \sqrt{n} -normalized partial sums of X to satisfies a CLT provided that $(\varepsilon_k)_{k \in \mathbb{Z}}$ are i.i.d. centered having finite second moments (see Merlevede et al. [6]). In this work, we investigate the rate of convergence in the CLT for X under the condition

$$\sum_{j \in \mathbb{Z}} |j|^\tau \|a_j\|_{\mathcal{L}(H)} < \infty \quad (2)$$

with $\tau = 1$ when $(\varepsilon_k)_{k \in \mathbb{Z}}$ are assumed to be i.i.d. centered and such that ε_0 belongs to L_H^∞ and $\tau = 1/2$ when $(\varepsilon_k)_{k \in \mathbb{Z}}$ are i.i.d. centered and such that ε_0 belongs to some Orlicz space $L_{H,\psi}$ (see section 2). This problem was previously studied (with $\tau = 1$ in Condition (2)) by Bosq [3] for (causal) Hilbert linear processes but a mistake in his proof was pointed out by V. Paulauskas [7]. However, in the particular case of Hilbertian autoregressive processes of order 1, Bosq [4] obtained the usual Berry-Esseen inequality provided that $(\varepsilon_k)_{k \in \mathbb{Z}}$ are i.i.d. centered with ε_0 in L_H^∞ .

2 Main result

In the sequel, C_{ε_0} is the autocovariance operator of ε_0 , $A := \sum_{j \in \mathbb{Z}} a_j$ and A^* is the adjoint of A . For any sequence $Z = (Z_k)_{k \in \mathbb{Z}}$ of random variables with values in H we denote

$$\Delta_n(Z) = \sup_{t \in \mathbb{R}} \left| \mathbb{P} \left(\left\| \frac{1}{\sqrt{n}} \sum_{k=1}^n Z_k \right\|_H \leq t \right) - \mathbb{P} (\|N\|_H \leq t) \right|$$

where $N \sim \mathcal{N}(0, AC_{\varepsilon_0}A^*)$.

For any $j \in \mathbb{Z}$, denote $c_{j,n} = \sum_{i=1}^n b_{i-j}$ where $b_i = a_i$ for any $i \neq 0$ and $b_0 = a_0 - A$.

Lemma 1 *For any positive integer n ,*

$$\sum_{k=1}^n X_k = A \left(\sum_{k=1}^n \varepsilon_k \right) + Q_n + R_n$$

where $Q_n = \sum_{k=1}^n \sum_{|j|>n} a_{k-j}(\varepsilon_j)$ and $R_n = \sum_{|j| \leq n} c_{j,n}(\varepsilon_j)$.

Recall that a Young function ψ is a real convex nondecreasing function defined on \mathbb{R}^+ which satisfies $\lim_{t \rightarrow +\infty} \psi(t) = +\infty$ and $\psi(0) = 0$. We define the Orlicz space $L_{H,\psi}$ as the space of H -valued random variables Z defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such

that $E[\psi(\|Z\|_H/c)] < +\infty$ for some $c > 0$. The Orlicz space $L_{H,\psi}$ equipped with the so-called Luxemburg norm $\|\cdot\|_\psi$ defined for any H -valued random variable Z by

$$\|Z\|_\psi = \inf\{c > 0; E[\psi(\|Z\|_H/c)] \leq 1\}$$

is a Banach space. In the sequel, $c(N)$ denotes a bound of the density of $\mathcal{N}(0, AC_{\varepsilon_0}A^*)$ (see Davydov et al. [5]). Our main result is the following.

Theorem 1 *Let $(\varepsilon_k)_{k \in \mathbb{Z}}$ be a sequence of i.i.d. centered H -valued random variables and let X be the Hilbertian linear process defined by (1).*

i) If ε_0 belongs to L_H^∞ and $\sum_{j \in \mathbb{Z}} |j| \|a_j\|_{\mathcal{L}(H)} < \infty$ then

$$\Delta_n(X) \leq \frac{c_1}{\sqrt{n}} \quad (3)$$

where $c_1 = c_2 + 14c(N)\|\varepsilon_0\|_\infty \sum_{j \in \mathbb{Z}} |j| \|a_j\|_{\mathcal{L}(H)}$ and c_2 is a positive constant which depend only on the distribution of ε_0 .

ii) If ψ is a Young function then

$$\Delta_n(X) \leq \Delta_n(A(\varepsilon)) + \varphi\left(\frac{c(N)\|Q_n + R_n\|_\psi}{\sqrt{n}}\right) \quad (4)$$

where $\varphi(x) = xh^{-1}(1/x)$ and $h(x) = x\psi(x)$ for any real $x > 0$.

The inequality (4) ensures a rate of convergence to zero for $\Delta_n(X)$ as n goes to infinity provided that $\Delta_n(A(\varepsilon_0))$ goes to zero as n goes to infinity and a bound for $\|Q_n + R_n\|_\psi$ exists. As just an illustration, we have the following corollary.

Corollary 1 *Assume that $(\varepsilon_k)_{k \in \mathbb{Z}}$ are i.i.d. centered H -valued random variables and that the condition (2) holds with $\tau = 1/2$.*

i) If ε_0 belongs to L_{H,ψ_1} then $\Delta_n(X) = O\left(\frac{\log n}{\sqrt{n}}\right)$ where ψ_1 is the Young function defined by $\psi_1(x) = \exp(x) - 1$.

ii) If ε_0 belongs to L_H^r for $r \geq 3$ then $\Delta_n(X) = O\left(n^{-\frac{r}{2(r+1)}}\right)$.

3 Proofs

Proof of Lemma 1. For any positive integer n , we have

$$\begin{aligned}
R_n &= \sum_{j=-n}^n c_{j,n}(\varepsilon_j) = \sum_{k=1}^n \sum_{j=-n}^n b_{k-j}(\varepsilon_j) \\
&= \sum_{k=1}^n \sum_{j \in [-n,n] \setminus \{k\}} a_{k-j}(\varepsilon_j) + (a_0 - A) \left(\sum_{k=1}^n \varepsilon_k \right) \\
&= \sum_{k=1}^n \sum_{j=-n}^n a_{k-j}(\varepsilon_j) - A \left(\sum_{k=1}^n \varepsilon_k \right) \\
&= -Q_n + \sum_{k=1}^n X_k - A \left(\sum_{k=1}^n \varepsilon_k \right).
\end{aligned}$$

The proof of Lemma 1 is complete.

Proof of Theorem 1. Let $\lambda > 0$ and $t > 0$ be fixed and denote $U = A(\sum_{k=1}^n \varepsilon_k / \sqrt{n})$ and $V = (Q_n + R_n) / \sqrt{n}$. So $U + V = \sum_{k=1}^n X_k / \sqrt{n}$ and

$$\mathbb{P}(\|U + V\|_H \leq t) \leq \mathbb{P}(\|U\|_H \leq t + \lambda) + \mathbb{P}(\|V\|_H \geq \lambda) \quad (5)$$

For $\lambda_0 = 2\|V\|_\infty$, we obtain

$$\mathbb{P}(\|U + V\|_H \leq t) - \mathbb{P}(\|N\|_H \leq t) \leq \mathbb{P}(\|U\|_H \leq t + \lambda_0) - \mathbb{P}(\|N\|_H \leq t).$$

If $c(N)$ denotes a bound for the density of $\|N\|_H$ (see Davydov et al. [5]) then

$$\Delta_n(X) \leq \Delta_n(A(\varepsilon)) + \frac{2c(N)\|Q_n + R_n\|_\infty}{\sqrt{n}}.$$

Noting that

$$Q_n = \sum_{j \geq n+2} a_j \left(\sum_{k=1-j}^{-n-1} \varepsilon_k \right) + \sum_{j < 0} a_j \left(\sum_{k=n+1}^{n-j} \varepsilon_k \right) \quad (6)$$

and

$$R_n = R'_n + R''_n \quad (7)$$

where

$$R'_n = - \sum_{j=-n}^{-1} a_j \left(\sum_{k=1}^{-j} \varepsilon_k \right) - \sum_{j < -n} a_j \left(\sum_{k=1}^n \varepsilon_k \right) - \sum_{j > 0} a_j \left(\sum_{k=n-j+1}^n \varepsilon_k \right)$$

and

$$R_n'' = \sum_{j=1}^n a_j \left(\sum_{k=-j+1}^0 \varepsilon_k \right) + \sum_{j=n+1}^{2n} a_j \left(\sum_{k=-n}^{n-j} \varepsilon_k \right),$$

we derive that $\|Q_n + R_n\|_\infty \leq 7\|\varepsilon_0\|_\infty \sum_{j \in \mathbb{Z}} |j| \|a_j\|_{\mathcal{L}(H)}$ and consequently

$$\Delta_n(X) \leq \Delta_n(A(\varepsilon)) + \frac{14c(N)\|\varepsilon_0\|_\infty \sum_{j \in \mathbb{Z}} |j| \|a_j\|_{\mathcal{L}(H)}}{\sqrt{n}}.$$

Combining the last inequality with the Berry-Esseen inequality for i.i.d. centered H -valued random variables (see Yurinski [9] or Bosq [2], Theorem 2.9) we obtain (3).

In the other part, if ψ is a Young function we have $\mathbb{P}(\|V\|_H \geq \lambda) \leq \frac{1}{\psi(\lambda/\|V\|_\psi)}$ and keeping in mind inequality (5), we derive

$$\Delta_n(X) \leq \Delta_n(A(\varepsilon)) + c(N)\lambda + \frac{1}{\psi(\lambda/\|V\|_\psi)}.$$

Noting that $c(N)\lambda = \frac{1}{\psi(\lambda/\|V\|_\psi)}$ if and only if $\lambda = \frac{\varphi(c(N)\|V\|_\psi)}{c(N)}$ where φ is defined by $\varphi(x) = xh^{-1}(1/x)$ and h by $h(x) = x\psi(x)$, we conclude

$$\Delta_n(X) \leq \Delta_n(A(\varepsilon)) + \varphi\left(\frac{c(N)\|Q_n + R_n\|_\psi}{\sqrt{n}}\right).$$

The proof of Theorem 1 is complete.

Proof of Corollary 1. Assume that $\|\varepsilon_0\|_{\psi_1} < \infty$ where ψ_1 is the Young function defined by $\psi_1(x) = \exp(x) - 1$. There exists $a > 0$ such that $E(\exp(a\|\varepsilon_0\|_H)) \leq 2$. So, there exist (see Arak and Zaizsev [1]) constants B and L such that

$$E\|\varepsilon_0\|_H^m \leq \frac{m!}{2} B^2 L^{m-2}, \quad m = 2, 3, 4, \dots$$

Applying Pinelis-Sakhanenko inequality (see Pinelis and Sakhanenko [?] or Bosq [2]), we obtain

$$\mathbb{P}\left(\left\|\sum_{k=p}^q \varepsilon_k\right\|_H \geq x\right) \leq \exp\left(-\frac{x^2}{2(q-p+1)B^2 + 2xL}\right), \quad x > 0$$

and using Lemma 2.2.10 in Van Der Vaart and Wellner [?], there exists a universal constant K such that

$$\left\|\sum_{k=p}^q \varepsilon_k\right\|_{\psi_1} \leq K \left(L + B\sqrt{q-p+1}\right) \quad (8)$$

Combining (6), (7) and (8), we derive $\|Q_n + R_n\|_{\psi_1} \leq C \sum_{j \in \mathbb{Z}} \sqrt{|j|} \|a_j\|_{\mathcal{L}(H)}$ where the constant C does not depend on n . Keeping in mind the Berry-Esseen's central limit theorem for i.i.d. centered H -valued random variables (see Yurinski [9] or Bosq [2], Theorem 2.9), we apply Theorem 1 with the Young function ψ_1 . Since the function φ defined by $\varphi(x) = xh^{-1}(1/x)$ with $h(x) = x\psi_1(x)$ satisfies

$$\lim_{x \rightarrow 0} \frac{\varphi(x)}{x \log(1 + \frac{1}{x})} = 0,$$

we derive $\Delta_n(X) = O\left(\frac{\log n}{\sqrt{n}}\right)$.

Now, assume that $\|\varepsilon_0\|_r < \infty$ for some $r \geq 3$. Applying Pinelis inequality (see Pinelis [8]), there exists a universal constant K such that

$$\left\| \sum_{k=p}^q \varepsilon_k \right\|_r \leq K \left(r \left(\sum_{k=p}^q E \|\varepsilon_k\|_H^r \right)^{1/r} + \sqrt{r} \left(\sum_{k=p}^q E \|\varepsilon_k\|_H^2 \right)^{1/2} \right)$$

and consequently

$$\left\| \sum_{k=p}^q \varepsilon_k \right\|_r \leq 2Kr \|\varepsilon_0\|_r \sqrt{q-p+1}. \quad (9)$$

Combining (6), (7) and (9), we derive $\|Q_n + R_n\|_r \leq C \sum_{j \in \mathbb{Z}} \sqrt{|j|} \|a_j\|_{\mathcal{L}(H)}$ where the constant C does not depend on n . Again, applying Berry-Esseen's CLT (see Yurinski [9] or Bosq [2], Theorem 2.9) and Theorem 1 with the Young function $\psi(x) = x^r$ and the function φ given by $\varphi(x) = x^{r/(r+1)}$, we obtain $\Delta_n(X) = O\left(n^{-\frac{r}{2(r+1)}}\right)$. The proof of Corollary 1 is complete.

References

- [1] T. V. Arak and A. Yu. Zaitsev. Uniform limit theorems for sums of independent random variables. *Proc. Steklov Inst. Math.*, (1(174)), 1988. A translation of Trudy Mat. Inst. Steklov. **174** (1986).
- [2] Denis Bosq. *Linear processes in function spaces*, volume 149 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 2000. Theory and applications.
- [3] Denis Bosq. Berry-Esseen inequality for linear processes in Hilbert spaces. *Statist. Probab. Lett.*, 63(3):243–247, 2003.

- [4] Denis Bosq. Erratum and complements to: “Berry-Esseen inequality for linear processes in Hilbert spaces” [Statist. Probab. Lett. **63** (2003), no. 3, 243–247; mr1986323]. *Statist. Probab. Lett.*, 70(2):171–174, 2004.
- [5] Yu. A. Davydov, M. A. Lifshits, and N. V. Smorodina. *Local properties of distributions of stochastic functionals*, volume 173 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1998. Translated from the 1995 Russian original by V. E. Nazaïkinskiĭ and M. A. Shishkova.
- [6] Florence Merlevède, Magda Peligrad, and Sergey Utev. Sharp conditions for the CLT of linear processes in a Hilbert space. *J. Theoret. Probab.*, 10(3):681–693, 1997.
- [7] V. Paulauskas. Personal communication, 2004.
- [8] Iosif Pinelis. Optimum bounds for the distributions of martingales in Banach spaces. *Ann. Probab.*, 22(4):1679–1706, 1994.
- [9] V. V. Yurinskii. On the accuracy of normal approximation of the probability of hitting a ball. *Teor. Veroyatnost. i Primenen.*, 27(2):270–278, 1982.

Mohamed EL MACHKOURI
 Laboratoire de Mathématiques Raphaël Salem
 UMR CNRS 6085, Université de Rouen
 Avenue de l’université
 76801 Saint-Etienne du Rouvray
 mohamed.elmachkouri@univ-rouen.fr