



HAL
open science

Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse

C.A. Castillo, J.L. Albasanz, D. León, J. Jordán, M. Pallàs, A. Camins, M.
Marti'N

► **To cite this version:**

C.A. Castillo, J.L. Albasanz, D. León, J. Jordán, M. Pallàs, et al.. Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse. *Experimental Gerontology*, 2009, 44 (6-7), pp.453. 10.1016/j.exger.2009.04.006 . hal-00493105

HAL Id: hal-00493105

<https://hal.science/hal-00493105>

Submitted on 18 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse

C.A. Castillo, J.L. Albasanz, D. León, J. Jordán, M. Pallàs, A. Camins, M. Martí ´n


PII: S0531-5565(09)00083-7
DOI: [10.1016/j.exger.2009.04.006](https://doi.org/10.1016/j.exger.2009.04.006)
Reference: EXG 8605

To appear in: *Experimental Gerontology*

Received Date: 2 February 2009
Revised Date: 3 April 2009
Accepted Date: 21 April 2009

Please cite this article as: Castillo, C.A., Albasanz, J.L., León, D., Jordán, J., Pallàs, M., Camins, A., Martí ´n, M., Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse, *Experimental Gerontology* (2009), doi: [10.1016/j.exger.2009.04.006](https://doi.org/10.1016/j.exger.2009.04.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



AGE-RELATED EXPRESSION OF ADENOSINE RECEPTORS IN BRAIN
FROM THE SENESENCE-ACCELERATED MOUSE

Castillo¹ C.A., Albasanz¹ J.L., León¹ D., Jordán² J., Pallàs³ M., Camins³ A. and
Martín¹, M*.

¹ Área de Bioquímica, Departamento de Química Inorgánica, Orgánica y Bioquímica,
Facultad de Química, Centro Regional de Investigaciones Biomédicas (CRIB),
Universidad de Castilla-La Mancha, Ciudad Real, Spain.

² Departamento de Ciencias Médicas, Facultad de Medicina, Centro Regional de
Investigaciones Biomédicas (CRIB), Universidad de Castilla-La Mancha, Albacete,
Spain.

³ Departamento de Farmacología y Química Terapéutica, Facultad de Farmacia, Centros
de Investigación Biomédica en Red de Enfermedades Neurodegenerativas
(CIBERNED), Universidad de Barcelona, Barcelona, Spain.

* Corresponding author: Dr. Mairena Martín López

Facultad de Químicas, Universidad de Castilla-La Mancha

Avenida Camilo José Cela, 10

13071 Ciudad Real, Spain

Tel: +34 92 629 5300 Ext 6279

Fax: +34 92 629 5318

e-mail: mairena.martin@uclm.es

Running title: Adenosine receptors in SAM

Keywords: Senescence-accelerated mouse, adenosine receptors, aging.

Abstract

1
2 Senescence-accelerated mice (SAM) are used as a model of aging and age-associated
3
4 diseases. SAMP8 are prone strains that show shortened life span and deficits in learning
5
6 and memory processes, while SAMR1 are strains of accelerated senescence-resistant,
7
8 long-lived mice. Due to their abnormal APP (amyloid precursor protein) metabolism in
9
10 brain, SAMP8 may be an Alzheimer-type model. Adenosine receptors are G-protein
11
12 coupled receptors which are altered in brain from Alzheimer Disease (AD) cases. The
13
14 analysis of adenosine receptors in brain from young (21 days old) and middle-aged (180
15
16 days old) SAMP8 as compared with SAMR1 mice revealed differences between these
17
18 strains associated with age. The age-related increase in mRNA coding A₁ and A_{2B}
19
20 receptors observed in SAMR1 was absent in SAMP8. A₁ receptors were significantly
21
22 decreased with age in SAMR1, while no differences were observed in SAMP8.
23
24 However, the levels of A₁ receptors in young SAMP8 were even lower than those
25
26 obtained in middle-aged SAMR1. In addition, A_{2A} receptors were significantly
27
28 increased only in aged SAMR1. A similar age-related decrease in A₁ receptors level was
29
30 also observed in brain from male Wistar rats. These results suggest different age-related
31
32 effects on adenosine receptors in SAMR1 and SAMP8 strains. Since A₁ receptors are
33
34 mainly neuroprotective, their important loss in very young SAMP8 strain suggests the
35
36 involvement of these receptors in the pathogenesis of neurodegenerative diseases
37
38 associated with aging.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

1
2 Senescence-accelerated prone mouse strains (SAMP) and resistant strains
3 (SAMR) were established by Takeda and colleagues at Kyoto University by selective
4 breeding from their AKR/J colony in the early 1970s (Takeda et al., 1981). Prone strains
5 of SAM (SAMP8) show shortened life span and early manifestation of senescence with
6 characteristic pathological phenotypes similar to those often observed in elderly
7 humans. These include senile osteoporosis, osteoarthritis, age-related deficits in
8 learning and memory with/without forebrain atrophy, senile amyloidosis, et al. SAMR1
9 are strains of accelerated senescence-resistant, long-lived mice. The mean life span in
10 the P series is about 9.7 months, 40% shorter than that of the R series (16.3 months).
11 Senescence accelerated mouse prone strain SAMP8 is one of the most appropriate
12 models to study aging and age-associated diseases because it shows an age-related
13 deterioration of learning and memory at an earlier age, compared with the control
14 mouse R1 strain in senescence-accelerated resistant mouse (SAMR1) (Miyamoto et al.,
15 1986; Miyamoto, 1997; Nomura and Okuma, 1999; Butterfield and Poom, 2005).
16 Alteration of learning and memory of SAMP8 have been associated with a decline in
17 neurotransmitter activity as well as decreased serotonin, increased GABA and decreased
18 cholinergic activity (Morley et al., 2002b). In addition, cognitive defect in these animals
19 can be due to overproduction of β -amyloid peptide (A β P) being reversed by specific
20 antibodies to β -amyloid. Furthermore, SAMP8 mice present lower mRNA levels of
21 apolipoprotein E and increased levels of presenilin-2 when compared to SAMR1 (Wei
22 et al., 1999). Therefore, these animals have been suggested as a good model for the
23 study of Alzheimer's disease (Morley, 2002; Morley et al., 2002a; Banks et al., 2007;
24 Pallas et al., 2008).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
SAMP8 mice also present other characteristics that contribute to their pathological phenotype. SAMP8 mice present a lower expression of neurotrophic genes as glial cell derived neurotrophic factor (GDNF) (Miyazaki et al., 2003), neurotrophin-3 (NT-3) and nerve growth factor (NGF) (Kaisho et al., 1994). Furthermore, the expression of proteins involved in reactive oxygen species (ROS) metabolism is altered in SAMP8 mice, which may be involved in the increased oxidative stress found in SAMP8 mouse brain. Compared to SAMR1 mice, the activity of enzymes involved in antioxidant processes are decreased in the cerebral cortex of aged SAMP8. This has been reported for manganese superoxide dismutase (Mn-SOD), glutamine synthase (GS), glutathione peroxidase (GPx) and peroxidase activities (Kurokawa et al., 2001; Sato et al., 1996; Okatani et al., 2002). Moreover, other enzymes which contribute to ROS generation, such as nitric oxide synthase (NOS), are increased in aged SAMP8 mice (Inada et al., 1996).

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Adenosine is a nucleoside widely distributed in central and peripheral nervous system that exerts its actions through four types of receptors named A₁, A_{2A}, A_{2B} and A₃, all of them being G-protein coupled receptors (GPCR). A₁ and A₃ receptors are coupled, through Gi/o proteins, to adenylyl cyclase activity inhibition, while A_{2A} and A_{2B} receptors are coupled to stimulation of the enzymatic activity, through Gs protein (Ralevic and Burnstock, 1998, Fredholm et al., 2001; 2005). Out of the four adenosine receptors, the A₁ subtype is the most abundant and widespread in the brain, where it plays a neuroprotective role because of its capacity to decrease the release of excitatory neurotransmitters, mainly glutamate (Dunwiddie and Masino, 2001). A_{2A} receptors are concentrated in the basal ganglia but they are also present throughout the brain, albeit in a considerably lower density. A_{2B} and A₃ receptors are the least abundant in the brain (Cunha, 2005).

1
2 Changes in adenosine receptors during aging have been demonstrated in several
3
4 animals (Cunha, 2005) and in humans (Meyer et al., 2007). Decreased expression and
5
6 density of A₁ in the cortical and hippocampal regions has been observed (Cheng et al.,
7
8 2000; Cunha et al., 1995, 2001a, b; Paganopoulou et al., 1992) whereas the density of
9
10 A_{2A} receptors increases with age (Cunha et al., 1995; Cunha, 2005; Lopes et al., 1999).
11
12 However, these age-related changes in adenosine receptor levels observed in some brain
13
14 areas have not been detected in the striatum, where these levels might be only slightly
15
16 affected by ageing (Cunha et al., 1995). Beyond this, adenosine receptors have not been
17
18 analyzed in SAM strains until now. The SAMP8 mouse has been proposed as an
19
20 excellent model for studying the pathogenesis of learning and memory disturbances
21
22 associated with AβP overproduction. The identification of senile plaques and the
23
24 increased expression of AβP in this mouse model suggest it may be an acceptable model
25
26 for Alzheimer's disease (Morley, 2002; Morley et al., 2002 a,b; Banks et al., 2007;
27
28 Pallas et al., 2008). We have previously reported that adenosine A₁ and A_{2A} receptors
29
30 are altered in frontal cortex brain from AD patients (Albasanz et al., 2008). The aim of
31
32 the present work was to determine the expression of different adenosine receptors,
33
34 mainly A₁ and A_{2A}, and their possible age-related changes in brains from SAMR1 and
35
36 SAMP8 strains, in order to use these animals as models to study the mechanism
37
38 involved in the neuropathogenesis of Alzheimer's disease.
39
40
41
42
43
44
45
46
47
48
49
50

51 **2. Materials and Methods**

52 **2.1. Materials**

53
54 Cyclopentyl-1,3-dipropylxanthine,8-[dipropyl-2,3-³H(N)] ([³H]DPCPX) 120 Ci/
55
56 mmol) and adenosine 3', 5'-cyclic phosphate [2,8-³H] ([³H]cAMP) 27.4 Ci/ mmol) were
57
58
59
60
61
62
63
64
65

1 purchased from PerkinElmer (Madrid, Spain). Anti-A₁ antibody was purchased from
2 Calbiochem (Bionova, Madrid, Spain), anti-A_{2A} antibody from Upstate (Millipore,
3 Madrid, Spain) and anti-β-actin from Abcam (Cambridge, UK). Guanosine-5'-O(3-
4 thiotriphosphate) tetralithium salt (GTP_γS) and calf intestine adenosine deaminase
5 (ADA) were purchased from Roche (Barcelona, Spain). Creatine kinase, Creatine
6 phosphate, Protein kinase A, 4-(3-butoxy-4-methoxybenzyl)imidazolidin-2-one (Ro 20-
7 1724), Forskolin, N⁶-cyclopentyladenosine (CPA), N⁶-cyclohexyladenosine (CHA),
8 Adenosine 5'-triphosphate (ATP) and Guanosine 5'-[β,γ-imido]triphosphate
9 (Gpp(NH)p) were from Sigma (Madrid, Spain). All other reagents were of analytical
10 grade and obtained from commercial sources.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **2.2. Animals**

28
29 Young (21 days old) and middle-aged (180 days old) SAMR1 and SAMP8 mice
30 or young male (3 months) and aged male (24 months) Wistar rats were kept at constant
31 temperature (25 ± 1 °C) and humidity, maintained on a 12 h light/12 h dark cycle with
32 *ad libitum* access to rodent food and water. All experiments followed the European
33 Union regulations about animal experimentation and those of the Experimentation
34 Animal Committee of the Castilla-La Mancha University. Special care was taken to
35 minimize animal suffering and to reduce the number of animals used.
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **2.3. Plasma membrane isolation**

51 Plasma membranes from mouse and rat brain were isolated as described
52 previously (León et al., 2002). Samples were homogenized in 20 volumes of isolation
53 buffer (50 mM Tris-HCl, pH 7.4 containing 10 mM MgCl₂ and protease inhibitors) in
54 Dounce homogenizer (10xA, 10xB). After homogenization, whole brain preparations
55
56
57
58
59
60
61
62
63
64
65

(excluding cerebellum and spinal cord) were centrifuged for 5 min at 1000 x g in a Beckman JA 21 centrifuge. Supernatant was centrifuged for 20 min at 27000 x g and the pellet was finally resuspended in isolation buffer.

2.4. [³H]DPCPX binding assays to plasma membranes

Binding assays to plasma membranes from mouse and rat brain were performed as described previously (León et al., 2002). Plasma membranes were incubated with 5 U/mg adenosine deaminase (ADA) in 50 mM Tris, 2 mM MgCl₂, pH 7.4, for 30 min at 25°C, in order to eliminate endogenous adenosine from membrane preparations. Then, plasma membranes (50-75 µg of protein) were incubated with [³H]DPCPX for 2 h at 25°C. Saturation assays were carried out at different [³H]DPCPX concentrations (0.1-20 nM) using CPA at a concentration 10⁵ times higher than the radioligand, in order to obtain non-specific binding. Binding assays were stopped by rapid filtration through Whatman GF/B filters, previously pre-incubated with 0.3% polyethylenimine using a FilterMate Harvester (Perkin-Elmer). Scintillation liquid mixture was added in order to measure radioactivity in a Microbeta Trilux (Perkin Elmer) liquid scintillation counter.

When the effect of Gpp(NH)p was studied [³H]DPCPX binding assays were performed at saturation concentration using the same methodology with minor modifications. Briefly, the buffer employed was 50 mM Tris, pH 7.4, containing 10 µM Gpp(NH)p, and the final concentration of [³H]DPCPX used was 20 nM.

2.5. Western blot assays

For Western blot assays, 50 µg of protein was mixed with loading buffer containing 0.125 M Tris (pH 6.8), 20% glycerol, 10% β-mercaptoethanol, 4% SDS and 0.002% bromophenol blue, and heated at 95°C for 5 min. Protein was electrophoresed

1 on a 10% SDS-PAGE gel using a mini-protean system (Bio-Rad) with molecular weight
2 standards (Bio-Rad). Protein transfer to nitrocellulose membranes was carried out in
3
4 iBlot™ Dry Blotting System (Invitrogen). Membranes were washed with PBS-tween-
5
6 20, blocked with PBS containing 5% skimmed milk, and then incubated with the
7
8 primary antibodies at 4°C overnight (1:1000 dilution for anti-A₁R, 1:500 dilution for
9
10 anti-A_{2A}R and 1:5000 dilution for anti-β-actin). After rinsing, the membranes were
11
12 incubated with the corresponding secondary antibody (Bio-Rad) at a dilution of 1:5000
13
14 in PBS containing 5% skimmed milk for 30 minutes. Antigen was visualized using the
15
16 ECL chemiluminescence detection kit (Amersham), and specific bands were quantified
17
18 by densitometry using Multianalyst 1.0 software (Bio-Rad). Membranes were stripped
19
20 for a second probe by incubating with 0.15% glycine, 0.1% SDS, and 1% tween-20, pH
21
22 2.2, for 30 minutes at room temperature. After removing the stripping buffer and
23
24 washing them twice with PBS, membranes were ready for the blocking stage.
25
26
27
28
29
30
31
32
33

34 **2.6. Preparation of Total RNA and cDNA**

35
36 Total RNA was extracted from whole brain (excluding cerebellum and spinal
37
38 cord) using an ABI 6100 Nucleic Acid PrepStation according to the manufacturer's
39
40 protocol. All chemicals for the ABI 6100 were purchased from Applied Biosystems
41
42 (Foster City, CA). Total RNA from mice was isolated and stored individually at -80°C.
43
44 Ratio of A₂₆₀/A₂₈₀ (purity of RNA) was in the range 1.9–2.1. RNA concentrations were
45
46 determined from the A₂₆₀. One microgram of total RNA was reverse transcribed using
47
48 Applied Biosystems High-Capacity cDNA Archive Kit.
49
50
51
52
53
54
55

56 **2.7. Quantitative Real Time RT-PCR Analysis**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Quantitative real time RT-PCR analysis (Higuchi et al, 1993) was performed with an Applied Biosystems Prism 7500 Fast Sequence Detection System using TaqMan® universal PCR master mix according to the manufacturer's specifications (Applied Biosystems Inc., Foster City, CA). The validated TaqMan probes and primers for A₁ (assay ID: Rn00567668_m1), A_{2A} (assay ID: Rn00583935_m1), A_{2B} (assay ID: Rn00567697_m1), A₃ (assay ID: Rn00563680_m1) and β-actin (Rn00667869_m1) were assay-on-demand gene expression products (Applied Biosystems). The TaqMan® primer and probe sequences are packaged together in a 20x solution. Rat β-actin gene was used as endogenous control. The thermal cycler conditions were as follows: hold for 20 s at 95 °C, followed by two-step PCR for 40 cycles of 95 °C for 3 s, followed by 60 °C for 30 s. Levels of RNA expression were determined using the 7500 Fast System SDS software version 1.3.1 (Applied Biosystems) according to the 2^{-ΔΔCt} method. Briefly, expression results for a gene were normalized to internal control β-actin relative to a calibrator, consisting of the mean expression level of the corresponding gene in control samples as follows: $2^{-\Delta\Delta C_t} = 2^{-((C_t \text{ receptor gene} - C_t \text{ actin})_{\text{sample}} - (C_t \text{ receptor gene} - C_t \text{ actin})_{\text{calibrator}})}$. The results from three independent repeat assays, performed on different plates each using different cDNAs from the animals analyzed, were averaged to produce a single mean quantity value for each mRNA for each animal.

2.8. Adenylyl cyclase activity determination

Adenylyl cyclase activity was determined in brain plasma membranes as previously described (León et al., 2005). Assays were performed with 10-20 μg of protein in a final volume of 0.25 mL of 50 mM Tris-HCl pH 7.4, 5 mM MgCl₂, 1 mM 1,4- Dithiothreitol, 1 mg/mL bovine serum albumin, 1 mg/mL creatine kinase, 10 mM creatine phosphate, and 0.1 mM Ro 20-1724 (a specific phosphodiesterase inhibitor).

1 Plasma membranes, previously incubated with adenosine deaminase (0.2 U/mg protein)
2 at 37°C for 30 min to remove endogenous adenosine, were incubated for 6 min at 30°C
3
4 in the absence or presence of 5 μ M GTP γ S, 10 μ M forskolin, or 1 mM CHA, followed
5
6 by a further incubation with 200 μ M ATP at 30°C for 10 min. Reaction was stopped by
7
8 the boiling of samples and then centrifugation at 12000 x g for 4 min. Fifty microliters
9
10 of supernatant were used to determine cAMP accumulation. Samples were incubated
11
12 with 0.25 pmol of [³H]cAMP and 6.25 μ g protein kinase A in a final volume of 200 μ L
13
14 of buffer assay (Tris-HCl 50 mM, pH 7.4, EDTA 4 mM) for 2 h at 4°C. Standard
15
16 samples were prepared in the same buffer in the range of concentration of 0-16 pmol.
17
18 Reaction was stopped by the addition of 100 μ L of active charcoal solution containing
19
20 bovine serum albumin and then centrifugation of the samples at 12000 x g for 4 min.
21
22 Radioactivity was measured in a Microbeta Trilux (Perkin Elmer) liquid scintillation
23
24 counter.
25
26
27
28
29
30
31
32
33

34 **2.9. Protein determination**

35
36 Protein concentration was measured by the method of Lowry et al. (1951), using
37
38 bovine serum albumin as standard.
39
40
41
42
43

44 **2.10. Statistical and data analysis**

45
46 Data are mean \pm SEM. Statistical analysis was performed using Student t-test.
47
48 Differences between mean values were considered statistically significant at $p < 0.05$.
49
50 Non-linear regression curve fit (one site binding hyperbola) was performed for binding
51
52 data using the GraphPad Prism 5.0 program for Windows (GraphPad Software, San
53
54 Diego, CA, USA).
55
56
57
58
59
60
61
62
63
64
65

3. Results

3.1. Expression of genes coding adenosine receptor types in SAMR1 and SAMP8 mice

To determine the age-related changes in expression of gene coding adenosine receptors in SAMP8 and SAMR1 mice, we isolated total RNA from both strains and performed quantitative real time PCR using selective oligonucleotides to adenosine A₁, A_{2A}, A_{2B} and A₃ adenosine receptors. The ages of animals were three weeks (21d) and six months (180d). Results show that mRNA coding adenosine A₁ receptors are significantly increased in middle-aged versus young SAMR1 animals, suggesting an age-associated up-regulation in resistant control mice. However, the level of A₁ mRNA in young and middle-aged SAMP8 animals was not significantly different—of the same order as that obtained in young SAMR1 (Figure 1, panel A). Similar results were obtained for A_{2B} receptors whose level was increased in middle-aged versus young resistant mice but was not altered in SAMP8 (Figure 1, panel D). Concerning A_{2A} receptors, no significant differences were obtained between young and middle-aged animals, either in SAMP8 or in SAMR1. However, the level of A_{2A} mRNA expression was significant lower in SAMP8 versus SAMR1 (Figure 1 panel C). Finally, an increase in mRNA coding A₃ receptor was observed in both middle-aged SAMR1 and SAMP8 animals without significant differences between strains (Figure 1, panel B).

3.2. Adenosine A₁ receptor in SAMR1 and SAMP8 mice

Adenosine A₁ receptors are widely distributed in the brain and they are involved in neuroprotection. We measured adenosine A₁ receptors in plasma membranes from SAMR1 and SAMP8 mice by binding assay using [³H]DPCPX, a selective A₁ receptor antagonist, as radioligand. In all cases assayed saturation curves were better fitted to one

1 binding site model. Results from saturation curves showed a significant decrease in total
2 adenosine A₁ receptor numbers in middle-aged SAMR1 (38%), suggesting a loss of
3 receptors associated with aging (Figure 2, panels A and C). This decrease was not
4 observed in the SAMP8 strain (Figure 2, panels B and C). However, the total adenosine
5 A₁ receptor number in SAMP8 was significantly lower than in SAMR1, with even
6 fewer receptors detected than in middle-aged SAMR1 mice, suggesting that A₁
7 receptors are already altered at a very early age in SAMP8. No significant age-
8 associated differences in K_d values were observed in any SAM strain. However, K_d
9 values in SAMP8 strain were lower than in SAMR1, suggesting a higher receptor
10 affinity in the former (Figure 2, panel D).
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Due to the discrepancy observed in data obtained by real time PCR and by
25 radioligand binding assays in these samples, additional binding assays were carried out
26 in the presence of Gpp(NH)p, a non-hydrolyzable GTP analogue, to rule out possible
27 [³H]DPCPX activity as an inverse agonist (Finlayson et al., 2003). Results obtained
28 (Figure 3) showed a significant decrease in [³H]DPCPX binding in middle-aged
29 SAMR1, young SAMP8 and middle-aged SAMP8 as compared to [³H]DPCPX binding
30 levels in young SAMR1 strain. These results correlate with binding data obtained in the
31 absence of Gpp(NH)p and confirm [³H]DPCPX as a selective antagonist.
32
33
34
35
36
37
38
39
40
41
42
43

44 To measure adenosine A_{2A} receptors, Western blot experiments were performed
45 with membranes from SAM animals. Results showed a significant age-related increase
46 in the level of A_{2A} receptors in SAMR1, while no significant difference was observed in
47 SAMP8 related to age. Moreover, levels of A_{2A} receptors observed in both young and
48 middle-aged SAMP8 were of the same order as those obtained from young SAMR1
49 (Figure 4). In addition, to confirm the variation of A₁ receptors detected by radioligand
50 binding assay, Western blot experiments were also performed using anti-A₁ receptor
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

antibody. Results show an aged-associated loss of adenosine A₁ receptors in SAMR1 mice (Figure 5). In addition, the level of A₁ receptors in very young SAMP8 mice was of the same order as that observed in middle-aged SAMR1 animals, confirming results obtained by saturation binding assays.

3.3. Adenosine A₁ receptors/AC activity in Wistar rats

To compare and confirm the effect of aging on adenosine A₁ receptor levels we also analyzed the status of these receptors in brain from Wistar rats. Two groups of animals were used, young (3 months) and old (24 months) male rats. Saturation curves obtained by radioligand binding assays showed a significant decrease in adenosine A₁ receptors in brain from old versus young rats (49.3% of young rats) without significant variation in receptor affinity (Figure 6).

To evaluate whether age-related variation in adenosine receptors from Wistar rats modulates the transduction pathway mediated by these receptors, we analyzed the status of adenylyl cyclase activity in young and old rats. Results show that basal and forskolin or forskolin plus GTP γ S-stimulated adenylyl cyclase activity were significantly lower in old than in young animals (Figure 7, panel A). Moreover, the ability of CHA, a selective A₁ receptor agonist, to inhibit this enzymatic activity, was also decreased, suggesting a desensitization of the response system associated with the loss of receptors in the plasma membrane (Figure 7, panel B).

4. Discussion

To our knowledge, this study is the first to deal with the expression of mRNA coding different adenosine receptors and to quantify adenosine A₁ and A_{2A} receptors in

1 brain from SAMR1 and SAMP8 mice. Results presented herein show an age-related
2 loss of adenosine A₁ receptors in SAMR1, associated with an increase in the rate of
3 synthesis of this receptor, probably as a compensatory mechanism to prevent the
4 important loss of receptor detected at the membrane surface. In contrast, an age-related
5 increase in adenosine A_{2A} receptors was observed in SAMR1 with no variation in
6 SAMP8. However, the most important finding in our study is that adenosine A₁
7 receptors, which have been described as being neuroprotective, are already severely
8 decreased in very young SAMP8 mice (three weeks old) suggesting a great alteration of
9 these receptors in this aging model.
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Morphological changes in the aging brain have been described and they depend
25 on the species, the strains and the ages included in the study, as well as on individual
26 differences. Changes in A₁ and A_{2A} receptors and their function during aging have been
27 demonstrated, with A₁ being decreased and A_{2A} increased, respectively (Cunha, 2005;
28 Rodrigues et al., 2008). In spite of the lack of anatomical resolution in our study, our
29 findings of age-related changes in A₁ receptors in normal aging mice (SAMR1) are
30 consistent with previously published data. Age-related changes in adenosine A₁ receptor
31 binding have been described in mouse brain using the selective agonist CHA. In the
32 cortex, hippocampus, and cerebellum of aged mice (28 months old), a significant
33 decrease of about 44%, 50% and 12%, respectively, was observed in CHA binding
34 compared to young animals (3 months old), without changes in K_d values and not
35 dependant on a general cell degeneration in old age (Pagonopoulou and Angelatou,
36 1992). Also in mice, there has been confirmed to be widespread age-related loss of A₁
37 receptors in various cortical and subcortical structures involved in the neuromodulatory
38 role of adenosine (Ekonomou et al., 2000).
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 The methodology employed to quantify A₁ receptors in SAM mice was
3 validated by our results in male Wistar rats, which agree with those published by Cunha
4 and co-workers in which the density of A₁ binding sites in aged rats (24 months) was
5 reduced by 42% in whole brain membranes (Cunha et al., 2001b) by 33% in the
6 hippocampus, and by 60% in the cortex, when compared with young adult rats (6
7 weeks), with no changes in K_d (Cunha et al., 1995). A significant decrease in DPCPX
8 binding to hippocampal membranes of aged versus young rats has also been observed
9 (Sperlágh et al., 1997). By immunoblotting analysis there was shown to be an age-
10 related decrease in A₁ receptor in cerebral cortex from Wistar rats. Compared to
11 preparations from 2-month-old animals, the levels of A₁ in 6, 12 and 24-month-old rats
12 were reduced by 14, 32 and 28% respectively (Cheng et al., 2000). In addition, the
13 pattern of binding of the A₁ receptor antagonist [³H]DPCPX decreased with aging in
14 membranes from whole brain of male Wistar rats while K_d values were not different
15 among age groups (Cunha et al., 2001b).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Loss of A₁ receptors in plasma membrane has been associated with a down-
40 regulation of adenosine receptor by ligand overexposure. Down-regulation and
41 desensitization of adenosine A₁ receptors in vivo has been suggested as being induced
42 by high adenosine levels (León et al., 2002; León et al., 2004; 2005) and it comprises
43 the classical GPCR response when exposed chronically to agonist (Hanyaloglu and von
44 Zastrow, 2008). In agreement, upon chronic noxious brain conditions, and also upon
45 aging, adenosine modulation is modified, with increased levels of adenosine, which
46 modifies the density of adenosine receptors (Lopes et al., 1999; reviewed in Cunha,
47 2005). Previous studies have shown that endogenous extracellular adenosine levels are
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 larger in hippocampal preparations (Cunha et al., 2001a) and basal forebrain (Murillo-
2 Rodriguez et al., 2004) from aged compared to young adult rats. This increase could be
3 related to increased activity of 5'-nucleotidases with age (Cunha et al., 2001a;
4 Mackiewicz et al., 2006), which transforms ATP released into adenosine at the synaptic
5 level (Zimmermann, 1996; Cunha, 2001). Other authors lend support to the idea that
6 adenosine is released as such, with only a minor contribution of extracellular ATP-
7 derived adenosine (Pedata et al., 1993), and the existence of non-concentrative
8 bidirectional adenosine transporters allows balancing of changes in the intracellular and
9 extracellular adenosine concentrations (Plagemann and Wohlhueter, 1984). It has been
10 described that the greater levels of extracellular adenosine in aged animals cause a
11 greater tonic A₁-mediated inhibition (Sebastião et al., 2000). Moreover, during aging,
12 the level of adenosine in the extracellular fluid is increased in the hippocampus,
13 associated with a down-regulation and reduced responsiveness of presynaptic adenosine
14 A₁ receptors (Spelágh et al., 1997). Further, in previous studies we have shown that
15 adenosine release during hypoxic conditions is responsible for adenosine A₁ receptor
16 down-regulation and adenosine A_{2A} receptor up-regulation in C6 cells (Castillo et al.,
17 2008). Therefore, we hypothesize that the age-related adenosine level increase could be
18 responsible for the down-regulation of A₁ receptors in SAMR1 mice.

19 A significant decrease in the A₁ mRNA level was obtained using Northern
20 blotting analysis, associated with age in male Wistar rats, probably due to a decrease in
21 transcription of the receptor gene and/or decrease in the stability of mRNA (Cheng et
22 al., 2000). This result contrasts with the significant increase in mRNA level coding A₁
23 receptor we detected by quantitative real time PCR in old SAMR1 animals, probably as
24 a compensatory mechanism for the loss of the adenosine A₁ receptor detected at the

1 membrane surface. This compensatory mechanism is absent in old SAMP8 mice, in
2 agreement with the non-variation in A_1 associated with aging.
3
4
5

6
7 SAMP8 have been proposed as a model for Alzheimer disease (AD). It has been
8 demonstrated that SAMP8 mice overproduce β -amyloid in the hippocampus, and late in
9 life they develop amyloid plaques (Morley et al., 2000). This is associated with an
10 increase in mRNA for the Amyloid Precursor Protein (APP). In addition, it has been
11 postulated that aged SAMP8 mice have higher levels of $A\beta$ P (Banks et al., 2007). The
12 12-month-old SAMP8 mouse has APP and $A\beta$ P levels that are about twice those of the
13 4-month-old SAMP8, and it has developed severe learning and memory problems
14 (Kumar et al., 2000; Morley et al., 2000), which can be reversed by treatment with
15 either antibody or antisense directed against $A\beta$ P (Kumar et al., 2000; Morley et al.,
16 2002a). Thus, the SAMP8 strain represents a model of age-related and $A\beta$ P-dependent
17 cognitive decline.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 Loss of adenosine receptors in SAMP8 mice observed in the present study could
35 be associated with the increased $A\beta$ P level detected in these animals, which is thereafter
36 related to degeneration characteristics of AD. However, there is no evidence supporting
37 increased $A\beta$ P levels in SAMP8 as young as three weeks old, at which time point we
38 observed an important loss of adenosine A_1 receptors. In addition, results obtained
39 previously by our group in frontal cortex brain from AD patients show a significant
40 increase in adenosine A_1 receptors already in early stages of the disease and then
41 maintained with the progression of neurodegeneration (Albasanz et al., 2008).
42 Moreover, up-regulation of A_1 receptors has also been observed in brain from patients
43 with other dementias such as Pick disease or argyrophilic grain disease, in which there
44 is no significant $A\beta$ P accumulation (Albasanz et al., 2007; Perez-Buira et al., 2007).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Therefore, variation in adenosine receptors seems not to be specific for neurodegenerative diseases associated with A β P accumulation. In other words, β -amyloid seems not to be responsible for variation in A₁ receptors. As postulated above, an increase in endogenous adenosine level could be responsible for the lower adenosine A₁ receptors observed in SAMP8 versus young SAMR1 animals. In agreement, the increase in adenosine receptors in human brain from AD patients has been related with the lower endogenous adenosine level, as detected in plasma (Selly, 2004). Moreover, adenosine and pharmaceutical agents that raise the extracellular adenosine concentration have been suggested as reinforcements to conventional treatment with acetylcholinesterase inhibitors in AD (Ferroni et al., 2002; Schubert et al., 2001). Furthermore, A_{2A} receptor blockade is also nowadays considered a promising target in AD treatment (Dall'Igna et al., 2007; Cunha et al., 2008).

Although no references are available about adenosine content in SAM brain, at least to our knowledge, changes in neurotransmitter levels in SAM animals have been reported as increasing in GABA and decreasing in acetylcholine and serotonin (Morley, 2002). In the hippocampus and/or cerebral cortex of P8 strain at early ages, the contents of Glutamate (Glu) and Glutamine (Gln) were higher than in R1, while the aspartate (Asp) and Alanine (Ala) content was lower, which suggests that a metabolic pathway from α -ketoglutarate to Glu predominates in SAMP8 brain (Kitamura et al., 1992; Nomura et al., 1991). Local accumulation of excitatory amino acids can lead to neuronal degeneration because of the neurotoxic effects at high concentrations of Glu and analogues in Glu receptive areas (Olney et al., 1990). Increased glutamate levels have been found in several brain areas in AD and in cerebrospinal fluid (CSF) from AD patients (Procter et al., 1988; Pomara et al., 1992). This excess of endogenous glutamate

1 would contribute to cell death in AD, possibly as the result of the failure to remove
2 glutamate from synapse (Greenamyre and Young, 1989). In agreement, a down-
3 regulation and desensitization of metabotropic glutamate receptors in the frontal cortex
4 in common forms of dementia with Lewy bodies (DLB) and in Alzheimer disease cases
5 have been detected (Albasanz et al., 2005). The neuroprotective role of adenosine is
6 mainly due to inhibition of glutamate release through adenosine A₁ receptor activation.
7 Therefore, the loss of A₁ receptors could be responsible for the increase in the glutamate
8 level in SAM mice.
9
10
11
12
13
14
15
16
17
18
19
20
21

22 In summary, in SAMR1 mice used here as controls, it was shown that there is an
23 age-related loss of adenosine A₁ receptor, as also observed in rats, but at early stages.
24 Moreover, the most important result in this work is that in brain from very young
25 SAMP8 mice there is a reduced A₁ receptor level similar to that obtained in aged
26 SAMR1, suggesting an important alteration of these receptors at only three weeks of
27 life which is maintained thereafter. In vivo PET study detected an age-dependent A₁
28 receptor loss in humans that may be of pathophysiological importance in various
29 neurological diseases associated with aging (Meyer et al., 2006). Therefore, SAM mice
30 may represent an important model in the study of alteration of neurotransmitter
31 receptors related to neurodegenerative diseases.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

1
2
3
4 This work was supported in part by the European Union through the Marie-
5 Curie Research Training Network PRAIRIES (contract MRTN-CT-2006-035810), by
6 the Consejería de Educación y Ciencia (PCI08-0125), the Consejería de Sanidad (PI-
7 2007/50 and G-2007-C/13) of the Junta de Comunidades de Castilla-La Mancha, the
8 Ministerio de Ciencia e Innovación (BFU2008-00138) and the Instituto de Salud Carlos
9 III (PI080400).
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

5. References

- 1
2
3
4
5
6
7 Albasanz, J.L., Perez, S., Barrachina, M., Ferrer I, Martín M. 2008. Up-regulation of adenosine
8 receptors in the frontal cortex in Alzheimer's disease. *Brain Pathol.* 18, 211-219.
9
10
11 Albasanz, J.L., Rodríguez, A., Ferrer, I., Martín, M. 2007. Up-regulation of adenosine A₁
12 receptors in frontal cortex from Pick's disease cases. *Eur J Neurosci.* 26, 3501-3508.
13
14
15 Banks, W.A., Farr, S.A., Morley, J.E., Wolf, K.M., Geylis, V., Steinitz, M. 2007. Anti-amyloid
16 beta protein antibody passage across the blood-brain barrier in the SAMP8 mouse
17 model of Alzheimer's disease: an age-related selective uptake with reversal of learning
18 impairment. *Exp Neurol.* 206, 248-256.
19
20
21
22
23
24
25
26 Butterfield, D.A., Poon, H.F. 2005. The senescence-accelerated prone mouse (SAMP8): a
27 model of age-related cognitive decline with relevance to alterations of the gene
28 expression and protein abnormalities in Alzheimer's disease. *Exp Gerontol.* 40, 774-
29 783.
30
31
32
33
34
35
36 Castillo C.A., León D., Ruiz M.A., Albasanz J. L. and Martín M. 2008. Modulation of
37 adenosine A(1) and A(2A) receptors in C6 glioma cells during hypoxia: involvement of
38 endogenous adenosine. *J Neurochem.* 105, 2315-2329.
39
40
41
42
43 Cheng, J.T., Liu, I.M., Juang, S.W., Jou, S.B. 2000. Decrease of adenosine A-1 receptor gene
44 expression in cerebral cortex of aged rats. *Neurosci Lett.* 283, 227-229.
45
46
47
48 Cunha G.M., Canas P.M., Melo C.S., Hockemeyer J., Muller C.E., Oliveira C.R. and Cunha
49 R.A. 2008. Adenosine A2A receptor blockade prevents memory dysfunction caused by
50 beta-amyloid peptides but not by scopolamine or MK-801. *Exp Neurol* 210, 776-781.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Cunha R.A. 2001. Adenosine as a neuromodulator and as a homeostatic regulator in the nervous system: different roles, different sources and different receptors. *Neurochem Int* 38, 107-125.
- Cunha, R.A. 2005. Neuroprotection by adenosine in the brain: From A₁ receptor activation to A_{2A} receptor blockade. *Purinergic Signal*. 1, 111-134.
- Cunha, R.A., Almeida, T., Ribeiro, J.A. 2001a. Parallel modification of adenosine extracellular metabolism and modulatory action in the hippocampus of aged rats. *J. Neurochem*. 76, 372-382.
- Cunha, R.A., Constantino, M.C., Sebastiao, A.M., Ribeiro, J.A. 1995. Modification of A₁ and A_{2a} adenosine receptor binding in aged striatum, hippocampus and cortex of the rat. *Neuroreport*. 6, 1583-1588.
- Cunha, R.A., Constantino, M.D., Fonseca, E., Ribeiro, J.A. 2001b. Age-dependent decrease in adenosine A₁ receptor binding sites in the rat brain. Effect of cis unsaturated free fatty acids. *Eur J Biochem*. 268, 2939-2947.
- Dall'Igna O.P., Fett P., Gomes M.W., Souza D.O., Cunha R.A. and Lara D.R. 2007. Caffeine and adenosine A_{2A} receptor antagonists prevent beta-amyloid (25-35)-induced cognitive deficits in mice. *Exp Neurol* 203, 241-245.
- Dunwiddie, T.V., Masino, S.A. 2001. The role and regulation of adenosine in the central nervous system. *Annu Rev Neurosci*. 24, 31-55.
- Ekonomou, A., Pagonopoulou, O., Angelatou, F. 2000. Age-dependent changes in adenosine A₁ receptor and uptake site binding in the mouse brain: an autoradiographic study. *J Neurosci Res*. 60, 257-265.
- Ferroni, S., Marchini, C., Ogata, T., Schubert, P., 2002. Recovery of deficient cholinergic calcium signaling by adenosine in cultured rat cortical astrocytes. *J Neurosci Res* 68, 615-621.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Finlayson K., Maemoto T., Butcher S.P., Sharkey J. and Olverman H.J. 2003. Comparison of effects of MgCl₂ and Gpp(NH)p on antagonist and agonist radioligand binding to adenosine A₁ receptors. *Acta Pharmacol Sin* 24, 729-740.
- Fredholm, B.B., Chen, J.F., Masino, S.A., Vaugeois, J.M. 2005. Actions of adenosine at its receptors in the CNS: insights from knockouts and drugs. *Annu Rev Pharmacol Toxicol*. 45, 385-412.
- Fredholm, B.B., IJzerman, A.P., Jacobson, K.A., Klotz, K.N., Linden, J. 2001. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev*. 53, 527-552.
- Greenamyre, J.T., Young, A.B. 1989. Excitatory amino acids and Alzheimer's disease. *Neurobiol Aging*. 10, 593-602.
- Hanyaloglu A.C. and von Zastrow M. 2008. Regulation of GPCRs by endocytic membrane trafficking and its potential implications. *Annu Rev Pharmacol Toxicol* 48, 537-568.
- Higuchi, R., Fockler, C., Dollinger, G., Watson, R. 1993. Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (NY)*. 11, 1026-1030.
- Inada K., Yokoi I., Kabuto H., Habu H., Mori A. and Ogawa N. 1996. Age-related increase in nitric oxide synthase activity in senescence accelerated mouse brain and the effect of long-term administration of superoxide radical scavenger. *Mech Ageing Dev* 89, 95-102.
- Kaisho Y., Miyamoto M., Shiho O., Onoue H., Kitamura Y. and Nomura S. 1994. Expression of neurotrophin genes in the brain of senescence-accelerated mouse (SAM) during postnatal development. *Brain Res* 647, 139-144.
- Kitamura, Y., Zhao, X.H., Ohnuki, T., Takei, M., Nomura, Y. 1992. Age-related changes in transmitter glutamate and NMDA receptor/channels in the brain of senescence-accelerated mouse. *Neurosci Lett*. 137, 69-72.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Kumar, V.B., Franko, M.W., Farr, S.A., Armbrrecht, H.J., Morley, J.E. 2000. Identification of age-dependent changes in expression of senescence-accelerated mouse (SAMP8) hippocampal proteins by expression array analysis. *Biochem Biophys Res Commun.* 272, 657-661.
- Kurokawa T., Asada S., Nishitani S. and Hazeki O. 2001. Age-related changes in manganese superoxide dismutase activity in the cerebral cortex of senescence-accelerated prone and resistant mouse. *Neurosci Lett* 298, 135-138.
- León, D., Albasanz, J.L., Fernández, M., Ruíz, M.A., Martín, M. 2004. Down-regulation of rat brain adenosine A1 receptors at the end of pregnancy. *J Neurochem.* 88, 993-1002.
- León, D., Albasanz, J.L., Ruíz, M.A., Fernández, M., Martín, M. 2002. Adenosine A1 receptor down-regulation in mothers and fetal brain after caffeine and theophylline treatments to pregnant rats. *J Neurochem.* 82, 625-634.
- León, D., Albasanz, J.L., Ruíz, M.A., Martín, M. 2005. Chronic caffeine or theophylline intake during pregnancy inhibits A1 receptor function in the rat brain. *Neuroscience.* 131, 481-489.
- León, D., Albasanz, J.L., Ruíz, M.A., Martín, M. 2005. Chronic caffeine or theophylline intake during pregnancy inhibits A1 receptor function in the rat brain. *Neuroscience.* 131, 481-489.
- Lopes, L.V., Cunha, R.A., Ribeiro, J.A. 1999. Cross talk between A(1) and A(2A) adenosine receptors in the hippocampus and cortex of young adult and old rats. *J Neurophysiol.* 82, 3196-3203.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. 1951. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 193, 265-275.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Mackiewicz, M., Nikonova, E.V., Zimmermann, J.E., Romer, M.A., Cater, J., Galante, R.J. 2006. Pack AI. Age-related changes in adenosine metabolic enzymes in sleep/wake regulatory areas of the brain. *Neurobiol Aging*. 27, 351-360.
- Meyer, P.T., Elmenhorst, D., Boy, C., Winz, O., Matusch, A., Zilles, K., Bauer, A. 2007. Effect of aging on cerebral A1 adenosine receptors: A [18F]CPFPX PET study in humans. *Neurobiol Aging*. 28, 1914-1924.
- Miyamoto, M. 1997. Characteristics of age-related behavioral changes in senescence-accelerated mouse SAMP8 and SAMP10. *Exp Gerontol*. 32, 139-148.
- Miyamoto, M., Kiyota, Y., Yamazaki, N., Nagaoka, A., Matsuo, T., Nagawa, Y., Takeda, T. 1986. Age-related changes in learning and memory in the senescence-accelerated mouse (SAM). *Physiol Behav*. 38, 399-406.
- Miyazaki H., Okuma Y., Nomura J., Nagashima K. and Nomura Y. 2003. Age-related alterations in the expression of glial cell line-derived neurotrophic factor in the senescence-accelerated mouse brain. *J Pharmacol Sci* 92, 28-34.
- Morley, J.E. 2002. The SAMP8 mouse: a model of Alzheimer disease? *Biogerontology*. 3, 57-60.
- Morley, J.E., Farr, S.A., Flood, J.F. 2002a. Antibody to amyloid beta protein alleviates impaired acquisition, retention, and memory processing in SAMP8 mice. *Neurobiol Learn Mem*. 78, 125-138.
- Morley, J.E., Farr, S.A., Kumar, V.B., Banks, W.A. 2002b. Alzheimer's disease through the eye of a mouse. Acceptance lecture for the 2001 Gayle A. Olson and Richard D. Olson prize. *Peptides*. 23, 589-599.
- Morley, J.E., Kumar, V.B., Bernardo, A.E., Farr, S.A., Uezu, K., Tumosa, N., Flood, J.F. 2000. Beta-amyloid precursor polypeptide in SAMP8 mice affects learning and memory. *Peptides*. 21, 1761-1767.

- 1
2 Murillo-Rodriguez, E., Blanco-Centurion, C., Gerashchenko, D., Salin-Pascual, R.J.,
3 Shiromani, P.J. 2004. The diurnal rhythm of adenosine levels in the basal forebrain of
4 young and old rats. *Neuroscience*. 123, 361-370.
5
6
7 Nomura, Y., Kitamura, Y., Zhao, X. 1991. Aging in the glutamatergic system with special
8 reference to the NMDA receptor/ion channel complex in the brains of senescence
9 accelerated mice. In T. Kameyama et al. (Eds), *NMDA Related Agents: Biochemistry,*
10 *Pharmacology and Behaviour*, NPP Books, Ann Arbor, pp- 287-298.
11
12
13
14
15
16
17 Nomura, Y., Okuma, Y. 1999. Age-related defects in lifespan and learning ability in SAMP8
18 mice. *Neurobiol Aging*. 20, 111-115.
19
20
21
22 Okatani Y., Wakatsuki A., Reiter R.J. and Miyahara Y. 2002. Melatonin reduces oxidative
23 damage of neural lipids and proteins in senescence-accelerated mouse. *Neurobiol Aging*
24 23, 639-644.
25
26
27
28
29 Olney, J.W. 1990. Excitotoxic amino acids and neuropsychiatric disorders. *Annu Rev*
30 *Pharmacol Toxicol*. 30, 47-71.
31
32
33
34 Pagonopoulou, O., Angelatou, F. 1992. Reduction of A1 adenosine receptors in cortex,
35 hippocampus and cerebellum in ageing mouse brain. *Neuroreport*. 3, 735-777.
36
37
38
39 Pallas M, Camins A, Smith MA, Perry G, Lee HG, Casadesus G. 2008. From aging to
40 Alzheimer's disease: unveiling "the switch" with the senescence-accelerated mouse
41 model (SAMP8). *J Alzheimers Dis*. 15, 615-624.
42
43
44
45
46 Pedata F., Latini S., Pugliese A.M. and Pepeu G. 1993. Investigations into the adenosine
47 outflow from hippocampal slices evoked by ischemia-like conditions. *J Neurochem* 61,
48 284-289.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7 Plagemann P.G. and Wohlhueter R.M. 1984. Nucleoside transport in cultured mammalian cells.
8
9 Multiple forms with different sensitivity to inhibition by nitrobenzylthioinosine or
10
11 hypoxanthine. *Biochim Biophys Acta* 773, 39-52.
- 12
13 Pomara, N., Singh, R., Deptula, D., Chou, J.C., Schwartz, M.B., LeWitt, P.A. 1992. Glutamate
14
15 and other CSF amino acids in Alzheimer's disease. *Am J Psychiatry*. 149, 251-254.
- 16
17 Procter, A.W., Palmer, A.M., Francis, P.T., Lowe, S.L., Neary, D., Murphy, E., Doshi, R.,
18
19 Bowen, D.M. 1988. Evidence of glutamatergic denervation and possible abnormal
20
21 metabolism in Alzheimer's disease. *J Neurochem*.50, 790-802.
- 22
23 Ralevic, V., Burnstock, G. 1998. Receptors for purines and pyrimidines. *Pharmacol Rev.* 50,
24
25 413-492.
- 26
27 Rodrigues, R.J., Canas, P.M., Lopes, L.V., Oliveira, C.R., Cunha, R.A. 2008. Modification of
28
29 adenosine modulation of acetylcholine release in the hippocampus of aged rats.
30
31 *Neurobiol Aging*. 29, 1597-1601.
- 32
33 Sato E., Oda N., Ozaki N., Hashimoto S., Kurokawa T. and Ishibashi S. 1996. Early and
34
35 transient increase in oxidative stress in the cerebral cortex of senescence-accelerated
36
37 mouse. *Mech Ageing Dev* 86, 105-114.
- 38
39 Schubert, P., Ogata, T., Marchini, C., Ferroni, S. 2001. Glia-related pathomechanisms in
40
41 Alzheimer's disease: a therapeutic target? *Mech Ageing Dev*. 123, 47-57.
- 42
43 Sebastião, A.M., Cunha, R.A., de Mendonça, A., Ribeiro, J.A. 2000. Modification of adenosine
44
45 modulation of synaptic transmission in the hippocampus of aged rats. *Br J Pharmacol*.
46
47 131, 1629-1634.
- 48
49
50 Selley, M.L. 2004. Increased homocysteine and decreased adenosine formation in Alzheimer's
51
52 disease. *Neurol Res*. 26, 554-547.
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Sperlágh, B., Zsilla, G., Baranyi, M., Kékes-Szabó, A., Vizi, E.S. 1997. Age-dependent changes of presynaptic neuromodulation via A1-adenosine receptors in rat hippocampal slices. *Int J Dev Neurosci.* 15, 739-747.

Takeda, T., Hosokawa, M., Takeshita, S., Irino, M., Higuchi, K., Matsushita, T., Tomita, Y., Yasuhira, K., Hammamoto H, Shimuzu, K, Ishii M and Yamamur T. 1981. A novel murine model of accelerated senescence, *Mech. Ageing Dev.*, 17, 183-194.

Wei X., Zhang Y. and Zhou J. 1999. Alzheimer's disease-related gene expression in the brain of senescence accelerated mouse. *Neurosci Lett* 268, 139-142.

Zimmermann H. 1996. Biochemistry, localization and functional roles of ecto-nucleotidases in the nervous system. *Prog Neurobiol* 49, 589-618.

FIGURE LEGENDS

Figure 1. Adenosine A₁, A_{2A}, A_{2B}, and A₃ receptor gene expression assayed by quantitative real time RT-PCR. After RNA isolation from 21-day-old SAMR1 (R1 21d), 180-day-old SAMR1 (R1 180d), 21-day-old SAMP8 (P8 21d) and 180-day-old SAMP8 (P8 180d) animals, quantitative real time RT-PCR assays for the determination of A₁ (panel A), A₃ (panel B) A_{2A} (panel C) and A_{2B} (panel D) gene expression levels were carried out using TaqMan® universal PCR following the protocol indicated in Materials and Methods. Histograms represent mean ± SEM values obtained from four separate experiments using different preparations. * p<0.05 and ** p<0.01 significantly different from indicated value.

1
2 **Figure 2. Adenosine A₁ receptor detection in SAM mice by radioligand binding**
3 **assay.** Saturation curves of [³H]DPCPX binding to plasma membranes were performed
4
5 by incubation of 50-75 μg of protein from SAMR1 (panel A) or SAMP8 (panel B) mice
6
7 with increasing concentrations of the radioligand, as described in Materials and
8
9 Methods, after pre-incubation with adenosine deaminase in order to remove endogenous
10
11 adenosine. Total receptor number (Bmax) and receptor affinity (Kd) determined by
12
13 Scatchard and nonlinear regression analysis are shown in panels C and D, respectively.
14
15 Data are mean ± SEM of independent experiments performed in triplicate using
16
17 different membranes isolated from four SAMR1 and four SAMP8 animals. *p<0.05 and
18
19 ***p<0.001 significantly different from indicated value.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Figure 3. Gpp(NH)p effect on radioligand binding assay in SAM mice.** [³H]DPCPX
3
4 binding assays were performed in plasma membranes (50-75 μg of protein) from
5 SAMR1 or SAMP8 mice. After pre-incubation with adenosine deaminase, binding
6
7 assays were performed with 20 nM [³H]DPCPX in the presence of 10 μM Gpp(NH)p,
8
9 without MgCl₂, and in the presence of MgCl₂, without Gpp(NH)p. CPA was used to
10
11 obtain nonspecific binding. Data are mean ± SEM of independent experiments
12
13 performed in duplicate using different membranes isolated from four animals of each
14
15 condition. No statistical differences were observed between binding data obtained in the
16
17 presence or the absence of Gpp(NH)p. *p < 0.05, **p < 0.01 significantly different from
18
19 control young SAMR1 mice in the absence of Gpp(NH)p; #p < 0.05 significantly
20
21 different from control young SAMR1 mice in the presence of Gpp(NH)p.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Figure 4. Western-blot of A_{2A}R in plasma membranes from SAM mice.** Fifty
3
4 micrograms of protein were subjected to SDS/PAGE, transferred electrophoretically to
5
6 nitrocellulose and probed with antisera anti-A_{2A}, as described in Materials and Methods.
7
8
9 Upper panel shows a representative autoradiography, showing the bands corresponding
10
11 to A_{2A}R and β-actin, which was used as gel loading control. Lower panel shows the
12
13 obtained histogram. Data is expressed as mean ± SEM using arbitrary units. **p < 0.01
14
15 significantly different from control young SAMR1 mice.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Figure 5. Western-blot of A₁R in plasma membranes from SAM mice.** Fifty
3
4 micrograms of protein were subjected to SDS/PAGE, transferred electrophoretically to
5
6 nitrocellulose and probed with antisera anti-A₁, as described in Materials and Methods.
7
8
9 Upper panel shows a representative autoradiography, showing the bands corresponding
10
11 to A₁R and β-actin, which was used as gel loading control. Lower panel shows the
12
13 obtained histogram from densitometric analysis of bands. Data is expressed as mean ±
14
15 SEM using arbitrary units. *p<0.05 and ***p<0.001 significantly different from control
16
17 young SAMR1 mice.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 6. Adenosine A₁ receptor detection in Wistar rats by radioligand binding

assay. Saturation curves of [³H]DPCPX binding to plasma membranes were performed by incubation of 50-75 µg of protein from young and old rats with increasing concentrations of the radioligand, as described in Materials and Methods, after pre-incubation with adenosine deaminase in order to remove endogenous adenosine. Total receptor number (B_{max}) and receptor affinity (K_d) determined by Scatchard and nonlinear regression analysis are shown in the inset. Data are mean ± SEM of number of experiments in brackets performed in triplicate using different membrane isolations from different animals. * p<0.01 significantly different from young rats.

1
2 **Figure 7. Adenylyl cyclase activity in brain plasma membranes from Wistar rats.**

3
4 **Panel A.** Ten to twenty micrograms of plasma membranes, previously incubated with
5 adenosine deaminase, were used to determine basal cAMP level and stimulation after
6 incubation with 10 μ M forskolin (Fsk) or 10 μ M Fsk plus 5 μ M GTP γ S. **Panel B.** Ten
7 to twenty micrograms of plasma membranes were incubated with Fsk plus GTP γ S in the
8 presence and in the absence of CHA, a selective A₁R agonist. cAMP accumulation was
9 measured as described in Materials and Methods. Data are means \pm SEM of at least four
10 experiments performed in triplicate with different membrane preparations from different
11 rats. * p<0.01 significantly different from young rats.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1




Figure 2




Figure 2





Figure 3




Figure 4




Figure 5





Figure 6



	Bmax (fmol/mg prot)	Kd (nM)
young rats	204.79 ± 26.22 (3)	4.42 ± 0.72 (3)
old rats	101.09 ± 10.10 (3)*	4.51 ± 1.25 (3)

Figure 7

