

A-infinity $GL(N)$ -equivariant matrix integrals-III.
Tree-level calculations and variations of nc-Hodge
structure on complex projective manifolds.

Serguei Barannikov

IMJ, CNRS

23/06/2010

Tree level BV on complex projective manifold (g=0 calculations)

- nc-BV on $Symm(C_\lambda[1+d])$, $C_\lambda = \bigoplus_{j=0}^{\infty} (U[1]^{\otimes j})^{\mathbb{Z}/j\mathbb{Z}}$

$$\hbar \Delta S + \frac{1}{2} \{S, S\} = 0, \quad S = \sum_{g \geq 0, i} \hbar^{2g-1+i} S_{g,i}, \quad S_{g,i} \in Symm^i(C_\lambda[1+d]),$$
$$\Leftrightarrow \Delta \exp(S/\hbar) = 0$$
$$\{S_{0,1}, S_{0,1}\} = 0, \text{ - } A_\infty - \text{CY algebra}$$

Tree level BV on complex projective manifold (g=0 calculations)

- nc-BV on $Symm(C_\lambda[1+d])$, $C_\lambda = \bigoplus_{j=0}^{\infty} (U[1]^{\otimes j})^{\mathbb{Z}/j\mathbb{Z}}$

$$\hbar \Delta S + \frac{1}{2} \{S, S\} = 0, \quad S = \sum_{g \geq 0, i} \hbar^{2g-1+i} S_{g,i}, \quad S_{g,i} \in Symm^i(C_\lambda[1+d]),$$
$$\Leftrightarrow \Delta \exp(S/\hbar) = 0$$
$$\{S_{0,1}, S_{0,1}\} = 0, \text{ - } A_\infty - \text{CY algebra}$$

- M - smooth projective/ \mathbb{C} , $c_1(T_M) = 0$, $\omega_0 \in \Gamma(M, K_M)$, \rightarrow BV operator on $\Omega^{0,*}(M, \Lambda^* T)$,
- $$(\Delta \alpha) \vdash \omega_0 = \partial(\alpha \vdash \omega_0)$$

Tree level BV on complex projective manifold (g=0 calculations)

- nc-BV on $Symm(C_\lambda[1+d])$, $C_\lambda = \bigoplus_{j=0}^{\infty} (U[1]^{\otimes j})^{\mathbb{Z}/j\mathbb{Z}}$

$$\hbar \Delta S + \frac{1}{2} \{S, S\} = 0, \quad S = \sum_{g \geq 0, i} \hbar^{2g-1+i} S_{g,i}, \quad S_{g,i} \in Symm^i(C_\lambda[1+d]),$$
$$\Leftrightarrow \Delta \exp(S/\hbar) = 0$$
$$\{S_{0,1}, S_{0,1}\} = 0, \text{ - } A_\infty - \text{CY algebra}$$

- M - smooth projective/ \mathbb{C} , $c_1(T_M) = 0$, $\omega_0 \in \Gamma(M, K_M)$, \rightarrow BV operator on $\Omega^{0,*}(M, \Lambda^* T)$,
$$(\Delta \alpha) \vdash \omega_0 = \partial(\alpha \vdash \omega_0)$$
- (tree level) BV equation on $\Omega^{0,*}(M, \Lambda^* T)$

$$u \Delta \gamma + \bar{\partial} \gamma + \frac{1}{2} [\gamma, \gamma] = 0,$$

$$\gamma = \sum_{i \geq 0} \gamma_i u^i, \quad \gamma_i \in \Omega^{0,*}(M, \Lambda^* T)$$

Tree level BV-equation on complex projective manifold

- (tree level) BV equation on $\Omega^{0,*}(M, \Lambda^* T)$

$$u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0,$$

Tree level BV-equation on complex projective manifold

- (tree level) BV equation on $\Omega^{0,*}(M, \Lambda^* T)$

$$u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0,$$

- the first order term $\gamma_0 \in \Omega^{0,*}(M, \Lambda^* T)$

$$\bar{\partial}\gamma_0 + \frac{1}{2}[\gamma_0, \gamma_0] = 0$$

γ_0 $-A_\infty$ -deformations of $D^b Coh(M)$

Tree level BV-equation on complex projective manifold

- (tree level) BV equation on $\Omega^{0,*}(M, \Lambda^* T)$

$$u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0,$$

- the first order term $\gamma_0 \in \Omega^{0,*}(M, \Lambda^* T)$

$$\bar{\partial}\gamma_0 + \frac{1}{2}[\gamma_0, \gamma_0] = 0$$

γ_0 – A_∞ -deformations of $D^b Coh(M)$

- The component

$$\gamma_0^{cl} \in \Omega^{0,1}(M, T)$$

$$\bar{\partial}\gamma_0^{cl} + \frac{1}{2}[\gamma_0^{cl}, \gamma_0^{cl}] = 0 \text{ - deformations of complex structure on } M$$

Noncommutative periods map (B[5])

•

$$u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0, \Leftrightarrow (u\Delta + \bar{\partial}) \exp\left(\frac{1}{u}\gamma\right) = 0$$

Noncommutative periods map (B[5])

-

$$u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0, \Leftrightarrow (u\Delta + \bar{\partial}) \exp\left(\frac{1}{u}\gamma\right) = 0$$

- Define r_u on components of $\Omega^{0,*}(M, \Lambda^* T)$

$$r_u \gamma = \sum_{p,q} u^{(q-p)/2} \gamma^{p,q}, \quad \gamma^{p,q} \in \Omega^{0,q}(M, \Lambda^p T)$$

$$(u\Delta + \bar{\partial}) \rightarrow \Delta + \bar{\partial}$$

-local system of $(u\Delta + \bar{\partial})$ -cohomology over $\mathbb{A}_u^1 \setminus \{0\}$

Noncommutative periods map (B[5])

- $u\Delta\gamma + \bar{\partial}\gamma + \frac{1}{2}[\gamma, \gamma] = 0 \Leftrightarrow (u\Delta + \bar{\partial}) \exp\left(\frac{1}{u}\gamma\right) = 0$
- Define r_u on components of $\Omega^{0,*}(M, \Lambda^* T)$

$$r_u \gamma = \sum_{p,q} u^{(q-p)/2} \gamma^{p,q}, \quad \gamma^{p,q} \in \Omega^{0,q}(M, \Lambda^p T)$$

$$(u\Delta + \bar{\partial}) \rightarrow \Delta + \bar{\partial}$$

-local system of $(u\Delta + \bar{\partial})$ -cohomology over $\mathbb{A}_u^1 \setminus \{0\}$

- $$\begin{aligned} \Omega(t, u) &= \int_M r_u \exp\left(\frac{1}{u}\gamma\right) \lrcorner \omega_0 \\ \Omega(t, u) &\in H_{DR}^*(M)((u)) \widehat{\otimes} \mathcal{O}_{\mathcal{M}_{\Lambda T}} \end{aligned}$$

$$t \in \mathcal{M}_{\Lambda T}, \quad T_0 \mathcal{M}_{\Lambda T} = H^*(M, \Lambda^* T)$$

Noncommutative periods map and g=0 Gromov-Witten of the mirror

-

$$\Omega(t, u) = \int_M r_u \exp\left(\frac{1}{u}\gamma\right) \lrcorner \omega_0$$

For γ^W , normalized using a filtration W opposite to $F^{Hodge}([B5], 1999)$:

$$\frac{\partial^2}{\partial t^i \partial t^j} \Omega^W = u^{-1} C_{ij}^k(t) \frac{\partial}{\partial t^k} \Omega^W$$

Noncommutative periods map and g=0 Gromov-Witten of the mirror



$$\Omega(t, u) = \int_M r_u \exp\left(\frac{1}{u}\gamma\right) \lrcorner \omega_0$$

For γ^W , normalized using a filtration W opposite to $F^{Hodge}([B5], 1999)$:

$$\frac{\partial^2}{\partial t^i \partial t^j} \Omega^W = u^{-1} C_{ij}^k(t) \frac{\partial}{\partial t^k} \Omega^W$$



$$C_{kij}(t) = \partial_{kij}^3 (\text{genus } = 0 \text{ GW-potential of } M^{\text{mirror}})$$

Semi-infinite (Noncommutative) Hodge structures ([B5]).

- The class $[\exp(\gamma^W)\omega] = \Omega(t, u)$ is obtained as intersection of moving subspace

$$\Omega(t, u) = \mathcal{L}(t) \cap (\text{Affine space}(W))$$

$$\mathcal{L}(t) \subset H_{DR}^*(M)((u)) \widehat{\otimes} \mathcal{O}_{\mathcal{M}_{\Lambda T}}$$

Semi-infinite (Noncommutative) Hodge structures ([B5]).

- The class $[\exp(\gamma^W)\omega] = \Omega(t, u)$ is obtained as intersection of moving subspace

$$\Omega(t, u) = \mathcal{L}(t) \cap (\text{Affine space}(W))$$

$$\mathcal{L}(t) \subset H_{DR}^*(M)((u)) \widehat{\otimes} \mathcal{O}_{\mathcal{M}_{\Lambda T}}$$

- The semi-infinite subspace $\mathcal{L}(t)$, $t \in \mathcal{M}_{\Lambda T}$, is defined for *arbitrary* projective manifold/ \mathbb{C}

$$\mathcal{L}(\gamma_0) : [r_u(\exp \frac{1}{u} i_{\gamma_0})(\varphi_0 + u\varphi_1 + \dots)], \quad \varphi_i \in \Omega_{DR}$$

where $\gamma_0 \in \Omega^{0,*}(M, \Lambda^* T)$, $\bar{\partial}\gamma_0 + \frac{1}{2}[\gamma_0, \gamma_0] = 0$

Semi-infinite (Noncommutative) Hodge structures ([B5]).

- The class $[\exp(\gamma^W)\omega] = \Omega(t, u)$ is obtained as intersection of moving subspace

$$\Omega(t, u) = \mathcal{L}(t) \cap (\text{Affine space}(W))$$

$$\mathcal{L}(t) \subset H_{DR}^*(M)((u)) \widehat{\otimes} \mathcal{O}_{\mathcal{M}_{\Lambda T}}$$

- The semi-infinite subspace $\mathcal{L}(t)$, $t \in \mathcal{M}_{\Lambda T}$, is defined for *arbitrary* projective manifold/ \mathbb{C}

$$\mathcal{L}(\gamma_0) : [r_u(\exp \frac{1}{u} i_{\gamma_0})(\varphi_0 + u\varphi_1 + \dots)], \quad \varphi_i \in \Omega_{DR}$$

where $\gamma_0 \in \Omega^{0,*}(M, \Lambda^* T)$, $\bar{\partial}\gamma_0 + \frac{1}{2}[\gamma_0, \gamma_0] = 0$

-

$$u\mathcal{L}(t) \subset \mathcal{L}(t), \quad \frac{\partial}{\partial t} \mathcal{L}(t) \subset u^{-1} \mathcal{L}(t)$$

$$\frac{\partial}{\partial u} \mathcal{L}(t) \subset u^{-2} \mathcal{L}(t), \quad \mathcal{L}(u) \oplus \overline{\mathcal{L}}(\bar{u}|_{\bar{u}=u^{-1}})$$

implies tt^* -equations, remarkably $\mathcal{D}_{\frac{\partial}{\partial u}}$ -modules over \mathbb{A}^1 with similar properties appeared in 70s in works of Birkhoff, Malgrange, K.Saito and M.Saito

Semi-infinite (Noncommutative) Hodge structures ([B5]).

- The class $[\exp(\gamma^W)\omega] = \Omega(t, u)$ is obtained as intersection of moving subspace

$$\Omega(t, u) = \mathcal{L}(t) \cap (\text{Affine space}(W))$$

$$\mathcal{L}(t) \subset H_{DR}^*(M)((u)) \widehat{\otimes} \mathcal{O}_{\mathcal{M}_{\Lambda T}}$$

- The semi-infinite subspace $\mathcal{L}(t)$, $t \in \mathcal{M}_{\Lambda T}$, is defined for *arbitrary* projective manifold/ \mathbb{C}

$$\mathcal{L}(\gamma_0) : [r_u(\exp \frac{1}{u} i_{\gamma_0})(\varphi_0 + u\varphi_1 + \dots)], \varphi_i \in \Omega_{DR}$$

where $\gamma_0 \in \Omega^{0,*}(M, \Lambda^* T)$, $\bar{\partial}\gamma_0 + \frac{1}{2}[\gamma_0, \gamma_0] = 0$

-

$$u\mathcal{L}(t) \subset \mathcal{L}(t), \frac{\partial}{\partial t}\mathcal{L}(t) \subset u^{-1}\mathcal{L}(t)$$

$$\frac{\partial}{\partial u}\mathcal{L}(t) \subset u^{-2}\mathcal{L}(t), \mathcal{L}(u) \oplus \overline{\mathcal{L}}(\bar{u}|_{\bar{u}=u^{-1}})$$

implies tt^* -equations, remarkably $\mathcal{D}_{\frac{\partial}{\partial u}}$ -modules over \mathbb{A}^1 with similar properties appeared in 70s in works of Birkhoff, Malgrange, K.Saito and M.Saito

- Over mod space of complex structures it reduces to VHS

Noncommutative Hodge structures ([B5]), cont'd

- $\mathcal{L}(t)$ corresponds via HKR and formality isomorphisms for $C^*(A, A) + k[\xi, \frac{\partial}{\partial \xi}]$ -module $C_*(A)$ to

$$\mathcal{L}(t) = HC^-(A_t) \subset HP(A_t)$$

Noncommutative Hodge structures ([B5]), cont'd

- $\mathcal{L}(t)$ corresponds via HKR and formality isomorphisms for $C^*(A, A) + k[\xi, \frac{\partial}{\partial \xi}]$ -module $C_*(A)$ to

$$\mathcal{L}(t) = HC^-(A_t) \subset HP(A_t)$$

- Recall $HP : C_*(A)((u))$, $b + uB$, $HC^- : C_*(A)[[u]]$, $b + uB$

Noncommutative Hodge structures ([B5]), cont'd

- $\mathcal{L}(t)$ corresponds via HKR and formality isomorphisms for $C^*(A, A) + k[\xi, \frac{\partial}{\partial \xi}]$ -module $C_*(A)$ to

$$\mathcal{L}(t) = HC^-(A_t) \subset HP(A_t)$$

- Recall $HP : C_*(A)((u))$, $b + uB$, $HC^- : C_*(A)[[u]]$, $b + uB$
- Let A be an arbitrary A_∞ -algebra, the $\frac{\infty}{2}$ subspace $HC^-(A) \rightarrow HP(A)$,

$$uHC^-(A) \subset HC^-(A)$$

$$\frac{\partial}{\partial u} HC^-(A) \subset u^{-2} HC^-(A)$$

$$\frac{\partial}{\partial t} HC^-(A_t) \subset u^{-1} HC^-(A_t), \quad \frac{\partial}{\partial t} - \text{Getzler flat connection on } HP(A_t)$$

where

$$rk_{C[[u]]} HC^-(A) = rk_{C((u))} HP$$

assumed, i.e. the degeneration of nc Hodge -to-De Rham spectral sequence, proven (Kaledin) for A -smooth and compact, Z_+ -graded, then $HC^- \subset HP$,

Noncommutative Hodge structures ([B5]), cont'd

- $\mathcal{L}(t)$ corresponds via HKR and formality isomorphisms for $C^*(A, A) + k[\xi, \frac{\partial}{\partial \xi}]$ -module $C_*(A)$ to

$$\mathcal{L}(t) = HC^-(A_t) \subset HP(A_t)$$

- Recall $HP : C_*(A)((u)), b + uB, HC^-(A) : C_*(A)[[u]], b + uB$
- Let A be an arbitrary A_∞ -algebra, the $\frac{\infty}{2}$ subspace $HC^-(A) \rightarrow HP(A)$,

$$uHC^-(A) \subset HC^-(A)$$

$$\frac{\partial}{\partial u} HC^-(A) \subset u^{-2} HC^-(A)$$

$$\frac{\partial}{\partial t} HC^-(A_t) \subset u^{-1} HC^-(A_t), \quad \frac{\partial}{\partial t} - \text{Getzler flat connection on } HP(A_t)$$

where

$$rk_{C[[u]]} HC^-(A) = rk_{C((u))} HP$$

- assumed, i.e. the degeneration of nc Hodge -to-De Rham spectral sequence, proven (Kaledin) for A -smooth and compact, Z_+ -graded, then $HC^- \subset HP$,
- Real structure on HP in the case of arbitrary A_∞ -algebra?

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \Sigma \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$

-critical value of the BCOV Kodaira-Spencer lagrangian

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$
- critical value of the BCOV Kodaira-Spencer lagrangian
- \mathcal{F}^{BCOV} is equal to the summation over trees with

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$

-critical value of the BCOV Kodaira-Spencer lagrangian

- \mathcal{F}^{BCOV} is equal to the summation over trees with

- edges $\rightarrow X = " \Delta \bar{\partial}^{-1} "$

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$

-critical value of the BCOV Kodaira-Spencer lagrangian

- \mathcal{F}^{BCOV} is equal to the summation over trees with

- edges $\rightarrow X = " \Delta \bar{\partial}^{-1} "$

- vertices \rightarrow product tensor on $\Omega^{0,*}(M, \Lambda^* T)$

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$
- critical value of the BCOV Kodaira-Spencer lagrangian
- \mathcal{F}^{BCOV} is equal to the summation over trees with
 - edges $\rightarrow X = " \Delta \bar{\partial}^{-1} "$
 - vertices \rightarrow product tensor on $\Omega^{0,*}(M, \Lambda^* T)$
 - legs \rightarrow elements of $H_*(\Lambda^* T)$

Summation over trees, BCOV lagrangian and BV cyclic operad.

- For $W = \overline{F^{Hodge}}$, $\gamma(t) = \sum \gamma_a t^a + \Delta\alpha(t)$, $\Delta(\gamma) = 0$

$$C_{kij}(t) = \partial_{kij}^3 \mathcal{F}^{BCOV}(t)$$

- $\mathcal{F}^{BCOV}(t) = \int -\frac{1}{2} \bar{\partial}\alpha(t) \wedge \Delta\alpha(t) + \frac{1}{3!} \gamma(t) \wedge \gamma(t) \wedge \gamma(t)$

-critical value of the BCOV Kodaira-Spencer lagrangian

- \mathcal{F}^{BCOV} is equal to the summation over trees with
 - edges $\rightarrow X = " \Delta \bar{\partial}^{-1} "$
 - vertices \rightarrow product tensor on $\Omega^{0,*}(M, \Lambda^* T)$
 - legs \rightarrow elements of $H_*(\Lambda^* T)$
- Meaning of this : the tree-level Feynman transform of the operad of BV algebras / $\Delta = 0 =$ operad of $H_*(\mathcal{M}_{0,n})$

References:

- [B1] S.Barannikov, *Modular operads and Batalin-Vilkovisky geometry*. IMRN, Vol. 2007, article ID rnm075. Preprint Max Planck Institute for Mathematics 2006-48 (04/2006),
- [B2] S.Barannikov, *Noncommutative Batalin-Vilkovisky geometry and matrix integrals*. «Comptes rendus Mathematique», presented for publication by M.Kontsevich in 05/2009, arXiv:0912.5484; Preprint NI06043 Newton Institute (09/2006), Preprint HAL, the electronic CNRS archive, hal-00102085 (09/2006)
- [B3] S.Barannikov, *Supersymmetry and cohomology of graph complexes*. Preprint hal-00429963; (11/2009).
- [B4] S.Barannikov, *Matrix De Rham complex and quantum A-infinity algebras*. arXiv:1001.5264, Preprint hal-00378776; (04/2009).
- [B5] S.Barannikov, *Quantum periods - I. Semi-infinite variations of Hodge structures*. Preprint ENS DMA-00-19. arXiv:math/0006193 (06/2000), Intern. Math. Res. Notices. 2001, No. 23
- [B6] S.Barannikov, *Solving the noncommutative Batalin-Vilkovisky equation*. Preprint hal-00464794 (03/2010). arXiv:1004.2253