

# Elimination of Hyperimaginaries and Stable Independence in simple CM-trivial theories

Daniel Palacin, Frank Olaf Wagner

### ▶ To cite this version:

Daniel Palacin, Frank Olaf Wagner. Elimination of Hyperimaginaries and Stable Independence in simple CM-trivial theories. 2011. hal-00495655v3

## HAL Id: hal-00495655 https://hal.science/hal-00495655v3

Preprint submitted on 10 Feb 2012 (v3), last revised 19 Mar 2012 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### ELIMINATION OF HYPERIMAGINARIES AND STABLE INDEPENDENCE IN SIMPLE CM-TRIVIAL THEORIES

#### D. PALACÍN AND F. O. WAGNER

ABSTRACT. In a simple CM-trivial theory every hyperimaginary is interbounded with a sequence of finitary hyperimaginaries. Moreover, such a theory eliminates hyperimaginaries whenever it eliminates finitary hyperimaginaries. In a supersimple CM-trivial theory, the independence relation is stable.

#### 1. INTRODUCTION

An important notion introduced by Shelah for a first-order theory is that of an *imaginary* element: the class of a finite tuple by a  $\emptyset$ -definable equivalence relation. The construction obtained by adding all imaginary elements to a structure does not change its basic model-theoretic properties, but introduces a convenient context and language to talk about quotients (by definable equivalence relations) and *canonical parameters* of definable sets. In the context of a stable theory it also ensures the existence of *canonical bases* for arbitrary complete types, generalizing the notion of a field of definition of an algebraic variety.

The generalization of stability theory to the wider class of simple theories necessitated the introduction of hyperimaginaries, classes of countable tuples modulo  $\emptyset$ -type-definable equivalence relations. Although the relevant model-theory for hyperimaginaries has been reasonably well understood [4], they cannot simply be added as extra sorts to the underlying structure, since inequality of two hyperimaginaries amounts to non-equivalence, and thus *a priori* is an open, but not a closed condition. While hyperimaginary elements are needed for the

The first author was partially supported by research project MTM 2008-01545 of the Spanish government and research project 2009SGR 00187 of the Catalan government. The second author was partially supported by the research project ANR-09-BLAN-0047 of the Agence Nationale de la Recherche.

This work was partially done while the first author was visiting the Institut Camille Jordan of Lyon. The first author wishes to express his gratitude to the members of Lyon logic group for their hospitality. He also would like to thank Enrique Casanovas for his valuable comments.

#### D. PALACÍN AND F. O. WAGNER

general theory, all known examples of a simple theory *eliminate* them in the sense that they are interdefinable (or at least interbounded) with a sequence of ordinary imaginaries; the latter condition is called *weak elimination*. It has thus been asked (and even been conjectured):

#### Question. Do all simple theories eliminate hyperimaginaries?

The answer is positive for stable theories [11], and for supersimple theories [1]. Among non-simple theories, the relation of being infinitely close in a non-standard real-closed field gives rise to non-eliminable hyperimaginaries; Casanovas and the second author have constructed non-eliminable hyperimaginaries in a theory without the strict order property [3].

A hyperimaginary is *finitary* if it is the class of a finite tuple modulo a type-definable equivalence relation. Kim [5] has shown that small theories eliminate finitary hyperimaginaries, and a result of Lascar and Pillay [8] states that bounded hyperimaginaries can be eliminated in favour of finitary bounded ones. We shall show that in a CM-trivial simple theory all hyperimaginaries are interbounded with sequences of finitary hyperimaginaries. We shall deduce that in such a theory hyperimaginaries can be eliminated in favour of finitary ones. In particular, a small CM-trivial simple theory eliminates hyperimaginaries. However, even the question whether all one-based simple theories eliminate hyperimaginaries is still open.

Elimination of hyperimaginaries is closely related to another question, the *stable forking conjecture*:

**Question.** In a simple theory, if  $a \not\perp_B M$  for some model M containing B, is there a stable formula in tp(a/M) which forks over B?

If we do not require M to be a model, nor to contain B, this is called *strong stable forking*. Every known simple theory has stable forking; Kim [6] has shown that one-based simple theories with elimination of hyperimaginaries have stable forking. Kim and Pillay [7] have strengthened this to show that one-based simple theories with weak elimination of imaginaries hyperimaginaries have strong stable forking; on the other hand pseudofinite fields (which are supersimple of SU-rank 1) do not. Conversely, stable forking implies weak elimination of hyperimaginaries (Adler).

While we shall not attack the stable forking conjecture as such, we shall show in the last section that the independence relation  $x 
ightarrow y_1 y_2$  is stable, meaning that it cannot order an infinite indiscernible sequence.

#### 2. Preliminaries

As usual, we shall work in the monster model  $\mathfrak{C}$  of a complete firstorder theory (with infinite models), and all sets of parameters and all sequences of elements will live in  $\mathfrak{C}^{eq}$ . Given any sequences a, b and any set of parameters A, we write  $a \equiv_A b$  whenever a and b have the same type over A. We shall write  $a \equiv_A^s b$  if in addition a and b lie in the same class modulo all A-definable finite equivalence relations (i.e. if a and b have the same strong type over A), and  $a \equiv_A^{Ls} b$  if they lie in the same class modulo all A-invariant bounded equivalence relations (i.e. if a and b have the same Lascar strong type over A). Recall that a theory is G-compact over a set A iff  $\equiv_A^{Ls}$  is type-definable over A (in which case it is the finest bounded equivalence relation type-definable over A). A theory T is G-compact whenever it is G-compact over any A. In particular, simple theories are G-compact [5].

**Definition 2.1.** A hyperimaginary h is finitary if  $h \in dcl^{heq}(a)$  for some finite tuple a of imaginaries, and quasi-finitary if  $h \in bdd(a)$  for some finite tuple a of imaginaries.

**Definition 2.2.** A hyperimaginary h is *eliminable* if it is interdefinable with a sequence  $e = (e_i : i \in I)$  of imaginaries, i.e. if there is such a sequence e with  $dcl^{heq}(e) = dcl^{heq}(h)$ . A theory T eliminates (finitary/quasi-finitary) hyperimaginaries if all (finitary/quasi-finitary) hyperimaginaries of T.

**Remark 2.3.** [8, Corollary 1.5] If  $h \in \operatorname{dcl}^{heq}(a)$ , then there is a typedefinable equivalence relation E on  $\operatorname{tp}(a)$  such that h and the class  $a_E$ of a modulo E are interdefinable.

**Lemma 2.4.** Let e be a finitary hyperimaginary. If T eliminates finitary hyperimaginaries, then T(e) eliminates finitary hyperimaginaries.

*Proof.* Let a be a finite tuple with  $e \in \operatorname{dcl}^{heq}(a)$ , and h a finitary hyperimaginary over e. So there is a finite tuple b with  $h \in \operatorname{dcl}^{heq}(eb) \subseteq \operatorname{dcl}^{heq}(ab)$ . Then there is a type-definable equivalence relation E over  $\emptyset$  such that e and  $a_E$  are interdefinable, and a type-definable equivalence relation  $F_a$  over a such that h and  $b_{F_a}$  are interdefinable. Moreover,  $F_a$  only depends on the E-class of a, that is, if a'Ea, then  $F_{a'} = F_a$ .

Type-define an equivalence relation by

 $xy\bar{E}uv \Leftrightarrow xEu \wedge yF_xv.$ 

It is easy to see that h is interdefinable with  $(ab)_{\overline{E}}$  over e. Moreover,  $(ab)_{\overline{E}}$  is clearly finitary, and hence eliminable in T. So h is eliminable in T(e).

The following fact appears in [8, Proof of Proposition 2.2], but was first stated as such in [1, Lemma 2.17].

**Fact 2.5.** Let *h* be a hyperimaginary and let *a* be a sequence of imaginaries such that  $a \in bdd(h)$  and  $h \in dcl^{heq}(a)$ . Then, *h* is eliminable.

**Fact 2.6.** [1, Lemma 2.18] Let h, e be hyperimaginaries with  $h \in bdd(e)$ . Then the set of *e*-conjugates of *h* is interdefinable with a hyperimaginary h'.

**Fact 2.7.** [8, Theorem 4.15] A bounded hyperimaginary is interdefinable with a sequence of finitary hyperimaginaries.

**Proposition 2.8.** If T eliminates finitary hyperimaginaries, then T eliminates quasi-finitary hyperimaginaries.

*Proof.* Let h be a quasi-finitary hyperimaginary and let a be a finite tuple of imaginaries such that  $h \in bdd(a)$ . Consider  $a' \equiv_h a$  with  $bdd(a) \cap bdd(a') = bdd(h)$ . Let h' be the hyperimaginary corresponding to the set of aa'-conjugates of h. Then h' is aa'-invariant, and hence finitary. It is thus interdefinable with a sequence e of imaginaries.

On the other hand,  $h \in bdd(a) \cap bdd(a')$ , as are all its aa'-conjugates. Thus  $h' \in bdd(a) \cap bdd(a') = bdd(h)$ . Hence  $e \in acl^{eq}(h)$  and  $h \in bdd(h') = bdd(e)$ . By Fact 2.7, there is a sequence h'' of finitary hyperimaginaries interdefinable with h over e. By Lemma 2.4 and elimination of finitary hyperimaginaries we see that h'' is interdefinable over e with a sequence e' of imaginaries. So  $h \in dcl^{heq}(ee')$  and  $e' \in dcl^{eq}(eh)$ . Moreover,  $ee' \in acl^{eq}(h)$  since  $e \in acl^{eq}(h)$ . Hence h is eliminable by Fact 2.5.

The following remarks and lemmata will need G-compactness.

**Remark 2.9.** Let T be G-compact over a set A. The following are equivalent:

- (1)  $a \equiv^{L_s}_A b$  iff  $a \equiv^s_A b$  for all sequences a, b.
- (2)  $\operatorname{Aut}(\mathfrak{C}/\operatorname{bdd}(A)) = \operatorname{Aut}(\mathfrak{C}/\operatorname{acl}^{eq}(A)).$
- (3)  $bdd(A) = dcl^{heq}(acl^{eq}(A)).$

*Proof.* Easy exercise.

**Remark 2.10.** Let *T* be a *G*-compact theory and assume further that  $a \equiv_A^{L_s} b \Leftrightarrow a \equiv_A^s b$  for all sequences a, b and for any set *A*. Let now *h* be a hyperimaginary and let *e* be a sequence of imaginaries such that *h* and *e* are interbounded. Then *h* is eliminable.

*Proof.* It follows from Remark 2.9 that  $bdd(e) = dcl^{heq}(acl^{eq}(e))$ . Fix an enumeration  $\bar{e}$  of  $acl^{eq}(e)$  and observe that  $h \in dcl^{heq}(\bar{e})$  and  $\bar{e} \in bdd(h)$ . Then apply Fact 2.5 to eliminate h.

4

It turns out for G-compact theories that elimination of hyperimaginaries can be decomposed as weak elimination of hyperimaginaries plus the equality between Lascar strong types and strong types over parameter sets.

**Fact 2.11.** [2, Proposition 18.27] Assume that T is G-compact. Then T eliminates all bounded hyperimaginaries iff  $a \equiv^{L_s} b \Leftrightarrow a \equiv^s b$  for all sequences a, b.

*Proof.* The proof in [2] is nice and intuitive; however, we will give another one using Remark 2.10. If T eliminates bounded hyperimaginaries, then  $\operatorname{Aut}(\mathfrak{C}/\operatorname{bdd}(\emptyset)) = \operatorname{Aut}(\mathfrak{C}/\operatorname{acl}^{eq}(\emptyset))$ . By Remark 2.9 we get  $\operatorname{Lstp} = \operatorname{stp.}$  For the other direction, let  $e \in \operatorname{bdd}(\emptyset)$  and let  $\overline{a}$  be an enumeration of  $\operatorname{acl}^{eq}(\emptyset)$ . It is clear that e and  $\overline{a}$  are interbounded. By Remark 2.10, e is eliminable.  $\Box$ 

**Lemma 2.12.** Suppose T is G-compact and assume further that T eliminates finitary hyperimaginaries. Then  $a \equiv_A^{Ls} b$  iff  $a \equiv_A^s b$  for all sequences a, b and for any set A.

*Proof.* Since T is G-compact, it is enough to check the condition for finite A. But then T(A) eliminates finitary hyperimaginaries by Remark 2.4, and hence all bounded hyperimaginaries by Fact 2.7. Now applying Fact 2.11 we obtain that  $a \equiv^{L_s} b$  iff  $a \equiv^s b$  in T(A).

#### 3. Elimination of hyperimaginaries in simple theories

In this section T will be a simple theory. Recall that the *canonical* base of a over b, denoted  $\operatorname{Cb}(a/b)$ , is the smallest definably closed subset C of  $\operatorname{bdd}(b)$  such that  $a \bigcup_{C} b$  and  $\operatorname{tp}(a/C)$  is Lascar strong.

**Lemma 3.1.** For any a and any  $h \in bdd(c)$  we have  $Cb(a/h) \subseteq dcl(ac) \cap bdd(h)$ . Therefore, the canonical base of the type of an imaginary finite tuple over a quasi-finitary hyperimaginary is finitary. Furthermore, if  $b \in Cb(a/c)$  then  $dcl(ab) \cap bdd(b) \subseteq Cb(a/c)$ . In particular, if  $c \in dcl(a)$  then  $Cb(a/c) = dcl(a) \cap bdd(c)$ .

*Proof.* Since  $h \in bdd(c)$ , equality of Lascar strong types over c refines equality of Lascar strong types over h, and the class of a modulo the former is clearly in dcl(ac). So the class of a modulo the latter is in dcl(ac), and  $Cb(a/h) \in dcl(ac) \cap bdd(h)$ . As a consequence, if a is a finite tuple and h is a quasi-finitary hyperimaginary bounded over some finite tuple c, then Cb(a/h) is definable over the finite tuple ac.

For the second assertion put  $b' = \operatorname{dcl}(ab) \cap \operatorname{bdd}(b)$ . Since  $b' \in \operatorname{dcl}(ab)$ there is an equivalence relation E on  $\operatorname{tp}(a/b)$  type-definable over b such that b' is interdefinable over b with  $a_E$ . As  $b' \in \operatorname{bdd}(b)$  and  $b \in \operatorname{Cb}(a/c)$ , the *E*-class of *a* is bounded over Cb(a/c); as tp(a/Cb(a/c)) is Lascarstrong,  $a_E \in Cb(a/c)$ .

The "in particular" clause is essentially [1, Remark 3.8]: If  $c \in dcl(a)$  then clearly  $c \in Cb(a/c)$ ; the assertion follows.

Recall the definition of CM-triviality.

**Definition 3.2.** A simple theory T is *CM-trivial* if for every tuple a and for any sets  $A \subseteq B$  with  $bdd(aA) \cap bdd(B) = bdd(A)$  we have  $Cb(a/A) \subseteq bdd(Cb(a/B))$ .

**Remark 3.3.** As in [10, Corollary 2.5], in the definition of CM-triviality we may take  $A \subseteq B$  to be models of the ambient theory and a to be a tuple from the home sort. Therefore, it makes no difference in the definition of CM-triviality whether we consider hyperimaginaries or just imaginaries.

Now we characterize canonical bases in simple CM-trivial theories in terms of finitary hyperimaginaries.

**Proposition 3.4.** Assume the theory is simple CM-trivial. If a is a finite imaginary tuple, then

$$bdd(Cb(a/B)) = bdd(Cb(a/b) : b \in X),$$

where X is the set of all finitary  $b \in bdd(Cb(a/B))$ .

*Proof.* Since  $\operatorname{Cb}(a/b) \subseteq \operatorname{bdd}(b) \subseteq \operatorname{bdd}(\operatorname{Cb}(a/B))$  for  $b \in X$ , we have

$$bdd(Cb(a/b): b \in X) \subseteq bdd(Cb(a/B)).$$

For the reverse inclusion, for every  $b \in X$  let  $\hat{b}$  be a real tuple with  $\operatorname{Cb}(a/b) \in \operatorname{dcl}(\hat{b})$ ; we choose them such that

$$(b: b \in X) \bigsqcup_{(\operatorname{Cb}(a/b): b \in X)} aB,$$

whence  $(\hat{b}: b \in X) igsquare$ \_B a.

Now, if  $a \not \perp_{(\hat{b}:b\in X)} B$  then there is a finite tuple  $b' \in B \cup \{\hat{b}: b \in X\}$ and a formula  $\varphi(x, b') \in \operatorname{tp}(a/B, \hat{b}: b \in X)$  which divides over  $(\hat{b}: b \in X)$ . Put  $\bar{b} = \operatorname{bdd}(ab') \cap \operatorname{bdd}(B, \hat{b}: b \in X)$ . Then  $\bar{b}$  is a quasi-finitary hyperimaginary, and by CM-triviality

$$\operatorname{Cb}(a/\overline{b}) \subseteq \operatorname{bdd}(\operatorname{Cb}(a/B, \hat{b} : b \in X)) = \operatorname{bdd}(\operatorname{Cb}(a/B)).$$

Since  $\operatorname{Cb}(a/\overline{b})$  is finitary by Lemma 3.1, it belongs to X. Note that  $b' \in \overline{b}$ ; but  $a \bigcup_{\operatorname{Cb}(a/\overline{b})} \overline{b}$ , so  $\varphi(x, b')$  cannot divide over  $\operatorname{Cb}(a/\overline{b})$ , and

even less over  $(\hat{b}: b \in X)$  as this contains  $Cb(a/\bar{b})$ . Thus,  $a \downarrow_{(\hat{b}:b\in X)} B$ , whence  $a \downarrow_{(Cb(a/b):b\in X)} B$  by transitivity. Therefore

$$\operatorname{Cb}(a/B) \subseteq \operatorname{bdd}(\operatorname{Cb}(a/b) : b \in X).$$

**Question.** The same proof will work without assuming CM-triviality if for every finite tuple  $b \in B$  there is some quasi-finitary hyperimaginary  $\bar{b} \in \text{bdd}(B)$  with  $b \in \text{dcl}(\bar{b})$  such that  $\text{Cb}(a/\bar{b}) \subseteq \text{bdd}(\text{Cb}(a/B))$ . Is this true in general?

We can now state (and prove) the main result.

**Theorem 3.5.** Let T be a simple CM-trivial theory. Then every hyperimaginary is interbounded with a sequence of finitary hyperimaginaries.

*Proof.* By Lemma 3.1 every hyperimaginary is interbounded with a canonical base. Since  $\operatorname{Cb}(A/B)$  is interdefinable with  $\bigcup \{\operatorname{Cb}(\bar{a}/B) : \bar{a} \in A \text{ finite}\}$ , it is enough to show that canonical bases of types of finite tuples are interbounded with sequences of finitary hyperimaginaries. This is precisely Proposition 3.4.

**Corollary 3.6.** A simple CM-trivial theory eliminates hyperimaginaries whenever it eliminates finitary ones.

*Proof.* By Theorem 3.5 every hyperimaginary is interbounded with a sequence of finitary hyperimaginaries and so with a sequence of imaginaries. Since T is simple, it is G-compact, whence Lstp = stp over any set by Lemma 2.12. We conclude that every hyperimaginary is eliminable by Remark 2.10.

**Corollary 3.7.** Every small simple CM-trivial theory eliminates hyperimaginaries.

*Proof.* A small simple theory eliminates finitary hyperimaginaries by [5]. Now apply Corollary 3.6.

#### 4. Stable independence for CM-trivial theories

Recall that an  $\emptyset$ -invariant relation R(x, y) is *stable* if there is no infinite indiscernible sequence  $(a_i, b_i : i < \omega)$  such that  $R(a_i, b_j)$  holds if and only if i < j. In this section, we shall show that independence is a stable relation, even with varying base set. We hope that this will help elucidate the stable forking problem.

**Theorem 4.1.** In a supersimple CM-trivial theory, the relation  $R(x; y_1y_2)$  given by  $x \downarrow_{y_1} y_2$  is stable.

*Proof.* Suppose not. Then there is an indiscernible sequence  $I = (a_i : a_i)$  $i \in \mathbb{Q}$ ) and tuples b, c such that

- $I^+ = (a_i : i > 0)$  is indiscernible over  $I^-bc$ ,
- $I^- = (a_i : i < 0)$  is indiscernible over  $I^+bc$ , and
- $a_i \, \bigcup_{a} b$  if and only if i > 0.

We consider limit types with respect to the cut at 0. Let  $p = \lim(I/I)$ ,  $p^+ = \lim(I^+/Ibc)$  and  $p^- = \lim(I^-/Ibc)$ . By finite satisfiability,  $p^+$ and  $p^-$  are both non-forking extensions of p, which is Lascar-strong. Let

$$A = \operatorname{Cb}(p) = \operatorname{Cb}(p^+) = \operatorname{Cb}(p^-) \in \operatorname{bdd}(I^+) \cap \operatorname{bdd}(I^-).$$

As  $p^+$  and  $p^-$  do not fork over A, we have

$$a_i \underset{A}{\bigcup} I^+ bc$$
 for all  $i < 0$ , and  $a_i \underset{A}{\bigcup} I^- bc$  for all  $i > 0$ .

We consider first  $e_0 = bdd(a_1c) \cap bdd(Ac)$ . Then

 $bdd(a_1e_0) \cap bdd(Ae_0) = e_0.$ 

Put  $A_0 = Cb(a_1/e_0)$ . By CM-triviality

$$A_0 \in e_0 \cap \mathrm{bdd}(\mathrm{Cb}(a_1/Ae_0)) \subseteq \mathrm{bdd}(a_1c) \cap \mathrm{bdd}(A),$$

since  $a_1 
int_A e_0$  implies  $\operatorname{Cb}(a_1/Ae_0) \subseteq \operatorname{bdd}(A)$ . Note that  $a_1 
int_c b$  yields  $a_1 
int_{A_0c} b$ . Moreover  $c \in e_0$ , so  $a_1 
int_{A_0} e_0$ implies  $a_1 
int_{A_0} c$ , whence  $a_1 
int_{A_0} cb$  by transitivity. On the other hand, suppose  $bc 
int_{A_0} a_{-1}$ . Then  $b 
int_{A_{0c}} a_{-1}$ ; as  $b 
int_c a_1$  implies  $b 
int_c A_0$ , we obtain  $b 
int_c A_0 a_{-1}$ , contradicting  $a_{-1} 
int_c b$ . Therefore  $bc 
int_{A_0} a_{-1}$ . Since I remains indiscernible over  $A_0$ , and both  $I^+$  and  $I^-$  remain indiscernible over  $A_0bc$ , we may add  $A_0$  to the parameters and suppose  $c = \emptyset$  (replacing b by bc).

Fact 4.2. [12, Theorem 5.2.18] In a supersimple theory, for any finitary a there are some  $B \mid a$  and a hyperimaginary finite tuple  $\bar{a}$  of independent realizations of regular types over B, such that  $\bar{a}$  is dominationequivalent with a over B.

By Fact 4.2 there are  $B \, \sqcup \, a_1$  and an independent tuple  $\bar{a}_1$  of realizations of regular types over B such that  $\bar{a}_1$  is domination-equivalent with  $a_1$  over B. Since  $B \, \sqcup \, a_1$  and I is indiscernible, we may assume by [12, Theorem 2.5.4] that  $Ba_i \equiv Ba_1$  for all  $i \in \mathbb{Q}$ , and  $B \downarrow I$ . So there are  $\bar{a}_i$  for  $i \in \mathbb{Q}$  with  $Ba_i\bar{a}_i \equiv Ba_1\bar{a}_1$ . We can also assume  $B imes_I b$ , whence  $B imes_I Ib$ . In particular  $b imes_{a_i} B$ , so for i > 0 we obtain  $b \perp Ba_i$  and thus  $b \perp_B a_i$ , while for i < 0 we have  $b \not\perp a_i B$  and  $b \perp B$ ,

whence  $b \not\perp_B a_i$ . By domination-equivalence,  $\bar{a}_i \perp_B b$  for i > 0 whereas  $\bar{a}_i \not \perp_B b$  for i < 0.

By compactness and Ramsey we may suppose in addition that  $\bar{I} =$  $(\bar{a}_i : i \in \mathbb{Q})$  is indiscernible over  $B, \bar{I}^+ = (\bar{a}_i : i > 0)$  is indiscernible over  $Bb\bar{I}^-$  and  $\bar{I}^- = (\bar{a}_i : i < 0)$  is indiscernible over  $Bb\bar{I}^+$ . We shall add B to the parameters and suppress it from the notation. We may further assume that  $\bar{a}'_{-1} \downarrow b$  for any proper subtuple  $\bar{a}'_{-1} \subseteq \bar{a}_{-1}$ . It follows that all the regular types in  $\bar{a}_i$  are non-orthogonal. Let  $w_{\mathcal{P}}(.)$ denote the weight with respect to that non-orthogonality class  $\mathcal{P}$  of regular types.

We again consider limit types with respect to the cut at 0. Let  $\bar{p} = \lim(\bar{I}/\bar{I}), \ \bar{p}^+ = \lim(\bar{I}^+/\bar{I}b) \text{ and } \ \bar{p}^- = \lim(\bar{I}^-/\bar{I}b).$  Once more,  $\bar{p}$  is Lascar-strong and  $\bar{p}^+$  and  $\bar{p}^-$  are non-forking extensions of  $\bar{p}$  by finite satisfiability; let

$$\bar{A} = \operatorname{Cb}(\bar{p}) = \operatorname{Cb}(\bar{p}^+) = \operatorname{Cb}(\bar{p}^-) \in \operatorname{bdd}(\bar{I}^+) \cap \operatorname{bdd}(\bar{I}^-).$$

As before,

$$\bar{a}_i \underset{\bar{A}}{\downarrow} \bar{I}^+ b$$
 for all  $i < 0$ , and  $\bar{a}_i \underset{\bar{A}}{\downarrow} \bar{I}^- b$  for all  $i > 0$ .

Put  $e_1 = bdd(\bar{a}_{-1}b) \cap bdd(Ab)$ . Then

$$bdd(\bar{a}_{-1}e_1) \cap bdd(\bar{A}e_1) = e_1.$$

Let  $A_1 = \operatorname{Cb}(\bar{a}_{-1}/e_1)$ . By CM-triviality

$$A_1 \in e_1 \cap \mathrm{bdd}(\mathrm{Cb}(\bar{a}_{-1}/\bar{A}e_1)) \subseteq \mathrm{bdd}(\bar{a}_{-1}b) \cap \mathrm{bdd}(\bar{A}),$$

since  $\bar{a}_{-1} \, \bigcup_{\bar{A}} e_1$  implies  $\operatorname{Cb}(a_{-1}/\bar{A}e_1) \subseteq \operatorname{bdd}(\bar{A})$ . As  $b \in e_1$  and  $\bar{a}_{-1} \, \bigcup_{A_1} e_1$  we obtain  $\bar{a}_{-1} \, \bigcup_{A_1} b$ . Moreover  $\bar{a}_1 \equiv_{A_1} \bar{a}_{-1}$ , since  $A_1 \subseteq \text{bdd}(\bar{A})$  and  $\bar{I}$  remains indiscernible over  $\bar{A}$ . Therefore

$$w_{\mathcal{P}}(\bar{a}_{-1}/A_1b) = w_{\mathcal{P}}(\bar{a}_{-1}/A_1) = w_{\mathcal{P}}(\bar{a}_1/A_1).$$

Recall that  $A_1 \subseteq \text{bdd}(\bar{a}_{-1}b)$ . Then

$$w_{\mathcal{P}}(\bar{a}_{-1}/b) = w_{\mathcal{P}}(\bar{a}_{-1}A_{1}/b) = w_{\mathcal{P}}(\bar{a}_{-1}/A_{1}b) + w_{\mathcal{P}}(A_{1}/b) = w_{\mathcal{P}}(\bar{a}_{1}/A_{1}) + w_{\mathcal{P}}(A_{1}/b) \geq w_{\mathcal{P}}(\bar{a}_{1}/A_{1}b) + w_{\mathcal{P}}(A_{1}/b) = w_{\mathcal{P}}(\bar{a}_{1}A_{1}/b) \geq w_{\mathcal{P}}(\bar{a}_{1}/b) = w_{\mathcal{P}}(a_{1}) = w_{\mathcal{P}}(a_{-1}) > w_{\mathcal{P}}(\bar{a}_{-1}/b),$$

since  $\bar{a}_1 \perp b$  and  $\bar{a}_{-1} \not\perp b$ . This final contradiction proves the theorem.

**Remark 4.3.** Note that the proof only uses the conclusion of Fact 4.2. The theorem thus still holds for simple CM-trivial theories with finite weights (strongly simple theories) and enough regular types, for instance CM-trivial simple theories without dense forking chains.

**Question.** By [9, Theorem 4.20] it is sufficient to assume that every regular type is CM-trivial, as this implies global CM-triviality. However, for a regular type p a more general notion of CM-triviality is often more appropriate, namely

$$\operatorname{cl}_p(aA) \cap \operatorname{cl}_p(B) = \operatorname{cl}_p(A) \quad \Rightarrow \quad \operatorname{Cb}(a/\operatorname{cl}_p(A)) \subseteq \operatorname{cl}_p(\operatorname{Cb}(a/\operatorname{cl}_p(B))).$$

If this holds for all regular types p, is independence still stable?

**Corollary 4.4.** An  $\omega$ -categorical supersimple CM-trivial theory has stable forking.

Let  $\Sigma$  be an  $\emptyset\text{-invariant}$  family of types. Recall the definition of  $\Sigma\text{-closure:}$ 

 $cl_{\Sigma}(A) = \{a : tp(a/A) \text{ is } \Sigma\text{-analysable}\}.$ 

**Fact 4.5.** [12, Lemma 3.5.3 and 3.5.5] If  $dcl(AB) \cap cl_{\Sigma}(A) \subseteq bdd(A)$ , then  $B \, \bigcup_{A} cl_{\Sigma}(A)$ . If  $A \, \bigcup_{B} C$ , then  $A \, \bigcup_{cl_{\Sigma}(B)} C$ .

**Corollary 4.6.** In a supersimple CM-trivial theory the relation  $R(x; y_1y_1)$  given by  $x \downarrow_{cly(y_1)} y_2$  is stable.

*Proof.* Suppose not. Then there is an indiscernible sequence  $I = (a_i : i \in \mathbb{Q})$  and tuples b, c such that

- $I^+ = (a_i : i > 0)$  is indiscernible over  $I^-bc$ ,
- $I^- = (a_i : i < 0)$  is indiscernible over  $I^+bc$ , and
- $a_i \perp_{\operatorname{cl}_{\Sigma}(c)} b$  if and only if i > 0.

Put  $c' = \operatorname{dcl}(bc) \cap \operatorname{cl}_{\Sigma}(c)$ . By Fact 4.5 we have  $b \bigcup_{c'} \operatorname{cl}_{\Sigma}(c)$ , so by transitivity  $a_i \bigcup_{c'} b$  for i > 0. Suppose  $a_i \bigcup_{c'} b$  for i < 0. Since  $\operatorname{cl}_{\Sigma}(c') = \operatorname{cl}_{\Sigma}(c)$ , Fact 4.5 yields  $a_i \bigcup_{\operatorname{cl}_{\Sigma}(c)} b$ , a contradiction. Thus  $a_i \bigcup_{c'} b$  if and only if i > 0, contradicting Theorem 4.1.

To conclude the paper we prove a version of Corollary 4.6 without the assumption of CM-triviality, but for a particular  $\emptyset$ -invariant family, namely the family  $\mathcal{P}$  of all non one-based types.

**Fact 4.7.** [9, Corollary 5.2] In a simple theory  $a \, {\rm bed}_{{\rm cl}_{\mathcal{P}}(a) \cap {\rm bdd}(b)} b$  for all tuples a and b.

**Theorem 4.8.** In a simple theory, the relation  $R(x; y_1y_2)$  given by  $x 
otin_{\operatorname{cl}_{\mathcal{P}}(y_1)} y_2$  is stable.

*Proof.* Suppose not. Then there is an indiscernible sequence  $I = (a_i : i \in \mathbb{Q})$  and tuples b, c such that

- $I^+ = (a_i : i > 0)$  is indiscernible over  $I^-bc$ ,
- $I^- = (a_i : i < 0)$  is indiscernible over  $I^+bc$ , and
- $a_i \perp_{cl_{\mathcal{P}}(c)} b$  if and only if i > 0.

As before, we consider limit types with respect to the cut at 0. Let  $p = \lim(I/I)$ ,  $p^+ = \lim(I^+/Ib)$  and  $p^- = \lim(I^-/Ib)$ . By finite satisfiability,  $p^+$  and  $p^-$  are both non-forking extensions of p, which is Lascar-strong. Let

$$A = \operatorname{Cb}(p) = \operatorname{Cb}(p^+) = \operatorname{Cb}(p^-) \in \operatorname{bdd}(I^+) \cap \operatorname{bdd}(I^-).$$

As in the proof of Theorem 4.1 we have

$$a_i \underset{A}{\bigcup} I^+ bc$$
 for all  $i < 0$ , and  $a_i \underset{A}{\bigcup} I^- bc$  for all  $i > 0$ .

We consider first  $e = cl_{\mathcal{P}}(a_1) \cap bdd(A)$ . Then  $a_1 \, \bigcup_e A$  by Fact 4.7; since I remains indiscernible over bdd(A) we have  $a_{-1} \equiv_{bdd(A)} a_1$ , whence  $e = cl_{\mathcal{P}}(a_{-1}) \cap bdd(A)$  and  $a_{-1} \, \bigcup_e A$ . On the other hand, since  $e \in bdd(A)$  and  $a_i \, \bigcup_A bc$  for  $i \in \mathbb{Q}$  we obtain

$$a_1 \bigsqcup_e bc$$
 and  $a_{-1} \bigsqcup_e bc$ .

Now put  $c' = \operatorname{dcl}(bc) \cap \operatorname{cl}_{\mathcal{P}}(c)$ . Then  $b \bigcup_{c'} \operatorname{cl}_{\mathcal{P}}(c)$  by Fact 4.5. Moreover,  $a_1 \bigcup_{\operatorname{cl}_{\mathcal{P}}(c)} b$  yields  $\operatorname{cl}_{\mathcal{P}}(a_1) \bigcup_{\operatorname{cl}_{\mathcal{P}}(c)} b$  by Fact 4.5, whence  $\operatorname{cl}_{\mathcal{P}}(a_1) \bigcup_{c'} b$ . Thus  $e \bigcup_{c'} b$  and hence  $e \bigcup_{c'} bc$  since  $c \subseteq c'$ . But now  $c' \subseteq \operatorname{dcl}(bc)$  and  $a_{-1} \bigcup_e bc$  imply that  $a_{-1} \bigcup_{c'} bc$ . Hence  $a_{-1} \bigcup_{\operatorname{cl}_{\mathcal{P}}(c)} b$  by Fact 4.5, as  $\operatorname{cl}_{\mathcal{P}}(c') = \operatorname{cl}_{\mathcal{P}}(c)$ . This contradiction finishes the proof.  $\Box$ 

**Remark 4.9.** If the theory is supersimple, we can take  $\mathcal{P}$  to be the family of all non one-based *regular* types.

**Remark 4.10.** Theorem 4.8 generalises the fact that independence is stable in a one-based theory. For a true generalisation of Theorem 4.1 to arbitrary theories, one should take  $\mathcal{P}$  to be the family of all 2-ample types. This is work in progress.

#### References

- [1] Steven Buechler, Anand Pillay, and Frank O. Wagner. *Supersimple theories*. Journal of the Amercian Mathematical Society. **14(1)**:109–124, 2000.
- [2] Enrique Casanovas. *Simple theories and hyperimaginaries*. Lecture Notes in Logic. Cambridge University Press, 2011.
- [3] Enrique Casanovas and Frank O. Wagner. The free roots of the complete graph. Proc. Amer. Math. Soc. 132(5):1543–1548, 2003.
- [4] Bradd Hart, Byunghan Kim, and Anand Pillay. Coordinatisation and canonical bases in simple theories. The Journal of Symbolic Logic. 65(1):293–309, 2000.
- [5] Byunghan Kim. A note on Lascar strong types in simple theories. The Journal of Symbolic Logic. 63(3):926–936, 1998.
- [6] Byunghan Kim. Simplicity, and stability in there. The Journal of Symbolic Logic. 66(2):822–836, 2001.
- [7] Byunghan Kim and Anand Pillay. Around stable forking. Fundamenta Mathematicæ. 170(1-2):107–118, 2001.
- [8] Daniel Lascar and Anand Pillay. Hyperimaginaries and automorphism groups. The Journal of Symbolic Logic. 66(1):127–143, 2001.
- [9] Daniel Palacín and Frank O. Wagner. Ample Thoughts. Preprint, 2011. HAL 00581795, arXiv 1104.0179.
- [10] Anand Pillay. The geometry of forking and groups of finite Morley rank. The Journal of Symbolic Logic. 60(4):1251–1259, 1995.
- [11] Anand Pillay and Bruno Poizat. Pas d'imaginaires dans l'infini!. The Journal of Symbolic Logic. 52(2):400–403, 1987.
- [12] Frank O. Wagner. Simple theories. Mathematics and Its Applications 503. Kluwer Academic Publishers, Dordrecht, 2000.

Universitat de Barcelona; Departament de Lògica, Història i Filosofia de la Ciència, Montalegre 6, 08001 Barcelona, Spain

UNIVERSITÉ DE LYON; CNRS; UNIVERSITÉ LYON 1; INSTITUT CAMILLE JOR-DAN UMR5208, 43 BOULEVARD DU 11 NOVEMBRE 1918, F-69622 VILLEUR-BANNE CEDEX, FRANCE

E-mail address: dpalacin@ub.edu

E-mail address: wagner@math.univ-lyon1.fr