



HAL
open science

Essential self-adjointness for combinatorial Schrödinger operators II- Metrically non complete graphs

Yves Colin de Verdière, Nabila Torki-Hamza, Françoise Truc

► **To cite this version:**

Yves Colin de Verdière, Nabila Torki-Hamza, Françoise Truc. Essential self-adjointness for combinatorial Schrödinger operators II- Metrically non complete graphs. *Mathematical Physics Analysis and Geometry*, 2011, 14 (1), pp.21-38. 10.1007/s11040-010-9086-7. hal-00496165v2

HAL Id: hal-00496165

<https://hal.science/hal-00496165v2>

Submitted on 9 Oct 2010 (v2), last revised 3 Mar 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Essential self-adjointness for combinatorial Schrödinger operators II- Metrically non complete graphs

Yves Colin de Verdière*
Nabila Torki-Hamza †
Françoise Truc‡

October 9, 2010

Abstract

We consider weighted graphs, we equip them with a metric structure given by a weighted distance, and we discuss essential self-adjointness for weighted graph Laplacians and Schrödinger operators in the metrically non complete case.

1 Introduction

This paper is a continuation of [To] which contains some statements about essential self-adjointness of Schrödinger operators on graphs. In [To], it was proved that on any metrically complete weighted graph with bounded degree, the Laplacian is essentially self-adjoint and the same holds for the Schrödinger operator provided the associated quadratic form is bounded from below. These results remind those in the context of Riemannian manifold in [Ol] and also in [B-M-S],

⁰ **Keywords:** metrically non complete graph, weighted graph Laplacian, Schrödinger operator, essential selfadjointness.

⁰ **Math Subject Classification (2000):** 05C63, 05C50, 05C12, 35J10, 47B25.

*Grenoble University, Institut Fourier, Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France); yves.colin-de-verdiere@ujf-grenoble.fr; <http://www-fourier.ujf-grenoble.fr/~ycolver/>

†Université du 7 Novembre à Carthage, Faculté des Sciences de Bizerte, Mathématiques et Applications (05/UR/15-02), 7021-Bizerte (Tunisie); nabila.torki-hamza@fsb.rnu.tn; torki@fourier.ujf-grenoble.fr

‡Grenoble University, Institut Fourier, Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France); françoise.truc@ujf-grenoble.fr; <http://www-fourier.ujf-grenoble.fr/~trucfr/>

[Shu1], [Shu2]. There are many recent independent researches in locally finite graphs investigating essential self-adjointness (see [Jor], [Go-Sch], [Ma]), and relations between stochastic completeness and essential self-adjointness (see [We], [Woj2] as well as the thesis [Woj1]). Similar results have been extended for arbitrary regular Dirichlet forms on discrete sets in [Ke-Le-2] which is mostly a survey of the original article [Ke-Le-1]. More recently the paper [Hu] is devoted to the stability of stochastic incompleteness, in almost the same setup as in [Ke-Le-1]. Here, we will investigate essential self-adjointness mainly on metrically non complete locally finite graphs.

Let us recall that a weighted graph G is a generalization of an electrical network where the set of vertices and the set of edges are respectively weighted with positive functions ω and c . For any given positive function p , a weighted distance d_p can be defined on G . Thus we have the usual notion of completeness for G as a metric space.

The main result of Section 3 states that the weighted graph Laplacian $\Delta_{\omega,c}$ (see the definition (1) below) is not essentially self-adjoint if the graph is of finite volume and metrically non complete (here the metric d_p is defined using the weights $p_{x,y} = c_{x,y}^{-\frac{1}{2}}$). The proof is derived from the existence of the solution for a Dirichlet problem at infinity, established in Section 2.

In Section 4, we establish some conditions implying essential self-adjointness. More precisely, defining the metric d_p with respect to the weights $p_{x,y}$ given by $p_{x,y} = (\min\{\omega_x, \omega_y\})c_{x,y}^{-\frac{1}{2}}$, and addressing the case of metrically non complete graphs, we get the essential self-adjointness of $\Delta_{\omega,c} + W$ provided the potential W is bounded from below by $N/2D^2$, where N is the maximal degree and D the distance to the boundary. We use for this result a technical tool deduced from Agmon-type estimates and inspired by the nice paper [Nen], see also [Col-Tr].

We discuss in Section 5 the case of star-like graphs. Under some assumptions on a , we prove that for any potential W , $\Delta_{1,a} + W$ is essentially self-adjoint using an extension of Weyl's theory to the discrete case. In the particular case of the graph \mathbb{N} , the same result had been proved in [Ber] (p.504) in the context of Jacobi matrices. We give some examples in Subsection 5.3 to illustrate the links between the previous results. Moreover we establish the sharpness of the conditions of Theorem 4.2.

The last Section is devoted to Appendix A dealing with Weyl's limit point-limit circle criteria (see [RS]) in the discrete case as well as in the continuous case, and to Appendix B including the unitary equivalence between Laplacians and Schrödinger operators [To] used repeatedly in Subsection 5.3.

Let us start with some definitions.

$G = (V, E)$ will denote an *infinite graph*, with $V = V(G)$ the set of *vertices* and $E = E(G)$ the set of *edges*. We write $x \sim y$ for $\{x, y\} \in E$.

The graph G is always assumed to be *locally finite*, that is any $x \in V$ has a finite number of neighbors, which we call the *degree* (or valency) of x . If the degree is

bounded independently of x in V , we say that the graph G is of *bounded degree*. The space of real functions on the graph G is denoted

$$C(V) = \{f : V \longrightarrow \mathbb{R}\}$$

and $C_0(V)$ is the subspace of functions with finite support.

We consider, for any weight $\omega : V \longrightarrow]0, +\infty[$, the space

$$l_\omega^2(V) = \{f \in C(V); \sum_{x \in V} \omega_x^2 f^2(x) < \infty\}.$$

It is a Hilbert space when equipped with the inner product:

$$\langle f, g \rangle_{l_\omega^2} = \sum_{x \in V} \omega_x^2 f(x) \cdot g(x) \quad .$$

For any $\omega : V \longrightarrow]0, +\infty[$, and $c : E \longrightarrow]0, +\infty[$, the *weighted graph Laplacian* $\Delta_{\omega,c}$ on the graph G weighted by the *conductance* c on the edges and by the *weight* ω on the vertices, is defined by:

$$(\Delta_{\omega,c}f)(x) = \frac{1}{\omega_x^2} \sum_{y \sim x} c_{x,y} (f(x) - f(y)) \quad (1)$$

for any $f \in C(V)$ and any $x \in V$. If $\omega \equiv 1$, we have

$$(\Delta_{1,a}f)(x) = \sum_{y \sim x} a_{x,y} (f(x) - f(y)).$$

Definition 1.1 Let $p : E \longrightarrow]0, +\infty[$ be given, the *weighted distance* $d_p(\leq +\infty)$ on the weighted graph G is defined by

$$d_p(x, y) = \inf_{\gamma \in \Gamma_{x,y}} L(\gamma)$$

where $\Gamma_{x,y}$ is the set of the paths $\gamma : x_1 = x, x_2, \dots, x_n = y$ from x to y . The length $L(\gamma)$ is computed as the sum of the p -weights for the edges of the path γ :

$$L(\gamma) = \sum_{1 \leq i \leq n} p_{x_i, x_{i+1}} \quad .$$

In particular, if x and y are in distinct connected components of G , $d_p(x, y) = \infty$. We say that the metric space (G, d_p) is *complete* when every Cauchy sequence of vertices has a limit in V .

Definition 1.2 We denote by \hat{V} the metric completion of (G, d_p) and by $V_\infty = \hat{V} \setminus V$ the *metric boundary* of V .

Definition 1.3 If G is a non finite graph and G_0 a finite sub-graph of G , the **ends of G relatively to G_0** are the non finite connected components of $G \setminus G_0$.

2 The Dirichlet problem at infinity

We will use in this section the distance d_p defined using the weights $p_{x,y} = c_{x,y}^{-\frac{1}{2}}$. Let us consider the quadratic form

$$Q(f) = \sum_{\{x,y\} \in E} c_{x,y} (f(x) - f(y))^2 + \sum_{x \in V} \omega_x^2 f(x)^2,$$

which is formally associated to the operator $\Delta_{\omega,c} + \text{Id}$ on l_ω^2 . We will need the following result which is close to lemma 2.5 in [Jo-Pe-2]:

Lemma 2.1 *For any $f : V \rightarrow \mathbb{R}$ so that $Q(f) < \infty$ and for any $a, b \in V$, we have*

$$|f(a) - f(b)| \leq \sqrt{Q(f)} d_p(a, b).$$

Proof.–

For any $\{x, y\} \in E$, $|f(x) - f(y)| \leq \sqrt{Q(f)} / \sqrt{c_{x,y}}$. For any path γ from a to b , defined by the vertices $x_1 = a, x_2, \dots, x_n = b$, we have $|f(a) - f(b)| \leq \sqrt{Q(f)} L(\gamma)$. Taking the infimum of the righthandside with respect to γ we get the result.

□

Remark 2.1 *Lemma 2.1 implies that any function f with $Q(f) < \infty$ extends to \hat{V} as a Lipschitz function \hat{f} . We will denote by f_∞ the restriction of \hat{f} to V_∞ .*

Theorem 2.1 *Let us assume that (V, d_p) is non complete. Let $f : V \rightarrow \mathbb{R}$ with $Q(f) < \infty$, then there exists a continuous function $F : \hat{V} \rightarrow \mathbb{R}$ which satisfies both conditions:*

(i) $(F - f)_\infty = 0$

(ii) $(\Delta_{\omega,c} + 1)(F|_V) = 0$.

Moreover, such an F satisfies $Q(F) < \infty$ and $F \in l_\omega^2$.

If \hat{V} is compact, such an F is unique.

Proof.–

We will denote by A_f the affine space of continuous functions $G : \hat{V} \rightarrow \mathbb{R}$ which satisfy $Q(G) < \infty$ and $(G - f)_\infty = 0$.

Q is lower semi-continuous for the pointwise convergence on V as defined by $Q = \sup Q_\alpha$ with $Q_\alpha(f) = \text{sum of a finite number of terms in } Q$.

Let $Q_0 := \inf_{G \in A_f} Q(G)$ and G_n be a corresponding minimizing sequence. The G_n 's are equicontinuous and pointwise bounded. From

Ascoli's Theorem, this implies the existence of a locally uniformly convergent subsequence $G_{n_k} \rightarrow F$. Using semi-continuity, we have $Q(F) = Q_0$.

If $x \in V$ and δ_x is the Dirac function at the vertex x , we have

$$\frac{d}{dt}\bigg|_{t=0} Q(F + t\delta_x) = 2\omega_x^2[(\Delta_{\omega,c} + 1)F(x)]$$

and this is equal to 0, because F is a minimum of Q restricted to A_f .

Uniqueness is proved using a maximum principle: let us assume that there exists a non zero continuous F with $F_\infty = 0$, then, changing, if necessary, F into $-F$, there exists $x_0 \in V$ with $F(x_0) = \max_{x \in V} F(x) > 0$. The identity (ii) evaluated at the vertex x_0 gives a contradiction. □

3 Not essentially self-adjoint Laplacians

Theorem 3.1 *Let $\Delta_{\omega,c}$ be a weighted graph Laplacian and assume the following conditions:*

- (i) (G, d_p) with $p_{x,y} = c_{x,y}^{-\frac{1}{2}}$ is NON complete,
- (ii) there exists a function $f : V \rightarrow \mathbb{R}$ with $Q(f) < \infty$ and $f_\infty \neq 0$.

Then $\Delta_{\omega,c}$ is not essentially self-adjoint.

Proof.–

Because $\Delta_{\omega,c}$ is ≥ 0 on $C_0(V)$, it is enough (see Theorem X.26 [RS]) to build a non zero function $F : V \rightarrow \mathbb{R}$ which is in $l_\omega^2(V)$ and satisfies

$$(\Delta_{\omega,c} + 1)F = 0 . \tag{2}$$

The function F given by Theorem 2.1 will be the solution of equation (2) the limit of which at infinity is f_∞ . □

Remark 3.1 *The assumptions of Theorem 3.1 are satisfied if (G, d_p) is non complete and $\sum \omega_y^2 < \infty$: it is enough to take $f \equiv 1$.*

They are already satisfied if G has a non complete “end” of finite volume.

Remark 3.2 *Theorem 3.1 is not valid for the Riemannian Laplacian: if X is a closed Riemannian manifold of dimension ≥ 4 , $x_0 \in X$ and $Y = X \setminus x_0$, the Laplace operator on Y is essentially self-adjoint (see [Col1]) and Y has finite volume.*

Question 3.1 *In Theorem 3.1, what is the deficiency index of $\Delta_{\omega,c}$ in terms of the geometry of the weighted graph?*

4 Schrödinger operators for metrically non complete graphs

We now discuss essential selfadjointness for Schrödinger operators of the type $H = \Delta_{\omega,c} + W$ on a graph G in the following setup: we define $\alpha_{x,y} = \min\{\omega_x, \omega_y\}$ and we assume that (G, d_p) , with $p_{x,y} = \frac{\alpha_{x,y}}{\sqrt{c_{x,y}}}$, is non complete as a metric space. It means that there exist Cauchy sequences of vertices without limit in the set V . We will assume that G is of bounded degree, and we denote the upper bound by N .

Definition 4.1 For a vertex $x \in V$, we denote by $D(x)$ the distance to the boundary V_∞ defined by

$$D(x) = \inf_{z \in V_\infty} d_p(x, z).$$

Lemma 4.1 We have, for any edge $\{x, y\}$,

$$|D(x) - D(y)| \leq d_p(x, y) \leq \frac{\min\{\omega_x, \omega_y\}}{\sqrt{c_{x,y}}}. \quad (3)$$

4.1 Agmon-type estimates

Lemma 4.2 Let $v, f \in C_0(V)$ be real valued and assume $Hv = 0$. Then

$$\langle fv, H(fv) \rangle_{l_\omega^2} = \sum_{\{x,y\} \in E} c_{x,y} v(x)v(y) (f(x) - f(y))^2. \quad (4)$$

Proof.–

In the case of positive v this type of formula is known as ground state transform (see [Hea-Kel] and references within). A particular case of this computation (for operators of the type $\Delta_{1,a} + W$) can be found in [To], let us recall the proof for the reader's convenience:

$$\langle fv, H(fv) \rangle_{l_\omega^2} = \sum_{x \in V} f(x)v(x) \left(\sum_{y \sim x} c_{x,y} (f(x) - f(y))v(y) \right)$$

where we used the fact that $Hv(x) = 0$. An edge $\{x, y\}$ contributes to the sum twice. The total contribution is

$$f(x)v(x) c_{x,y} (f(x) - f(y))v(y) + f(y)v(y) c_{y,x} (f(y) - f(x))v(x)$$

so

$$\langle fv, H(fv) \rangle_{l_\omega^2} = \sum_{\{x,y\} \in E} c_{x,y} (f(x) - f(y)) (f(x)v(x)v(y) - f(y)v(y)v(x)).$$

□

Theorem 4.1 *Let v be a solution of $(H - \lambda)v = 0$. Assume that v belongs to $l_\omega^2(V)$ and that there exists a constant $c > 0$ such that, for all $u \in C_0(V)$,*

$$\langle u | (H - \lambda)u \rangle_{l_\omega^2} \geq \frac{N}{2} \sum_{x \in V} \max \left(\frac{1}{D(x)^2}, 1 \right) \omega_x^2 |u(x)|^2 + c \|u\|_{l_\omega^2}^2, \quad (5)$$

then $v \equiv 0$.

Proof.–

This theorem is based on Lemma 4.2 applied to $H - \lambda$. Let us consider ρ and R satisfying $0 < \rho < \frac{1}{2}$ and $1 < R < +\infty$. For any $\varepsilon > 0$, we define the function $f_\varepsilon : V \rightarrow \mathbb{R}$ by $f_\varepsilon = F_\varepsilon(D)$ where D denotes the distance associated to the metric d_p , and $F_\varepsilon : \mathbb{R}^+ \rightarrow \mathbb{R}$ is the continuous piecewise affine function defined by

$$F_\varepsilon(u) = \begin{cases} 0 & \text{for } u \leq \varepsilon \\ \rho(u - \varepsilon)/(\rho - \varepsilon) & \text{for } \varepsilon \leq u \leq \rho \\ u & \text{for } \rho \leq u \leq 1 \\ 1 & \text{for } 1 \leq u \leq R \\ R + 1 - u & \text{for } R \leq u \leq R + 1 \\ 0 & \text{for } u \geq R + 1 \end{cases}$$




Figure 1: The function F

Using Lemma 4.2, Lemma 4.1 and the inequalities

$$v(x)v(y) \leq \frac{1}{2}(v(x)^2 + v(y)^2),$$

the lefthandside of Equation (4) is bounded as follows

$$\langle f_\varepsilon v | (H - \lambda)(f_\varepsilon v) \rangle_{l_\omega^2} \leq \frac{1}{2} \sum_{x \in V} v(x)^2 \omega_x^2 \Phi_\varepsilon(x),$$

with

$$\Phi_\varepsilon(x) = \sum_{y \sim x} \frac{(f_\varepsilon(x) - f_\varepsilon(y))^2}{d_p(x, y)^2} \leq \frac{N\rho^2}{(\rho - \varepsilon)^2}$$

where the second inequality uses the fact that f_ε is $\frac{\rho}{\rho - \varepsilon}$ -Lipshitz.

This implies

$$\langle f_\varepsilon v | (H - \lambda)(f_\varepsilon v) \rangle_{l_\omega^2} \leq \frac{N\rho^2}{2(\rho - \varepsilon)^2} \|v\|_{l_\omega^2}^2. \quad (6)$$

On the other hand, due to assumption (5) the lefthandside of Equation (4) is bounded from below as follows:

$$\langle f_\varepsilon v | (H - \lambda)(f_\varepsilon v) \rangle_{l_\omega^2} \geq \frac{N}{2} \sum_{\rho \leq D(x) \leq R} \omega_x^2 v(x)^2 + c \|f_\varepsilon v\|_{l_\omega^2}^2. \quad (7)$$

Putting together Equations (6) and (7) we get

$$\frac{N}{2} \sum_{\rho \leq D(x) \leq R} \omega_x^2 v(x)^2 + c \|f_\varepsilon v\|_{l_\omega^2}^2 \leq \frac{N\rho^2}{2(\rho - \varepsilon)^2} \|v\|_{l_\omega^2}^2. \quad (8)$$

Then we do $\varepsilon \rightarrow 0$. After that, we do $\rho \rightarrow 0$ and $R \rightarrow \infty$. We get $v \equiv 0$.

□

Remark 4.1 *The previous result is inspired by a nice idea from [Nen], so following the terminology of [Nen] we call Agmon-type estimates Equation (8).*

4.2 Essential self-adjointness

Theorem 4.2 *Consider the Schrödinger operator $H = \Delta_{\omega, c} + W$ on a graph G , define $\alpha_{x, y} = \min\{\omega_x, \omega_y\}$ and assume that (G, d_p) , with $p_{xy} = \frac{\alpha_{x, y}}{\sqrt{c_{x, y}}}$, is non complete as a metric space. For a vertex $x \in V$, we denote by $D(x)$ the distance from x to the boundary V_∞ . We assume the following conditions:*

- (i) G is of bounded degree and we denote the upper bound by N ,
- (ii) there exists $M < \infty$ so that

$$\forall x \in V, W(x) \geq \frac{N}{2D(x)^2} - M. \quad (9)$$

Then the Schrödinger operator H is essentially self-adjoint.

Remark 4.2 *In the particular case when $\sum \omega_x^2 < \infty$, the Laplacian $H = \Delta_{\omega,c}$ does not satisfy the assumption (9) so this result is coherent with Theorem 3.1.*

Remark 4.3 *The exponent of $D(x)$ in (9) is sharp. In fact, one can find a potential W such that $W(x) \geq \frac{k}{D(x)^2}$ where $k < \frac{N}{2}$ and weights ω and c such that $H = \Delta_{\omega,c} + W$ is non essentially self-adjoint. See Example 5.3.2 .*

Remark 4.4 *In the case where $\omega \equiv 1$ the result is an immediate consequence of [Ke-Le-1] (Theorem 5).*

Proof.–

We have, for any $u \in C_0(V)$

$$\langle u | Hu \rangle_{l_\omega^2} \geq \sum_{x \in V} W(x) \omega_x^2 |u(x)|^2,$$

so using assumption (9) we get:

$$\langle u | (H - \lambda)u \rangle_{l_\omega^2} - \frac{N}{2} \sum_{x \in V} \frac{1}{D(x)^2} \omega_x^2 |u(x)|^2 \geq \sum_{x \in V} -(M + \lambda) |u(x)|^2.$$

Then choosing for example

$$\lambda = -M - 1$$

we get the inequality (5) with $c = 1$, and the proof follows from Theorem 4.1.

□

5 Schrödinger operators on “star-like” graphs

5.1 Introduction

Definition 5.1 *The graph \mathbb{N} is the graph defined by $V = \{0, 1, 2, \dots\}$ and $E = \{\{n, n + 1\} \mid n = 0, 1, \dots\}$.*

Definition 5.2 *We will call an infinite graph $G = (V, E)$ star-like if there exists a finite sub-graph G_0 of G so that $G \setminus G_0$ is the union of a finite number of disjoint copies G_α of the graph \mathbb{N} (the ends of G relatively to G_0 according to Definition 1.3).*

For example, the graph \mathbb{Z} , defined similarly to \mathbb{N} , is star-like.

Let us consider a Laplace operator $L = \Delta_{1,a}$ on G . On each end G_α of G , L will be given by

$$L^\alpha f_n = -a_{n,n+1}^\alpha f_{n+1} + (a_{n-1,n}^\alpha + a_{n,n+1}^\alpha) f_n - a_{n-1,n}^\alpha f_{n-1} ,$$

where the $a_{n-1,n}^\alpha$'s are > 0 . If $W : V \rightarrow \mathbb{R}$, we will consider Schrödinger operators H on $C_0(G)$ defined by $H = \Delta_{1,a} + W$.

Lemma 5.1 *Let G_0 be a finite sub-graph of G . The operator $H = \Delta_{1,a} + W$ on G is essentially self-adjoint if and only if it is essentially self-adjoint on each end of G relatively to G_0 . More precisely, the deficiency indices n_\pm are the sum of the corresponding deficiency indices of the ends.*

We will need the following Lemma which is a consequence of Kato-Rellich Theorem, see [Go-Sch], Proposition 2.1:

Lemma 5.2 *If A and B are 2 symmetric operators with the same domains and $R = B - A$ is bounded, then the deficiency indices of A and B are the same.*

Proof.–

We give here an alternative proof to this result. Let us define, for $t \in \mathbb{R}$, $A_t = A + tR$ so that $A_0 = A$ and $A_1 = B$. The domains of the closures of the A_t 's coincide: the “graph-norms” $\|A_t u\|_{l^2} + \|u\|_{l^2}$ are equivalent. The domains of the adjoints coincide too. Let $K = D(A^*)/D(\bar{A})$ and $Q_t(u, v) = -i(\langle A_t^* u | v \rangle - \langle u | (A_t^* v \rangle)$ which is well defined on K . We know that these bounded Hermitian forms are non degenerate on K with the graph norm and continuous w.r. to t . Hence the Morse index $n_-(t)$ is locally constant: take a decomposition $K = K_+ \oplus K_-$ where $q = Q_{t_0}$ satisfies $q|_{K_+} \geq C > 0$ and $q|_{K_-} \leq -C < 0$.

□

Using Lemma 5.2, we can prove Lemma 5.1:

Proof.–

We will consider the operator H_{red} where we replace the entries $a_{x,y}$ of H with $\{x, y\} \in E(G_0)$ by 0. The claim of the Lemma is clear for H_{red} because it is the direct orthogonal sum of the Schrödinger operators of the ends and a finite rank l^2 -bounded matrix. We can then use Lemma 5.2 because $H - H_{\text{red}}$ is bounded.

□

Remark 5.1 *It follows from Lemma 5.1 that, concerning essential self-adjointness questions for star-like graphs, it is enough to work on the graph \mathbb{N} . We have*

$$(Hf)_0 = -a_{0,1}f_1 + a_{0,1}f_0 + W_0f_0.$$

This implies that the space of solutions of $(H - \lambda)u = 0$ on \mathbb{N} is of dimension 1 and any solution so that f_0 vanishes is $\equiv 0$. We will consider also solutions “near infinity”, i.e. $(f_n)_{n \geq 0}$ satisfies $((H - \lambda)f)_n = 0$ for $n \geq 1$; this space is of dimension 2.

5.2 Main result

It is known ([Dod]) that $H = \Delta_{1,a} + W$ is essentially self-adjoint provided $\Delta_{1,a}$ is bounded as an operator on $l^2(G)$ and W bounded from below. For star-like graphs, we have the following result, which holds for *any* potential W :

Theorem 5.1 *If G is star-like and if for each end G_α ,*

$$1/a_{n-1,n}^\alpha \notin l^1(\mathbb{N}) \quad (10)$$

then $H = \Delta_{1,a} + W$ with domain $C_0(V)$ is essentially self-adjoint for any potential W .

Remark 5.2 *The condition (10) is sufficient but not necessary. See Example 5.3.2 .*

Proof.–

Due to Remark 5.1 we only have to prove the following

Theorem 5.2 *If*

$$\frac{1}{a_{n-1,n}} \notin l^1(\mathbb{N}) , \quad (11)$$

the Schrödinger operator $H = \Delta_{1,a} + W$ with domain $C_0(\mathbb{N})$ is essentially self-adjoint for any potential W .

This result is contained in the book [Ber] (p. 504). We propose here a short proof, obtained by contradiction using Corollary 6.1 which is an analog of Weyl’s limit point-limit circle criteria in the discrete case.

Let us consider an operator $\Delta_{1,a}$ such that (10) is fulfilled. We assume that any sequence u , such that $(H - i)u = 0$ near infinity, is in $l^2(\mathbb{N})$. In particular, there exists a basis f, g of solutions of $(H - i)f = 0$ with $f \in l^2(\mathbb{N})$ and $g \notin l^2(\mathbb{N})$.

We have

$$-a_{n,n+1} f_{n+1} + (a_{n-1,n} + a_{n,n+1} + (W_n - i)) f_n - a_{n-1,n} f_{n-1} = 0 ,$$

and the same holds for g . The Wronskian of f and g is the sequence $\mathcal{W}_n = f_n g_{n-1} - f_{n-1} g_n$. We have, for any $n \in \mathbb{N}$:

$$\mathcal{W}_{n+1} = \frac{a_{n-1,n}}{a_{n,n+1}} \mathcal{W}_n ,$$

which implies

$$\mathcal{W}_n = \frac{a_{0,1}}{a_{n-1,n}} \mathcal{W}_1 .$$

But since the Wronskian is in $l^1(\mathbb{N})$ according to the assumption that f and g are in $l^2(\mathbb{N})$, we get a contradiction with the hypothesis (10).

□

5.3 Examples of Schrödinger operators

5.3.1 Example 1

Let us consider the Laplacian $\Delta_{\omega,c}$ on \mathbb{N} , with, $\forall n > 0$, $c_{n-1,n} = n^3$ and, $\forall n \geq 0$, $\omega_n = \frac{1}{n+1}$. Since $\sum \omega_n^2 < \infty$ and $\sum c_{n-1,n}^{-1/2} < \infty$ we deduce from Theorem 3.1 (due to Rem 3.1) that $\Delta_{\omega,c}$ is not essentially self-adjoint.

Applying a result of [To] (see Proposition 7.1 in Appendix B) we get that this Laplacian is unitarily equivalent to the Schrödinger operator $H = \Delta_{1,a} + W$ with $a_{n-1,n} = \frac{c_{n-1,n}}{\omega_{n-1}\omega_n} \sim n^5$ and

$$W_n = \frac{1}{\omega_n} \left[c_{n,n+1} \left(\frac{1}{\omega_n} - \frac{1}{\omega_{n+1}} \right) + c_{n-1,n} \left(\frac{1}{\omega_n} - \frac{1}{\omega_{n-1}} \right) \right] \sim -3n^3 ,$$

which is therefore not essentially self-adjoint.

According to Theorem 5.1, such an operator must verify $\frac{1}{a_{n-1,n}} \in l^1(\mathbb{N})$, which is indeed the case.

5.3.2 Example 2: Discretization of a Schrödinger operator on \mathbb{R}^+

Let us consider the Schrödinger operator on $]0, +\infty[$ defined on smooth compactly supported functions by $Lf := -f'' + \frac{A}{x^2}f$. This operator is essentially self-adjoint if and only if $A > 3/4$ (see [RS] theorem X 10). We discretize this operator in the following way: let us consider the graph $\Gamma = (V, E)$ resulting of the following dyadic subdivision of the interval $(0, 1)$: the vertices are the $x_n = 2^{-n}$ and the edges are the pairs $\{2^{-n}, 2^{-n+1}\}$ which correspond to the intervals $[2^{-n}, 2^{-n+1}]$ of length $\omega_n^2 = 2^{-n}$.

Then we define, for any

$$f \in l_\omega^2(V) = \left\{ f \in C(V) \mid \sum_{n \in \mathbb{N}} 2^{-n} f_n^2 < +\infty \right\}$$

where we set $f = (f_n)$, the quadratic form

$$Q(f) = \sum_{n \in \mathbb{N}} 2^{-n} \left[\left(\frac{f_{n+1} - f_n}{2^{-n}} \right)^2 + A 2^{2n} f_n^2 \right].$$

According to the previous definitions and if we set $c_{n,n+1} = 2^n$, this quadratic form is associated to the Schrödinger operator $H = \Delta_{\omega,c} + W$ on \mathbb{N} with the potential $W_n := A 2^{2n}$.

Let us set $a_{n,n+1} = \frac{c_{n,n+1}}{\omega_n \omega_{n+1}} = 2^{2n+\frac{1}{2}}$. Applying Proposition 7.1 we get that H is unitarily equivalent to

$$\widehat{H} = \Delta_{1,a} + \widehat{W} + W$$

with

$$\widehat{W}_n = \frac{1}{\omega_n} \left[c_{n,n+1} \left(\frac{1}{\omega_n} - \frac{1}{\omega_{n+1}} \right) + c_{n-1,n} \left(\frac{1}{\omega_n} - \frac{1}{\omega_{n-1}} \right) \right] = 2^{2n} \left(\frac{3}{2} - \frac{5\sqrt{2}}{4} \right).$$

We have $\widehat{H} = \Delta_{1,a} + (A - A_0)4^n$ with $A_0 = \frac{5\sqrt{2}}{4} - \frac{3}{2} (> 0)$. The metric graph (\mathbb{N}, d_p) with $p_{n,n+1} = a_{n,n+1}^{-1/2}$ is non complete. The solutions u of $Hu = 0$ verify

$$4u_{n+1} - \left(5 + 2\sqrt{2}(A - A_0) \right) u_n + u_{n-1} = 0.$$

The solutions are generated by α_1^n and α_2^n where α_1 and α_2 are the roots of

$$4\alpha^2 - \left(5 + 2\sqrt{2}(A - A_0) \right) \alpha + 1 = 0.$$

We have $|\alpha_1| < 1$ and $|\alpha_2| < 1$ if and only if $A_0 - \frac{5}{\sqrt{2}} < A < A_0$.

Using Proposition 6.1, with $d = 2$ and $U_n = \begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix}$, we get, for any $\lambda \in \mathbb{C}$, the exponential decay of all solutions near infinity of $(\widehat{H} - \lambda)u = 0$ if $A_0 - \frac{5}{\sqrt{2}} < A < A_0$, and the existence of a solution of $(\widehat{H} - \lambda)u = 0$ with

exponential growth in the case when $A > \frac{5\sqrt{2}}{4} - \frac{3}{2}$ or $A < -\frac{5\sqrt{2}}{4} - \frac{3}{2}$.

Hence (by Corollary 6.1) we get the following result:

Proposition 5.1 1. If $-\frac{5\sqrt{2}}{4} - \frac{3}{2} < A < \frac{5\sqrt{2}}{4} - \frac{3}{2}$, then the discretized operator H is not essentially self-adjoint.

2. If $A > \frac{5\sqrt{2}}{4} - \frac{3}{2}$ (*) or $A < -\frac{5\sqrt{2}}{4} - \frac{3}{2}$, then H is essentially self-adjoint.

From this result we can deduce several informations:

1. The condition (*) is analogous to the condition $A > 3/4$ in the continuous case.
2. Proposition 5.1 implies that for $A = 0$ the operator $H = \Delta_{\omega,c}$ is not essentially self-adjoint, which is a result predicted by Theorem 3.1.
3. This gives examples of essentially self-adjoint operators with $1/a_n \in l^1$.
4. Sharpness of the assumption (9) in Theorem 4.2

In this context, the distance d_p is associated to

$$p_{x,y} = \frac{\alpha_{x,y}}{\sqrt{c_{x,y}}}$$

with $\alpha_{x,y} = \min\{\omega_x, \omega_y\}$ so we get

$$D(n) = \sum_{p \geq n} \frac{\alpha_{p,p+1}}{\sqrt{c_{p,p+1}}} = \sum_{p \geq n} \left(\frac{2^{-p-1}}{2^p} \right)^{1/2} = 2^{-\frac{1}{2}} 2^{-n} 2$$

so

$$\frac{1}{D(n)^2} = 2^{2n-1} .$$

If the operator $H = \Delta_{\omega,c} + A4^n$ satisfies the assumption (9), then

$A > \frac{1}{2}$ which involves condition (*), since $\frac{1}{2} > \frac{5\sqrt{2}}{4} - \frac{3}{2}$, so Theorem 4.2 is coherent with proposition 5.1. Moreover the operator $H = \Delta_{\omega,c} + A4^n$ with $A = \frac{5\sqrt{2}}{4} - \frac{3}{2}$ is not essentially self-adjoint, which implies that the estimate (9) on the growth of the potential in Theorem 4.2 is sharp.

5.3.3 Example 3

Let us consider the Laplacian $\Delta_{\omega,c}$ on \mathbb{N} , where the coefficients verify $c_{n-1,n} = n^\gamma$ with $\gamma > 2$ and $\omega_n = (n+1)^{-\beta}$ with $\beta > \frac{1}{2}$. Since $\sum \omega_n^2 < \infty$ and $\sum c_{n-1,n}^{-1/2} < \infty$ we deduce from Theorem 3.1 (due to Remark 3.1) that $\Delta_{\omega,c}$ is not essentially self-adjoint.

Applying one more time Proposition 7.1, we see that this operator is unitarily equivalent to the Schrödinger operator $H = \Delta_{1,a} + W$, with $a_{n-1,n} \sim n^{\gamma+2\beta}$ and the potential $W_n \sim -\beta(\beta + \gamma - 1)n^{2\beta+\gamma-2}$, which is therefore also not essentially self-adjoint. We emphasize that W is not bounded from below, which is predicted in [To], Theorem 3.2.

Furthermore, according to Theorem 5.1, such an operator must verify the condition $\frac{1}{a_{n-1,n}} \in l^1(\mathbb{N})$, which is indeed the case. Following the terminology of the previous sections, it means the non completeness of (\mathbb{N}, d_p) with the weights $p_{n-1,n} = a_{n-1,n}^{-1/2}$.

5.3.4 Example 4

Let us consider the Laplacian $H = \Delta_{\omega,c}$ on a spherically homogeneous rooted tree $G = (V, E)$ (see [Bre] and references within). For any vertex x , we denote by $\delta(x)$ the distance from x to the root 0 and define $\omega_x = 2^{-\delta(x)}$, and $c_{x,y} = 2^{\delta(x)}$, for any $y \sim x$ so that $\delta(y) = n + 1$. We assume that the graph G has a uniform degree $N + 1$.

Let us set $a_{x,y} = \frac{c_{x,y}}{\omega_x \omega_y}$. We have $a_{x,y} = 2^{3n+1}$ for any edge x, y , so that $\delta(x) = n$ and $\delta(y) = n + 1$. Then, due to Proposition 7.1, the operator H is unitarily equivalent to

$$\widehat{H} = \Delta_{1,a} + W$$

with

$$W(x) = 2^{3n} \left(-N + \frac{1}{4} \right)$$

for any x such that $\delta(x) = n$.

The radial solutions u of $Hu = 0$ can be seen as sequences (u_n) which satisfy the equation:

$$-2Nu_{n+1} + \left(N + \frac{1}{2} \right) u_n - \frac{1}{4}u_{n-1} = 0 .$$

The solutions are generated by α_1^n and α_2^n where α_1 and α_2 are the roots of

$$\alpha^2 - \left(\frac{1}{2} - \frac{1}{4N} \right) \alpha + \frac{1}{8N} = 0 .$$

We have $|\alpha_1| < 1$ and $|\alpha_2| < 1$ for any $N > 0$.

The radial solutions of $(\widehat{H} - \lambda)u = 0$ satisfy

$$-2Nu_{n+1} + \left(N + \frac{1}{2} \right) u_n - \frac{1}{4}u_{n-1} = 2N\lambda 2^{-(3n+1)}u_n .$$

Using Proposition 6.1, with $d = 2$ and $U_n = \begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix}$, we get the exponential decay of all solutions near infinity of $(\widehat{H} - \lambda)u = 0$.

Hence (by Corollary 6.1) we get the following result:

Proposition 5.2 *For any $N \geq 1$ H is not essentially self-adjoint .*

Remark 5.3 *We have*

$$\sum_x \omega_x^2 = \sum_n \omega_n^2 N^n = \sum_n \left(\frac{N}{4}\right)^n .$$

If $N < 4$, then $\sum_x \omega_x^2 < \infty$ so Theorem 3.1 can also be applied to get the result since the graph is non complete with respect to the metric d_p , with $p_{x,y} = c_{x,y}^{-\frac{1}{2}}$.

6 Appendix A: Weyl’s “limit point-limit circle” criteria

6.1 The discrete case

The goal of this section is to prove the discrete version of the Weyl’s “limit point-limit circle” criterium. Our presentation is simpler than the classical presentation for the continuous case (see [RS], Appendix to section X.1).

Let us consider the Hilbert space $\mathcal{H} := l^2(\mathbb{N}, \mathbb{C}^N)$ and the formally symmetric differential operator P defined by

$$Pf(0) = P_{0,0}f(0) + P_{0,1}f(1), \forall l \geq 1, Pf(l) = P_{l,l-1}f(l-1) + P_{l,l}f(l) + P_{l,l+1}f(l+1)$$

where

1. $\forall l \geq 1, P_{l-1,l}^* = P_{l,l-1}$
2. $\forall l \geq 0, P_{l,l}^* = P_{l,l}$
3. $\forall l \geq 0, P_{l,l+1}$ is invertible
4. $\exists M \in \mathbb{R}$ so that for any $f \in C_0(\mathbb{N}, \mathbb{C}^N)$, $Q_P(f) = \langle Pf | f \rangle \geq -M \|f\|^2$.

Let us define the subspace \mathcal{E} of \mathcal{H} as the set of l^2 sequences f so that, for all $l \geq 1$, $(P - i)f(l) = 0$; the space \mathcal{E} is isomorphic to the space of germs at infinity of l^2 solutions of $(P - i)f = 0$. Assumption 3. implies that $\dim \mathcal{E} \leq 2N$. Let us denote by $\mathcal{K} = \ker(P - i) \cap l^2$ and consider the following sequence

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{E} \rightarrow \mathbb{C}^N \rightarrow 0, \quad (12)$$

where the non trivial arrow is given by $f \rightarrow (P - i)f(0)$. We have the

Theorem 6.1 *The sequence (12) is exact and the deficiency indices $n_{\pm} = \dim \mathcal{K}$ of P are given by $n_{\pm} = \dim \mathcal{E} - N$.*

Proof.–

Assumption 4. implies (using Corollary of Theorem X.1 in [RS]) that the deficiency indices are equal. The only non trivial point is to prove that the arrow $p : \mathcal{E} \rightarrow \mathbb{C}^N$ is surjective. Let us consider \tilde{P} a self-adjoint extension of P which exists because $n_+ = n_-$. Let us consider the map $\rho : \mathbb{C}^N \rightarrow \mathcal{E}$ defined by $\rho(x) = (\tilde{P} - i)^{-1}(x, 0, 0, \dots)$. Then $p \circ \rho = \text{Id}_{\mathbb{C}^N}$.

□

Corollary 6.1 *The Schrödinger operator $H = \Delta_{1,a} + W$ defined on $C_0(\mathbb{N})$ is essentially self-adjoint if and only if there exists a sequence u such that $(H - i)u = 0$ near infinity (i.e. $((H - i)u)_n = 0$ for n large enough) which is not in $l^2(\mathbb{N})$.*

6.2 Asymptotic behavior of perturbed hyperbolic iterations

In order to apply Corollary 6.1, the following results will be useful

Proposition 6.1 *Let us consider the following linear dynamical system on \mathbb{C}^d :*

$$\forall n \geq 0, U_{n+1} = AU_n + R(n)U_n \quad (13)$$

where

1. *A is hyperbolic: all eigenvalues λ_j of A satisfy $|\lambda_j| \neq 1$*
2. *$R(n) \rightarrow 0$ as $n \rightarrow \infty$.*

Then

- **case A:** *If all eigenvalues λ_j of A satisfy $|\lambda_j| < 1$, all solutions (U_n) of Equation (13) are exponentially decaying.*
- **case B:** *If m eigenvalues satisfy $|\lambda_j| > 1$, then there exists an m -dimensional vector space F of solutions of Equation (13) whose non-zero vectors have exponential growth.*

Proof.–

Case A: There exists a norm $\|\cdot\|$ on \mathbb{C}^d so that the operator norm of A satisfies $\|A\| = k < 1$. For n large enough, we have $\|A + R(n)\| \leq k' < 1$. The conclusion follows.

Case B: There exists a splitting $\mathbb{C}^d = Y \oplus Z$, denoted $x = y + z$, with $\dim Y = m$, stable by A , norms on Y and Z and 2 constants $\mu < 1 < \sigma$, so that

$$\begin{aligned} \forall y \in Y, \|Ay\| &\geq \sigma \|y\| , \\ \forall z \in Z, \|Az\| &\leq \mu \|z\| . \end{aligned}$$

Let us choose $\varepsilon > 0$ so that $1 < \sigma - 2\varepsilon$ and N so that $\|R(n)\| \leq \varepsilon$ for $n \geq N$. We have, for $n \geq N$,

$$\|y_{n+1}\| \geq \sigma \|y_n\| - \varepsilon(\|y_n\| + \|z_n\|), \|z_{n+1}\| \leq \mu \|z_n\| + \varepsilon(\|y_n\| + \|z_n\|) ,$$

so that

$$\|y_{n+1}\| - \|z_{n+1}\| \geq (\sigma - 2\varepsilon)(\|y_n\| - \|z_n\|) .$$

Any solution which satisfies $\|y_N\| > \|z_N\|$ will have exponential growth. Take for F the space of solutions for which $z_N = 0$.

□

6.3 The continuous case

A similar method works for the continuous case. Let $H = -\frac{d^2}{dx^2} + A(x)$ be a system of differential operators where $A(x)$ is Hermitian for every x and is continuous on $[0, a[$ as a function of x . The differential operator H is L^2 -symmetric on the Dirichlet domain

$$D = C_0^\infty([0, a[, \mathbb{C}^N) \cap \{u \mid u(0) = 0\} .$$

We denote H_D the closure of (H, D) . Let us assume that $n_+(H_D) = n_-(H_D)$ which is true for example if A is bounded from below or if A is real-valued. Then

Theorem 6.2 *If \mathcal{E} is the space of solutions u of the differential equation $(H - i)u = 0$ which are L^2 near a , then $n_\pm(H_D) = \dim \mathcal{E} - N$.*

Proof.–

Let us consider the sequence

$$0 \rightarrow \ker(H_D - i) \rightarrow \mathcal{E} \rightarrow \mathbb{C}^N \rightarrow 0 , \quad (14)$$

where the only non trivial arrow is given by $u \rightarrow u(0)$. This sequence is exact: we have only to prove the surjectivity of the non trivial arrow. Let \tilde{H} be a self-adjoint extension of H_D and $\chi \in C_0^\infty([0, a[, \mathbb{R})$ with $\chi(0) = 1$. For any $X \in \mathbb{C}^N$, let us consider

$$u = \chi X - (\tilde{H} - i)^{-1}((H - i)(\chi X)) .$$

Then $(H - i)u = 0$, $u(0) = X$ and u is L^2 near a .

□

7 Appendix B: Unitary equivalence between Laplacians and Schrödinger operators

In this section, we recall the following results (see [To] Proposition 2.1 and Theorem 5.1): the first one states that a Laplacian is always unitarily equivalent to a Schrödinger operator, and the second result asserts that a Schrödinger operator with a strictly positive quadratic form is unitarily equivalent to a Laplacian.

For a weighted graph G by the weight ω on its vertices, let

$$U_\omega : l_\omega^2(V) \longrightarrow l^2(V)$$

the unitary operator defined by

$$U_\omega(f) = \omega f .$$

This operator preserves the set of functions on V with finite support.

Proposition 7.1 *The operator*

$$\widehat{\Delta} = U_\omega \Delta_{\omega,c} U_\omega^{-1} ,$$

is a Schrödinger operator on G . More precisely:

$$\widehat{\Delta} = \Delta_{1,a} + W$$

where a is a strictly positive weight on E given by:

$$a_{x,y} = \frac{c_{x,y}}{\omega_x \omega_y}$$

and the potential $W : V \longrightarrow \mathbb{R}$ is given by:

$$W = -\frac{1}{\omega} \Delta_{1,a} \omega .$$

The following Theorem uses the existence of a strictly positive harmonic function (see [To], section 4).

Theorem 7.1 *Let P a Schrödinger operator on a graph G . We assume that $\langle Pf, f \rangle_{l^2} > 0$ for any function f in $C_0(V) \setminus \{0\}$. Then there exist weights: ω on V and c on E such that P is unitarily equivalent to the Laplacian $\Delta_{\omega,c}$ on the graph G .*

Acknowledgments

The second author is greatly indebted to the research unity "Mathématiques et Applications" (05/UR/15-02) of Faculté des Sciences de Bizerte (Tunisie) for the financial support, and would like to present special thanks to Institut Fourier where this work was carried on.

Thanks to D. Lenz for giving notes on some references.

All the authors would like to thank the anonymous referee for careful reading, numerous remarks, useful suggestions and valuable references.

References

- [Ber] Ju.M. Berezans'kii. *Expansions in eigenfunctions of selfadjoint operators*. Translations of Mathematical Monographs **17**, American Mathematical Society, Providence, R.I. (1968).
- [B-M-S] M. Braverman, O. Milatovic & M. Shubin. *Essential self-adjointness of Schrödinger-type operators on manifolds*, Russian Math. Surveys **57** (2002) 641-692 .
- [Bre] J. Breuer. *Singular continuous spectrum for the Laplacian on certain sparse trees*, Commun. Math. Phys. **269** (3) (2007) 851-857.
- [Col1] Y. Colin de Verdière. *Pseudos-Laplaciens I*, Ann. Inst. Fourier (Grenoble), **32** (1982) 275-286 .
- [Col2] Y. Colin de Verdière. *Spectre de graphes*, Cours spécialisés **4**, Société mathématique de France (1998).
- [Col-Tr] Y. Colin de Verdière & F. Truc. *Confining quantum particles with a purely magnetic field*, Ann. Inst. Fourier (Grenoble) (to appear).
- [Dod] J. Dodziuk. *Elliptic operators on infinite graphs*, Analysis geometry and topology of elliptic operators, 353-368, World Sc. Publ., Hackensack NJ. (2006).
- [Du-Sc] N. Dunford & J. T. Schwartz. *Linear operator II, Spectral Theory*, John Wiley & Sons, New York (1971).
- [Go-Sch] S. Golénia & C. Schumacher. *The problem of deficiency indices for discrete Schrödinger operators on locally finite graphs*, [arXiv:1005.0165](https://arxiv.org/abs/1005.0165) (2010).
- [Hea-Kel] S. Haeseler & M. Keller. *Generalised solutions and spectrum for Dirichlet forms on graphs*, [arXiv:1002.1040](https://arxiv.org/abs/1002.1040) (2010).

- [Hu] X. Huang. *A note on stochastic incompleteness for graphs and weak Omori-Yau maximum principle*, [arXiv:1009.2579](#) (2010).
- [Jor] P.E.T. Jorgensen. *Essential self-adjointness of the graph-Laplacian*, J. Math. Phys. **49** (7) (2008) 073510, 33pp.
- [Jo-Pe-1] P.E.T. Jorgensen & E.P.J. Pearse. *Spectral reciprocity and matrix representations of unbounded operators*, [arXiv:0911.0185](#) (2009)
- [Jo-Pe-2] P.E.T. Jorgensen & E.P.J. Pearse. *A discrete Gauss-Green identity for unbounded Laplace operators, and the transience of Random walks*, [arXiv:0906.1586](#) (2010).
- [Ke-Le-1] M. Keller & D. Lenz. *Dirichlet forms and stochastic completeness of graphs and subgraphs*, [arXiv:0904.2985](#) (2009).
- [Ke-Le-2] M. Keller & D. Lenz. *Unbounded Laplacians on graphs: basic spectral properties and the heat equation*, Math. Nat. Phenomena, **5** (2010), n 4.
- [Nen] G. Nenciu & I. Nenciu. *On confining potentials and essential self-adjointness for Schrödinger operators on bounded domains in \mathbb{R}^n* , Ann. Henri Poincaré, **10** (2009), 377-394.
- [Ma] J. Masamune. *A Liouville property and its application to the Laplacian of an infinite graph*, Contemp. Math., **484** (2009), 103-115,
- [Ol] I.M. Oleinik. *On the essential self-adjointness of the operator on complete Riemannian manifolds*, Mathematical Notes **54** (1993), 934-939 .
- [RS] M.Reed & B.Simon. *Methods of Modern mathematical Physics, I-Functional analysis*,(1980), II-*Fourier analysis, Self-adjointness*, (1975). New York Academic Press.
- [Shu1] M. Shubin. *The essential self-adjointness for semi-bounded magnetic Schrödinger operators on non-compact manifolds*, J. Func. Anal. **186** (2001), 92-116.
- [Shu2] M. Shubin. *Classical and quantum completeness for the Schrödinger operators on non-compact manifolds*, Geometric Aspects of Partial Differential Equations (Proc. Sympos., Roskilde, Denmark (1998)) Amer. Math. Soc. Providence, RI, 257-269 (1999).
- [To] N. Toriki-Hamza. *Laplaciens de graphes infinis I Graphes métriquement complets*, Confluentes Mathematici **2** (2010), n 3.
- [We] A. Weber. *Analysis of the physical Laplacian and the heat flow on a locally finite graph*, J. Math. Anal. Appl. **370** (2010), 146-158.

- [Woj1] R.K. Wojciechowski. *Stochastic completeness of graphs*, Ph.D. Thesis, The graduate Center of the University of New-York (2008).
- [Woj2] R.K. Wojciechowski. *Heat kernel and essential spectrum of infinite graphs*, Univ. Math. J. **58** (2009), n 3, 1419-1442.