



HAL
open science

Realizability algebras II: new models of ZF + DC

Jean-Louis Krivine

► **To cite this version:**

Jean-Louis Krivine. Realizability algebras II: new models of ZF + DC. Logical Methods in Computer Science, 2012, 8 (1), pp.10. 10.2168/LMCS-8(1:10)2012 . hal-00497587

HAL Id: hal-00497587

<https://hal.science/hal-00497587>

Submitted on 5 Jul 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Realizability algebras II : new models of ZF + DC

Jean-Louis Krivine

July 1, 2010

Introduction

The technology of *classical realizability* was developed in [13, 16] in order to extend the proof-program correspondence (*Curry-Howard correspondence*) to the whole of mathematical proofs, with excluded middle, axioms of ZF, dependent choice, existence of a well ordering on $\mathcal{P}(\mathbb{N})$, ...

We show here that this technology is also a new method in order to build models of ZF and to obtain relative consistency results.

The main tools are :

- The structure of *standard realizability algebra* [16], which plays a role similar to a set of forcing conditions.
- The theory ZF_ε [11] which is a conservative extension of ZF, with a notion of *strong membership*, denoted as ε .

The theory ZF_ε is essentially ZF without the extensionality axiom. We note an analogy with the Fraenkel-Mostowski models with “urelements” : we obtain a non well orderable set, which is a Boolean algebra denoted $\mathfrak{J}2$, all elements of which except the unity are empty. But we also notice two important differences :

- The final model of $\text{ZF} + \neg \text{AC}$ is obtained directly, without taking a suitable submodel.
- There exists an injection from the “pathological set” $\mathfrak{J}2$ into \mathbb{R} , and therefore \mathbb{R} is *also not well orderable*.

We show the consistency, relatively to the consistency of ZF, of the theory $\text{ZF} + \text{DC}$ (dependent choice) with the following properties :

- there exists a sequence \mathcal{X}_n of infinite subsets of \mathbb{R} , the “cardinals” of which are strictly decreasing (this means that there is an injection but no surjection from \mathcal{X}_{n+1} to \mathcal{X}_n) ;
- there exists a sequence \mathcal{X}_n of infinite subsets of \mathbb{R} , the “cardinals” of which are strictly increasing, and such that $\mathcal{X}_m \times \mathcal{X}_n$ is equipotent with \mathcal{X}_{mn} .

More detailed properties of \mathbb{R} in this model are given in theorems 35 and 39.

As far as I know, these consistency results are new, and cannot be obtained by forcing. But, in any case, the fact that the simplest non trivial realizability model has a real line with so unusual properties, is of interest in itself. Another aspect of these results, which

is interesting from the point of view of computer science, is the following : in [16], we introduce *read* and *write* instructions in a global memory, in order to realize a form of the axiom of choice (well ordering of \mathbb{R}). Therefore, what we show here, is that these instructions are *indispensable* : without them, we can build a realizability model in which \mathbb{R} is not well ordered.

Standard realizability algebras

The notion of *realizability algebra*, and the particular case of *standard realizability algebra* are defined in [16]. They are variants of the usual notion of *combinatory algebra*. Here, we only need the *standard* realizability algebras, the definition of which we recall below :

We have a countable set Π_0 which is the set of *stack constants*.

We define recursively two sets : Λ (the set of *terms*) and Π (the set of *stacks*). Terms and stacks are finite sequences of elements of the set :

$$\Pi_0 \cup \{B, C, E, I, K, W, \mathbf{cc}, \zeta, \mathbf{k}, (,), [,], \bullet\}$$

which are obtained by the following rules :

- $B, C, E, I, K, W, \mathbf{cc}, \zeta$ are terms ;
- each element of Π_0 is a stack (*empty stacks*) ;
- if ξ, η are terms, then $(\xi)\eta$ is a term (this operation is called *application*) ;
- if ξ is a term and π a stack, then $\xi \bullet \pi$ is a stack (this operation is called *push*) ;
- if π is a stack, then $\mathbf{k}[\pi]$ is a term.

A term of the form $\mathbf{k}[\pi]$ is called a *continuation*. It will also be denoted as \mathbf{k}_π .

A term which does not contain any continuation (i.e. in which the symbol \mathbf{k} does not appear) is called *proof-like*.

Every stack has the form $\pi = \xi_1 \bullet \dots \bullet \xi_n \bullet \pi_0$, where $\xi_1, \dots, \xi_n \in \Lambda$ and $\pi_0 \in \Pi_0$, i.e. π_0 is a stack constant.

If $\xi \in \Lambda$ and $\pi \in \Pi$, the ordered pair (ξ, π) is called a *process* and denoted as $\xi \star \pi$; ξ and π are called respectively the *head* and the *stack* of the process $\xi \star \pi$.

The set of processes $\Lambda \times \Pi$ will also be written $\Lambda \star \Pi$.

Notation. The term $(\dots(((\xi)\eta_1)\eta_2)\dots)\eta_n$ will be also denoted by $(\xi)\eta_1\eta_2\dots\eta_n$ or $\xi\eta_1\eta_2\dots\eta_n$.

For example : $\xi\eta\zeta = (\xi)\eta\zeta = (\xi\eta)\zeta = ((\xi)\eta)\zeta$.

We now choose a recursive bijection from Λ onto \mathbb{N} , which is written $\xi \longmapsto \mathbf{n}_\xi$.

We put $0 = \lambda f \lambda x x$, $\sigma = \lambda n \lambda f \lambda x (f)(n)fx$ (the successor in λ -calculus). For each $n \in \mathbb{N}$, we define $\underline{n} \in \Lambda$ recursively, by putting : $\underline{0} = 0$; $\underline{n+1} = (\sigma)\underline{n}$.

We define a preorder relation \succ , on $\Lambda \star \Pi$. It is the least reflexive and transitive relation such that, for all $\xi, \eta, \zeta \in \Lambda$ and $\pi, \varpi \in \Pi$, we have :

- $(\xi)\eta \star \pi \succ \xi \star \eta \bullet \pi$.
- $I \star \xi \bullet \pi \succ \xi \star \pi$.
- $K \star \xi \bullet \eta \bullet \pi \succ \xi \star \pi$.
- $E \star \xi \bullet \eta \bullet \pi \succ (\xi)\eta \star \pi$.
- $W \star \xi \bullet \eta \bullet \pi \succ \xi \star \eta \bullet \eta \bullet \pi$.
- $C \star \xi \bullet \eta \bullet \zeta \bullet \pi \succ \xi \star \zeta \bullet \eta \bullet \pi$.

$B \star \xi \cdot \eta \cdot \zeta \cdot \pi \succ (\xi)(\eta)\zeta \star \pi.$
 $cc \star \xi \cdot \pi \succ \xi \star k_\pi \cdot \pi.$
 $k_\pi \star \xi \cdot \varpi \succ \xi \star \pi.$
 $\varsigma \star \xi \cdot \eta \cdot \pi \succ \xi \star \underline{n}_\eta \cdot \pi.$

Finally, we have a subset \perp of $\Lambda \star \Pi$ which is a final segment for this preorder, which means that : $\mathfrak{p} \in \perp, \mathfrak{p}' \succ \mathfrak{p} \Rightarrow \mathfrak{p}' \in \perp.$

In other words, we ask that \perp has the following properties :

$(\xi)\eta \star \pi \notin \perp \Rightarrow \xi \star \eta \cdot \pi \notin \perp.$
 $I \star \xi \cdot \pi \notin \perp \Rightarrow \xi \star \pi \notin \perp.$
 $K \star \xi \cdot \eta \cdot \pi \notin \perp \Rightarrow \xi \star \pi \notin \perp.$
 $E \star \xi \cdot \eta \cdot \pi \notin \perp \Rightarrow (\xi)\eta \star \pi \notin \perp.$
 $W \star \xi \cdot \eta \cdot \pi \notin \perp \Rightarrow \xi \star \eta \cdot \eta \cdot \pi \notin \perp.$
 $C \star \xi \cdot \eta \cdot \zeta \cdot \pi \notin \perp \Rightarrow \xi \star \zeta \cdot \eta \cdot \pi \notin \perp.$
 $B \star \xi \cdot \eta \cdot \zeta \cdot \pi \notin \perp \Rightarrow (\xi)(\eta)\zeta \star \pi \notin \perp.$
 $cc \star \xi \cdot \pi \notin \perp \Rightarrow \xi \star k_\pi \cdot \pi \notin \perp.$
 $k_\pi \star \xi \cdot \varpi \notin \perp \Rightarrow \xi \star \pi \notin \perp.$
 $\varsigma \star \xi \cdot \eta \cdot \pi \notin \perp \Rightarrow \xi \star \underline{n}_\eta \cdot \pi \notin \perp.$

Remark. Thus, the only arbitrary part in a standard realizability algebra is the set \perp of processes.

c-terms and λ -terms

We call *c-term* a term which is built with variables, the elementary combinators $B, C, E, I, K, W, cc, \varsigma$ and the application (binary function). A closed c-term is exactly what we have called a proof-like term.

Given a c-term t and a variable x , we define inductively on t , a new c-term denoted by $\lambda x t$. To this aim, we apply the first possible case in the following list :

1. $\lambda x t = (K)t$ if t does not contain x .
2. $\lambda x x = I$.
3. $\lambda x tu = (C\lambda x(E)t)u$ if u does not contain x .
4. $\lambda x tx = (E)t$ if t does not contain x .
5. $\lambda x tx = (W)\lambda x(E)t$ (if t contains x).
6. $\lambda x(t)(u)v = \lambda x(B)tuv$ (if uv contain x).

In [16], it is shown that this definition is correct. This allows us to translate every λ -term into a c-term. In the following, almost every c-term will be written as a λ -term. The fundamental property of this translation is given by theorem 1, which is proved in [16] :

Theorem 1. *Let t be a c-term with the only variables x_1, \dots, x_n ; let $\xi_1, \dots, \xi_n \in \Lambda$ and $\pi \in \Pi$. Then $\lambda x_1 \dots \lambda x_n t \star \xi_1 \cdot \dots \cdot \xi_n \cdot \pi \succ t[\xi_1/x_1, \dots, \xi_n/x_n] \star \pi.$*

The formal system

We write formulas and proofs in the language of first order logic. This formal language consists of :

- *individual variables* x, y, \dots ;
- *function symbols* f, g, \dots ; each one has an arity, which is an integer ; function symbols of arity 0 are called *constant symbols*.
- *relation symbols* ; each one has an arity ; relation symbols of arity 0 are called *propositional constants*. We have two particular propositional constants \top, \perp and three particular binary relation symbols \neq, \notin, \subseteq .

The *terms* are built in the usual way with individual variables and function symbols.

Remark. We use the word “term” with two different meanings : here as a term in a first order language, and also as an element of the set Λ of a realizability algebra. I think that, with the help of the context, no confusion is possible.

The *atomic formulas* are the expressions $R(t_1, \dots, t_n)$, where R is a n -ary relation symbol, and t_1, \dots, t_n are terms.

Formulas are built as usual, from atomic formulas, *with the only logical symbols* \rightarrow, \forall :

- each atomic formula is a formula ;
- if A, B are formulas, then $A \rightarrow B$ is a formula ;
- if A is a formula and x an individual variable, then $\forall x A$ is a formula.

Notations.

The formula $A_1 \rightarrow (A_2 \rightarrow (\dots (A_n \rightarrow B) \dots))$ will be written $A_1, A_2, \dots, A_n \rightarrow B$.

The usual logical symbols are defined as follows :

$\neg A \equiv A \rightarrow \perp$; $A \vee B \equiv (A \rightarrow \perp), (B \rightarrow \perp) \rightarrow \perp$; $A \wedge B \equiv (A, B \rightarrow \perp) \rightarrow \perp$; $\exists x F \equiv \forall x (F \rightarrow \perp) \rightarrow \perp$.

More generally, we shall write $\exists x \{F_1, \dots, F_k\}$ for $\forall x (F_1, \dots, F_k \rightarrow \perp) \rightarrow \perp$.

We shall sometimes write \vec{F} for a finite sequence of formulas F_1, \dots, F_k ;

Then, we shall also write $\vec{F} \rightarrow G$ for $F_1, \dots, F_k \rightarrow G$ and $\exists x \{\vec{F}\}$ for $\forall x (\vec{F} \rightarrow \perp) \rightarrow \perp$.

$A \leftrightarrow B$ is the pair of formulas $\{A \rightarrow B, B \rightarrow A\}$.

The rules of natural deduction are the following (the A_i 's are formulas, the x_i 's are variables of \mathbf{c} -term, t, u are \mathbf{c} -terms, written as λ -terms) :

1. $x_1 : A_1, \dots, x_n : A_n \vdash x_i : A_i$.
2. $x_1 : A_1, \dots, x_n : A_n \vdash t : A \rightarrow B, \quad x_1 : A_1, \dots, x_n : A_n \vdash u : A$
 $\Rightarrow x_1 : A_1, \dots, x_n : A_n \vdash tu : B$.
3. $x_1 : A_1, \dots, x_n : A_n, x : A \vdash t : B \Rightarrow x_1 : A_1, \dots, x_n : A_n \vdash \lambda x t : A \rightarrow B$.
4. $x_1 : A_1, \dots, x_n : A_n \vdash t : A \Rightarrow x_1 : A_1, \dots, x_n : A_n \vdash t : \forall x A$ where x is an individual variable which does not appear in A_1, \dots, A_n .
5. $x_1 : A_1, \dots, x_n : A_n \vdash t : \forall x A \Rightarrow x_1 : A_1, \dots, x_n : A_n \vdash t : A[\tau/x]$ where x is an individual variable and τ is a term.
6. $x_1 : A_1, \dots, x_n : A_n \vdash \mathbf{cc} : ((A \rightarrow B) \rightarrow A) \rightarrow A$ (law of Peirce).

The theory \mathbf{ZF}_ε

We write below a set of axioms for a theory called \mathbf{ZF}_ε . Then :

- We show that \mathbf{ZF}_ε is a conservative extension of \mathbf{ZF} .
- We define the realizability models and we show that each axiom of \mathbf{ZF}_ε is realized by a proof-like \mathbf{c} -term, in every realizability model.

It follows that the axioms of ZF are also realized by proof-like c-terms in every realizability model.

We write the axioms of ZF_ε with the predicate constants $\notin, \notin, \subseteq$. Of course, $x \varepsilon y$ and $x \in y$ are the formulas $x \notin y \rightarrow \perp$ and $x \notin y \rightarrow \perp$.

The notation $x \simeq y \rightarrow F$ means $x \subseteq y, y \subseteq x \rightarrow F$. Thus $x \simeq y$, which represents the usual (extensional) equality of sets, is the pair of formulas $\{x \subseteq y, y \subseteq x\}$.

We use the notations $(\forall x \varepsilon a)F(x)$ for $\forall x(\neg F(x) \rightarrow x \notin a)$ and $(\exists x \varepsilon a)\vec{F}(x)$ for $\forall x(\vec{F}(x) \rightarrow x \notin a) \rightarrow \perp$.

For instance, $(\exists x \varepsilon y)t \simeq u$ is the formula $\forall x(t \subseteq u, u \subseteq t \rightarrow x \notin y) \rightarrow \perp$.

The axioms of ZF_ε are the following :

0. Extensionality axioms.

$$\forall x \forall y [x \in y \leftrightarrow (\exists z \varepsilon y)x \simeq z] ; \forall x \forall y [x \subseteq y \leftrightarrow (\forall z \varepsilon x)z \in y].$$

1. Foundation scheme.

$$\forall x((\forall y \varepsilon x)F[y, x_1, \dots, x_n] \rightarrow F[x, x_1, \dots, x_n]) \rightarrow \forall x F[x, x_1, \dots, x_n]$$

for every formula $F[x, x_1, \dots, x_n]$.

The intuitive meaning of axioms 0 and 1 is that ε is a well founded relation, and that the relation \in is obtained by “collapsing” ε into an extensional binary relation.

The following axioms essentially express that the relation ε satisfies the axioms of Zermelo-Fraenkel *except extensionality*.

2. Comprehension scheme.

$$\forall x_1 \dots \forall x_n \forall a \exists b \forall x (x \varepsilon b \leftrightarrow (x \varepsilon a \wedge F[x, x_1, \dots, x_n]))$$

for every formula $F[x, x_1, \dots, x_n]$.

3. Pairing axiom.

$$\forall a \forall b \exists x \{a \varepsilon x, b \varepsilon x\}.$$

4. Union axiom.

$$\forall a \exists b (\forall x \varepsilon a)(\forall y \varepsilon x) y \varepsilon b.$$

5. Power set axiom.

$$\forall a \exists b \forall x (\exists y \varepsilon b) \forall z (z \varepsilon y \leftrightarrow (z \varepsilon a \wedge z \varepsilon x)).$$

6. Collection scheme.

$$\forall x_1 \dots \forall x_n \forall a \exists b (\forall x \varepsilon a) (\exists y F[x, y, x_1, \dots, x_n] \rightarrow (\exists y \varepsilon b) F[x, y, x_1, \dots, x_n])$$

for every formula $F[x, y, x_1, \dots, x_n]$.

7. Infinity scheme.

$$\forall x_1 \dots \forall x_n \forall a \exists b \{a \varepsilon b, (\forall x \varepsilon b) (\exists y F[x, y, x_1, \dots, x_n] \rightarrow (\exists y \varepsilon b) F[x, y, x_1, \dots, x_n])\}$$

for every formula $F[x, y, x_1, \dots, x_n]$.

The usual Zermelo-Fraenkel set theory is obtained from ZF_ε by identifying the predicate symbols \notin and \notin . Thus, the axioms of ZF are written as follows, with the predicate symbols \notin, \subseteq (recall that $x \simeq y$ is the conjunction of $x \subseteq y$ and $y \subseteq x$) :

0. Equality and extensionality axioms.

$$\forall x \forall y [x \in y \leftrightarrow (\exists z \in y)x \simeq z] ; \forall x \forall y [x \subseteq y \leftrightarrow (\forall z \in x)z \in y].$$

1. Foundation scheme.

$$\forall x((\forall y \in x)F[y, x_1, \dots, x_n] \rightarrow F[x, x_1, \dots, x_n]) \rightarrow \forall x F[x, x_1, \dots, x_n]$$

for every formula $F[x, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

2. Comprehension scheme.

$$\forall a \exists b \forall x (x \in b \leftrightarrow (x \in a \wedge F[x, x_1, \dots, x_n]))$$

for every formula $F[x, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

3. Pairing axiom.

$$\forall a \forall b \exists x \{a \in x, b \in x\}.$$

4. Union axiom.

$$\forall a \exists b (\forall x \in a) (\forall y \in x) y \in b.$$

5. Power set axiom.

$$\forall a \exists b \forall x (\exists y \in b) \forall z (z \in y \leftrightarrow (z \in a \wedge z \in x)).$$

6. Collection scheme.

$$\forall a \exists b (\forall x \in a) (\exists y F[x, y, x_1, \dots, x_n] \rightarrow (\exists y \in b) F[x, y, x_1, \dots, x_n])$$

for every formula $F[x, y, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

7. Infinity scheme.

$$\forall a \exists b \{a \in b, (\forall x \in b) (\exists y F[x, y, x_1, \dots, x_n] \rightarrow (\exists y \in b) F[x, y, x_1, \dots, x_n])\}$$

for every formula $F[x, y, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

Remark. The usual statement of the axiom of infinity is the particular case of this scheme, where $a = \emptyset$, and $F(x, y)$ is the formula $y = x \cup \{x\}$.

Let us show that ZF_ε is a conservative extension of ZF. First, it is clear that, if $ZF_\varepsilon \vdash F$, where F is a formula of ZF (i.e. written only with \notin and \subseteq), then $ZF \vdash F$; indeed, it is sufficient to replace \notin with \notin in any proof of $ZF_\varepsilon \vdash F$.

Conversely, we must show that each axiom of ZF is a consequence of ZF_ε .

Theorem 2.

i) $ZF_\varepsilon \vdash \forall a (a \subseteq a)$ (and thus $a \simeq a$).

ii) $ZF_\varepsilon \vdash \forall a \forall x (x \varepsilon a \rightarrow x \in a)$.

i) Using the foundation axiom, we assume $\forall x (x \varepsilon a \rightarrow x \subseteq x)$, and we must show $a \subseteq a$; therefore, we add the hypothesis $x \varepsilon a$. It follows that $x \subseteq x$, then $x \simeq x$, and therefore: $\exists y \{x \simeq y, y \varepsilon a\}$, that is to say $x \in a$. Thus, we have $\forall x (x \varepsilon a \rightarrow x \in a)$, and therefore $a \subseteq a$.

ii) Just shown.

Q.E.D.

Corollary 3. $ZF_\varepsilon \vdash \forall x (x \in a \rightarrow x \in b) \rightarrow a \subseteq b$.

We must show $x \varepsilon a \rightarrow x \in b$, which follows from $x \in a \rightarrow x \in b$ and $x \varepsilon a \rightarrow x \in a$ (theorem 2(ii)).

Q.E.D.

Lemma 4. $ZF_\varepsilon \vdash a \subseteq b, \forall x (x \in b \rightarrow x \in c) \rightarrow a \subseteq c$.

We must show $x \varepsilon a \rightarrow x \in c$, which follows from $x \varepsilon a \rightarrow x \in b$ and $x \in b \rightarrow x \in c$.

Q.E.D.

Theorem 5. $ZF_\varepsilon \vdash \forall y \forall z (y \simeq a, a \varepsilon z \rightarrow y \varepsilon z)$; $ZF_\varepsilon \vdash \forall y \forall z (a \subseteq y, z \varepsilon a \rightarrow z \in y)$.

Call $F(a)$, $F'(a)$ these two formulas. We show $F(a)$ by foundation :

thus, we suppose $(\forall x \varepsilon a)F(x)$ and we first show $F'(a)$: by hypothesis, we have $a \subseteq y$, $z \in a$; thus, there exists a' such that $z \simeq a'$ and $a' \varepsilon a$, and thus $F(a')$. From $a' \varepsilon a$ and $a \subseteq y$, we deduce $a' \in y$. From $z \simeq a'$ and $a' \in y$, we deduce $z \in y$ by $F(a')$.

Then, we show $F(a)$: by hypothesis, we have $y \simeq a$, $a \in z$, thus $a \simeq y'$ and $y' \varepsilon z$ for some y' . In order to show $y \in z$, it is sufficient to show $y \simeq y'$.

Now, we have $y \simeq a$, $a \simeq y'$, and thus $y' \subseteq a$, $a \subseteq y$. From $F'(a)$, we get $\forall z(z \in a \rightarrow z \in y)$; from $y' \subseteq a$, we deduce $y' \subseteq y$ by lemma 4.

We have also $y \subseteq a$, $a \subseteq y'$. From $F'(a)$, we get $\forall z(z \in a \rightarrow z \in y')$; from $y \subseteq a$, we deduce $y \subseteq y'$ by lemma 4.

Q.E.D.

With corollary 3, we obtain :

Corollary 6. $ZF_\varepsilon \vdash b \subseteq c \leftrightarrow \forall x(x \in b \rightarrow x \in c)$.

It is now easy to deduce the equality and extensionality axioms of ZF :

$\forall x(x \simeq x)$; $\forall x \forall y(x \simeq y \rightarrow y \simeq x)$; $\forall x \forall y \forall z(x \simeq y, y \simeq z \rightarrow x \simeq z)$;
 $\forall x \forall x' \forall y \forall y'(x \simeq x', y \simeq y', x \notin y \rightarrow x' \notin y')$; $\forall x \forall y(\forall z(z \notin x \leftrightarrow z \notin y) \rightarrow x \simeq y)$;
 $\forall x \forall y(x \subseteq y \leftrightarrow \forall z(z \notin y \rightarrow z \notin x))$.

Remark. This shows that \simeq is an equivalence relation which is compatible with the relations \in and \subseteq ; but, in general, it is *not compatible with* ε . It is the equality relation for ZF ; it will be called *extensional equivalence*.

Notation. The formula $\forall z(z \notin y \rightarrow z \notin x)$ will be written $x \subset y$. The ordered pair of formulas $x \subset y, y \subset x$ will be written $x \sim y$.

By theorem 2, we get $ZF_\varepsilon \vdash \forall x \forall y(x \subset y \rightarrow x \subseteq y)$. Thus \subset will be called *strong inclusion*, and \sim will be called *strong extensional equivalence*.

- Foundation scheme.

Let $F[x]$ be written with only \notin, \subseteq and let $G[x]$ be the formula $\forall y(y \simeq x \rightarrow F[y])$. Clearly, $\forall x G[x]$ is equivalent to $\forall x F[x]$. Therefore, from axiom scheme 1 of ZF_ε , it is sufficient to show : $\forall b(\forall x(x \in b \rightarrow F[x]) \rightarrow F[b]) \rightarrow (\forall x(x \varepsilon a \rightarrow G[x]) \rightarrow G[a])$, i.e. :

$\forall b(\forall x(x \in b \rightarrow F[x]) \rightarrow F[b]), \forall x \forall y(x \varepsilon a, y \simeq x \rightarrow F[y]), a \simeq b \rightarrow F[b]$.

Therefore, it is sufficient to prove : $\forall x \forall y(x \varepsilon a, y \simeq x \rightarrow F[y]), a \simeq b \rightarrow \forall x(x \in b \rightarrow F[x])$.

From $x \in b, a \simeq b$, we deduce $x \in a$ and therefore (by axiom 0), $x' \varepsilon a$ for some $x' \simeq x$. Finally, we get $F[x]$ from $\forall x \forall y(x \varepsilon a, y \simeq x \rightarrow F[y])$.

- Comprehension scheme : $\forall a \exists b \forall x(x \in b \leftrightarrow (x \in a \wedge F[x]))$

for every formula $F[x, x_1, \dots, x_n]$ written with \notin, \subseteq .

From the axiom scheme 2 of ZF_ε , we get $\forall x(x \varepsilon b \leftrightarrow (x \varepsilon a \wedge F[x]))$. If $x \in b$, then $x \simeq x'$, $x' \varepsilon b$ for some x' . Thus $x' \varepsilon a$ and $F[x']$. From $x \simeq x'$ and $x' \varepsilon a$, we deduce $x \in a$. Since \subseteq and \in are compatible with \simeq , it is the same for F ; thus, we obtain $F[x]$.

Conversely, if we have $F[x]$ and $x \in a$, we have $x \simeq x'$ and $x' \varepsilon a$ for some x' . Since F is compatible with \simeq , we get $F[x']$, thus $x' \varepsilon b$ and $x \in b$.

- Pairing axiom : $\forall x \forall y \exists z \{x \in z, y \in z\}$.

Trivial consequence of axiom 3 of ZF_ε , and theorem 2(ii).

- Union axiom : $\forall a \exists b \forall x \forall y(x \in a, y \in x \rightarrow y \in b)$.

From $x \in a$ we have $x \simeq x'$ and $x' \varepsilon a$ for some x' ; we have $y \in x$, therefore $y \in x'$, thus $y \simeq y'$ and $y' \varepsilon x'$. From axiom 4 of ZF_ε , $x' \varepsilon a$ and $y' \varepsilon x'$, we get $y' \varepsilon b$; therefore $y \in b$, by $y \simeq y'$.

- Power set axiom : $\forall a \exists b \forall x \exists y \{y \in b, \forall z (z \in y \leftrightarrow (z \in a \wedge z \in x))\}$

Given a , we obtain b by axiom 5 of ZF_ε ; given x , we define x' by the condition : $\forall z (z \varepsilon x' \leftrightarrow (z \varepsilon a \wedge z \in x))$ (comprehension scheme of ZF_ε). By definition of b , there exists $y \varepsilon b$ such that $\forall z (z \varepsilon y \leftrightarrow z \varepsilon a \wedge z \varepsilon x')$, and therefore $\forall z (z \varepsilon y \leftrightarrow z \varepsilon a \wedge z \in x)$. It follows easily that $\forall z (z \in y \leftrightarrow z \in a \wedge z \in x)$.

- Collection scheme : $\forall a \exists b (\forall x \in a) (\exists y F[x, y] \rightarrow (\exists y \in b) F[x, y])$

for every formula $F[x, y, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

From $x \in a$ and $\exists y F[x, y]$, we get $x \simeq x'$, $x' \varepsilon a$ for some x' , and thus $\exists y F[x', y]$ since F is compatible with \simeq . From axiom scheme 6 of ZF_ε , we get $\exists y (y \varepsilon b \wedge F[x', y])$, and therefore :

$\exists y (y \in b \wedge F[x, y])$, because $y \varepsilon b \rightarrow y \in b$ and F is compatible with \simeq .

- Infinity scheme : $\forall a \exists b \{a \varepsilon b, (\forall x \in b) (\exists y F[x, y] \rightarrow (\exists y \in b) F[x, y])\}$

for every formula $F[x, y, x_1, \dots, x_n]$ written with the only relation symbols \notin, \subseteq .

Same proof.

Q.E.D.

Realizability models of ZF_ε

We start with an (ordinary) model \mathcal{M} of ZFC, called *the ground model* or *the standard model*. In particular, the integers of \mathcal{M} are called *the standard integers*.

The elements of \mathcal{M} will be called *individuals*.

We define a *realizability model* \mathcal{N} , with the same set of individuals. But \mathcal{N} is not a model in the usual sense, because its truth values are subsets of Π instead of $\{0, 1\}$. Therefore, although \mathcal{M} and \mathcal{N} have the same domain (the quantifier $\forall x$ describes the same domain for both), the model \mathcal{N} may (and will, in all non trivial cases) have much more individuals than \mathcal{M} , because it has individuals which are *not named*. In particular, it will have *non standard integers*.

Remark. This is a great difference between *realizability* and *forcing* models of ZF. In a forcing model, each individual is named in the ground model; it follows that integers, and even ordinals, are not changed.

For each closed formula F with parameters in \mathcal{M} , we define two truth values :

$\|F\| \subseteq \Pi$ and $|F| \subseteq \Lambda$.

$|F|$ is defined immediately from $\|F\|$ as follows :

$$\xi \in |F| \Leftrightarrow (\forall \pi \in \|F\|) \xi \star \pi \in \perp.$$

Notation. We shall write $\xi \Vdash F$ for $\xi \in |F|$.

$\|F\|$ is now defined by recurrence on the length of F :

- F is atomic ;

then F has one of the forms $\top, \perp, a \notin b, a \subseteq b, a \notin b$ where a, b are parameters in \mathcal{M} .

We set :

$\|\top\| = \emptyset$; $\|\perp\| = \Pi$; $\|a \neq b\| = \{\pi \in \Pi; (a, \pi) \in b\}$.

$\|a \subseteq b\|$, $\|a \not\subseteq b\|$ are defined simultaneously by induction on $(\text{rk}(a) \cup \text{rk}(b), \text{rk}(a) \cap \text{rk}(b))$ ($\text{rk}(a)$ being the rank of a).

$\|a \subseteq b\| = \bigcup_c \{\xi \cdot \pi; \xi \in \Lambda, \pi \in \Pi, (c, \pi) \in a, \xi \Vdash c \not\subseteq b\}$;

$\|a \not\subseteq b\| = \bigcup_c \{\xi \cdot \xi' \cdot \pi; \xi, \xi' \in \Lambda, \pi \in \Pi, (c, \pi) \in b, \xi \Vdash a \subseteq c, \xi' \Vdash c \subseteq a\}$.

- $F \equiv A \rightarrow B$; then $\|F\| = \{\xi \cdot \pi; \xi \Vdash A, \pi \in \|B\|\}$.
- $F \equiv \forall x A$: then $\|F\| = \bigcup_a \|A[a/x]\|$.

The following theorem is an essential tool :

Theorem 7 (Adequacy lemma).

Let A_1, \dots, A_n, A be closed formulas of ZF_ε , and suppose that $x_1 : A_1, \dots, x_n : A_n \vdash t : A$.

If $\xi_1 \Vdash A_1, \dots, \xi_n \Vdash A_n$ then $t[\xi_1/x_1, \dots, \xi_n/x_n] \Vdash A$.

In particular, if $\vdash t : A$, then $t \Vdash A$.

We need to prove a (seemingly) more general result, that we state as a lemma :

Lemma 8. Let $A_1[\vec{z}], \dots, A_n[\vec{z}], A[\vec{z}]$ be formulas of ZF_ε , with $\vec{z} = (z_1, \dots, z_k)$ as free variables, and suppose that $x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash t : A[\vec{z}]$.

If $\xi_1 \Vdash A_1[\vec{a}], \dots, \xi_n \Vdash A_n[\vec{a}]$ for some parameters (i.e. individuals in \mathcal{M})

$\vec{a} = (a_1, \dots, a_k)$, then $t[\xi_1/x_1, \dots, \xi_n/x_n] \Vdash A[\vec{a}]$.

Proof by recurrence on the length of the derivation of $x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash t : A[\vec{z}]$.

We consider the last used rule.

1. $x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash x_i : A_i[\vec{z}]$. This case is trivial.

2. We have the hypotheses :

$x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash u : B[\vec{z}] \rightarrow A[\vec{z}]$; $x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash v : B[\vec{z}]$;
 $t = uv$.

By the induction hypothesis, we have $u[\vec{\xi}/\vec{x}] \Vdash B[\vec{a}/\vec{z}] \rightarrow A[\vec{a}/\vec{z}]$ and $v[\vec{\xi}/\vec{x}] \Vdash B[\vec{a}/\vec{z}]$.

Therefore $(uv)[\vec{\xi}/\vec{x}] \Vdash A[\vec{a}/\vec{z}]$ which is the desired result.

3. We have the hypotheses :

$x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}], y : B[\vec{z}] \vdash u : C[\vec{z}]$; $A[\vec{z}] \equiv B[\vec{z}] \rightarrow C[\vec{z}]$; $t = \lambda y u$.

We want to show that $(\lambda y u)[\vec{\xi}/\vec{x}] \Vdash B[\vec{a}/\vec{z}] \rightarrow C[\vec{a}/\vec{z}]$. Thus, let :

$\eta \Vdash B[\vec{a}/\vec{z}]$ and $\pi \in \|C[\vec{a}/\vec{z}]\|$. We must show :

$(\lambda y u)[\vec{\xi}/\vec{x}] \star \eta \cdot \pi \in \perp$ or else $u[\vec{\xi}/\vec{x}, \eta/y] \star \pi \in \perp$.

Now, by the induction hypothesis, we have $u[\vec{\xi}/\vec{x}, \eta/y] \Vdash C[\vec{a}/\vec{z}]$,

which gives the result.

4. We have the hypotheses :

$x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash t : B[\vec{z}]$; $A[\vec{z}] \equiv \forall z_1 B[\vec{z}]$; $\xi_i \Vdash A_i[a_1/z_1, a_2/z_2, \dots, a_k/z_k]$;
the variable z_1 is not free in $A_1[\vec{z}], \dots, A_n[\vec{z}]$.

We have to show that $t[\vec{\xi}/\vec{x}] \Vdash \forall z_1 B[\vec{a}/\vec{z}]$ i.e. $t[\vec{\xi}/\vec{x}] \Vdash \forall z_1 B[a_2/z_2, \dots, a_k/z_k]$. Thus,

we take an arbitrary set b in \mathcal{M} and we show $t[\vec{\xi}/\vec{x}] \Vdash B[b/z_1, a_2/z_2, \dots, a_k/z_k]$.

By the induction hypothesis, it is sufficient to show that $\xi_i \Vdash A_i[b/z_1, a_2/z_2, \dots, a_k/z_k]$. But this follows from the hypothesis on ξ , because z_1 is not free in the formulas A_i .

5. We have the hypotheses :

$x_1 : A_1[\vec{z}], \dots, x_n : A_n[\vec{z}] \vdash t : \forall y B[y, \vec{z}]$; $A[\vec{z}] \equiv B[\tau[\vec{z}]/y, \vec{z}]$.

By the induction hypothesis, we have $t[\vec{\xi}/\vec{x}] \Vdash \forall y B[y, \vec{a}/\vec{z}]$; therefore $t[\vec{\xi}/\vec{x}] \Vdash B[b, \vec{a}/\vec{z}]$ for every parameter b . We get the desired result by taking $b = \tau[\vec{a}]$.

6. The result follows from the following :

Theorem 9. *For every formulas A, B , we have $\text{cc} \Vdash ((A \rightarrow B) \rightarrow A) \rightarrow A$.*

Let $\xi \Vdash (A \rightarrow B) \rightarrow A$ and $\pi \in \|A\|$. Then $\text{cc} \star \xi \cdot \pi \succ \xi \star k_\pi \cdot \pi$ which is in \perp , because $k_\pi \Vdash A \rightarrow B$ by lemma 10.

Q.E.D.

Lemma 10. *If $\pi \in \|A\|$, then $k_\pi \Vdash A \rightarrow B$.*

Indeed, let $\xi \Vdash A$; then $k_\pi \star \xi \cdot \pi' \succ \xi \star \pi \in \perp$ for every stack $\pi' \in \|B\|$.

Q.E.D.

This completes the proof of lemma 8 and theorem 7.

Q.E.D.

Realized formulas and coherent models

In the ground model \mathcal{M} , we interpret the formulas of the *language of ZF* : this language consists of $\in, =$; we add some function symbols, but these functions are always defined, in \mathcal{M} , by some formulas written with $\in, =$. We suppose that this ground model satisfies ZFC. The value, in \mathcal{M} , of a closed formula F of the language of ZF, with parameters in \mathcal{M} , is of course 1 or 0. In the first case, we say that \mathcal{M} *satisfies* F , and we write $\mathcal{M} \models F$.

In the realizability model \mathcal{N} , we interpret the formulas of the *language of ZF_ε* , which consists of $\notin, \subseteq, \notin$ and the same function symbols as in the language of ZF. The domain of \mathcal{N} and the interpretation of the function symbols are the same as for the model \mathcal{M} .

The value, in \mathcal{N} , of a closed formula F of ZF_ε with parameters (in \mathcal{M} or in \mathcal{N} , which is the same thing) is an element of $\mathcal{P}(\Pi)$ which is denoted as $\|F\|$, the definition of which has been given above.

Thus, we can no longer say that \mathcal{N} satisfies (or not) a given closed formula F . But we shall say that \mathcal{N} *realizes* F (and we shall write $\mathcal{N} \Vdash F$), if there exists a proof-like term θ such that $\theta \Vdash F$. We say that two closed formulas F, G are *interchangeable* if $\mathcal{N} \Vdash F \leftrightarrow G$.

Notice that, if $\|F\| = \|G\|$, then F, G are interchangeable (indeed $I \Vdash F \rightarrow G$), but the converse is far from being true.

The model \mathcal{N} allows us to make relative consistency proofs, since it is clear, from the adequacy lemma (theorem 7), that the class of formulas which are realized in \mathcal{N} is closed by deduction in classical logic. Nevertheless, we must check that the realizability model \mathcal{N} is *coherent*, i.e. that it does not realize the formula \perp . We can express this condition in the following form :

For every proof-like term θ , there exists a stack $\pi \in \Pi$ such that $\theta \star \pi \notin \perp$.

When the model \mathcal{N} is coherent, it is not *complete*, except in trivial cases. This means that there exist closed formulas F of ZF_ε such that $\mathcal{N} \not\models F$ and $\mathcal{N} \not\models \neg F$.

The axioms of ZF_ε are realized in \mathcal{N}

- Extensionality axioms.

We have $\|\forall z(z \notin b \rightarrow z \notin a)\| = \bigcup \{\xi \cdot \pi; \xi \Vdash c \notin b, \pi \in \|c \notin a\|\}$

by definition of the value of $\|\forall z(z \notin b \rightarrow z \notin a)\|$;

and $\|a \subseteq b\| = \bigcup \{\xi \cdot \pi; (c, \pi) \in a, \xi \Vdash c \notin b\}$ by definition of $\|a \subseteq b\|$.

Therefore, we have $\|a \subseteq b\| = \|\forall z(z \notin b \rightarrow z \notin a)\|$, so that :

$I \Vdash \forall x \forall y (x \subseteq y \rightarrow \forall z (z \notin y \rightarrow z \notin x))$ and $I \Vdash \forall x \forall y (\forall z (z \notin y \rightarrow z \notin x) \rightarrow x \subseteq y)$.

In the same way, we have :

$\|\forall z(a \subseteq z, z \subseteq a \rightarrow z \notin b)\| = \bigcup \{\xi \cdot \xi' \cdot \pi; \xi \Vdash a \subseteq c, \xi' \Vdash c \subseteq a; \pi \in \|c \notin b\|\}$

by definition of the value of $\|\forall z(a \subseteq z, z \subseteq a \rightarrow z \notin b)\|$;

and $\|a \notin b\| = \bigcup \{\xi \cdot \xi' \cdot \pi; (c, \pi) \in b, \xi \Vdash a \subseteq c, \xi' \Vdash c \subseteq a\}$ by definition of $\|a \notin b\|$.

Therefore, we have $\|a \notin b\| = \|\forall z(a \subseteq z, z \subseteq a \rightarrow z \notin b)\|$, so that :

$I \Vdash \forall x \forall y (x \notin y \rightarrow \forall z (x \subseteq z, z \subseteq x \rightarrow z \notin y))$;

$I \Vdash \forall x \forall y (\forall z (x \subseteq z, z \subseteq x \rightarrow z \notin y) \rightarrow x \notin y)$.

Notation. We shall write $\vec{\xi}$ for a finite sequence (ξ_1, \dots, ξ_n) of terms. Therefore, we shall write $\vec{\xi} \Vdash \vec{A}$ for $\xi_i \Vdash A_i$ ($i = 1, \dots, n$).

In particular, the notation $\vec{\xi} \Vdash a \simeq b$ means $\xi_1 \Vdash a \subseteq b, \xi_2 \Vdash b \subseteq a$;

the notation $\vec{\xi} \Vdash A \leftrightarrow B$ means $\xi_1 \Vdash A \rightarrow B, \xi_2 \Vdash B \rightarrow A$.

- Foundation scheme.

Theorem 11. $\mathbf{Y} \Vdash \forall x (\forall y (\vec{F}[y] \rightarrow y \notin x), \vec{F}[x] \rightarrow \perp) \rightarrow \forall x (\vec{F}[x] \rightarrow \perp)$

for every finite sequence $\vec{F}[x, x_1, \dots, x_n]$ of formulas.

Let $\xi \Vdash \forall x (\forall y (\vec{F}[y] \rightarrow y \notin x), \vec{F}[x] \rightarrow \perp)$ and $\vec{\eta} \Vdash \vec{F}[a]$. We show that $\mathbf{Y} \star \xi \cdot \vec{\eta} \cdot \pi \in \perp$, for every $\pi \in \Pi$, by induction on the rank of a . It suffices to show $\xi \star \mathbf{Y} \xi \cdot \vec{\eta} \cdot \pi \in \perp$.

Now, $\xi \Vdash \forall y (\vec{F}[y] \rightarrow y \notin a), \vec{F}[a] \rightarrow \perp$, so that it suffices to show $\mathbf{Y} \xi \Vdash \forall y (\vec{F}[y] \rightarrow y \notin a)$, in other words $\mathbf{Y} \xi \Vdash \vec{F}[b] \rightarrow b \notin a$ for every b . Let $\vec{\zeta} \Vdash \vec{F}[b]$ and $\varpi \in \|b \notin a\|$. Thus, we have $(b, \varpi) \in a$, therefore $\text{rk}(b) < \text{rk}(a)$ so that $\mathbf{Y} \star \xi \cdot \vec{\zeta} \cdot \varpi \in \perp$ by induction hypothesis. It follows that $\mathbf{Y} \xi \star \vec{\zeta} \cdot \varpi \in \perp$, which is the desired result.

Q.E.D.

It follows from theorem 11 that the axiom scheme 1 of ZF_ε (foundation) is realized.

- Comprehension scheme.

Let a be a set, and $F[x]$ a formula with parameters. We put :

$b = \{(x, \xi \cdot \pi); (x, \pi) \in a, \xi \Vdash F[x]\}$; then, we have trivially $\|x \notin b\| = \|F(x) \rightarrow x \notin a\|$.

Therefore $I \Vdash \forall x [x \notin b \rightarrow (F(x) \rightarrow x \notin a)]$ and $I \Vdash \forall x [(F(x) \rightarrow x \notin a) \rightarrow x \notin b]$.

- Pairing axiom.

We consider two sets a and b , and we put $c = \{a, b\} \times \Pi$. We have $\|a \notin c\| = \|b \notin c\| = \|\perp\|$, thus $I \Vdash a \varepsilon c$ and $I \Vdash b \varepsilon c$.

- Union axiom.

Given a set a , let $b = \text{Cl}(a)$ (the transitive closure of a , i.e. the least transitive set which contains a). We show $\|y \notin x \rightarrow x \notin a\| \supset \|y \notin b \rightarrow x \notin a\|$: indeed, let $\xi \cdot \pi \in \|y \notin b \rightarrow x \notin a\|$, i.e. $\xi \Vdash y \notin b$ and $(x, \pi) \in a$. Therefore, $\text{Cl}(a) \supset x$, i.e. $b \supset x$ and thus $\|y \notin b\| \supset \|y \notin x\|$.

Thus, we have $\xi \Vdash y \notin x$, which gives the result.

It follows that $I \Vdash \forall x \forall y [(y \notin x \rightarrow x \notin a) \rightarrow (y \notin b \rightarrow x \notin a)]$.

- Power set axiom.

Given a set a , let $b = \mathcal{P}(\text{Cl}(a) \times \Pi) \times \Pi$. For every set x , we put :

$y = \{(z, \xi \cdot \pi); (z, \pi) \in a, \xi \Vdash F[z, x]\}$, where $F[z, x]$ is a formula with parameters. Then, as we have seen above (comprehension scheme), we have $\|z \notin y\| = \|F[z, x] \rightarrow z \notin a\|$ and therefore $(I, I) \Vdash \forall z (z \notin y \leftrightarrow (F[z, x] \rightarrow z \notin a))$.

Now, it is obvious that $y \in \mathcal{P}(\text{Cl}(a) \times \Pi)$, and therefore $(y, \pi) \in b$ for every $\pi \in \Pi$.

Thus, we have $\|y \notin b\| = \Pi = \|\perp\|$. Therefore $I \Vdash y \varepsilon b$, and finally :

$(I, (I, I)) \Vdash y \varepsilon b \wedge \forall z (z \notin y \leftrightarrow (F[z, x] \rightarrow z \notin a))$.

The power set axiom is the particular case when the formula $F[z, x]$ is $z \varepsilon x$.

- Collection scheme.

Given a set a , and a formula $F[x, y]$ with parameters, let :

$b = \bigcup \{\Phi(x, \xi) \times \text{Cl}(a); x \in \text{Cl}(a), \xi \in \Lambda\}$ with

$\Phi(x, \xi) = \{y \text{ of minimum rank}; \xi \Vdash F[x, y]\}$ or $\Phi(x, \xi) = \emptyset$ if there is no such y .

We show that $\|\forall y (F[x, y] \rightarrow y \notin b)\| \supset \|\forall y (F[x, y] \rightarrow x \notin a)\|$:

Suppose indeed that $\xi \cdot \pi \in \|\forall y (F[x, y] \rightarrow x \notin a)\|$, i.e. $(x, \pi) \in a$ and $\xi \Vdash F[x, y]$ for some y . By definition of $\Phi(x, \xi)$, there exists $y' \in \Phi(x, \xi)$. Moreover, we have :

$x \in \text{Cl}(a)$, $\pi \in \text{Cl}(a)$, and therefore $(y', \pi) \in b$; it follows that $\|y' \notin b\| \supset \|x \notin a\|$. But $y' \in \Phi(x, \xi)$, and therefore $\xi \Vdash F[x, y']$; thus, we have $\xi \cdot \pi \in \|F[x, y'] \rightarrow y' \notin b\|$, which gives the result.

We have proved that $I \Vdash \forall y (F[x, y] \rightarrow y \notin b) \rightarrow \forall y (F[x, y] \rightarrow x \notin a)$.

- Infinity scheme.

Given a set a , we define b as the least set such that :

$\{a\} \times \Pi \subseteq b$ and $\forall x (\forall \pi \in \Pi) (\forall \xi \in \Lambda) ((x, \pi) \in b \Rightarrow \Phi(x, \xi) \times \{\pi\} \subseteq b)$

where $\Phi(x, \xi)$ is defined as above.

We have $\{a\} \times \Pi \subseteq b$, thus $\|a \notin b\| = \|\perp\|$, and therefore $I \Vdash a \varepsilon b$.

We now show that $\|\forall y (F[x, y] \rightarrow y \notin b)\| \supset \|\forall y (F[x, y] \rightarrow x \notin b)\|$:

Suppose indeed that $\xi \cdot \pi \in \|\forall y (F[x, y] \rightarrow x \notin b)\|$, i.e. $(x, \pi) \in b$ and $\xi \Vdash F[x, y]$ for some y . By definition of $\Phi(x, \xi)$, there exists $y' \in \Phi(x, \xi)$. By definition of b , we have $(y', \pi) \in b$, i.e. $\pi \in \|y' \notin b\|$. Now, $y' \in \Phi(x, \xi)$, and therefore $\xi \Vdash F[x, y']$; thus, we have :

$\xi \cdot \pi \in \|F[x, y'] \rightarrow y' \notin b\|$, which gives the result.

It follows that $I \Vdash \forall y (F[x, y] \rightarrow y \notin b) \rightarrow \forall y (F[x, y] \rightarrow x \notin b)$ and therefore :

$(I, I) \Vdash \{a \varepsilon b, \forall x (\forall y (F[x, y] \rightarrow y \notin b) \rightarrow \forall y (F[x, y] \rightarrow x \notin b))\}$.

Function symbols and equality

Following our needs, we shall add to the language of ZF_ε , some *function symbols* f, g, \dots of any arity. A k -ary function symbol f will be interpreted, in the realizability model \mathcal{N} , by a *functional relation*, which is defined in the ground model \mathcal{M} by a formula $F[x_1, \dots, x_k, y]$ of ZF. Thus, we assume that $\mathcal{M} \models \forall x_1 \dots \forall x_k \exists! y F[x_1, \dots, x_k, y]$ ($\exists! y F[y]$ is the conjunction of $\forall y \forall y' (F[y], F[y'] \rightarrow y = y')$ and $\exists y F[y]$).

The axiom schemes of ZF_ε , written in the extended language, are still realized in the model \mathcal{N} , because the above proofs remain valid.

On the other hand, in order to make sure that the axiom schemes of ZF, which use a k -ary function symbol f , are still realized, one must check that this symbol is *compatible with* \simeq , i.e. that the following formula is realized in \mathcal{N} :

$$\forall x_1 \dots \forall x_k (x_1 \simeq y_1, \dots, x_k \simeq y_k \rightarrow f x_1 \dots x_k \simeq f y_1 \dots y_k).$$

We now add a new rule to build formulas of ZF_ε :

If t, u are two terms and F is a formula of ZF_ε , then $t = u \mapsto F$ is a formula of ZF_ε .

The formula $t = u \mapsto \perp$ is denoted $t \neq u$.

The formula $t \neq u \rightarrow \perp$, i.e. $(t = u \mapsto \perp) \rightarrow \perp$ is denoted $t = u$.

The truth value of these new formulas is defined as follows, assuming that t, u, F are closed, with parameters in \mathcal{N} :

$$\|t = u \mapsto F\| = \emptyset \text{ if } t \neq u ; \|t = u \mapsto F\| = \|F\| \text{ if } t = u.$$

It follows that :

$$\begin{aligned} \|t \neq u\| &= \emptyset = \|\top\| \text{ if } t \neq u ; \|t \neq u\| = \Pi = \|\perp\| \text{ if } t = u ; \\ \|t = u\| &= \|\top \rightarrow \perp\| \text{ if } t \neq u ; \|t = u\| = \|\perp \rightarrow \perp\| \text{ if } t = u. \end{aligned}$$

Proposition 12 shows that $t = u \mapsto F$ and $t = u \rightarrow F$ are interchangeable.

Proposition 12.

- i) $\lambda x(x)I \Vdash (t = u \rightarrow F) \rightarrow (t = u \mapsto F)$;
- ii) $\lambda x \lambda y (\text{cc}) \lambda k(y)(k)x \Vdash (t = u \mapsto F), t = u \rightarrow F$.

i) Let $\xi \Vdash t = u \rightarrow F$ and $\pi \in \|t = u \mapsto F\|$. Thus, we have $t = u$ and $\pi \in \|F\|$.

We must show $\lambda x(x)I \star \xi \cdot \pi \in \perp$, that is $\xi \star I \cdot \pi \in \perp$. This is immediate, by hypothesis on ξ , since $I \Vdash t = u$.

ii) Let $\xi \Vdash t = u \mapsto F$, $\eta \Vdash t = u$ and $\pi \in \|F\|$. We must show that :

$$\lambda x \lambda y (\text{cc}) \lambda k(y)(k)x \star \xi \cdot \eta \cdot \pi \in \perp, \text{ soit } \eta \star \kappa_\pi \xi \cdot \pi \in \perp.$$

If $t \neq u$, then $\eta \Vdash \top \rightarrow \perp$, hence the result.

If $t = u$, then $\xi \Vdash F$, thus $\xi \star \pi \in \perp$, therefore $\kappa_\pi \xi \Vdash \perp$.

But we have $\eta \Vdash \perp \rightarrow \perp$, and therefore $\eta \star \xi \cdot \pi \in \perp$.

Q.E.D.

Proposition 13 shows that the formulas $t = u$ and $\forall x (u \notin x \rightarrow t \notin x)$ (*Leibniz equality*) are interchangeable.

Proposition 13.

- i) $I \Vdash t = u \mapsto \forall x (u \notin x \rightarrow t \notin x)$;
- ii) $I \Vdash \forall x (x \notin u \rightarrow x \notin t) \rightarrow t = u$.

i) It suffices to check that $I \Vdash \forall x(u \notin x \rightarrow t \notin x)$ when $t = u$, which is obvious.
ii) We must show that $I \Vdash \forall x(u \notin x \rightarrow t \notin x), t \neq u \rightarrow \perp$. Thus let $\xi \Vdash \forall x(u \notin x \rightarrow t \notin x)$, $\eta \Vdash t \neq u$ and $\pi \in \Pi$; we must show that $\xi \star \eta \bullet \pi \in \perp$.
We have $\xi \Vdash u \notin a \rightarrow t \notin a$ for every a ; we take $a = \{t\} \times \Pi$, thus $\|t \notin a\| = \Pi$, hence $\pi \in \|t \notin a\|$.
If $t = u$, we have $\eta \Vdash \perp$, thus $\eta \Vdash u \notin a$, hence the result.
If $t \neq u$, we have $\|u \notin a\| = \emptyset = \|\top\|$, thus $\eta \Vdash u \notin a$, hence the result.
Q.E.D.

We now show that the axioms of equality are realized.

Proposition 14. $I \Vdash \forall x(x = x)$; $I \Vdash \forall x \forall y(x = y \mapsto y = x)$;

$I \Vdash \forall x \forall y \forall z(x = y \mapsto (y = z \mapsto x = z))$;

$I \Vdash \forall x \forall y(x = y \mapsto (F[x] \rightarrow F[y]))$ for every formula F with one free variable, with parameters.

Trivial, by definition of \mapsto .

Q.E.D.

Conservation of well-foundedness

Theorem 15 says that every well founded relation in the ground model \mathcal{M} , gives a well founded relation in the realizability model \mathcal{N} .

Theorem 15. Let f be a binary function such that $f(x, y) = 1$ is a well founded relation in the ground model \mathcal{M} . Then, for every formula $F[x]$ of ZF_ε with parameters in \mathcal{M} :
 $\mathbf{Y} \Vdash \forall x(\forall y(f(y, x) = 1 \mapsto F[y]) \rightarrow F[x]) \rightarrow \forall x F[x]$.

Let us fix a and let $\xi \Vdash \forall x(\forall y(f(y, x) = 1 \mapsto F[y]) \rightarrow F[x])$. We show, by induction on a , following the well founded relation $f(x, y) = 1$, that $\mathbf{Y} \star \xi \bullet \pi \in \perp$ for every $\pi \in \|F[a]\|$. Thus, suppose that $\pi \in \|F[a]\|$; we need to show that $\xi \star \mathbf{Y} \xi \bullet \pi \in \perp$. By hypothesis, we have :

$\xi \Vdash \forall y(f(y, a) = 1 \mapsto y \notin z) \rightarrow F[a]$; thus, it suffices to show that :

$\mathbf{Y} \xi \Vdash f(y, a) = 1 \mapsto F[y]$ for every y . This is clear if $f(y, a) \neq 1$, by definition of \mapsto .

If $f(y, a) = 1$, we must show $\mathbf{Y} \xi \Vdash F[y]$, i.e. $\mathbf{Y} \star \xi \bullet \rho \in \perp$ for every $\rho \in \|F[y]\|$. But this follows from the induction hypothesis.

Q.E.D.

Sets in \mathcal{M} give type-like sets in \mathcal{N}

We define a unary function symbol \mathbf{J} by putting $\mathbf{J}(a) = a \times \Pi$ for every individual a (element of the ground model \mathcal{M}).

For each set E of the ground model \mathcal{M} , we also introduce the unary function 1_E with values in $\{0, 1\}$, defined as follows :

$1_E(a) = 1$ if $a \in E$; $1_E(a) = 0$ if $a \notin E$.

The formula $1_E(x) = 1 \mapsto A$ will also be denoted as $x \varepsilon \mathbf{J}E \mapsto A$.

In particular, $a \notin \mathbf{J}E$ is identical with $a \varepsilon \mathbf{J}E \mapsto \perp$ that is $1_E(a) \neq 1$.

We shall write $(\forall x \varepsilon \mathbf{J}E) A[x]$ for $\forall x(x \varepsilon \mathbf{J}E \mapsto A[x])$.

Proposition 12 shows that $x \varepsilon \mathfrak{J}E \mapsto A$ and $x \varepsilon \mathfrak{J}E \rightarrow A$ are interchangeable. Therefore $(\forall x \varepsilon \mathfrak{J}E) A[x]$ and $\forall x(x \varepsilon \mathfrak{J}E \rightarrow A[x])$ are also interchangeable. We have :

$$\|(\forall x \varepsilon \mathfrak{J}E) A[x]\| = \bigcup_{a \in E} \|A[a/x]\| \quad \text{and} \quad |(\forall x \varepsilon \mathfrak{J}E) A[x]| = \bigcap_{a \in E} |A[a/x]|.$$

As already said, we shall add to the language of ZF_ε , some function symbols of any arity, which will be interpreted in the ground model \mathcal{M} by some functional relations. Then every formula of the form $\forall \vec{x}(t_1[\vec{x}] = u_1[\vec{x}], \dots, t_k[\vec{x}] = u_k[\vec{x}] \rightarrow t[\vec{x}] = u[\vec{x}])$ which is satisfied in the model \mathcal{M} , is *realized* in the model \mathcal{N} ($t_1, u_1, \dots, t_k, u_k, t, u$ are terms of the language).

Indeed, we verify immediately that :

$$I \Vdash \forall \vec{x}(t_1[\vec{x}] = u_1[\vec{x}] \mapsto (\dots \mapsto (t_k[\vec{x}] = u_k[\vec{x}] \mapsto t[\vec{x}] = u[\vec{x}])) \dots).$$

It follows that if, for instance, $t[x, x']$ sends $E \times E'$ into D in the model \mathcal{M} , then it sends $\mathfrak{J}E \times \mathfrak{J}E'$ into $\mathfrak{J}D$ in the model \mathcal{N} . Indeed, we have then :

$\mathcal{M} \models \forall x \forall x'(1_E(x) = 1, 1_{E'}(x') = 1 \rightarrow 1_D(t[x, x']) = 1)$ and therefore, we have :

$$I \Vdash \forall x \forall x'(1_E(x) = 1 \mapsto (1_{E'}(x') = 1 \mapsto 1_D(t[x, x']) = 1)), \text{ in other words :}$$

$$I \Vdash (\forall x \varepsilon \mathfrak{J}E)(\forall x' \varepsilon \mathfrak{J}E')(t[x, x'] \varepsilon \mathfrak{J}D).$$

Notice, in particular, that the characteristic function 1_E , which takes its values in the set $2 = \{0, 1\}$ in the model \mathcal{M} , takes its values in $\mathfrak{J}2$ in the realizability model \mathcal{N} .

We shall denote \wedge, \vee, \neg the (trivial) Boolean algebra operations in $\{0, 1\}$ (they should not be confused with the logical connectives \wedge, \vee, \neg). In this way, we have defined three function symbols of the language of ZF_ε ; thus, in the realizability model \mathcal{N} , they define a *Boolean algebra structure* on the set $\mathfrak{J}2$.

The set $\tilde{\mathfrak{N}}$ of integers in \mathcal{N}

We add to the language of ZF_ε a constant symbol 0 and a unary function symbol s . Their interpretation in the model \mathcal{M} is as follows :

0 is \emptyset ; $s(a)$ is $\{a\} \times \Pi$ for every set a , in other words $s(a) = \mathfrak{J}(\{a\})$.

Remark. In the definition of the set of integers in the realizability model \mathcal{N} , we are using the singleton as the successor function s , instead of the usual one $x \mapsto x \cup \{x\}$, which is more complicated to define in the realizability model. It would give :

$$s(a) = \{(a, \underline{1} \cdot \pi); \pi \in \Pi\} \cup \{(x, \underline{0} \cdot \pi); x \in a, \pi \in \Pi\}.$$

Theorem 16. *The following formulas are realized in \mathcal{N} :*

- i) $\forall x \forall y (sx = sy \mapsto x = y)$;
- ii) $\forall x (sx \neq 0)$;
- iii) $\forall x \forall y (x \simeq y \rightarrow sx \simeq sy)$;
- iv) $\forall x \forall y (sx \simeq sy \rightarrow x \simeq y)$.

This shows, in particular, that the function s is *compatible with the extensional equivalence* \simeq .

i) We check that $I \Vdash sa = sb \mapsto a = b$. We may suppose $sa = sb$, because $\|sa = sb \mapsto a = b\| = \emptyset$ if $sa \neq sb$. But, in this case, we have $a = b$, by definition of sa, sb .

ii) We have $\|a \notin sa\| = \|\perp\|$, thus $I \Vdash a \varepsilon sa$. Since $\mathcal{N} \Vdash \forall x \forall y (y \notin x \rightarrow y \notin x)$, we have : $\mathcal{N} \Vdash a \in sa$. But $I \Vdash a \notin 0$, and therefore $\mathcal{N} \Vdash (a \notin 0 \rightarrow a \notin sa) \rightarrow \perp$; thus $\mathcal{N} \Vdash \forall x (x \notin 0 \rightarrow x \notin sa) \rightarrow \perp$, i.e. $\mathcal{N} \Vdash sa \not\subseteq 0$. Therefore, $\mathcal{N} \Vdash sa \neq 0$.

iii) We show that the formula $a \simeq b \rightarrow sa \simeq sb$ is realized ; it suffices to realize the formula $a \simeq b \rightarrow sa \subseteq sb$. We prove it by means of already realized sentences.

We need to prove $a \simeq b, x \notin sb \rightarrow x \notin sa$. But $x \notin sa$ has the same truth value as $x \neq a$. Thus, we simply have to prove $a \simeq b \rightarrow a \in sb$. But $a \in sb$ follows from $b \varepsilon sb$ and $a \simeq b$.

iv) In the same way, we prove the formula $sa \simeq sb \rightarrow a \simeq b$ and, in fact $sa \subseteq sb \rightarrow a \simeq b$. The formula $sa \subseteq sb$ is $\forall x (x \notin sb \rightarrow x \notin sa)$; but $x \notin sa$ is the same as $x \neq a$. Thus, from $sa \subseteq sb$ we obtain $a \in sb$, i.e. $(\exists x \varepsilon sb) x \simeq a$. But $x \varepsilon sb$ is the same as $x = b$, so that we obtain $a \simeq b$.

Q.E.D.

The individuals $s^n 0$ are obviously distinct, for $n \in \mathbb{N}$. Therefore, we can define :

$$\tilde{\mathbb{N}} = \{(s^n 0, \underline{n} \cdot \pi) ; n \in \mathbb{N}, \pi \in \Pi\}$$

and we have :

$\|a \notin \tilde{\mathbb{N}}\| = \emptyset$ if a is not of the form $s^n 0$, with $n \in \mathbb{N}$;

$\|s^n 0 \notin \tilde{\mathbb{N}}\| = \{\underline{n} \cdot \pi ; \pi \in \Pi\}$.

The formula $x \varepsilon \tilde{\mathbb{N}}$ will also be written $\text{ent}(x)$.

In the sequel, we shall use the restricted quantifier $\forall x \varepsilon \tilde{\mathbb{N}}$, which we also write $\forall x^{\text{ent}}$, with the following meaning :

$\|\forall x^{\text{ent}} F[x]\| = \|(\forall x \varepsilon \tilde{\mathbb{N}}) F[x]\| = \{\underline{n} \cdot \pi ; n \in \mathbb{N}, \pi \in \|F[s^n 0]\|\}$.

The restricted existential quantifier $\exists x \varepsilon \tilde{\mathbb{N}}$ or $\exists x^{\text{ent}}$ is defined as :

$\exists x^{\text{ent}} F[x] \equiv (\exists x \varepsilon \tilde{\mathbb{N}}) F[x] \equiv \neg \forall x^{\text{ent}} \neg F[x]$.

Proposition 17 shows that these quantifiers have indeed the intended meaning : the formulas $\forall x^{\text{ent}} F[x]$ and $\forall x (x \varepsilon \tilde{\mathbb{N}} \rightarrow F[x])$ are interchangeable.

Proposition 17.

i) $\lambda x \lambda y \lambda z (y)(x)z \Vdash \forall x^{\text{ent}} F[x] \rightarrow \forall x (\neg F[x] \rightarrow x \notin \tilde{\mathbb{N}})$;

ii) $\lambda x \lambda y (\text{cc}) \lambda k(x) k y \Vdash \forall x (\neg F[x] \rightarrow x \notin \tilde{\mathbb{N}}) \rightarrow \forall x^{\text{ent}} F[x]$.

i) Let $\xi \Vdash \forall x^{\text{ent}} F[x]$, $\eta \Vdash \neg F[a]$ and $\varpi \in \|a \notin \tilde{\mathbb{N}}\|$. Thus, we have $a = s^n 0$ for some $n \in \mathbb{N}$ (else $\|a \notin \tilde{\mathbb{N}}\| = \emptyset$) and $\varpi = \underline{n} \cdot \pi$. We must show that $\eta \star \xi \underline{n} \cdot \pi \in \perp$.

Now, by hypothesis on ξ , we have $\xi \star \underline{n} \cdot \rho \in \perp$ for any $\rho \in \|F[s^n 0]\|$; thus $\xi \underline{n} \Vdash F[s^n 0]$. Since $\eta \Vdash \neg F[s^n 0]$, we have $\eta \star \xi \underline{n} \cdot \pi \in \perp$, which is the desired result.

ii) Let $\xi \Vdash \forall x (\neg F[x] \rightarrow x \notin \tilde{\mathbb{N}})$ and $\underline{n} \cdot \pi \in \|\forall x^{\text{ent}} F[x]\|$, with $n \in \mathbb{N}$ and $\pi \in \|F[s^n 0]\|$.

We have : $\lambda x \lambda y (\text{cc}) \lambda k(x) k y \star \xi \cdot \underline{n} \cdot \pi \succ \xi \star k_\pi \cdot \underline{n} \cdot \pi$.

Now, we have $k_\pi \Vdash \neg F[s^n 0]$ and $\underline{n} \cdot \pi \in \|s^n 0 \notin \tilde{\mathbb{N}}\|$. Therefore $\xi \star k_\pi \cdot \underline{n} \cdot \pi \in \perp$.

Q.E.D.

Theorem 18 (Recurrence scheme). *For every formula $F[\vec{x}, y]$:*

i) $I \Vdash \forall \vec{x} (\forall y (F[\vec{x}, y] \rightarrow F[\vec{x}, sy]), F[\vec{x}, 0] \rightarrow (\forall n \varepsilon \tilde{\mathbb{N}}) F[\vec{x}, n])$.

ii) $I \Vdash \forall \vec{x} (\forall n \varepsilon \tilde{\mathbb{N}}) (\forall y (F[\vec{x}, sy] \rightarrow F[\vec{x}, y]), F[\vec{x}, n] \rightarrow F[\vec{x}, 0])$.

i) This can be written $I \Vdash (\forall n \varepsilon \tilde{\mathbb{N}}) \forall \vec{x} (\forall y (F[\vec{x}, y] \rightarrow F[\vec{x}, sy]), F[\vec{x}, 0] \rightarrow F[\vec{x}, n])$.

Thus, let $n \in \mathbb{N}$, \vec{a} a sequence of d'individuals, $\xi \Vdash \forall y (F[\vec{a}, y] \rightarrow F[\vec{a}, sy])$, $\alpha \Vdash F[\vec{a}, 0]$.

We must show that, for every $\pi \in \|F[\vec{a}, n]\|$, we have $I \star \underline{n} \cdot \xi \cdot \alpha \cdot \pi \in \perp$.

In fact, we show, by recurrence on n , that $\underline{n} \star \xi \cdot \alpha \cdot \pi \in \perp$.

This is immediate if $n = 0$. In order to go from n to $n+1$, we suppose now $\pi \in \|F[\vec{a}, sn]\|$; we have $\underline{n+1} \star \xi \cdot \alpha \cdot \pi \succ \underline{n+1} \star \xi \cdot \alpha \cdot \pi \succ \sigma \star \underline{n} \cdot \xi \cdot \alpha \cdot \pi \succ \xi \star \underline{n} \xi \alpha \cdot \pi$.

But, by the recurrence hypothesis, we have $\underline{n} \xi \alpha \Vdash F[\vec{a}, n]$; by hypothesis on ξ , we have : $\xi \Vdash F[\vec{a}, n] \rightarrow F[\vec{a}, sn]$. Hence the result, since $\pi \in \|F[\vec{a}, sn]\|$.

ii) Almost the same proof : take now $\xi \Vdash \forall y (F[\vec{a}, sy] \rightarrow F[\vec{a}, y])$, $\pi \in \|F[\vec{a}, 0]\|$ and show, by recurrence on n , that $\underline{n} \star \xi \cdot \alpha \cdot \pi \in \perp$, for every $\alpha \Vdash F[\vec{a}, n]$.

Q.E.D.

Definition. We denote by $\text{int}(n)$ the formula $\forall x (\forall y (sy \not\leq x \rightarrow y \not\leq x), n \not\leq x \rightarrow 0 \not\leq x)$.

Theorem 20 shows that the formulas $\text{int}(n)$ and $n \in \tilde{\mathbb{N}}$ are interchangeable, i.e. the formula $\forall n (\text{int}(n) \leftrightarrow n \in \tilde{\mathbb{N}})$ is realized by a proof-like term : this is the *storage theorem for integers*.

Lemma 19. $\lambda g \lambda x (g)(\sigma)x \Vdash \forall y (sy \not\leq \tilde{\mathbb{N}} \rightarrow y \not\leq \tilde{\mathbb{N}})$.

We show that $\lambda g \lambda x (g)(\sigma)x \Vdash sb \not\leq \tilde{\mathbb{N}} \rightarrow b \not\leq \tilde{\mathbb{N}}$ for every individual b .

This is obvious if b is not of the form $s^n 0$, since then $\|b \not\leq \tilde{\mathbb{N}}\| = \emptyset$. Thus, it remains to show :

$\lambda g \lambda x (g)(\sigma)x \Vdash s^{n+1} 0 \not\leq \tilde{\mathbb{N}} \rightarrow s^n 0 \not\leq \tilde{\mathbb{N}}$. Thus, let $\xi \Vdash s^{n+1} 0 \not\leq \tilde{\mathbb{N}}$; we must show :

$\lambda g \lambda x (g)(\sigma)x \star \xi \cdot \underline{n} \cdot \pi \in \perp$, i.e. $\xi \star \sigma \underline{n} \cdot \pi \in \perp$, which is clear, since $\sigma \underline{n} = \underline{n+1}$.

Q.E.D.

Theorem 20 (Storage theorem).

i) $I \Vdash (\forall x \in \tilde{\mathbb{N}}) \text{int}(x)$.

ii) $T \Vdash \forall x (\text{int}(x), x \not\leq \tilde{\mathbb{N}} \rightarrow \perp)$ with $T = \lambda n \lambda f ((n) \lambda g \lambda x (g)(\sigma)x) f 0$.

i) It is theorem 18(ii), if we take for $F[x, y]$ the formula $y \not\leq x$.

ii) Let $\nu \Vdash \text{int}(a)$, $\phi \Vdash a \not\leq \tilde{\mathbb{N}}$ and $\pi \in \Pi$. We must show $T \star \nu \cdot \phi \cdot \pi \in \perp$, that is : $\nu \star \lambda g \lambda x (g)(\sigma)x \cdot \phi \cdot \underline{0} \cdot \pi \in \perp$.

By hypothesis, we have $\nu \Vdash \forall y (sy \not\leq \tilde{\mathbb{N}} \rightarrow y \not\leq \tilde{\mathbb{N}}), a \not\leq \tilde{\mathbb{N}} \rightarrow 0 \not\leq \tilde{\mathbb{N}}$.

But we have $\underline{0} \cdot \pi \in \|0 \not\leq \tilde{\mathbb{N}}\|$ by definition of $\tilde{\mathbb{N}}$ and, by lemma 19 :

$\lambda g \lambda x (g)(\sigma)x \Vdash \forall y (sy \not\leq \tilde{\mathbb{N}} \rightarrow y \not\leq \tilde{\mathbb{N}})$. Hence the result.

Q.E.D.

From theorem 18(i), it follows immediately that the *recurrence scheme of ZF* is realized in \mathcal{N} ; it is the scheme :

$\forall \vec{x} (\forall y (F[\vec{x}, y] \rightarrow F[\vec{x}, sy]), F[\vec{x}, 0] \rightarrow (\forall n \in \tilde{\mathbb{N}}) F[\vec{x}, n])$ for every formula $F[\vec{x}, y]$ of ZF (i.e. written with $\not\leq, \subseteq, 0, s$).

Then, indeed, the formula F is compatible with the extensional equivalence \simeq .

Since the function s is compatible with \simeq , we deduce from lemma 19 that the formula : $\forall y (y \in \tilde{\mathbb{N}} \rightarrow sy \in \tilde{\mathbb{N}})$ is realized in \mathcal{N} ; the formula $0 \in \tilde{\mathbb{N}}$ is also obviously realized.

From the recurrence scheme just proved, we deduce that :

$\tilde{\mathbb{N}}$ is the set of integers of the model \mathcal{N} , considered as a model of ZF.

Theorem 21.

i) Let $f : \mathbb{N}^k \rightarrow \mathbb{N}$ be a recursive function. Then, the formula :

$(\forall x_1 \in \tilde{\mathbb{N}}) \dots (\forall x_k \in \tilde{\mathbb{N}}) f(x_1, \dots, x_k) \in \tilde{\mathbb{N}}$ is realized in \mathcal{N} .

ii) Let $g : \mathbb{N}^k \rightarrow 2$ be a recursive function. Then, the formula :

$(\forall x_1 \in \tilde{\mathbb{N}}) \dots (\forall x_k \in \tilde{\mathbb{N}}) (g(x_1, \dots, x_k) = 1 \vee g(x_1, \dots, x_k) = 0)$ is realized in \mathcal{N} .

i) This can be written $\forall x_1^{\text{ent}} \dots \forall x_k^{\text{ent}} \text{ent}(f(x_1, \dots, x_k))$. The proof is done in [16, 13].

ii) We have $\mathcal{N} \Vdash (\forall x_1 \in \mathbb{J}\mathbb{N}) \dots (\forall x_k \in \mathbb{J}\mathbb{N}) g(x_1, \dots, x_k) \in \mathbb{J}2$.

Now, since g is recursive, we have, by (i) :

$\mathcal{N} \Vdash (\forall x_1 \in \tilde{\mathbb{N}}) \dots (\forall x_k \in \tilde{\mathbb{N}}) g(x_1, \dots, x_k) \in \tilde{\mathbb{N}}$.

Hence the result, by lemma 22.

Q.E.D.

Lemma 22. $\lambda x \lambda y \lambda f(f)xy \Vdash (\forall x \in \mathbb{J}2)(x \neq 1, x \neq 0 \rightarrow x \notin \tilde{\mathbb{N}})$.

We have to show :

$\lambda x \lambda y \lambda f(f)xy \Vdash \top, \perp \rightarrow 0 \notin \tilde{\mathbb{N}}$ and $\lambda x \lambda y \lambda f(f)xy \Vdash \perp, \top \rightarrow 1 \notin \tilde{\mathbb{N}}$.

Thus let $\xi \Vdash \top$ (i.e. $\xi \in \Lambda$ arbitrary) and $\eta \Vdash \perp$. We have to show :

$\lambda x \lambda y \lambda f(f)xy \star \xi \cdot \eta \cdot \underline{0} \cdot \pi \in \perp$ and $\lambda x \lambda y \lambda f(f)xy \star \eta \cdot \xi \cdot \underline{1} \cdot \pi \in \perp$

which is trivial.

Q.E.D.

Remarks. i) In the present paper, theorem 21 is used only in trivial particular cases.

ii) Let us recall the difference between $\mathbb{J}\mathbb{N}$ and $\tilde{\mathbb{N}}$ (the set of integers in the model \mathcal{N}) ; we have :

$\xi \Vdash (\forall x \in \mathbb{J}\mathbb{N}) F[x]$ iff $(\forall n \in \mathbb{N})(\forall \pi \in \|F[s^n 0]\|) \xi \star \pi \in \perp$.

$\xi \Vdash (\forall x \in \tilde{\mathbb{N}}) F[x]$ iff $(\forall n \in \mathbb{N})(\forall \pi \in \|F[s^n 0]\|) \xi \star \underline{n} \star \pi \in \perp$.

Notice that we have $K \Vdash \forall x(x \notin \mathbb{J}\mathbb{N} \rightarrow x \notin \tilde{\mathbb{N}})$, in other words $K \Vdash \tilde{\mathbb{N}} \subset \mathbb{J}\mathbb{N}$. This means that, in \mathcal{N} , the set $\tilde{\mathbb{N}}$ of integers is strongly included in $\mathbb{J}\mathbb{N}$. In the particular realizability model considered below (and, in fact, in every non trivial realizability model) the formula $\mathbb{J}\mathbb{N} \not\subset \tilde{\mathbb{N}}$ is realized.

Non extensional and dependent choice

For each formula $F(x, y_1, \dots, y_m)$ of ZF_ε , we add a function symbol f_F of arity $m + 1$,

with the axiom : $\forall \vec{y}((\forall k \in \tilde{\mathbb{N}}) F[f_F(k, \vec{y}), \vec{y}] \rightarrow \forall x F[x, \vec{y}])$

or else : $\forall \vec{y}(\forall k^{\text{ent}} F[f_F(k, \vec{y}), \vec{y}] \rightarrow \forall x F[x, \vec{y}])$.

It is the *axiom scheme of non extensional choice*, in abbreviated form NEAC.

Remarks. i) The axiom scheme NEAC does not imply the axiom of choice in ZF, because we do not suppose that the symbol f_F is compatible with the extensional equivalence \simeq . It is the reason why we speak about *non extensional* axiom of choice. On the other hand, as we show below, it implies DC (the axiom of dependent choice).

ii) It seems that we could take for f_F a m -ary function symbol and use the following simpler (and logically equivalent) axiom scheme NEAC' : $\forall \vec{y}(F[f_F(\vec{y}), \vec{y}] \rightarrow \forall x F[x, \vec{y}])$.

But this axiom scheme cannot be realized, even though the axiom scheme NEAC is realized by a very simple proof-like term (theorem 23), *provided the instruction ς is present*.

More precisely, we can define a function f_F in \mathcal{M} , such that NEAC is realized in \mathcal{N} , but this is impossible for NEAC'.

Theorem 23 (NEAC).

For each closed formula $\forall x \forall \vec{y} F$, we can define a $(m+1)$ -ary function symbol f_F such that :

$$\lambda x(\zeta)xx \Vdash \forall \vec{y} (\forall k^{\text{ent}} F[f_F(k, \vec{y})/x, \vec{y}] \rightarrow \forall x F[x, \vec{y}]).$$

For each $k \in \mathbb{N}$ we put $P_k = \{\pi \in \Pi; \xi \star \underline{k} \cdot \pi \notin \perp, k = n_\xi\}$.

For each individual x , we have : $\|\forall x F[x, \vec{y}]\| = \bigcup^a \|F[a, \vec{y}]\|$.

Therefore, there exists a function f_F such that, given $k \in \mathbb{N}$ and \vec{y} such that $P_k \cap \|\forall x F[x, \vec{y}]\| \neq \emptyset$, we have $P_k \cap \|F[f_F(k, \vec{y}), \vec{y}]\| \neq \emptyset$.

Now, we want to show $\lambda x(\zeta)xx \Vdash \forall k^{\text{ent}} F[f_F(k, \vec{y}), \vec{y}] \rightarrow F[x, \vec{y}]$, for every individuals x, \vec{y} .

Thus, let $\xi \Vdash \forall k^{\text{ent}} F[f_F(k, \vec{y}), \vec{y}]$ and $\pi \in \|F[a, \vec{y}]\|$; we must show $\lambda x(\zeta)xx \star \xi \cdot \pi \in \perp$.

If this is false, we have $\zeta \star \xi \cdot \xi \cdot \pi \notin \perp$ and therefore $\xi \star \underline{j} \cdot \pi \notin \perp$ with $j = n_\xi$.

It follows that $\pi \in P_j \cap \|F[a, \vec{y}]\|$; thus, there exists $\pi' \in P_j \cap \|F[f_F(j, \vec{y}), \vec{y}]\|$.

Now, we have $\underline{j} \cdot \pi' \in \|\forall k^{\text{ent}} F[f_F(k, \vec{y}), \vec{y}]\|$, and therefore, by hypothesis on ξ , we have : $\xi \star \underline{j} \cdot \pi' \in \perp$. This is a contradiction.

Q.E.D.

NEAC implies DC

Let us call DCS (dependent choice scheme) the following axiom scheme :

$$\forall \vec{z} (\forall x \exists y F[x, y, \vec{z}] \rightarrow \forall n^{\text{ent}} \exists ! y S_F[n, y, \vec{z}] \wedge \forall n^{\text{ent}} \exists y \exists y' \{S_F[n, y, \vec{z}], S_F[sn, y', \vec{z}], F[y, y', \vec{z}]\}).$$

where F is a formula of ZF_ε with free variables x, y, \vec{z} ; the formula S_F is written below.

In the following, we omit the variables \vec{z} (the parameters), for sake of simplicity.

The usual axiom of dependent choice DC is obtained by taking for $F[x, y, z_0, z_1]$ the formula $y \varepsilon z_0 \wedge (x \varepsilon z_0 \rightarrow \langle x, y \rangle \varepsilon z_1)$.

We now show how to define the formula S_F , so that $\text{ZF}_\varepsilon, \text{NEAC} \vdash \text{DCS}$; we shall conclude that DC is realized.

So, let us assume $\forall x \exists y F[x, y]$. By NEAC, there is a function symbol f such that :

$\forall x \exists k^{\text{ent}} F[x, f(k, x)]$. We define the formula $R_F[x, y]$ as follows :

$$R_F[x, y] \equiv \exists k^{\text{ent}} \{F[x, f(k, x)], \forall i^{\text{ent}} (i < k \rightarrow \neg F[x, f(i, x)]), y = f(k, x)\}.$$

This means : “ $y = f(k, x)$ for the first integer k such that $F[x, f(k, x)]$ ”.

Therefore, R_F is functional, i.e. we have $\forall x \exists ! y R_F(x, y)$.

S_F is defined so as to represent a sequence obtained by iteration of the function given by R_F , beginning (arbitrarily) at 0 :

$$S_F(n, x) \equiv \forall z [\forall m \forall y \forall y' (\langle m, y \rangle \varepsilon z, R_F(y, y') \rightarrow \langle sm, y' \rangle \varepsilon z), \langle 0, 0 \rangle \varepsilon z \rightarrow \langle n, x \rangle \varepsilon z].$$

It should be clear that, with this definition of S_F , we obtain :

$$\forall n^{\text{ent}} \exists ! y S_F[n, y] \text{ and } \forall n^{\text{ent}} \exists y \exists y' \{S_F[n, y], S_F[sn, y'], F[y, y']\}.$$

Thus, DCS is provable from ZF_ε and NEAC.

Remark. We have used the binary function symbol $\langle x, y \rangle$ which is defined, in the ground model \mathcal{M} , in the usual way : $\langle a, b \rangle = \{\{a\}, \{a, b\}\}$. Then, the formulas :

$$\forall x \forall x' \forall y \forall y' (\langle x, y \rangle = \langle x', y' \rangle \rightarrow x = x'), \quad \forall x \forall x' \forall y \forall y' (\langle x, y \rangle = \langle x', y' \rangle \rightarrow y = y'),$$

are trivially realized by I .

Properties of the Boolean algebra $\mathfrak{J}2$

Let $(x < y)$ be the binary recursive function defined as follows in \mathcal{M} :
 $(m < n) = 1$ if $m, n \in \mathbb{N}$, $m < n$; else $(m < n) = 0$.

Theorem 24. *For every choice of \perp , the relation $(x < y) = 1$ is, in \mathcal{N} , a strict well founded partial order, which is the usual order on integers (i.e. on $\tilde{\mathbb{N}}$).*

Indeed, the formulas $\forall x((x < x) \neq 1)$; $\forall x \forall y((x < y) = 1 \leftrightarrow (y < x) \neq 1)$ and $\forall x \forall y \forall z((x < y) = 1 \leftrightarrow ((y < z) = 1 \leftrightarrow (x < z) = 1))$ are trivially realized.

Moreover, since the relation $(x < y) = 1$ is well founded, we have (theorem 15) :

$\mathbf{Y} \Vdash \forall x(\forall y((y < x) = 1 \leftrightarrow F[y]) \rightarrow F[x]) \rightarrow \forall x F[x]$

for every formula $F[x]$ with parameters and one free variable.

By theorem 21(ii), the binary recursive function $(x < y)$ sends $\tilde{\mathbb{N}}^2$ into $\{0, 1\}$, in the model \mathcal{N} .

Therefore, it suffices to check that the following formulas are realized in \mathcal{N} :

$(\forall x, y \in \tilde{\mathbb{N}})(y \leq x \rightarrow (x < y) \neq 1)$; $(\forall x, y \in \tilde{\mathbb{N}})(x < y \rightarrow (x < y) = 1)$.

Now the following formulas are trivially realized :

$(\forall x, y, z \in \mathfrak{J}\mathbb{N})(x = y + z \rightarrow (x < y) \neq 1)$; $(\forall x, y, z \in \mathfrak{J}\mathbb{N})(y = x + z + 1 \rightarrow (x < y) = 1)$.

Q.E.D.

Theorem 25.

The following formulas are realized in \mathcal{N} :

i) $(\forall x \in \mathfrak{J}\mathbb{N})(\forall m \in \mathfrak{J}\mathbb{N})((x < m) = 1 \leftrightarrow x \in \mathfrak{J}m)$;

ii) $(\forall m \in \mathfrak{J}\mathbb{N})(\forall n \in \mathfrak{J}\mathbb{N})((m < n) = 1 \rightarrow \mathfrak{J}m \subset \mathfrak{J}n)$.

iii) $(\forall x \in \mathfrak{J}\mathbb{N})(\forall m \in \mathfrak{J}\mathbb{N})((x < m) = 1 \leftrightarrow (\exists y \in \mathfrak{J}\mathbb{N})(m = x + y + 1))$.

Recall that $x \subset y$ is the formula $\forall z(z \notin y \rightarrow z \notin x)$.

i) We have trivially $\|(a < m) \neq 1\| = \|a \notin \mathfrak{J}m\|$ for every $a, m \in \mathbb{N}$.

ii) By transitivity of the relation $(m < n) = 1$ (theorem 24).

iii) We observe that $\|(a < m) \neq 1\| = \|(\forall y \in \mathfrak{J}\mathbb{N})(m \neq a + y + 1)\|$ for every $a, m \in \mathbb{N}$.

Q.E.D.

For each $n \in \mathfrak{J}\mathbb{N}$ (and, in particular, for each $n \in \tilde{\mathbb{N}}$, i.e. for each integer of \mathcal{N}), the set defined, in \mathcal{N} , by $(x < n) = 1$ (the strict initial segment defined by n) is therefore extensionally equivalent to $\mathfrak{J}n$.

Theorem 26. *In \mathcal{N} , the application $(x, y) \mapsto my + x$ is a bijection from $\mathfrak{J}m \times \mathfrak{J}n$ onto $\mathfrak{J}(mn)$. Indeed, the following formulas are realized in \mathcal{N} by I :*

i) $(\forall m, n \in \mathfrak{J}\mathbb{N})(\forall x \in \mathfrak{J}m)(\forall y \in \mathfrak{J}n)(my + x \in \mathfrak{J}(mn))$;

ii) $(\forall m, n \in \mathfrak{J}\mathbb{N})(\forall x, x' \in \mathfrak{J}m)(\forall y, y' \in \mathfrak{J}n)(my + x = my' + x' \leftrightarrow x = x')$;

$(\forall m, n \in \mathfrak{J}\mathbb{N})(\forall x, x' \in \mathfrak{J}m)(\forall y, y' \in \mathfrak{J}n)(my + x = my' + x' \leftrightarrow y = y')$;

iii) $(\forall m, n \in \mathfrak{J}\mathbb{N})(\forall z \in \mathfrak{J}(mn))(\exists x \in \mathfrak{J}m)(\exists y \in \mathfrak{J}n) z = my + x$.

i) and ii) We simply have to replace $(\forall m \in \mathfrak{J}\mathbb{N})$ and $(\forall x \in \mathfrak{J}m)$ with their definitions, which are : $(\forall m \in \mathfrak{J}\mathbb{N}) F \equiv \forall m(1_{\mathbb{N}}(m) = 1 \leftrightarrow F)$; $(\forall x \in \mathfrak{J}m) F \equiv \forall x((x < m) = 1 \leftrightarrow F)$.

We see immediately that these two formulas are realized by I .

iii) We show that :

$$I \Vdash (\forall m, n, z \in \mathbb{N})((\forall x, y \in \mathbb{N})((x < m) = 1 \mapsto ((y < n) = 1 \mapsto z \neq my + x)) \rightarrow (z < mn) \neq 1).$$

Thus, we consider :

$m, n, z_0 \in \mathbb{N}$; $\xi \in \Lambda, \xi \Vdash \forall x \forall y ((x < m) = 1 \mapsto ((y < n) = 1 \mapsto z \neq my + x))$
and $\pi \in \|(z_0 < mn) \neq 1\|$. We must show $I \star \xi \cdot \pi \in \perp$, that is $\xi \star \pi \in \perp$.

We have $\|(z_0 < mn) \neq 1\| \neq \emptyset$, therefore $z_0 < mn$.

Thus, there exists $x_0, y_0 \in \mathbb{N}, x_0 < m, y_0 < n$ such that $z_0 = mx_0 + y_0$. Now, by hypothesis on ξ , we have :

$\xi \Vdash (x_0 < m) = 1 \mapsto ((y_0 < n) = 1 \mapsto z \neq my_0 + x_0)$, in other words $\xi \Vdash \perp$.

Q.E.D.

Injection of \mathbb{N} into $\mathcal{P}(\tilde{\mathbb{N}})$

Recall that we have fixed a recursive bijection : $\xi \mapsto n_\xi$ from Λ onto \mathbb{N} . The inverse bijection will be denoted $n \mapsto \xi_n$.

This bijection is used in the execution rule of the instruction ς , which is as follows :

$$\varsigma \star \xi \cdot \eta \cdot \pi \succ \xi \star \eta \cdot \pi.$$

We define, in \mathcal{M} , a function $\Delta : \mathbb{N} \rightarrow 2$ by putting $\Delta(n) = 0 \Leftrightarrow \xi_n \Vdash \perp$.

Thus, we have defined a function symbol Δ , in the language of ZF_ε . In the realizability model \mathcal{N} , the symbol Δ represents a function from \mathbb{N} into $\mathbb{2}$. In particular, the function Δ sends the set $\tilde{\mathbb{N}}$ of integers of the model \mathcal{N} into the Boolean algebra $\mathbb{2}$.

Lemma 27. *For every $n \in \mathbb{N}$, we have $\xi_n \Vdash \Delta(n) \neq 0$.*

We write this as $\Delta(n) = 0 \Rightarrow \xi_n \Vdash \perp$, which follows from the definition of Δ .

Q.E.D.

Theorem 28. *Let us put $\theta = \lambda x \lambda y (\varsigma) y x x$; then, we have :*

$$\theta \Vdash (\forall x \in \mathbb{2})(x \neq 0 \rightarrow \exists n^{\text{ent}} \{\Delta(n) \neq 0, \Delta(n) \leq x\})$$

where \leq is the order relation of the Boolean algebra $\mathbb{2}$: $y \leq x$ is the formula $x = (y \vee x)$.

We must show $\theta \Vdash (\forall x \in \mathbb{2})(x \neq 0, \forall n^{\text{ent}} (\Delta(n) \neq 0 \rightarrow x \neq \Delta(n) \vee x) \rightarrow \perp)$.

Thus, let $a \in \{0, 1\}$, $\xi \Vdash a \neq 0$, $\eta \Vdash \forall n^{\text{ent}} (\Delta(n) \neq 0 \rightarrow a \neq \Delta(n) \vee a)$ and $\pi \in \Pi$.

We must show $\theta \star \xi \cdot \eta \cdot \pi \in \perp$ that is $\varsigma \star \eta \cdot \xi \cdot \xi \cdot \pi \in \perp$, or else $\eta \star \xi \cdot \xi \cdot \pi \in \perp$.

By hypothesis on η , it suffices to show $n_\xi \cdot \xi \cdot \pi \in \|\forall n^{\text{ent}} (\Delta(n) \neq 0 \rightarrow a \neq \Delta(n) \vee a)\|$.

By lemma 27, we have $\xi \Vdash \Delta(n_\xi) \neq 0$. By definition of the quantifier $\forall n^{\text{ent}}$, it remains to show $\pi \in \|a \neq \Delta(n_\xi) \vee a\|$ or else $a = \Delta(n_\xi) \vee a$.

This is obvious if $a = 1$; if $a = 0$, then $\xi \Vdash \perp$, by hypothesis on ξ .

Therefore $\Delta(n_\xi) = 0$ by definition of Δ , hence the result.

Q.E.D.

By theorem 28, the set $\{\Delta(n); n \in \tilde{\mathbb{N}}, \Delta(n) \neq 0\}$ is, in the realizability model \mathcal{N} , a countable dense subset of the Boolean algebra $\mathbb{2}$: this means that each element $\neq 0$ of this Boolean algebra has a lower bound of the form $\Delta(n) \neq 0$ with $n \in \tilde{\mathbb{N}}$.

It follows that the application of $\mathbb{2}$ into $\mathcal{P}(\tilde{\mathbb{N}})$ given by :

$$x \mapsto \{n \in \tilde{\mathbb{N}}; \Delta(n) \leq x, \Delta(n) \neq 0\}$$

is one to one : indeed, if $a, b \in \mathbb{J}2$ with $a \neq b$, then $a + b \neq 0$; thus, there exists an integer $n \in \tilde{\mathbb{N}}$ such that $\Delta(n) \neq 0$ and $\Delta(n) \leq a + b$. Therefore, we have $\Delta(n) \leq a$ iff $(b \wedge \Delta(n)) = 0$.

But, since $\Delta(n) \neq 0$, we get : $\Delta(n) \leq a$ iff $\Delta(n) \not\leq b$.

We have shown :

Theorem 29.

The formula : “ there exists an injection of $\mathbb{J}2$ into $\mathcal{P}(\tilde{\mathbb{N}})$ ” is realized in the model \mathcal{N} .

Corollary 30. *The formula : “ for every integer n there exists an injection of $\mathbb{J}n$ into $\mathcal{P}(\tilde{\mathbb{N}})$ ” is realized in the model \mathcal{N} .*

Using theorem 26 we see, by recurrence on m , that the model \mathcal{N} realizes the formula : “ $(\forall m \in \tilde{\mathbb{N}}) (\mathbb{J}2)^m$ is equipotent to $\mathbb{J}(2^m)$ ” ; and therefore also the formula :

“ $(\forall m \in \tilde{\mathbb{N}})$ there exists an injection of $\mathbb{J}(2^m)$ into $\mathcal{P}(\tilde{\mathbb{N}})$ ”.

Finally, by theorem 25(ii), we see that the following formula is realized :

“ $(\forall n \in \tilde{\mathbb{N}})$ there exists an injection of $\mathbb{J}n$ into $\mathcal{P}(\tilde{\mathbb{N}})$ ”.

Q.E.D.

Realizability models in which \mathbb{R} is not well ordered

$\mathbb{J}2$ atomless

Theorem 31. *We suppose there exist two proof-like terms ω_0, ω_1 such that, for every $\pi \in \Pi$, we have $\omega_0 k_\pi \Vdash \perp$ or $\omega_1 k_\pi \Vdash \perp$. Then, the Boolean algebra $\mathbb{J}2$ is non trivial. Indeed :*

$\theta \Vdash \forall x (x \neq 1, x \neq 0 \rightarrow x \notin \mathbb{J}2) \rightarrow \perp$ with $\theta = \lambda f (cc) \lambda k ((f)(\omega_1)k)(\omega_0)k$.

Let $\xi \Vdash \forall x (x \neq 1, x \neq 0 \rightarrow x \notin \mathbb{J}2)$ and $\pi \in \Pi$. We must show :

$\theta \star \xi \cdot \pi \in \perp$, that is $\xi \star \omega_1 k_\pi \cdot \omega_0 k_\pi \cdot \pi \in \perp$.

But, by hypothesis on ξ , we have $\xi \Vdash \top, \perp \rightarrow \perp$ and $\xi \Vdash \perp, \top \rightarrow \perp$. Hence the result, by hypothesis on ω_1, ω_0 .

Q.E.D.

Theorem 32. *We suppose that there exists three proof-like terms $\alpha_0, \alpha_1, \alpha_2$ such that, for every $\xi \in \Lambda$ and $\pi \in \Pi$, we have $k_\pi \xi \alpha_0 \Vdash \perp$ or $k_\pi \xi \alpha_1 \Vdash \perp$ or $k_\pi \xi \alpha_2 \Vdash \perp$.*

Then, the Boolean algebra $\mathbb{J}2$ is atomless. Indeed :

$\theta \Vdash \forall x [\forall y (xy \neq 0, xy \neq x \rightarrow y \notin \mathbb{J}2), x \neq 0 \rightarrow x \notin \mathbb{J}2]$

with $\theta = \lambda x \lambda y (cc) \lambda k ((x)(k)y\alpha_0)((x)(k)y\alpha_1)(k)y\alpha_2$.

By a simple computation, we see that we must show :

i) $\theta \Vdash (\perp, \perp \rightarrow \perp), \perp \rightarrow \perp$.

ii) $\theta \Vdash |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|, \top \rightarrow \perp$.

Proof of (i) : let $\eta \in |\perp, \perp \rightarrow \perp|$ and $\xi \in |\perp|$. We must show $\theta \star \eta \cdot \xi \cdot \pi \in \perp$, that is : $\eta \star k_\pi \xi \alpha_0 \cdot ((\eta)(k_\pi)\xi \alpha_1)(k_\pi)\xi \alpha_2 \cdot \pi \in \perp$.

But, from $\xi \Vdash \perp$, we deduce $k_\pi \xi \zeta \Vdash \perp$ for every $\zeta \in \Lambda_c$.

Since $\eta \Vdash \perp, \perp \rightarrow \perp$, we have $((\eta)(\mathbf{k}_\pi)\xi\alpha_1)(\mathbf{k}_\pi)\xi\alpha_2 \Vdash \perp$ and therefore :
 $\eta \star \mathbf{k}_\pi\xi\alpha_0 \cdot ((\eta)(\mathbf{k}_\pi)\xi\alpha_1)(\mathbf{k}_\pi)\xi\alpha_2 \cdot \pi \in \perp$.

Proof of (ii) : let $\eta \in |\top, \perp \rightarrow \perp| \cap |\perp, \top \rightarrow \perp|$ and $\xi \in \Lambda_c$. Again, we must show that :
 $\eta \star \mathbf{k}_\pi\xi\alpha_0 \cdot ((\eta)(\mathbf{k}_\pi)\xi\alpha_1)(\mathbf{k}_\pi)\xi\alpha_2 \cdot \pi \in \perp$. If this is false, then :

$\mathbf{k}_\pi\xi\alpha_0 \not\Vdash \perp$ (because $\eta \Vdash \perp, \top \rightarrow \perp$) and

$((\eta)(\mathbf{k}_\pi)\xi\alpha_1)(\mathbf{k}_\pi)\xi\alpha_2 \not\Vdash \perp$ (because $\eta \Vdash \top, \perp \rightarrow \perp$).

But, since $\eta \Vdash \perp, \top \rightarrow \perp$ (resp. $\top, \perp \rightarrow \perp$), we have $\mathbf{k}_\pi\xi\alpha_1 \not\Vdash \perp$ (resp. $\mathbf{k}_\pi\xi\alpha_2 \not\Vdash \perp$).

This contradicts the hypothesis of the theorem.

Q.E.D.

\mathbb{R} not well orderable

Theorem 33.

We suppose that there exists a proof-like term ω such that, for every $\xi, \xi' \in \Lambda$, $\xi \neq \xi'$ and $\pi \in \Pi$, we have $\omega\mathbf{k}_\pi\xi \Vdash \perp$ or $\omega\mathbf{k}_\pi\xi' \Vdash \perp$.

Then we have, for every formula F with three free variables :

$\theta \Vdash (\forall m, n \in \mathbb{N}) \forall z [(m < n) = 1 \mapsto$

$(\forall x \forall y \forall y' (F(x, y, z), F(x, y', z), y \neq y' \rightarrow \perp), (\forall y \varepsilon \mathbb{J}n) \neg (\forall x \varepsilon \mathbb{J}m) \neg F(x, y, z) \rightarrow \perp)]$

with $\theta = \lambda x \lambda x' (\mathbf{cc}) \lambda k (x') \lambda z (x z z) (\omega) k z$.

Remark. This shows that, if $(m < n) = 1$, then $(\mathbb{J}m \subset \mathbb{J}n)$ and there is no surjection of $\mathbb{J}m$ onto $\mathbb{J}n$: indeed, it suffices to take, as $F(x, y, z)$, the formula $\langle x, y \rangle \varepsilon z$.

Assume this is false ; then, there exist $m, n \in \mathbb{N}$ with $m < n$, an individual c , two terms $\xi, \xi' \in \Lambda$ and a stack $\pi \in \Pi$ such that :

$\theta \star \xi \cdot \xi' \cdot \pi \notin \perp$;

$\xi \Vdash \forall x \forall y \forall y' [F(x, y, c), F(x, y', c), y \neq y' \rightarrow \perp]$;

$\xi' \Vdash (\forall y \varepsilon \mathbb{J}n) \neg (\forall x \varepsilon \mathbb{J}m) \neg F(x, y, z)$.

Therefore, we have $\xi' \star \eta \cdot \pi \notin \perp$ with $\eta = \lambda z (\xi z z) (\omega) \mathbf{k}_\pi z$. By hypothesis on ξ' we have, for every integer $i < n$: $\eta \not\Vdash (\forall x \varepsilon \mathbb{J}m) \neg F(x, i, c)$. Thus, there exists an integer $m_i < m$ such that $\eta \not\Vdash \neg F(m_i, i, c)$. It follows that there exist $\xi_i \in \Lambda$ and $\pi_i \in \Pi$ such that $\xi_i \Vdash F(m_i, i, c)$ and $\eta \star \xi_i \cdot \pi_i \notin \perp$. By definition of η , we get $\xi \star \xi_i \cdot \xi_i \cdot \omega \mathbf{k}_\pi \xi_i \cdot \pi_i \notin \perp$. By hypothesis on ξ , it follows that $\omega \mathbf{k}_\pi \xi_i \not\Vdash i \neq i$; in other words, we have $\omega \mathbf{k}_\pi \xi_i \not\Vdash \perp$ for every integer $i < n$.

By the hypothesis of the theorem, it follows that we have $\xi_i = \xi_j$ for every $i, j < n$.

But, since $m_i < m < n$ and $i < n$, there exist $i, j < n$, $i \neq j$ such that $m_i = m_j = k$. Then, $\xi_i = \xi_j \Vdash F(k, i, c), F(k, j, c)$ and $\omega \mathbf{k}_\pi \xi_i \Vdash i \neq j$ since $\|i \neq j\| = \emptyset$.

Therefore, by hypothesis on ξ , we have $\xi \star \xi_i \cdot \xi_i \cdot \omega \mathbf{k}_\pi \xi_i \cdot \pi_i \in \perp$, which is a contradiction.

Q.E.D.

Now, we see that, with the hypothesis of theorem 33, there is no surjection from $\mathbb{J}2$ onto $\mathbb{J}2 \times \mathbb{J}2$. Indeed, by theorem 26, there exists a bijection from $\mathbb{J}2 \times \mathbb{J}2$ onto $\mathbb{J}4$ and, by theorem 33, there is no surjection from $\mathbb{J}2$ onto $\mathbb{J}4$. It follows that $\mathbb{J}2$ cannot be well ordered.

Now, by theorem 28, $\mathbb{J}2$ is equipotent with a subset of $\mathcal{P}(\tilde{\mathbb{N}})$, and this subset is infinite, by theorem 32. Therefore, the hypothesis of theorems 32 and 33 are sufficient in order that the following formula be realized in the model \mathcal{N} :

There is no well ordering on the set of reals.

In fact, the hypothesis of theorem 33 is sufficient : this follows from theorem 34.

Theorem 34.

Same hypothesis as theorem 33 : there exists a proof-like term ω such that, for every $\pi \in \Pi$ and $\xi, \xi' \in \Lambda$, $\xi \neq \xi'$, we have $\omega k_\pi \xi \Vdash \perp$ or $\omega k_\pi \xi' \Vdash \perp$.

Then we have, for every formula F with three free variables :

$\theta \Vdash \forall z \{ \forall x [\forall n^{ent} F(n, x, z) \rightarrow x \notin \mathbb{N}], \forall n \forall x \forall y [\neg F(n, x, z) \neg F(n, y, z), x \neq y \rightarrow \perp] \rightarrow \perp \}$
with $\theta = \lambda x \lambda x' (cc) \lambda k(x) \lambda n(cc) \lambda h(x'hh) (\omega k) \lambda f(f) hn$.

Remark. This formula means that, in the realizability model \mathcal{N} , there is no surjection from the set of integers $\tilde{\mathbb{N}}$ onto \mathbb{N} : it suffices to take for $F(x, y, z)$ the formula $\langle x, y \rangle \notin z$ (the graph of an hypothetical surjection being $\langle x, y \rangle \varepsilon z$).

Reasoning by contradiction, we suppose that there is an individual c , a stack $\pi \in \Pi$, and two terms ξ, ξ' such that :

$\xi \Vdash \forall x [\forall n^{ent} F(n, x, c) \rightarrow x \notin \mathbb{N}]$; $\xi' \Vdash \forall n \forall x \forall y [\neg F(n, x, z) \neg F(n, y, z), x \neq y \rightarrow \perp]$ and $\theta \star \xi \cdot \xi' \cdot \pi \notin \perp$.

Therefore, we have $\xi \star \eta \cdot \pi \notin \perp$, with $\eta = \lambda n(cc) \lambda h(\xi'hh) (\omega k_\pi) \lambda f(f) hn$.

By hypothesis on ξ , we have $\eta \not\Vdash \forall n^{ent} F(n, 0, c)$ and $\eta \not\Vdash \forall n^{ent} F(n, 1, c)$. Thus, we see that there exist $n_0, n_1 \in \mathbb{N}$, $\pi_0 \in \|F(n, 0, c)\|$ and $\pi_1 \in \|F(n, 1, c)\|$ such that $\eta \star n_0 \cdot \pi_0 \notin \perp$ and $\eta \star n_1 \cdot \pi_1 \notin \perp$. By performing these two processes, we obtain :

$\xi' \star k_{\pi_0} \cdot k_{\pi_0} \cdot \zeta_0 \cdot \pi_0 \notin \perp$ et $\xi' \star k_{\pi_1} \cdot k_{\pi_1} \cdot \zeta_1 \cdot \pi_1 \notin \perp$,

with $\zeta_0 = (\omega k_\pi) \lambda f(f) k_{\pi_0} n_0$ and $\zeta_1 = (\omega k_\pi) \lambda f(f) k_{\pi_1} n_1$.

By hypothesis on ξ' , we have $\xi' \Vdash \neg F(n_0, 0, c), \neg F(n_0, 0, c), 0 \neq 0 \rightarrow \perp$.

Since $k_{\pi_0} \Vdash \neg F(n_0, 0, c)$, we see that $\zeta_0 \not\Vdash \perp$ and, in the same way, $\zeta_1 \not\Vdash \perp$.

Thus, by the hypothesis of the theorem, we have :

$\lambda f(f) k_{\pi_0} n_0 = \lambda f(f) k_{\pi_1} n_1$, and therefore $n_0 = n_1$ and $\pi_0 = \pi_1$.

But, we have $\xi' \Vdash \neg F(n_0, 0, c), \neg F(n_0, 1, c), 0 \neq 1 \rightarrow \perp$. Moreover, we have :

$\pi_0 \in \|F(n_0, 0, c)\|$ and $\pi_1 \in \|F(n_1, 1, c)\|$, thus $\pi_0 \in \|F(n_0, 1, c)\|$ since $n_0 = n_1$, $\pi_0 = \pi_1$.

Therefore $k_{\pi_0} \Vdash \neg F(n_0, 0, c)$ and $\neg F(n_0, 1, c)$. Moreover, we have obviously $\zeta_0 \Vdash 0 \neq 1$, since $\|0 \neq 1\| = \emptyset$. Therefore, we have $\xi' \star k_{\pi_0} \cdot k_{\pi_0} \cdot \zeta_0 \cdot \pi_0 \in \perp$, which is a contradiction.

Q.E.D.

Theorems 33 and 34 show that \mathbb{N} is infinite and not equipotent with $\mathbb{N} \times \mathbb{N}$, thus not well orderable. Since \mathbb{N} is equipotent with a subset of $\mathcal{P}(\tilde{\mathbb{N}})$ (theorem 29), we have shown that $\mathcal{P}(\tilde{\mathbb{N}})$ is not well orderable, with the hypothesis of theorem 33.

More precisely, by corollary 30, we know that \mathbb{N} is equipotent with a subset of $\mathcal{P}(\tilde{\mathbb{N}})$ for each integer n . Therefore, we have :

Theorem 35. With the hypothesis of theorem 33, the following formula is realized :

“ There exists a sequence \mathcal{X}_n of infinite subsets of $\mathcal{P}(\tilde{\mathbb{N}})$ such that, for every integers $m, n \geq 2$:

- there is an injection from \mathcal{X}_n into \mathcal{X}_{n+1} ;
- there is no surjection from \mathcal{X}_n onto \mathcal{X}_{n+1} ;
- $\mathcal{X}_m \times \mathcal{X}_n$ and \mathcal{X}_{mn} are equipotent ”.

For each integer n , the set $\{0, 1, \dots, n-1\}$ is a ring : the ring of integers modulo n ; the Boolean algebra $\{0, 1\}$ is a set of idempotents in this ring. These ring operations extend to the realizability model, giving a ring structure on $\mathbb{J}n$, and $\mathbb{J}2$ is a set of idempotents in $\mathbb{J}n$.

For each $a \in \mathbb{J}2$, the equation $ax = x$ defines an ideal in $\mathbb{J}n$, which we denote as $a\mathbb{J}n$. The application $x \mapsto ax$ is a retraction from $\mathbb{J}n$ onto $a\mathbb{J}n$.

Proposition 36. *The following formulas are realized in \mathcal{N} :*

- i) $(\forall n \in \mathbb{J}\mathbb{N})(\forall a \in \mathbb{J}2)$ (the application $x \mapsto (ax, (1-a)x)$ is a bijection
from $\mathbb{J}n$ onto $a\mathbb{J}n \times (1-a)\mathbb{J}n$).
- ii) $(\forall m, n \in \mathbb{J}\mathbb{N})(\forall a \in \mathbb{J}2)$ (the application $(x, y) \mapsto my + x$ is a bijection
from $a\mathbb{J}m \times a\mathbb{J}n$ onto $a\mathbb{J}(mn)$).

i) Trivial : the inverse is $(y, y') \mapsto y + y'$.

ii) By theorem 26, this application is injective ; clearly, it sends $a\mathbb{J}m \times a\mathbb{J}n$ into $a\mathbb{J}(mn)$. Conversely, if $z \in a\mathbb{J}(mn)$, then there exists $x \in \mathbb{J}m$ and $y \in \mathbb{J}n$ such that $z = my + x$; thus, we have $z = az = may + ax$ with $ax \in a\mathbb{J}m$ and $ay \in a\mathbb{J}n$.

Q.E.D.

Theorem 37. *We suppose that, for each $\alpha \in \Lambda$, $\pi \in \Pi$, and every distinct $\zeta_0, \zeta_1, \zeta_2 \in \Lambda$, we have $k_\pi \alpha \zeta_0 \Vdash \perp$ or $k_\pi \alpha \zeta_1 \Vdash \perp$ or $k_\pi \alpha \zeta_2 \Vdash \perp$.*

Then, for each formula $F(x, y, z)$ with three free variables, we have :

$\theta \Vdash \forall z (\forall m, n \in \mathbb{J}\mathbb{N})(\forall a \in \mathbb{J}2)[(2m < n) = 1 \mapsto$
 $(a \neq 0, \forall x \forall y \forall y' (F(x, y, z), F(x, y', z), y \neq y' \rightarrow \perp), (\forall y \in \mathbb{J}n)(\exists x \in \mathbb{J}m) F(x, ay, z) \rightarrow \perp)]$
with $\theta = \lambda a \lambda x \lambda y (\text{cc}) \lambda k (y) \lambda z (x z z) (k) a z$.

Remark. This formula means that, if $n > 2m$, $a \in \mathbb{J}2$, $a \neq 0$, then there is no surjection from $\mathbb{J}m$ onto $a\mathbb{J}n$: it suffices to take $F(x, y, z) \equiv \langle x, y \rangle \varepsilon z$.

Reasoning by contradiction, let us consider $m, n \in \mathbb{N}$ with $n > 2m$, $a \in \{0, 1\}$, an individual c , three terms $\alpha, \xi, \eta \in \Lambda$ and $\pi \in \Pi$ such that :

$\theta \star \alpha \cdot \xi \cdot \eta \cdot \pi \notin \perp$, $\alpha \Vdash a \neq 0$, $\xi \Vdash \forall x \forall y \forall y' (F(x, y, c), F(x, y', c), y \neq y' \rightarrow \perp)$,
 $\eta \Vdash (\forall y \in \mathbb{J}n) \neg (\forall x \in \mathbb{J}m) \neg F(x, ay, c)$.

We have $\theta \star \alpha \cdot \xi \cdot \eta \cdot \pi \succ \eta \star \theta' \cdot \pi$ and therefore $\eta \star \theta' \cdot \pi \notin \perp$ with $\theta' = \lambda z (\xi z z) (k_\pi) \alpha z$. It follows that, for every $y \in \{0, \dots, n-1\}$, we have $\theta' \not\Vdash (\forall x \in \mathbb{J}m) \neg F(x, ay, c)$.

Thus, there exist two functions $y \mapsto x_y$ (resp. $y \mapsto \zeta_y$) from $\{0, \dots, n-1\}$ into $\{0, \dots, m-1\}$ (resp. into Λ), such that $\zeta_y \Vdash F(x_y, ay, c)$ and $\theta' \star \zeta_y \cdot \varpi_y \notin \perp$ (for some suitable stacks ϖ_y).

Now, we have $\theta' \star \zeta_y \cdot \varpi_y \succ \xi \star \zeta_y \cdot \zeta_y \cdot \kappa_y \cdot \varpi_y$ with $\kappa_y = k_\pi \alpha \zeta_y$; therefore, we have : $\xi \star \zeta_y \cdot \zeta_y \cdot \kappa_y \cdot \varpi_y \notin \perp$ for each $y \in \{0, \dots, n-1\}$.

By hypothesis on ξ (with $y = y'$), it follows that $\kappa_y \not\Vdash \perp$ for every $y < n$.

It follows first that $\alpha \not\Vdash \perp$ and therefore, we have $a = 1$; thus $\zeta_y \Vdash F(x_y, y, c)$.

Moreover, since $n > 2m$, there exist $y_0, y_1, y_2 < n$ distinct, such that $x_{y_0} = x_{y_1} = x_{y_2}$.

But, following the hypothesis of the theorem, the terms $\zeta_{y_0}, \zeta_{y_1}, \zeta_{y_2}$ cannot be distinct, because $\kappa_{y_0}, \kappa_{y_1}, \kappa_{y_2} \not\Vdash \perp$. Therefore we have, for instance, $\zeta_{y_0} = \zeta_{y_1}$; then, we apply the hypothesis on ξ with $y = y_0, y' = y_1$, which gives $\xi \star \zeta_{y_0} \cdot \zeta_{y_1} \cdot \kappa \cdot \varpi \in \perp$ for every $\kappa \in \Lambda$ and $\varpi \in \Pi$. But it follows that $\xi \star \zeta_{y_0} \cdot \zeta_{y_0} \cdot \kappa_{y_0} \cdot \varpi_{y_0} \in \perp$ which is a contradiction.

Q.E.D.

Corollary 38. *With the hypothesis of theorem 37, the following formulas are realized :*

- i) $(\forall n \in \tilde{\mathbb{N}})(\forall a \in \mathbb{J}2)(a \neq 0 \rightarrow \text{there is no surjection from } \mathbb{J}n \text{ onto } a\mathbb{J}(n+1)).$
- ii) $(\forall n \in \tilde{\mathbb{N}})(\forall a, b \in \mathbb{J}2)(ab = 0, b \neq 0 \rightarrow \text{there is no surjection from } a\mathbb{J}n \text{ onto } b\mathbb{J}2).$
- iii) $(\forall n \in \tilde{\mathbb{N}})(\forall a, b \in \mathbb{J}2)(ab = a, a \neq b \rightarrow \text{there is no surjection from } a\mathbb{J}n \text{ onto } b\mathbb{J}2).$

i) We prove this formula by contradiction : if there is a surjection from $a\mathbb{J}n$ onto $a\mathbb{J}(n+1)$ then, for each integer $k \in \tilde{\mathbb{N}}$, there exists a surjection from $(\mathbb{J}n)^k$ onto $(a\mathbb{J}(n+1))^k$; and thus also a surjection from $\mathbb{J}(n^k)$ onto $a\mathbb{J}((n+1)^k)$ (theorem 36). But, for $k > n$, we have $(n+1)^k > 2n^k$ and this contradicts theorem 37.

ii) Indeed, we have $(a+b)\mathbb{J}n = a\mathbb{J}n \times b\mathbb{J}n$. Reasoning by contradiction, there would exist a surjection from $(a+b)\mathbb{J}n$ onto $b\mathbb{J}2 \times b\mathbb{J}n$, thus also onto $b\mathbb{J}(2n)$ (theorem 36), thus a surjection from $\mathbb{J}n$ onto $b\mathbb{J}(2n)$, which contradicts (i).

iii) Otherwise, it would exist a surjection from $a\mathbb{J}n$ onto $(b-a)\mathbb{J}2$, which contradicts (ii).

Q.E.D.

Application. By DC, since $\mathbb{J}2$ is atomless, there exists in $\mathbb{J}2$ a strictly decreasing sequence. By corollary 38(iii) and theorem 29, there exists a sequence of infinite subsets of $\mathcal{P}(\tilde{\mathbb{N}})$, the “cardinals” of which are strictly decreasing. More precisely, let \mathcal{B} be the image of $\mathbb{J}2$ by the injection in $\mathcal{P}(\tilde{\mathbb{N}})$ given by theorem 29 ; then we have :

Theorem 39. *With the hypothesis of theorem 37, the following formula is realized in \mathcal{N} :*

“There exists a subset \mathcal{B} of $\mathcal{P}(\tilde{\mathbb{N}})$ (the real line of the model \mathcal{N}), such that \mathcal{B} is an atomless Boolean algebra for the usual order \subseteq on $\mathcal{P}(\tilde{\mathbb{N}})$, with $\emptyset, \mathbb{N} \in \mathcal{B}$; $a, b \in \mathcal{B} \Rightarrow a \cap b \in \mathcal{B}$.

If $a \in \mathcal{B}, a \neq \emptyset$ then $a\mathcal{B}$ is infinite and there is no surjection from \mathcal{B} onto $a\mathcal{B} \times a\mathcal{B}$ (where $a\mathcal{B}$ means $\{x \in \mathcal{B}; x \subseteq a\}$).

If $a, b \in \mathcal{B}, a, b \neq \emptyset$ and $a \cap b = \emptyset$, then there is no surjection from $a\mathcal{B}$ onto $b\mathcal{B}$ (the “cardinals” of $a\mathcal{B}, b\mathcal{B}$ are incomparable).

If $a, b \in \mathcal{B}, a \subseteq b$ and $a \neq b$, then there is no surjection from $a\mathcal{B}$ onto $b\mathcal{B}$ (the “cardinal” of $a\mathcal{B}$ is strictly less than the “cardinal” of $b\mathcal{B}$).”.

In other words, for $a, b \in \mathcal{B}$, we have : $a \subseteq b \Leftrightarrow$ there exists a surjection from $b\mathcal{B}$ onto $a\mathcal{B}$. The order, in the atomless Boolean algebra \mathcal{B} , is the order on the “cardinals” of its initial segments, like in a finite Boolean algebra.

The model of threads

This model is the canonical instance of a non trivial coherent realizability model. It is defined as follows :

Let $n \mapsto \pi_n$ be an enumeration of the *stack constants* and let $n \mapsto \theta_n$ be a recursive enumeration of the *proof-like terms*. For each $n \in \mathbb{N}$, the *thread of number n* is the set of processes which appear during the execution of the process $\theta_n \star \pi_n$.

Note that every term which appears in the n -th thread contains the only stack constant π_n .

We define \perp^c (the complement of \perp) as the union of the threads.

Therefore, we have $\xi \star \pi \in \perp$ iff the process $\xi \star \pi$ never appears in any thread.

For every term ξ , we have $\xi \Vdash \perp$ iff ξ never appears in head position in any thread.

If ξ is a proof-like term, we have $\xi = \theta_n$ for some integer n , and therefore $\xi \star \pi_n \notin \perp$, by definition of \perp . It follows that *the model of threads is coherent*.

If $\xi \in \Lambda$, $\xi \not\perp$ then ξ appears in head position in at least one thread. This thread is unique, unless ξ is a proof-like term, because it is determined by the number of any stack constant which appears in ξ .

Theorem 40. *The hypothesis of theorems 31, 32, 33 and 37 are satisfied in the model of threads.*

The hypothesis of theorems 33 and 31 are trivially satisfied if we take :

$\omega = (\lambda x xx)\lambda x xx$, $\omega_0 = \omega 0$, and $\omega_1 = \omega 1$.

Moreover, the hypothesis of theorem 37 is obviously stronger than the hypothesis of theorem 32.

We check the hypothesis of theorem 37 by contradiction : we suppose $k_\pi \alpha \zeta_0 \not\perp$, $k_\pi \alpha \zeta_1 \not\perp$ and $k_\pi \alpha \zeta_2 \not\perp$. Therefore, these three terms appear in head position, and moreover in the same thread : indeed, since they contain the stack π , this thread has the same number as the stack constant of π .

Let us consider their first appearance in head position, for instance with the order 0, 1, 2.

Therefore we have, in this thread : $k_\pi \alpha \zeta_0 \star \rho_0 \succ \alpha \star \pi \succ \dots \succ k_\pi \alpha \zeta_1 \star \rho_1 \succ \alpha \star \pi \succ \dots$

But, at the second appearance of $\alpha \star \pi$, the thread enters into a loop, and the term $k_\pi \alpha \zeta_2$ can never arrive in head position, since $\zeta_1 \neq \zeta_2$.

Q.E.D.

References

- [1] S. Berardi, M. Bezem, T. Coquand. *On the computational content of the axiom of choice*. J. Symb. Log. 63 (1998), p. 600-622.
- [2] H.B. Curry, R. Feys. *Combinatory Logic*. North-Holland (1958).
- [3] W. Easton. *Powers of regular cardinals*. Ann. Math. Logic 1 (1970), p. 139-178.
- [4] J.Y. Girard. *Une extension de l'interprétation fonctionnelle de Gödel à l'analyse*. Proc. 2nd Scand. Log. Symp. (North-Holland) (1971) p. 63-92.
- [5] T. Griffin. *A formulæ-as-type notion of control*. Conf. record 17th A.C.M. Symp. on Principles of Progr. Languages (1990).
- [6] S. Grigorieff. *Combinatorics on ideals and forcing*. Ann. Math. Logic 3(4) (1971), p. 363-394.
- [7] W. Howard. *The formulas-as-types notion of construction*. Essays on combinatory logic, λ -calculus, and formalism, J.P. Seldin and J.R. Hindley ed., Acad. Press (1980) p. 479-490.
- [8] J. M. E. Hyland. *The effective topos*. The L.E.J. Brouwer Centenary Symposium (Noordwijkerhout, 1981), 165-216, Stud. Logic Foundations Math., 110, North-Holland, Amsterdam-New York, 1982.

- [9] G. Kreisel. *On the interpretation of non-finitist proofs I.*
J. Symb. Log. 16 (1951) p. 248-26.
- [10] G. Kreisel. *On the interpretation of non-finitist proofs II.*
J. Symb. Log. 17 (1952), p. 43-58.
- [11] J.-L. Krivine. *Typed lambda-calculus in classical Zermelo-Fraenkel set theory.*
Arch. Math. Log., 40, 3, p. 189-205 (2001).
http://www.pps.jussieu.fr/~krivine/articles/zf_epsilon.pdf
- [12] J.-L. Krivine. *Dependent choice, 'quote' and the clock.*
Th. Comp. Sc., 308, p. 259-276 (2003).
<http://hal.archives-ouvertes.fr/hal-00154478>
<http://www.pps.jussieu.fr/~krivine/articles/quote.pdf>
- [13] J.-L. Krivine. *Realizability in classical logic.*
In *Interactive models of computation and program behaviour.*
Panoramas et synthèses, Société Mathématique de France, 27 (2009).
<http://hal.archives-ouvertes.fr/hal-00154500>
Version mise à jour à :
<http://www.pps.jussieu.fr/~krivine/articles/Luminy04.pdf>
- [14] J.-L. Krivine. *Realizability : a machine for Analysis and set theory.*
Geocal'06 (fevrier 2006 - Marseille); Mathlogaps'07 (juin 2007 - Aussois).
<http://cel.archives-ouvertes.fr/cel-00154509>
Version mise à jour à :
<http://www.pps.jussieu.fr/~krivine/articles/Mathlog07.pdf>
- [15] J.-L. Krivine. *Structures de réalisabilité, RAM et ultrafiltre sur \mathbb{N} .* (2008)
<http://hal.archives-ouvertes.fr/hal-00321410>
<http://www.pps.jussieu.fr/~krivine/articles/Ultrafiltre.pdf>
- [16] J.-L. Krivine. *Realizability algebras : a program to well order \mathbb{R} .* (2010)
<http://hal.archives-ouvertes.fr/hal-00483232>
http://www.pps.jussieu.fr/~krivine/articles/Well_order.pdf