



**HAL**  
open science

## Thermodynamic and spectroscopic study for the interaction of dimethyltin(IV) with L-cysteine in aqueous solution

Paola Cardiano, Concetta de Stefano, Ottavia Giuffrè, Silvio Sammartano

► **To cite this version:**

Paola Cardiano, Concetta de Stefano, Ottavia Giuffrè, Silvio Sammartano. Thermodynamic and spectroscopic study for the interaction of dimethyltin(IV) with L-cysteine in aqueous solution. *Biophysical Chemistry*, 2008, 133 (1-3), pp.19. 10.1016/j.bpc.2007.11.005 . hal-00501692

**HAL Id: hal-00501692**

**<https://hal.science/hal-00501692>**

Submitted on 12 Jul 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript

Thermodynamic and spectroscopic study for the interaction of dimethyltin(IV) with L-cysteine in aqueous solution

Paola Cardiano, Concetta De Stefano, Ottavia Giuffrè, Silvio Sammartano


PII: S0301-4622(07)00278-5  
DOI: doi: [10.1016/j.bpc.2007.11.005](https://doi.org/10.1016/j.bpc.2007.11.005)  
Reference: BIOCHE 5048

To appear in: *Biophysical Chemistry*

Received date: 24 October 2007  
Revised date: 15 November 2007  
Accepted date: 15 November 2007

Please cite this article as: Paola Cardiano, Concetta De Stefano, Ottavia Giuffrè, Silvio Sammartano, Thermodynamic and spectroscopic study for the interaction of dimethyltin(IV) with L-cysteine in aqueous solution, *Biophysical Chemistry* (2007), doi: [10.1016/j.bpc.2007.11.005](https://doi.org/10.1016/j.bpc.2007.11.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Submitted to: Biophysical Chemistry

**Thermodynamic and spectroscopic study for the interaction of  
dimethyltin(IV) with L-cysteine in aqueous solution**

**Paola Cardiano, Concetta De Stefano, Ottavia Giuffrè, Silvio Sammartano\***

Dipartimento di Chimica Inorganica, Chimica Analitica e Chimica Fisica. Università di Messina.

Salita Sperone 31, Villaggio S. Agata, 98166 Messina (Italy)

---

\* Author for correspondence: [ssammartano@unime.it](mailto:ssammartano@unime.it)

**Abstract**

Thermodynamic and spectroscopic properties of the species formed by dimethyltin(IV) cation with L-cysteine (cys) were studied by potentiometric, calorimetric, UV and NMR investigations in aqueous solution. The resulting speciation model showed the formation of five complex species:  $(\text{CH}_3)_2\text{Sn}(\text{cys})\text{H}^+$ ,  $(\text{CH}_3)_2\text{Sn}(\text{cys})^0$ ,  $(\text{CH}_3)_2\text{Sn}(\text{cys})\text{OH}^-$ ,  $(\text{CH}_3)_2\text{Sn}(\text{cys})_2\text{H}^-$ ,  $(\text{CH}_3)_2\text{Sn}(\text{cys})_2^{2-}$ . The stability and the formation percentages, for the mononuclear mixed species in particular, are very high, in a wide pH range. Thermodynamic parameters indicate that the enthalpy values are exothermic and the enthalpic contribution to the stability is higher than entropic one. Individual UV spectra of cys and dimethyltin(IV)-cys species were calculated. Spectroscopic results of UV and  $^1\text{H}$  NMR investigations fully confirm the speciation model. The structures calculated from NMR investigations show that all the species have an eq- $(\text{CH}_3)_2$ -tbp structure.

**Keywords:** dimethyltin(IV) complexes; thermodynamic parameters; potentiometry; calorimetry; spectroscopy.

## Introduction

The study of the interactions between metal cations and amino acids is of fundamental relevance for the knowledge of natural occurring metalloprotein chelation as well as for human and environmental toxicology. The most relevant amino acids, able to bind metal cations, are histidine, cysteine, glutamic and aspartic acids [1]. **Cys** is the only naturally occurring sulfur-containing amino acid. It is a component of the antioxidant glutathione (L- $\gamma$ -glutamyl-L cysteinyl-glycine). A living organism is able to synthesize **cys** from the essential amino acid methionine under normal physiologic conditions; furthermore, among the amino acids considered in blood plasma models it is present at the highest concentration, with a value  $\geq 100 \mu\text{mol L}^{-1}$  [2]. **Cys** and its derivatives aroused particular interest owing to their involvement in many important biological processes and to be an active site both in the catalytic function of the enzymes cysteine proteases and in several peptides and proteins [3]. It is recognized that **cys** can coordinate metal cations through sulfur sites in several proteins and metalloenzymes [4-6].

In several investigations alkyltin(IV) cations, as metals, were used for studying their interaction with biological ligands. This choice is induced by the widespread use of organotin(IV) compounds in industrial, agricultural and biological field, which caused their release and accumulation into the environment and consequently in biological systems. It is well known that these alkyltin(IV) compounds are generally toxic and their toxicity depends on the nature of the alkyl groups bound to the tin(IV) atom and it is directly proportional to their number. In contrast, some others organotin compounds have been found active *in vitro* against human tumour cell lines [7]. For this reason the solution and structural studies on the

interaction of the organotin moieties with several biologically relevant compounds such as carbohydrates, amino acids, peptides, nucleic acids and DNA has been carried out mostly in the last decade [8-21]. From some of these cited studies, the interaction of dimethyltin(IV) and trialkyltin(IV) (with R = methyl, ethyl, and butyl groups) in aqueous solution, at physiological pH, with thiol groups of model molecules, as well as feline and rat haemoglobin, have been investigated by  $^{119}\text{Sn}$  Mössbauer spectroscopy. The results showed that the above species form tetrahedral or trigonal bipyramidal tin sites, with covalent Sn-S bonds and further coordination by nitrogen donors, from **cys** and histidine side chain, respectively [14, 15]. Since for understanding biological action of alkyltin chemicals, their speciation in biological systems must be known, it seems of interest to study quantitatively the interactions between alkyltin(IV) cation and **cys**. Thermodynamic data on metal-**cys** complexes are extremely poor. In particular, no calorimetric study has been performed. The only available data obtained by the van't Hoff equation, regard several metals, such as  $\text{Pb}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Hg}^{2+}$  and others, but there are no enthalpic data on tin or alkyltin cations [21]. Spectroscopic studies on the characterization of several organotin(IV) derivatives of **cys** [17, 18]; potentiometric [19, 20] and Mössbauer [19] investigations on the coordination of **cys** and its derivatives towards trialkyltin(IV) cations can be found in literature.

This work represents a further step in our systematic investigations on the speciation of mono-, di- and tri-methyltin(IV) cations and their interactions with several carboxylic ligands in aqueous solution [22-30]. In this paper we give a complete picture of the thiolic group influence on the interaction of dimethyltin(IV) cation with **cys** by determining thermodynamic parameters by potentiometric and calorimetric measurements, describing the speciation in aqueous solution and confirming findings by spectroscopic (UV and  $^1\text{H}$  NMR) investigations.

## 1. Experimental

### 1.1. Chemicals

Fresh dimethyltin(IV) solutions were prepared every day by weighing dichloride salt (Fluka product), twice re-crystallised before use. Cys (microselect >99.5%) (Fluka product) was used without further purification and its purity, which was always higher than 99.5%, was checked by potentiometric titrations with standard solutions of NaOH. Sodium hydroxide and hydrochloric acid solutions were prepared by diluting concentrated Fluka ampoules and were standardised against potassium biphthalate and sodium carbonate, respectively. All the solutions were prepared using analytical grade water (resistivity = 18 M $\Omega$  cm) and grade A glassware.

### 1.2. Potentiometric apparatus and procedure

Potentiometric titrations were carried out (at 25.0 $\pm$ 0.1 $^{\circ}$ C) using an apparatus consisting of a Metrohm Model 713 potentiometer, equipped with a combined glass electrode (Ross type 8102, from Orion) and a Metrohm Model 765 motorized burette. Estimated accuracy was  $\pm$ 0.2 mV and  $\pm$ 0.003 mL for e.m.f. and titrant volume readings, respectively. The apparatus was connected to a PC, and automatic titrations were performed using a suitable computer program to control titrant delivery, data acquisition and to check for e.m.f. stability. All titrations were carried out under magnetic stirring and presaturated N<sub>2</sub> was bubbled through

the purified solution in order to exclude O<sub>2</sub> and CO<sub>2</sub> inside. A volume of 25 mL of the solution containing *cys*, dimethyltin(IV) dichloride and NaCl was titrated with standard NaOH up to 80-90% neutralization. Details of experimental measurements are reported in Table 1. Separate titrations of HCl at the same ionic strength as the sample under study, were carried out to determine standard electrode potential  $E^0$  and to obtain pH = -log [H<sup>+</sup>] readings. The reliability of the calibration in the alkaline range was checked by calculating p*K*<sub>w</sub> values.

### *1.3. Calorimetric apparatus and procedure*

The measurements were carried out using a Tronac model 450 Iso-peribolic Titration calorimeter, coupled with a Keithley 196 system Dmm digital multimeter. A volume of 50 mL of solution, containing dimethyltin(IV) dichloride at 25.000±0.001 °C was titrated with a solution of Na<sub>3</sub>*cys*. Details of experimental measurements are reported in Table 1. The titrant was delivered by a 2.5 mL capacity Hamilton syringe, model 1002TLL. A computer program was used for the acquisition of calorimetric data. Accuracy was checked by titrating a THAM (tris-(hydroxymethyl)amino-methane) buffer with HCl. The heat of dilution was measured before each experiment. The accuracy of calorimetric apparatus was Q±0.008 J and v±0.001 cm<sup>3</sup>.

### *1.4. UV spectra*

The UV spectra were recorded using a Varian Cary 50 UV-VIS spectrophotometer with an optic fibre probe having a fixed 1 cm path length, from 220 to 280 nm. The spectrophotometer was connected to a PC for the acquisition of the spectra. The measurement

cell and potentiometric apparatus are the same of those already described in the *potentiometric apparatus and procedure* paragraph. A volume of 25 mL of the solutions containing ligand or metal-ligand were prepared and analysed in a wide pH range values. Small increments of standard NaOH solution were added to each solution in order to obtain the maximum absorbance of each predominant species, according to the preliminarily determined speciation diagrams of the system. Details of experimental measurements are reported in Table 1.

### 1.5. NMR measurements

$^1\text{H}$  NMR spectra were recorded on a Bruker AMX R-300 spectrometer. The proton chemical shifts were measured with respect to 1,4-dioxane as reference and converted relative to TMS, using  $\delta_{\text{dioxane}} = 3.70$  ppm. Measurements were generally made in a 9:1  $\text{H}_2\text{O}:\text{D}_2\text{O}$  solution. The ligand and metal concentrations were varied in the range 0.005-0.01 mol  $\text{L}^{-1}$ . In order to obtain the individual NMR parameters ( $\delta$ ,  $^2\text{J}$ ) of the various species, the protonic spectra were recorded at different pH ranging from 2 to 11. The individual chemical shifts as well as  $^2\text{J}(^{119}\text{Sn}-^1\text{H})$  coupling constants belonging to the dimethyltin(IV)-cys complexes were calculated assuming fast mutual exchange [12]. The heteronuclear couplings of the tin-bound methyl groups  $^2\text{J}(^{119}\text{Sn}-^1\text{H})$  obtained in such a way were “converted” into C-Sn-C angles by using the published equation [31].

### 1.6. Calculations

All the parameters relative to alkalimetric purity determination were refined using the nonlinear least squares computer program ESAB2M. Formation constants were refined using the nonlinear least squares computer programs STACO and BSTAC. Speciation profiles were obtained using the computer program ES4EC. Details of calculation methods and programs have already been reported [32].

The interactions of dimethyltin(IV) with small amounts of  $\text{Cl}^-$  from the dimethyltin(IV) dichlorides and from NaCl used as ionic medium were taken into account in the calculations. Formation constants were extrapolated to zero ionic strength as already reported [33, 34]. Both BSTAC and STACO computer programs can deal with potentiometric data obtained in variable ionic strength conditions. The dependence on ionic strength of formation constants was taken into account using the Debye-Hückel type equation:

$$\log\beta = \log^T\beta - z^* \sqrt{I} / (2+3\sqrt{I}) + CI \quad (1)$$

where

$$C = (0.025 \pm 0.015) z^*, \text{ at } I \leq 0.15$$

$$z^* = \sum z_{\text{reactants}}^2 - \sum z_{\text{products}}^2$$

( $\beta$  = stoichiometric formation constant;  ${}^T\beta$  = formation constant at infinite dilution;  $z$  is the component charge).

Calorimetric titration data were analysed by the computer program ES5CM [35]. UV spectra were analysed by the Hyperquad 2000 program [36], which allows to calculate stability constants and molar absorbance spectrum of each absorbing species, using experimental absorbances, analytical concentrations of reagents and the proposed chemical model as input. NMR calculations were performed using the general linear and nonlinear least squares computer program LIANA [32]. Concentrations and thermodynamic parameters are given in the molar scale. Errors are given as standard deviations. The overall formation equilibria are expressed as (charges omitted)



Where  $r > 0$  for protonated species and  $r < 0$  for hydrolytic species.

Thermodynamic parameters of protonation of **cys** used in the calculations, showed in Table 2 at  $I = 0 \text{ mol L}^{-1}$  and  $T=25^\circ\text{C}$ , have been published in the ref. 37. Dimethyltin(IV) cation has a strong tendency to hydrolyze in aqueous solution, forming five hydrolytic species  $[(\text{CH}_3)_2\text{Sn}(\text{OH})^+; (\text{CH}_3)_2\text{Sn}(\text{OH})_2^0; (\text{CH}_3)_2\text{Sn}(\text{OH})_3^-; [(\text{CH}_3)_2\text{Sn}]_2(\text{OH})_2^{2+}; [(\text{CH}_3)_2\text{Sn}]_2(\text{OH})_3^+$ , whose thermodynamic parameters are reported in Table 3 together with those regarding the complexes with  $\text{Cl}^-$ .

## 2. Results and discussion

### 2.1. Potentiometric and calorimetric measurements and speciation model

The stoichiometries and the stability constants of the species formed in the system dimethyltin(IV)-cys were obtained by performing on the potentiometric data several trials with different sets of complex species. The selected speciation model was that which had the best statistical fit, without giving systematic drifts in the residuals. This model very well fits experimental data, see for example Fig. s.1 in Supporting information. The results showed the formation of five complex species between  $(\text{CH}_3)_2\text{Sn}^{2+}$  (M) and cys (L): three mononuclear, such as MLH, ML, MLOH and two binuclear, such as  $\text{ML}_2\text{H}$  and  $\text{ML}_2$ . The overall thermodynamic formation parameters values referring to reaction 2 are reported in Table 4. The thermodynamic formation parameter values obtained on the basis of the most probable reaction were collected in Table 5. The analysis of the thermodynamic results reported in Tables 4, 5 indicates that all enthalpy values are exothermic and in some cases the enthalpic contribution is higher than the entropic one. The errors associated to the enthalpic values are acceptable, except for  $\text{ML}_2\text{H}$  species, which is affected by a high error, so it seems convenient to indicate approximately this value. LogK values, referring mainly to the mononuclear species, show a very high stability, that allows to hypothesize high formation percentages in a wide pH range.

The formation of MLH species in the acidic pH range ( $2 \leq \text{pH} \leq 4$ ) is confirmed by the results obtained for the same system by Buzás et al., Silvestri et al. [18, 19] and in the system

containing trialkyltin(IV) studied by Hynes and O'Dowd [20]. Further deprotonation of **cys**, promoted by metal cation, occurs at higher pH values ( $5 \leq \text{pH} \leq 7$ ), where the ML species is formed. At  $6 \leq \text{pH} \leq 7$ , when the L/M ratio increases, the species  $\text{ML}_2\text{H}$  becomes more significant. In the alkaline range ( $8 \leq \text{pH} \leq 11$ ) the hydroxo-complex MLOH is predominant, while the  $\text{ML}_2$  species, present only in our speciation model, prevails when M/L concentration ratio is at least equal to 0.75 or lower than this value.

By analysing the distribution of the species  $(\text{CH}_3)_2\text{Sn}^{2+}$ -**cys** vs. pH, it arises that the formation percentages of the mononuclear mixed species are very high in all the pH range investigated ( $2 \leq \text{pH} \leq 11$ ), suppressing almost completely the hydrolysis of dimethyltin(IV) cation, as confirmed by speciation diagrams (Fig. 1a-c). When  $C_{(\text{CH}_3)_2\text{Sn}} = 5 \text{ mmol L}^{-1}$  and  $C_{\text{cys}} = 10 \text{ mmol L}^{-1}$  (Fig. 1a), in the acidic range  $2.1 \leq \text{pH} \leq 4.1$  the predominant species MLH achieves a maximum percentage of 82.9 at  $\text{pH} = 3.05$ , while in the same range the hydrolytic MOH species has a maximum percentage of 6.0. In the wide range  $4.2 \leq \text{pH} \leq 7.0$  the predominant species is ML and at  $\text{pH} = 7.0$  its formation percentage is 92.9. In the alkaline range  $8.0 \leq \text{pH} \leq 9.5$ , where the  $\text{ML}_2$  and MLOH species are present in high percentages: the former reaches 59.0 at  $\text{pH} = 9.2$  and this percentage lowers to 8.6 and 3.1 at  $C_{(\text{CH}_3)_2\text{Sn}} = C_{\text{cys}} = 1$  and  $0.1 \text{ mmol L}^{-1}$ , respectively (Figs. 1b-c). The decreasing of the concentration of the metal and the ligand induces a lowering of MLH as well as an increase of the hydrolytic species, such as MOH and  $\text{M}(\text{OH})_2$  formation percentages; on the contrary, the percentages referring to MLOH species remain very high throughout alkaline range. For example, at dimethyltin(IV)/**cys** concentration ratio = 1 and  $8 \leq \text{pH} \leq 11$ , the MLOH species is absolutely predominant: at  $\text{pH} = 9.5$  its formation percentage is 87.3 at  $1 \text{ mmol L}^{-1}$  and 76.8 at  $0.1 \text{ mmol L}^{-1}$ .

In Table s.1 of Supporting information we report the percentages of the species in a pH range, which includes the physiological one (pH = 7.4) and those of interest for many natural waters. The pH values reported are: 7; 7.5; 8.0 and the concentration ratios dimethyltin(IV)/*cys*: 0.5 and 1.

## 2.2. UV measurements

Before studying the UV properties of a complicated system containing metal complexes of an absorbing ligand it was necessary to determine the molar absorbance spectrum of each protonated ligand form. The only paper [17] on the UV properties of complexes formed by *cys* with dimethyltin(IV) merely reports  $\lambda_{\max}$  and  $\epsilon_{\max}$  values, obtained for solutions containing *cys* and dimethyltin(IV)-*cys*, at a specific pH value, without any hypothesis of speciation model and consequently without calculating the individual spectra of the species.

The molar absorbance spectra of protonated and unprotonated species of the ligand were calculated by using aqueous solutions of *cys* at different concentrations by means of spectrophotometric investigations in the 220-280 nm spectral range, by varying the pH through alkalimetric titration. While preliminary tests revealed that solutions containing dimethyltin(IV) cation do not absorb in the spectral range investigated, the UV measurements on the *cys* solutions showed that the absorbance intensity and the position of absorption maximum are clearly affected by the degree of the ligand protonation (Fig. 2). Experimental spectra referring to solutions containing *cys*, at  $C = 0.4 \text{ mmol L}^{-1}$ , show very low absorbance intensity from pH = 1.5 (%  $\text{LH}_3 = 81.4$  and %  $\text{LH}_2 = 18.6$ ) to pH = 6.5. On the contrary in the  $8.9 \leq \text{pH} \leq 11.5$  range they show large spectral variations: 1. by increasing both pH and ligand deprotonation, absorbance values increase; 2. with the predominating of L species on the LH

one, a batocromic shift from  $\lambda_{\max} = 230$  nm to  $\lambda_{\max} = 235$  nm occurs. As examples: at pH = 8.9, where the formation percentages of the species are 18.9 e 75.8 for LH<sub>2</sub> and LH respectively, the absorbance intensity observed is  $A = 1.14$  at the  $\lambda_{\max} = 231$  nm; at pH = 9.5, where % LH<sub>2</sub> = 4.6, % LH = 74.2 and % L = 21.2,  $A = 1.35$ , at the  $\lambda_{\max} = 232$  nm; at pH = 11.0, % LH = 9.3 and % L = 90.6,  $A = 1.74$  at the  $\lambda_{\max} = 235$  nm. Individual UV spectra of the four species of **cys**, namely L<sup>2-</sup>, HL<sup>-</sup>, H<sub>2</sub>L<sup>0</sup>, H<sub>3</sub>L<sup>+</sup> are obtained by deconvolution of experimental spectra into the component gaussian peaks, as shown in Fig. 3. Those relating to the L<sup>2-</sup> and HL<sup>-</sup> species display a higher molar absorbances with respect to LH<sub>2</sub> and LH<sub>3</sub> species, to attribute to the ionisation of the SH group, *i.e.* n→σ\* transition of the –S– chromophore, while the deprotonation of carboxylic and aminic group produces a little spectral variation in molar absorbance in the range investigated. Precisely, at  $\lambda_{\max} = 235$  nm, L and LH species, reach  $\epsilon_{\max} = 4279, 4345$  L mol<sup>-1</sup> cm<sup>-1</sup>, respectively; while, at  $\lambda_{\max} = 220$  nm, LH<sub>2</sub> and LH<sub>3</sub> species, reach  $\epsilon_{\max} = 402, 77$  L mol<sup>-1</sup> cm<sup>-1</sup>, respectively. In literature [17] it was reported a value of  $\epsilon_{\max} = 4300$  L mol<sup>-1</sup> cm<sup>-1</sup>, at  $\lambda_{\max} = 236$  nm and pH≈12, which is very close to our result for L species ( $\epsilon_{\max} = 4279$  L mol<sup>-1</sup> cm<sup>-1</sup>), at  $\lambda_{\max} = 235$  nm.

With the aim to know the UV spectral properties of the complexes with alkyltin(IV) cations and to confirm the reliability of their formation constant values obtained by potentiometry, a wide number of spectra were performed by varying the pH, in the 1.5≤pH≤11.0 range, and the stoichiometric ratio between the **cys** and the dimethyltin(IV). In Fig. s.2 of the Supporting Information some experimental spectra registered on solutions containing dimethyltin(IV) and **cys** at  $C_M = C_L = 0.5$  mmol L<sup>-1</sup> are reported. They show slight absorptions up to pH = 6.5 that increase to a great extent in the 8.7≤pH≤11.5 range. Individual UV spectra of the MLH, ML and MLOH complex species are reported in Fig. 3 together with those of **cys**. The comparison

of these individual spectra with those of **cys** species reveals: 1. a batocromic shift of molar absorbance maximum; 2. a decrease of molar absorbances; as an example, at  $\lambda_{\max} = 220$  nm,  $\epsilon_{\max} = 1666, 2038, 2591$  L mol<sup>-1</sup> cm<sup>-1</sup> for MLH, ML and MLOH species, respectively. These results totally confirm the complex formation between the absorbing ligand **cys** and the dimethyltin(IV) cation.

The stability constants evaluation of these complexes can be performed by peak heights of UV spectra. By using the experimental spectra obtained on dimethyltin(IV)-**cys** solutions at several concentrations we refined the formation constants of the three main species (MLH, ML, MLOH) at  $I = 0.013$  mol L<sup>-1</sup>. The results are shown in Table 6 together with those obtained from the potentiometric measurements. The comparison shows that the spectroscopic technique is less sensitive than the potentiometric one, consequently the latter allows to determinate the stabilities of all the species examined with lower standard deviations; nevertheless, it represents a further method of validation of the speciation model calculated on the basis of potentiometric evidences.

### 2.3. NMR measurements

The spectroscopic properties of the dimethyltin(IV)-**cys** complexes were studied by means of <sup>1</sup>H NMR investigations in H<sub>2</sub>O/D<sub>2</sub>O solutions. A wide number of spectra were performed by varying the pH in the range 3 - 11 and the stoichiometric ratio between the **cys** and dimethyltin(IV) as well. The evidence that the spectra show a single set of peaks despite the experimental conditions suggests that although several species may be present in solution, they are involved in fast mutual exchange; as a consequence, the resonances of the bound and the free ligands cannot be directly observed from the spectra. Nevertheless, some spectra of

**cys** have been carried out in experimental conditions corresponding to the maxima of speciation curves, thus allowing the NMR characterization of LH<sub>2</sub> and L species, in particular conditions of concentration and pH.

As a common feature of all the spectra, on increasing the pH, upfield chemical shifts of all the signals occurred regardless of the dimethyltin(IV)/**cys** ratio employed. It is worth to mention that Silvestri *et al.* [18] studied the system dimethyltin(IV)-**cys** in aqueous solution by means of <sup>1</sup>H NMR, not considering the fast mutual exchange occurring in the experimental conditions. Furthermore some small signals clearly associated to the presence of low amount of cystine were wrongly ascribed to **cys**-containing species.

As far as the CH<sub>3</sub> signals are concerned, the <sup>1</sup>H NMR spectra of both dimethyltin(IV) and dimethyltin(IV)-**cys** solutions showed a single sharp signal featured by the satellite peaks of heteronuclear couplings, namely <sup>2</sup>J(<sup>117</sup>Sn-<sup>1</sup>H) and <sup>2</sup>J(<sup>119</sup>Sn-<sup>1</sup>H), with the two NMR active isotopes of tin. Fig. 4 shows the variations of δCH<sub>3</sub> vs pH for dimethyltin(IV) and dimethyltin(IV)-**cys** solutions in different stoichiometric ratio. The strong differences showed by the curves obtained with and without **cys** suggest that the presence of **cys** involves the formation of complex species in a wide pH range rather than simple hydrolysis. Furthermore, the curves corresponding to dimethyltin(IV)-**cys** solutions show a different behavior depending on the stoichiometric ratio employed, starting at pH~6. These conclusions fully support the potentiometric data interpretation that suggests the formation of complexes such as ML<sub>2</sub> or ML<sub>2</sub>H whenever the measurements are performed in excess of **cys**.

As can be inferred from Fig. 5, the δCH of the **cys** alone shows a first variation at low pH supporting the deprotonation of COOH group in this range; furthermore, on increasing the pH, the CH resonance keeps almost constant in a wide region (namely 3.4-7.2) while after neutral pH it decreases steadily. The dimethyltin(IV) containing mixtures are featured by a

CH signal very close to the one showed by **cys** solutions up to 4.5, whilst at higher pH the signals behavior is different; this evidence may be ascribed to the formation of dimethyltin(IV)-**cys** containing species [8]. Furthermore, while the  $^1\text{H}$  NMR spectra of **cys** registered in the presence of dimethyltin(IV) at  $\text{pH} < 4.5$  show that the resonance associated to CH is comparable with the one obtained for the free ligand in the same pH region, some differences can be detected in the above range in the  $\text{CH}_2$  signals suggesting an interaction between **cys** and dimethyltin(IV) at low pH by means of a distant group from the CH, i.e. the thiol. The absence of interactions between the two species present in solution must be discarded on the basis of potentiometric analysis.

Even though the direct measurements of NMR parameters cannot be carried out owing to fast mutual exchange among the formed species, it is possible to obtain both  $\delta$  and  $^2\text{J}$  for each supposed complex formed starting from speciation study. In fact, since the concentration of each species over the whole pH range is known from potentiometric data, individual chemical shifts and coupling constants can be easily calculated [31]. The aim of obtaining these calculated values is twofold: on one hand these individual “indirect”  $\delta$  and  $^2\text{J}$  were used to recalculate the mean corresponding to experimental pHs (simulating a fast mutual exchange) in order to validate the model used for speciation (by comparing the average recalculated and experimental values); on the other hand individual  $^2\text{J}$  calculated for all the complexes allows some structural considerations to be discussed.

For example, Fig. 6 shows the good agreement between calculated and observed chemical shifts of the tin-bound methylic groups in the mixture with  $C_{(\text{CH}_3)_2\text{Sn}} = 0.00869 \text{ mol L}^{-1}$  and  $C_{\text{cys}} = 0.01114 \text{ mol L}^{-1}$ . As a result the NMR data are consistent with the model proposed for the interpretation of potentiometric investigations. Furthermore, the  $^2\text{J}(\text{Sn}-^1\text{H})$  values can be conveniently converted into C-Sn-C angles by means of the empiric relation proposed by

Lockhart and Manders [31]. The  $^2J(\text{Sn}-^1\text{H})$  relative to the single species were calculated (Table 7) and the corresponding angles obtained show that all the complexes are characterized by an angle ranging from  $123^\circ$  to  $130^\circ$  suggesting a pentacoordinated trigonal bipyramidal arrangement around the tin with equatorial  $\text{CH}_3$  groups for all the species formed in solution, confirming literature evidences [38].

#### 2.4. Sequestering power of *L*-cysteinate towards dimethyltin(IV) cation

The sequestering capacity of a ligand towards a metal can be expressed by the function  $\Sigma(\%)$  vs.  $\text{pL}$ , where  $\Sigma(\%)$  is the total percentage of alkyltin(IV) cation complexed and  $\text{pL} = -\log [\text{L}]_{\text{tot}}$ . Since this function is a typically sigmoidal curve (or a dose response curve), increasing rapidly over a relatively small change in concentration, we can use the Boltzmann type equation (with asymptotes of 100 for  $\text{pL} \rightarrow -\infty$  and 0 for  $\text{pL} \rightarrow +\infty$ ):

$$\Sigma (\%) = 100 \times \left[ \frac{1}{1 + e^{(\text{pL} - \text{pL}_{50})/S}} - 1 \right] \quad (3)$$

where  $\text{pL}_{50}$  and  $S$  are empirical parameters, which define the ligand concentration necessary to sequester 50% of metal ion, while  $S$  is a measure of the slope in flex of the function ( $\Sigma\%$ ) vs  $\text{pL}$ . The  $\text{pL}_{50}$  parameter is very useful because gives a representation of the binding ability of a ligand (L) toward a specific cation in the investigated conditions.

Fig. 7 shows sequestration diagram of **cys** towards dimethyltin(IV) cation, at  $C_{\text{RSn}} = 10^{-3} \mu\text{mol L}^{-1}$  and  $T=25^\circ\text{C}$  and at several pH values.  $\text{pL}_{50}$  values are 4.082, 4.394, 4.970 at pH = 5, 6, 8, respectively; while  $S = 0.434, 0.433, 0.428$ . With increasing pH value, increases the binding ability of the ligand, too. These results are interesting for the toxicological

implications, because they indicate that the sequestering ability of **cys** is very high; for example at pH = 8 when  $C_{(\text{CH}_3)_2\text{Sn}} = 10^{-3} \mu\text{mol L}^{-1}$ , the concentration of the ligand necessary to sequester 50% of the dimethyltin(IV) is equal to  $10 \mu\text{mol L}^{-1}$ , i.e. 10 times lower than **cys** concentration in blood plasma models ( $\geq 100 \mu\text{mol L}^{-1}$ ) [2].

### 3. Literature comparisons

In Table 8 the speciation model and the stability constants of the species formed between **cys** and dimethyltin(IV) at  $I = 0.1 \text{ mol L}^{-1}$  in NaCl are compared with those referring to the system containing the same ligand and diethyltin(IV) at  $I = 0.1 \text{ mol L}^{-1}$  in NaClO<sub>4</sub>. The two speciation models, proposed by us and by Buzás et al. [19], respectively, are different only for the ML<sub>2</sub> species, present in our model only. The stability constants referring to MLH, ML, MLOH, and ML<sub>2</sub>H species are similar, precisely the species of the dimethyltin(IV) with **cys** are slightly higher than those of diethyltin(IV) (the highest difference is 0.53 and refers to the ML<sub>2</sub>H species), as expected for higher steric crowding of the ethyl groups with respect to methyl ones in the interaction with the ligand. The results obtained by Hynes and O'Dowd on the triethyltin(IV)-**cys** system at  $I = 0.3 \text{ mol L}^{-1}$  in NaClO<sub>4</sub> are reported in Table 8, too [20]. The only species proposed is the MLH and the corresponding stability is much lower than that referring to dimethyltin(IV)-**cys** system.

The comparisons of the complex stabilities of dimethyltin(IV)-**cys** system with those formed by dimethyltin(IV) with a simple amino acid, such as alanine, are reported in Table 9. The high extra stability obtained for the dimethyltin(IV)-**cys** system, with respect to dimethyltin(IV)-alanine (from 4.1 for MLOH to 7.6 for MLH species), is to attribute

unequivocally to the –SH group of the *cys*. From the same Table, the little differences between the stabilities of dimethyltin(IV)-*cys* and diethyltin(IV)-tiolactate species confirm what was already discussed.

#### 4. Conclusions and biological implications

The results obtained by different equilibrium (pH-metric, calorimetric and spectrophotometric) and structural ( $^1\text{H}$  NMR) methods for the study of the interaction of dimethyltin(IV) cation with *cys*, can be summarized as follows.

- The formation of five complex species, with very high stabilities and formation percentages (mainly for mononuclear species), in a wide pH range, was found.
- Exothermic enthalpy values and in some cases with an enthalpic contribution higher than entropic one, are observed.
- The speciation model proposed on the basis of potentiometric results was also validated by means of spectroscopic technique.
- Pentacoordinated trigonal arrangement around the tin, with equatorial  $\text{CH}_3$ , for all the species, was found by NMR data.
- Very high sequestering ability of *cys* toward dimethyltin(IV), was observed.
- The high extra stability of dimethyltin(IV)-*cys* species with respect to those of dimethyltin(IV) with simple amino acids, such as alanine, confirms that –S– interaction leads to more stable coordination than  $\text{COO}^-$  or  $\text{NH}_2$ .

As pointed up by Pellerito and Nagy in a recent review [9] on the interaction of organotin(IV) cations with biologically active ligands, this field is still open. Equilibrium data on the

different systems are largely missing. Therefore the study of the interaction of dimethyltin(IV) with a biological ligand such as cysteine, is useful to understand the biological action of dimethyltin(IV) complexes. The information obtained on these complexes has biological implications, because cys is a sulfur containing amino acid, present in the blood, which plays an important role in the metabolism of living cells (it participates in many biochemical processes, e.g. in the synthesis of tripeptides such as glutathione).

### **Acknowledgements**

The authors thank University of Messina for financial support.

**References**

- [1] R. Spezia, G. Tournois, T. Cartailier, J. Tortajada and Y. Jeanvoine,  $\text{Co}^{2+}$  binding Cysteine and selenocysteine: a DFT study, *J. Phys. Chem. A* 110 (2006) 9727-9735 and references therein.
- [2] E. Bottari and M.R. Festa, On the behaviour of cysteine as ligand of cadmium(II), *Talanta* 44 (1997) 1705-1718.
- [3] S.S. Khaloo, M.K. Amini, S. Tangestaninejad, S. Shahrokhian and R. Kia, Voltammetric and Potentiometric study of cysteine at Cobalt(II) Phtalocyanine modified carbon-paste electrode, *J. Ir. Chem. Soc.* 1 (2004) 128-135.
- [4] C. Mathé, T.A. Mattioli, O. Horner, M. Lombard, J.M. Latour, M. Fontecave and V. Nivière, Identification of Iron(III) peroxo species in the active site of the superoxide reductase SOR from *Desulfoarculus baarsi*, *J. Am. Chem. Soc.* 124 (2002) 4966-4967.
- [5] I. Endo, M. Nojiri, M. Tsujimura, M. Nakasako, S. Nagashima, M. Yohda and M. Odaka, Fe-type nitrile hydratase, *J. Inorg. Biochem* 83 (2001) 247-253.
- [6] A. Miyanaga, S. Fushinobu, K. Ito and T Wakagi, Crystal structure of Cobalt-containing nitrile hydratase, *Biochem. Biophys Res. Commun.* 288 (2001) 1169-1174.
- [7] M. Gielen, Tin-based antitumour drugs, *Coord. Chem. Rev.* 151 (1996) 41-51.
- [8] M. Nath, S. Pokharia and Y. Rakesh, Organotin(IV) complexes of amino acids and peptides, *Coord. Chem. Rev.* 215 (2001) 99-149.

- [9] L. Pellerito and L. Nagy, Organotin(IV)<sup>n+</sup> complexes formed with biologically active ligands: equilibrium and structural studies, and some biological aspects, *Coord. Chem. Rev.* 224 (2002) 111-150.
- [10] G. Arena, R. Cali, A. Contino, A. Musumeci, S. Musumeci and R. Purrello, Coordination properties of dialkyltin (IV) in aqueous solution. Thermodynamic study of dimethyltin (IV) complexes with L-amino acids, *Inorg. Chim. Acta* 237 (1995) 187-191.
- [11] M.M. Shoukry, Equilibrium studies of the diorganotin(IV) complexes with some amino acids and related compounds, *Talanta* 43 (1996) 177-183.
- [12] P. Surdy, P. Rubini, N. Buzás, B. Henry, L. Pellerito and T. Gajda, Interaction of Dimethyltin(IV)<sup>2+</sup> Cation with Gly-Gly, Gly-His, and Some Related Ligands, A New Case of a Metal Ion Able To Promote Peptide Nitrogen Deprotonation in Aqueous Solution, *Inorg. Chem.* 38 (1999) 346-352.
- [13] A. Atkinson, M.D. Rodriguez, T.E. Shewmaker and J.E. Walmsley, Synthesis and characterization of compounds of di- and tributyltin chloride with adenine and guanine mononucleotides, *Inorg. Chim. Acta* 285 (1999) 60-69.
- [14] G. Barone, R. Barbieri, G. La Manna and M.H. Koch, The interaction of deoxyribonucleic acid with methyltin(IV) moieties in solution studied by small-angle X-ray scattering, circular dichroism and UV spectroscopy, *Appl. Organomet. Chem.* 14 (2000) 189-196.
- [15] R. Barbieri, A. Silvestri, M.T. Lo Giudice, G. Ruisi and M.T. Musmeci, The binding of trialkyltin(IV) moieties to Rat Haemoglobin, and the structure of Model Systems, studied by Tin-119 Mössbauer spectroscopy, *J. Chem. Soc., Dalton Trans.* (1989) 519-525.

- [16] M.T. Musmeci, G. Madonia, M.T. Lo Giudice, A. Silvestri, G. Ruisi and R. Barbieri, Interactions of organotins with biological systems, *Appl. Organomet. Chem.* 6 (1992) 127-138.
- [17] G. Domazetis, R.J. Magee and B.D. James, Organotin(IV) derivatives of L-cysteine and DL-penicillamine, *J. Organomet. Chem.* 162 (1978) 239-252.
- [18] A. Silvestri, D. Duca and F. Huber, A study of dimethyltin(IV)-L-cysteinate in aqueous solution, *Appl. Organomet. Chem.* 2 (1988) 417-425.
- [19] N. Buzás, T. Gajda, E. Kuzmann, L. Nagy, A. Vértes and K. Burger, Coordination properties of L-cysteine and its derivatives towards diethyltin(IV) in aqueous solution, *Main Group Metal Chemistry* 18(11) (1995) 641-649.
- [20] M.J. Hynes and M. O'Dowd, Interactions of the trimethyltin(IV) cation with carboxylic acids, aminoacids and related ligands, *J. Chem. Soc. Dalton Trans.* (1987) 563-566.
- [21] G. Berthon, The Stability Constants of Metal Complexes of Amino Acids with Polar Side Chains, *Pure & Applied Chem.* 67(7) (1995) 1117-1240.
- [22] C. De Stefano, C. Foti, A. Gianguzza, M. Martino, L. Pellerito and S. Sammartano, Hydrolysis of  $(\text{CH}_3)_2\text{Sn}^{2+}$  in Different Ionic Media: Salt Effects and Complex Formation, *J. Chem. Eng. Data* 41 (1996) 511-515.
- [23] C. De Stefano, C. Foti, A. Gianguzza, F. J. Millero and S. Sammartano, Hydrolysis of  $(\text{CH}_3)_3\text{Sn}^+$  in various salt media, *J. Solution Chem.* 28 (7) (1999) 959-972.
- [24] C. Foti, A. Gianguzza, D. Piazzese and G. Trifiletti, Inorganic speciation of organotin(IV) cations in natural waters with particular references to seawater, *Chem. Spec. Bioavail.* 12(2) (2000) 41-52.

- [25] C. De Stefano, C. Foti, A. Gianguzza and S. Sammartano, in: Hydrolysis processes of organotin(IV) compounds in sea water, eds. A. Gianguzza, E. Pellizzetti and S. Sammartano, *Chemical Processes in the Marine Environment* (Environmental Sciences Library, Springer, Berlin, 2000) pp. 213-228.
- [26] C. De Stefano, A. Gianguzza, F. Marrone and D. Piazzese, Interaction of Alkyltin(IV) Compounds with Ligands of Interest in the Speciation of Natural Fluids: Complexes of  $(\text{CH}_3)_2\text{Sn}^{2+}$  with carboxylates, *Appl. Organomet. Chem.* 11 (1997) 683-691.
- [27] C. De Stefano, C. Foti and A. Gianguzza, Interactions of Alkyltin(IV) Compounds with Ligands of Interest in the Speciation of Natural Fluids: Complex of  $(\text{CH}_3)_3\text{Sn}^+$  with Carboxylates, *Ann. Chim. (Rome)* 89 (1999) 147-155.
- [28] C. Foti, A. Gianguzza and S. Sammartano, Interaction of alkyltin(IV) compounds with ligands of interest in the speciation of natural fluids: carboxylate and hydroxycarboxylate complexes of monomethyltin(IV) trichloride, *Ann. Chim. (Rome)* 92(7-8) (2002) 705-715.
- [29] P. Cardiano, O. Giuffrè, L. Pellerito, A. Pettignano, S. Sammartano and M. Scopelliti, Thermodynamic and spectroscopic study of the binding of dimethyltin(IV) by citrate at 25°C, *Appl. Organomet. Chem.* 20 (2006) 425-435.
- [30] A. De Robertis, A. Gianguzza, O. Giuffrè, A. Pettignano, S. Sammartano, Interaction of methyltin(IV) compounds with carboxylate ligands. Part 1: Formation and stability of methyltin(IV)-carboxylate complexes and their relevance in the speciation of natural waters, *Appl. Organomet. Chem.* 20 (2006), 89-98.

- [31] T.P. Lockhart and W.F. Manders, Structure determination by NMR spectroscopy. Correlation of  $|^2J(^{119}\text{Sn}, ^1\text{H})|$  and the Me-Sn-Me angle in methyltin(IV) compounds, *Inorg. Chem.* 25 (1986) 892-895.
- [32] C. De Stefano, S. Sammartano, P. Mineo and C. Rigano, in: *Computer Tools for the Speciation of Natural Fluids*, eds. A. Gianguzza, E. Pellizzetti and S. Sammartano, *Marine Chemistry - An Environmental Analytical Chemistry Approach* (Kluwer Academic Publishers, Amsterdam, 1997) pp. 71-83.
- [33] P. G. Daniele, A. De Robertis, C. De Stefano, S. Sammartano and C. Rigano, On the Possibility of Determining the Thermodynamic Parameters for the Formation of Weak Complexes Using a Simple Model for the Dependence on Ionic Strength of Activity Coefficients.  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$  Complexes of Low Molecular Weight Ligands in Aqueous Solution, *J. Chem. Soc. Dalton Trans.* 85 (1985) 2353-2361.
- [34] P. G. Daniele, C. De Stefano, C. Foti and S. Sammartano, The Effect of Ionic Strength and Ionic Medium on the Thermodynamic Parameters of Protonation and Complex Formation, *Curr. Top. Sol. Chem.* 2 (1997) 253-274.
- [35] A. De Robertis, C. De Stefano and C. Rigano, Computer Analysis of Equilibrium Data in Solution. ES5CM Fortran and Basic Programs for Computing Formation Enthalpies from Calorimetric Measurements, *Thermochim. Acta* 138 (1986) 141-146.
- [36] Hyperquad 2000, P.Gans, A. Sabatini, A. Vacca, version 2.1.
- [37] V.K. Sharma, F. Casteran, F.J. Millero and C. De Stefano, Dissociation Constants of Protonated Cysteine Species in NaCl Media, *J. Solution Chem.* 31 (2002) 783-792.

- [38] R. Barbieri and M.T. Musmeci, A  $^{119}\text{Sn}$  mössbauer spectroscopic study on the interaction of dimethyltin(IV) derivatives with rat hemoglobin, and of related model systems in aqueous solution, *J. Inorg. Biochem.* 1988, 32, 89-108.
- [39] C. Foti, A. Gianguzza, F.J. Millero and S. Sammartano, The speciation of  $(\text{CH}_3)_2\text{Sn}^{2+}$  in electrolyte solution containing the major components of natural waters, *Aquatic Geochem.* 5 (1999) 381-398.
- [40] K. Gajda-Schranz, L. Nagy, T. Fiore, L. Pellerito and T. Gajda, Equilibrium and spectroscopic studies of diethyltin(IV) complexes formed with hydroxymono- and dicarboxylic acids and their thioanalogues. *J. Chem. Soc., Dalton Trans.*, 2002, 152-158 and references therein.

Table 1

Experimental conditions for potentiometric, calorimetric and spectrophotometric measurements at T=25°C

potentiometric measurements	$C_{\text{NaCl}}^a$	$C_{(\text{CH}_3)_2\text{SnCl}_2}^a$	$C_{\text{eys}}^a$	$I^{a,b}$	$N_{\text{tit}}^c$	$N_{\text{pts}}^d$
		0.002	0.003	0.008	2	125
		0.001	0.002	0.004	2	148
		0.002	0.006	0.009	2	136
		0.004	0.002	0.009	2	226
	0.14	0.002	0.003	0.142	2	146
	0.15	0.001	0.002	0.146	2	205
	0.14	0.002	0.006	0.141	2	163
	0.14	0.004	0.002	0.137	2	147
calorimetric measurements	$C_{\text{NaCl}}^a$	$C_{(\text{CH}_3)_2\text{SnCl}_2}^a$	$C_{\text{Na}_2\text{eys}}^a$	$I^{a,b}$	$N_{\text{tit}}^c$	$N_{\text{pts}}^d$
	0.1	0.0025	0.2507	0.104	2	60
	0.1	0.005	0.2507	0.108	2	60
spectrophotometric measurements	$C_{\text{HCl}}^a$	$C_{(\text{CH}_3)_2\text{SnCl}_2}^a$	$C_{\text{eys}}^a$	$I^{a,b}$	$N_{\text{tit}}^c$	$N_{\text{spectra}}^e$
	0.012		0.0004	0.019	1	10
	0.020		0.0006	0.031	1	22
	0.101		0.002	0.129	1	15
	0.040		0.005	0.072	1	13
	0.004	0.002	0.0004	0.015	1	10
		0.0004	0.0008	0.004	1	10
	0.004	0.005	0.001	0.027	1	9
	0.003	0.0005	0.0005	0.009	1	32

<sup>a</sup> Concentrations in mol L<sup>-1</sup>; <sup>b</sup> mean value of ionic strength; <sup>c</sup> number of titrations; <sup>d</sup> number of points; <sup>e</sup> number of spectra.

Table 2

Thermodynamic parameters of protonation of **cys** at

T=25°C and I = 0 mol L<sup>-1</sup>

pqr	log $\beta^a$	$\Delta H^{0\ a, b}$
011	10.21	-42
012	18.39	-77
013	20.53	-81

<sup>a</sup> Rif. 37; <sup>b</sup> expressed in kJ mol<sup>-1</sup>.

Table 3

Thermodynamic parameters for hydrolysis and  $\text{Cl}^-$  complex formation of  $(\text{CH}_3)_2\text{Sn}^{2+}$  at  $T=25^\circ\text{C}$  and  $I = 0$  mol  $\text{L}^{-1}$

pqr	$\log\beta^{a, b}$	$\Delta H^{0 c, d}$
10-1	-2.86	33.1
10-2	-8.16	62.1
10-3	-19.35	97.7
20-2	-4.99	60.0
20-3	-9.06	85.0
11 0	0.78	—
11-1	-3.17	—

<sup>a</sup> reaction:  $p\text{M}^{2+} + q\text{Cl}^- + r\text{H}_2\text{O} = \text{M}_p\text{Cl}_q(\text{OH})_r^{2p-(r+1)} + r\text{H}^+$ ; <sup>b</sup> refs. 22, 24; <sup>c</sup> expressed in  $\text{kJ mol}^{-1}$ ; <sup>d</sup> ref. 39.

Table 4

Overall thermodynamic formation parameters of  $(\text{CH}_3)_2\text{Sn-cys}$  complexes at  $T=25^\circ\text{C}$  and  $I = 0$  mol  $\text{L}^{-1}$

pqr <sup>a</sup>	$\log\beta \pm 3s^b$	$\log\beta^c$	$-\Delta G^{0d}$	$\Delta H^{0 \pm s^{b,d}}$	$T\Delta S^{0d}$
1 1 1	$18.81 \pm 0.04$	18.03	107.4	$-61.5 \pm 0.6$	46
1 1 0	$14.67 \pm 0.03$	13.89	83.7	$-44.1 \pm 0.3$	40
1 1 -1	$7.29 \pm 0.03$	6.71	41.6	$-10 \pm 1$	32
1 2 1	$27.77 \pm 0.04$	26.60	158.5	$\sim -134$	$\sim 25$
1 2 0	$20.27 \pm 0.04$	19.49	115.7	$-48.7 \pm 0.7$	67

<sup>a</sup> Reaction:  $pM + qL + rH = M_pL_qH_r$ ; <sup>b</sup> standard deviation; <sup>c</sup> calculated at  $I = 0.15$  mol  $\text{L}^{-1}$  with eq. 1; <sup>d</sup> expressed in  $\text{kJ mol}^{-1}$ .

Table 5

Thermodynamic formation parameters of  $(\text{CH}_3)_2\text{Sn-cys}$  complexes at  $T=25^\circ\text{C}$  and  $I = 0 \text{ mol L}^{-1}$

Equilibrium	logK	$-\Delta G^{0a}$	$\Delta H^{0a}$	$T\Delta S^{0a}$
$\text{HL} + \text{M} = \text{MLH}$	8.6	49.1	-19.5	30
$\text{L} + \text{M} = \text{ML}$	14.67	83.7	-44.1	40
$\text{L} + \text{MOH} = \text{MLOH}$	10.1	57.7	-43	15
$\text{ML} + \text{HL} = \text{ML}_2\text{H}$	2.9	16.6	~-48	-31
$\text{ML} + \text{L} = \text{ML}_2$	5.6	32.0	-4.6	27

<sup>a</sup> Expressed in  $\text{kJ mol}^{-1}$ .

Table 6

Formation constants of dimethyltin(IV)-cys species obtained with two different techniques and calculation programs

species	$\log\beta\pm 3s^a$	
	from potentiometry <sup>b</sup>	from UV spectroscopy <sup>c</sup>
MLH	18.81±0.04	18.5±0.3
ML	14.67±0.03	15.4±0.2
MLOH	7.29±0.03	7.7±0.3

<sup>a</sup> s = standard deviations; <sup>b</sup> at I = 0 mol L<sup>-1</sup>; <sup>c</sup> at I = 0.013 mol L<sup>-1</sup>.

Table 7

Calculated individual  $^1\text{H}$  NMR parameters

species	$\delta(\text{CH}_3)$ (ppm)	$^2\text{J}(\text{Sn-H})$ (Hz)	$\angle\text{C-Sn-C}$ ( $^\circ$ )
MLH	$0.871\pm 0.005$	$79.5\pm 0.2$	130
ML	$0.853\pm 0.001$	$75.5\pm 0.1$	125
MLOH	$0.617\pm 0.005$	$75.2\pm 0.2$	125
$\text{ML}_2\text{H}$	$1.19\pm 0.04$	$74\pm 1$	123
$\text{ML}_2$	$0.67\pm 0.01$	$74.9\pm 0.2$	124

Table 8

Literature comparisons at T=25°C

species	log $\beta$		
	M = (CH <sub>3</sub> ) <sub>2</sub> Sn <sup>2+</sup> <sup>a</sup> this work	M = (C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> Sn <sup>2+</sup> <sup>b</sup> ref. 19	M = (C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub> Sn <sup>+</sup> <sup>b</sup> ref. 20
M(cys)H	19.43	19.43	15.21
M(cys)	14.86	14.77	—
M(cys)OH	7.29	6.98	—
M(cys) <sub>2</sub> H	28.88	28.35	—
M(cys) <sub>2</sub>	20.83	—	—

<sup>a</sup> At I = 0.1 mol L<sup>-1</sup> in NaCl; <sup>b</sup> at I = 0.1 mol L<sup>-1</sup> in NaClO<sub>4</sub>.

Table 9

Literature comparisons at T=25°C

species	log $\beta$		
	M = (CH <sub>3</sub> ) <sub>2</sub> Sn <sup>2+</sup> L = <b>cys</b> <sup>a</sup> this work	M = (CH <sub>3</sub> ) <sub>2</sub> Sn <sup>2+</sup> L = alanine <sup>b</sup> ref. 10	M = (C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> Sn <sup>2+</sup> L = tiolactic acid <sup>c</sup> ref. 40
MLH	18.81	11.20	—
ML	14.67	8.25	14.13
MLOH	7.29	3.23	7.48
ML(OH) <sub>2</sub>	—	-5.95	—
ML <sub>2</sub> H	27.77	—	—
ML <sub>2</sub>	20.27	—	—

<sup>a</sup> At I = 0 mol L<sup>-1</sup>; <sup>b</sup> at I = 0.1 mol L<sup>-1</sup> in KNO<sub>3</sub>; <sup>c</sup> at I = 0.3 mol L<sup>-1</sup> in NaClO<sub>4</sub>.

## Supporting Information

Table s.1

Percentages of the dimethyltin(IV)-cys species in the  $7 \leq \text{pH} \leq 8$  range and at two concentration ratios

M/L ratio	I <sup>a</sup>	pH	%ML	%MLOH	%ML <sub>2</sub> H	%ML <sub>2</sub>	Σ%
0.5 <sup>b</sup>	0.05	7.0	55.9	25.4	13.1	4.9	99.3
		7.5	26.4	39.3	14.9	19.1	99.7
		8.0	8.6	41.9	9.3	40.0	99.8
1 <sup>c</sup>	0.065	7.0	61.9	28.3	6.1	2.3	98.6
		7.5	33.5	50.3	6.5	8.5	98.8
		8.0	13.4	65.1	3.9	16.6	99.0

<sup>a</sup> Expressed in mol L<sup>-1</sup>; <sup>b</sup> C<sub>M</sub> = 5 mmol L<sup>-1</sup>; C<sub>L</sub> = 10 mmol L<sup>-1</sup>; <sup>c</sup> C<sub>M</sub> = C<sub>L</sub> = 10 mmol L<sup>-1</sup>.

Table s.2

Molar absorbances of protonated and unprotonated species of **cys** and of complexes with dimethyltin(IV) cation at T=25°C

species	$\lambda^a$						
	L	LH	LH <sub>2</sub>	LH <sub>3</sub>	MLH	ML	MLOH
280	31	44	2	0	18	17	14
275	90	97	3	0	28	24	21
270	217	237	4	0	44	37	31
265	489	513	5	1	70	66	57
260	998	1052	8	1	132	132	110
255	1828	1928	15	1	246	256	225
250	2875	3004	26	2	444	492	463
245	3757	3864	48	2	735	843	865
240	4177	4211	80	2	1092	1302	1419
235	4279	4345	151	4	1417	1737	1981
230	4157	4289	230	17	1581	1964	2303
225	4002	4141	317	41	1584	1990	2416
220	3883	3999	402	77	1666	2038	2591

<sup>a</sup> Expressed in nm; <sup>b</sup> expressed in L mol<sup>-1</sup> cm<sup>-1</sup>.




Fig. s.1 Comparison between experimental potentiometric data and theoretical fit recorded on aqueous solutions of dimethyltin(IV) (M) and **cys** (L) at  $C_M = 0.001007 \text{ mol L}^{-1}$ ,  $C_L = 0.002014 \text{ mmol L}^{-1}$ ,  $C_{\text{NaOH}} = 0.1975 \text{ mol L}^{-1}$ ,  $v_0 = 25 \text{ mL}$  and  $T=25^\circ\text{C}$




Fig. s.2 Experimental UV spectra recorded on aqueous solutions of dimethyltin(IV) (M) and **cys** (L) at  $C_M = C_L = 0.5 \text{ mmol L}^{-1}$  and  $T=25^\circ\text{C}$




Fig. 1a-c.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Fig. 7.

Fig. 1a-c. Speciation diagrams of  $(\text{CH}_3)_2\text{Sn}^{2+}(\text{M})$ -**cys** (L) vs. pH at  $I = 0 \text{ mol L}^{-1}$  and  $T=25^\circ\text{C}$ . Indexes refer to reaction (2). Hydrolytic species are shown by dashed lines. **a**:  $C_M = 5 \text{ mmol L}^{-1}$ ;  $C_L = 10 \text{ mmol L}^{-1}$ . **b**:  $C_M = C_L = 1 \text{ mmol L}^{-1}$ . **c**:  $C_M = C_L = 0.1 \text{ mmol L}^{-1}$ .

Fig. 2. Experimental UV spectra recorded on aqueous solutions of **cys** at  $C = 0.4 \text{ mmol L}^{-1}$  and  $T=25^\circ\text{C}$ .

Fig. 3. Molar absorbances of protonated and unprotonated species of **cys** and of complexes with  $(\text{CH}_3)_2\text{Sn}^{2+}$  cation at  $T=25^\circ\text{C}$ .

Fig. 4. Measured chemical shifts, at  $T=25^\circ\text{C}$ , of  $\text{CH}_3$  vs. pH for:

- solutions containing dimethyltin(IV) only;
- dimethyltin(IV):**cys** 1:1 mixtures;
- △— dimethyltin(IV):**cys** 1.5:2 mixtures;

Fig. 5. Measured chemical shifts, at  $T=25^\circ\text{C}$ , of CH vs. pH for:

- solutions containing **cys** only;
- dimethyltin(IV):**cys** 1.5:2 mixtures;
- △— dimethyltin(IV):**cys** 1:1 mixtures;

Fig. 6. Chemical shifts of  $\text{CH}_3$  vs. pH in dimethyltin(IV)-**cys** solutions at  $C_{(\text{CH}_3)_2\text{Sn}} = 0.00869 \text{ mol L}^{-1}$ ;  $C_{\text{cys}} = 0.01114 \text{ mol L}^{-1}$  and  $T=25^\circ\text{C}$ .

Fig. 7. Sequestration diagram of  $(\text{CH}_3)_2\text{Sn}^{2+}$  cation by **cys** at  $T=25^\circ\text{C}$  and  $C_{(\text{CH}_3)_2\text{Sn}} = 10^{-3} \mu\text{mol L}^{-1}$ . □: pH = 5; ○: pH = 6; △: pH = 8.