

Stationary Solutions to the Vlasov-Poisson System

Fahd Karami, Simon Labrunie, Bruno Pinçon

▶ To cite this version:

Fahd Karami, Simon Labrunie, Bruno Pinçon. Stationary Solutions to the Vlasov-Poisson System. 2010. hal-00501842v1

HAL Id: hal-00501842 https://hal.science/hal-00501842v1

Preprint submitted on 12 Jul 2010 (v1), last revised 19 Jun 2012 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stationary Solutions to the Vlasov-Poisson System

Fahd KARAMI * Sime

Simon Labrunie †

Bruno PINÇON[‡]

july 12, 2010

Abstract

In this paper, we present an existence and uniqueness result for stationary Vlasov Poisson system and we study the asymptotic behavior of the solution near the corner.

Keywords: Vlasov-Poisson, Plasma physics, Polygonal domains, Asymptotic analysis.

1 Introduction.

We consider a gas of charged particles. This particles are described by their distribution function f(x, v) where $x \in \Omega \subset \mathbb{R}^N$ is the position of the particle and v its velocity. We consider the Stationary Vlasov-Poisson system which governs the motion of charged particles under the influence of an electrostatic field. The stationary Vlasov equation reads

$$v \cdot \nabla_x f - E(x) \cdot \nabla_v f = 0, \quad (x, v) \in \Omega \times \mathbb{R}^N, \tag{1}$$

N 7

where the electric field E(x) is given by the equations

$$E(x) = -\nabla_x (\phi[f] - \phi_e), \quad -\Delta\phi[f] = \int f(x, v) \, dv := \rho[f] \quad \text{and} \quad -\Delta\phi_e = \rho_e.$$

where $\rho[f]$ is a spatial density of particles, ϕ the self-consistent electrostatic potential, ρ_e is the density of the fixed charges and ϕ_e is the external potential. Denoted by $\nu(x)$ the outward normal vector of Ω at a point $x \in \partial \Omega$ and we assume that $\partial \Omega := \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$ where

$$\begin{aligned}
\phi[f] &= \phi_e = 0 & \text{on } \Gamma_1 \\
\phi[f] &= 0, \ \phi_e = \phi_{in} & \text{on } \Gamma_2 \\
\frac{\partial \phi[f]}{\partial \nu} &= \frac{\partial \phi_e}{\partial \nu} = 0 & \text{on } \Gamma_3.
\end{aligned}$$
(2)

^{*}Email : karami@iecn.u-nancy.fr

[†]Email : labrunie@iecn.u-nancy.fr

[‡]Email : pincon@iecn.u-nancy.fr

1.1 Assumptions

Throughout this paper, we assume that

$$(H_1) \ \rho_e \in L^p(\Omega) \text{ and } \phi_{in} \in L^{\infty}(\Gamma_2) \cap W^{1-1/p, p}(\Gamma_2) \text{ for some } p > N.$$

 $(H_2) \ \gamma \ge 0$ is a strictly decreasing function such that $\forall \delta > 0$

$$s^{\frac{N}{2}-1}\gamma(s-\delta) \in L^1(0,+\infty).$$

(*H*₃) Assume that $\sigma(r) = -\int_0^r \gamma^{-1}(s) \, ds$ with $\sigma \in \mathcal{C}^2(\mathbb{R}^+) \cup \mathcal{C}^0(\mathbb{R}^+)$ bounded from below and strictly convex

$$\sup_{x \in \Omega} \int_0^\infty \sigma(h(x,s)) ds < \infty \quad \text{and} \quad \lim_{s \to \infty} \frac{\sigma(s)}{s} = +\infty$$

where $h(x, s) = s^{\frac{N}{2}-1} \gamma(s - \phi_e(x)).$

1.2 Preliminaries

We consider the spaces V and W of functions defined by:

 $V = \{ w \in H^1(\Omega) \quad \text{such that} \ w = 0 \text{ on } \Gamma_1 \cup \Gamma_2 \}$

 $W = \{ w \in H^1(\Omega) \text{ such that } w = 0 \text{ on } \Gamma_1 \text{ and } w = \phi_{in} \text{ on } \Gamma_2 \}.$

Let $f \in L^1(\Omega \times \mathbb{R}^N)$. If $\rho[f] \in V'$ then there exists $\phi \in V$ solution of $-\Delta \phi = \rho[f]$ satisfying

$$\int_{\Omega} \nabla \phi \cdot \nabla \xi \, dx = \int_{\Omega} \rho[f] \, \xi \, dx, \quad \text{ for any } \xi \in V.$$

We define a mapping Φ and ϕ by $\Phi[\rho] = \Phi[\rho[f]] = \phi$ and $\phi[f] = \phi$, we have

$$\int_{\Omega} |\nabla \phi[f]|^2 < \infty.$$

If $\rho[f] \in L^1(\Omega) \setminus V'$ then, the solution $\phi \in V$ does not exist, and in this case, we agree that

$$\int_{\Omega} |\nabla \phi[f]|^2 = \infty.$$

Indeed, by density there exists a sequence $\rho_n \in L^1(\Omega) \cap V'$ such that ρ_n converges to ρ in $L^1(\Omega)$. Then, as $n \to \infty$ we have

$$\int_{\Omega} |\nabla \Phi[\rho_n]|^2 \sim ||\rho_n||_{V'}^2 \to \infty,$$

otherwise there exists a subsequence will be noted by ρ_n which converges weakly to ρ in V' and this is contradictory because $\rho \notin V'$. Similarly, we agree that

 $\int_{\Omega \times \mathbb{R}^N} \sigma(f) = \infty \quad \text{if} \quad \sigma(f) \notin L^1(\Omega \times \mathbb{R}^N).$

It is clear that the operator ϕ is a linear and satisfies

$$\int_{\Omega \times \mathbb{R}^N} f\phi[g] \, d(x,v) \ = \ \int_{\Omega \times \mathbb{R}^N} g\phi[f] \, d(x,v) \quad \text{and} \quad \int_{\Omega \times \mathbb{R}^N} f\phi[f] \, d(x,v) \ = \int_{\Omega} |\nabla \phi[f]|^2 \, dx.$$

In the next section, we prove existence and uniqueness of the solution of the stationary problem (1-2). For that, we use some techniques for nonlinear optimization in L^1 and we give also some preliminary properties that will be used afterwards. In the third section, we study the asymptotic of the solution near the corner. Then, we establish some interesting properties for the singularity coefficient. The last section is devoted to the asymptotic of the solution of Maxwell-Boltzmann problem with respect to the total mass.

2 Existence and uniqueness

Thanks to (H_1) , there exists a unique $\phi_e \in W^{1, p}(\Omega)$ solution to problem

$$\begin{cases} -\Delta \phi_e = \rho_e & \text{in } \Omega \\ \phi_e = 0 & \text{on } \Gamma_1 \\ \phi_e = \phi_{in} & \text{on } \Gamma_2 \\ \frac{\partial \phi_e}{\partial \nu} = 0 & \text{on } \Gamma_3. \end{cases}$$
(3)

Now, any function f satisfying

$$f(x,v) = \gamma\left(\frac{1}{2}|v|^2 + \phi[f](x) - \phi_e(x) - \beta\right) \quad \text{and} \qquad \int_{\Omega \times \mathbb{R}^N} f(x,v) = M \tag{4}$$

is a stationary solution to problem (1-2), where β is unknown constant and M is the total mass. Now, we consider the space $L^1_M(\Omega \times \mathbb{R}^N)$ defined by:

$$L^1_M(\Omega \times \mathbb{R}^N) := \{ f \in L^1(\Omega \times \mathbb{R}^N) : \int_{\Omega \times \mathbb{R}^N} f \, dx \, dv = M \}.$$

To prove the existence of a solution to problem (4), we use some similar argument in [1], [2] and [3].

Proposition 2.1 Assume that the Hypothesis $(H_1 - H_3)$ are satisfied. There exists a unique function f solution to problem (4). The function f is the minimum on $L^1_M(\Omega \times \mathbb{R}^N)$ of the functional:

$$J[f] = \int_{\Omega \times \mathbb{R}^N} \left(\sigma(f) + \left(\frac{1}{2}|v|^2 - \phi_e\right) f \right) dx dv + \frac{1}{2} \int_{\Omega} |\nabla \phi[f]|^2 dx.$$

Proof: It is not difficult to see that \mathcal{J} is convex and lower semi-continuous, then \mathcal{J} is weakly *l.s.c.* Taking $g(x,v) = \gamma(\frac{1}{2}|v|^2 - \phi_e(x))$ by (H_2) , we have $g(x,v) \in L^1(\Omega \times \mathbb{R}^N)$. Thanks to the fact that $\sigma(s) - \sigma(r) - (s-r)\sigma'(r) \ge 0$ for any $r, s \ge 0$, we have

$$\begin{aligned} \mathcal{J}[f] &\geq \int_{\Omega \times \mathbb{R}^{N}} \left(\sigma(g) + (\frac{1}{2}|v|^{2} - \phi_{e}) g \right) d(x,v) \\ &\geq 2^{\frac{N}{2} - 1} |S^{N-1}| \int_{\Omega} \int_{0}^{\infty} \sigma(s^{\frac{N}{2} - 1} \gamma(s - \phi_{e})) d(s,x) - ||\phi_{e}||_{L^{\infty}(\Omega)} ||g||_{L^{1}(\Omega \times \mathbb{R}^{N})} \end{aligned}$$

where $|S^{N-1}|$ is the measure of the unit sphere in \mathbb{R}^N . Using (H_2) and (H_3) we have $\mathcal{J}[f]$ is bounded from below on $L^1_M(\Omega \times \mathbb{R}^N)$. Let $(f_n) \in L^1_M(\Omega \times \mathbb{R}^N)$ is a minimizing sequence for \mathcal{J} . Now, using the Dunford-Pettis compactness criterion, we can extract a subsequence still noted (f_n) such that

$$f_n \rightharpoonup f$$
 weakly in $L^1(\Omega \times \mathbb{R}^N)$ as $n \to \infty$

Since \mathcal{J} is weakly *l.s.c*, then

$$\mathcal{J}[f] \le \liminf_{n \to \infty} \mathcal{J}[f_n] = \inf_{L^1_M} \mathcal{J}$$

Then the function f is a global minimum of \mathcal{J} , that satisfies $\int_{\Omega} |\nabla \phi[f]|^2 dx < \infty$ and

$$-\gamma^{-1}(f) + \phi[f] + \frac{1}{2}|v|^2 - \phi_e(x) - \beta = 0,$$

where β is the Lagrange multiplier associated to the constraint $\int_{\Omega \times \mathbb{R}^N} f(x, v) = M$. Finally, the uniqueness of the solution follows by the strict convexity of the function σ .

Now, let the function G defined by

$$G(r) = \int_{\mathbb{R}^N} \gamma(\frac{1}{2}|v|^2 + r) \, dv = 2^{\frac{N}{2}-1} |S^{N-1}| \int_0^{+\infty} s^{\frac{N}{2}-1} \gamma(s+r) \, ds$$

then the problem (4) is equivalent to solving:

$$\begin{aligned} & & -\Delta\phi = \rho, \ \rho = G(\phi - \phi_e - \beta) & \text{in } \Omega \\ & & \phi = 0 & & \text{on } \Gamma_1 \cup \Gamma_2 \\ & & \frac{\partial\phi}{\partial\nu} = 0 & & \text{on } \Gamma_3 \end{aligned}$$
(5)

where β is defined implicitly by:

$$\int_{\Omega} G(\phi - \phi_e - \beta) = M.$$

Proposition 2.2 For any fixed β , there exists a unique solution to Problem (5) satisfying

$$\int_{\Omega} \nabla \phi \cdot \nabla \xi = \int_{\Omega} G(\phi - \phi_e - \beta) \xi \quad \text{for all} \quad \xi \in V.$$

Poof: We consider the nonlinear operator $\mathcal{B}: V \longrightarrow V'$ defined by

$$\langle \mathcal{B}\phi, v \rangle = \int_{\Omega} \nabla \phi \cdot \nabla v - \int_{\Omega} G(\phi - \phi_e - \beta) v + \int_{\Omega} G(-\phi_e - \beta) v \quad \text{for all} \quad v \in V$$

where $\langle .,. \rangle$ is the duality pairing between V and its dual V'. The operator \mathcal{B} is bounded and weakly continuous. Moreover, since G is a decreasing function then $\int_{\Omega} (G(-\phi_e - \beta) - G(\phi - \phi_e - \beta)) \phi \ge 0$ and

$$\frac{\langle \mathcal{B}\phi, \phi \rangle}{||\phi||_V} \ge ||\phi||_V \to \infty \quad \text{as} \quad ||\phi||_V \to \infty.$$

Then the operator \mathcal{B} is coercive. Thanks to [4], for any element of V' (in particular for $G(-\phi_e - \beta) \in L^{\infty}(\Omega)$) there exists $\phi \in V$, such that $\langle \mathcal{B}\phi, v \rangle = \langle G(-\phi_e - \beta), v \rangle$ for all $v \in V$.

2.1 Dependence of the solution on the total mass

Now, we study the dependence of the solution with respect to the total mass M, and we derive a comparison principle between M and the Lagrange multiplier β .

Proposition 2.3 Let ϕ_1 and ϕ_2 be two solutions to Problem (5) corresponding to β_1 and β_2 respectively.

If
$$\beta_1 \ge \beta_2$$
 then $\phi_1 \ge \phi_2$ in Ω .

Poof: Let ϕ_1 et ϕ_2 be two solutions corresponding respectively to β_1 and β_2 in the sense of the proposition 2.2, we have

$$\int_{\Omega} \nabla(\phi_1 - \phi_2) \cdot \nabla \xi = \int_{\Omega} \left(G(\phi_1 - \phi_e - \beta_1) - G(\phi_2 - \phi_e - \beta_2) \right) \xi.$$

Taking $\xi = (\phi_2 - \phi_1)^+$ as a test function in this formulation, then

$$\int_{\Omega} \nabla(\phi_1 - \phi_2) \cdot \nabla(\phi_2 - \phi_1)^+ = \int_{\Omega} \left(G(\phi_1 - \phi_e - \beta_1) - G(\phi_2 - \phi_e - \beta_2) \right) (\phi_2 - \phi_1)^+.$$

We have

$$\int_{\Omega} \nabla (\phi_1 - \phi_2) \cdot \nabla (\phi_2 - \phi_1)^+ = -\int_{\Omega} |\nabla (\phi_2 - \phi_1)^+|^2 \le 0,$$

and

$$G(\phi_1 - \phi_e - \beta_1) - G(\phi_2 - \phi_e - \beta_2) = \left(G(\phi_1 - \phi_e - \beta_2) - G(\phi_2 - \phi_e - \beta_2)\right) + \left(G(\phi_1 - \phi_e - \beta_1) - G(\phi_1 - \phi_e - \beta_2)\right)$$

The second term on the right-hand side is positive because $\beta_1 \ge \beta_2$ and G is a decreasing function. Using again the monotonicity of G, we have

$$\int_{\Omega} \left(G(\phi_1 - \phi_e - \beta_2) - G(\phi_2 - \phi_e - \beta_2) \right) (\phi_2 - \phi_1)^+ \ge 0,$$

then

$$\int_{\Omega} \left(G(\phi_1 - \phi_e - \beta_1) - G(\phi_2 - \phi_e - \beta_2) \right) (\phi_2 - \phi_1)^+ \ge 0$$

Hence

$$-\int_{\Omega} |\nabla(\phi_2 - \phi_1)^+|^2 \ge 0$$

and therefore

$$\nabla(\phi_2 - \phi_1)^+ = 0$$

then $(\phi_2 - \phi_1)^+ = const$. Since $(\phi_2 - \phi_1)^+ = 0$ on $\Gamma_1 \cup \Gamma_2$, we deduce that $(\phi_2 - \phi_1)^+ = 0$ and $\phi_1 \ge \phi_2$.

Theorem 2.4 Let ϕ and ϕ_e be given by the solution of problem (5). Define the mapping

$$\mu : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$$
$$\beta \longmapsto M = \int_{\Omega} G(\phi - \phi_e - \beta) dx.$$

Then, μ is a nondecreasing, one to one and onto mapping.

Proof: Let $\beta_1, \beta_2 \in \mathbb{R}^+$, with $\beta_1 \ge \beta_2$ then for i = 1, 2, we have

$$-\Delta \phi_i = \rho_i$$

$$\rho_i = G(\phi_i - \phi_e - \beta_i) \quad \text{and} \quad M_i = \int_{\Omega} G(\phi_i - \phi_e - \beta_i) dx.$$

We subtract two equations and taking $w = \phi_1 - \phi_2$, then

$$\begin{cases} -\Delta w = \rho_1 - \rho_2 & \text{in } \Omega \\ w = 0 & \text{on } \Gamma_1 \cup \Gamma_2 \\ \frac{\partial w}{\partial \nu} = 0 & \text{on } \Gamma_3, \end{cases}$$

Stationary Solutions to the Vlasov-Poisson

in other words, for any $\xi \in V$, we have

$$\int_{\Omega} \nabla w \cdot \nabla \xi = \int_{\Omega} (\rho_1 - \rho_2) \, \xi$$

Let us take $\xi = \mathcal{H}_{\varepsilon}^{+}(w)$ as a test function in the previous equality, where $\mathcal{H}_{\varepsilon}^{+}(.)$ is the approximation of the Heaviside function $\mathcal{H}^{+}(.)$ defined as

$$\mathcal{H}_{\varepsilon}^{+}(r) = \begin{cases} 1 & \text{if} \quad r > \varepsilon \\ \frac{1}{\varepsilon}r & \text{if} \quad 0 \le r \le \varepsilon \\ 0 & \text{if} \quad r < 0, \end{cases}$$

then

$$-\int_{\Omega} |\nabla w|^2 \mathcal{H}_{\varepsilon}'^+(w) = \int_{\Omega} (\rho_2 - \rho_1) \mathcal{H}_{\varepsilon}^+(w).$$

Now, letting $\varepsilon \to 0$, we get

$$\int_{\Omega} (\rho_2 - \rho_1) \mathcal{H}^+(w) \le 0.$$

Using the proposition 2.3, we have $\beta_1 \ge \beta_2$ then $w = \phi_1 - \phi_2 \ge 0$ and $\mathcal{H}^+(w) = 1$. Finally, we have

$$M_1 := \int_{\Omega} \rho_1 \ge \int_{\Omega} \rho_2 := M_2$$

Let us introduce the spaces Φ_p defined by

 $\Phi_p = \{ v \in W^{1,p}(\Omega); \ \Delta v \in L^p(\Omega), v = 0 \ \text{ on } \ \Gamma_1 \cup \Gamma_2 \}$

endowed with the norm $||.||_{\Phi_p}$ where

$$||u||_{\Phi_p} \sim ||\Delta u||_{L^p(\Omega)}$$
 for all $u \in \Phi_p$.

Lemma 2.5 Let f the solution of the problem (4). Then $\rho[f] \in L^{\infty}(\Omega)$ and $\phi[f] \in \Phi_p$.

Proof: We know that the function f satisfying

$$f(x,v) = \gamma \left(\frac{1}{2} |v|^2 + \phi[f](x) - \phi_e(x) - \beta\right)$$
 and $\int_{\Omega \times \mathbb{R}^2} f(x,v) = M.$

Using the weak maximum principle, we find $\phi[f](x) \ge 0$ in Ω . On the other hand, thanks to (H_1) we have $\rho_e \in L^p(\Omega)$ and $\phi_{in} \in W^{1-1/p, p}(\Gamma_2) \cap L^{\infty}(\Gamma_2)$, then $\phi_e \in W^{1, p}(\Omega) \subset L^{\infty}(\Omega)$. Using the fact that γ is a decreasing function, we have

$$f(x,v) \le \gamma \left(\frac{1}{2} |v|^2 - \phi_e(x) - \beta\right) \le \gamma \left(\frac{1}{2} |v|^2 - ||\phi_e||_{\infty} - \beta\right)$$

Integrating this inequality of the variable v and using the monotonicity of γ , we have

$$\rho[f](x) \le 2^{\frac{N}{2}-1} |S^{N-1}| \int_0^{+\infty} s^{\frac{N}{2}-1} \gamma \left(s - ||\phi_e||_{\infty} - \beta\right) ds.$$

According to (H_2) , we deduce that $\rho[f] \in L^{\infty}(\Omega)$ and $\phi[f] \in \Phi_p$.

3 Corner behavior (N=2)

In this section we consider the case where Ω is a polygonal domain of \mathbb{R}^2 and denote by S the vertex of the re-entrant corner in Γ_1 , let $\frac{\pi}{\alpha}$ be the interior angle of Ω at the vertex and (r, θ) the polar coordinates centered at S. Using Grisvard's results [5], then for all $\phi \in \Phi_p$ with $1 , there exists <math>\lambda_m$, such that

$$\phi - \sum_{0 < m\alpha < 2-2/p} \lambda_m \chi(r) r^{m\alpha} \sin(m\alpha\theta) \in W^{2,p}(\Omega)$$

with $m \ge 1$ and $\chi \in \mathcal{C}^{\infty}(\mathbb{R}^+)$ is a cut-off function defined by

$$\chi(r) = \begin{cases} 1 & \text{if } 0 \le r < \eta \\ \\ 0 & \text{if } r \ge \eta, \end{cases}$$

where η is small enough. In particular, for $p \in \left[\frac{2}{2-\alpha}, \frac{1}{1-\alpha}\right]$ we have $\phi, \phi_e \in \Phi_p$ then its can be decomposed into the regular part $W^{2,p}(\Omega)$ and a singular part as follows:

$$\phi = \phi_{\mathcal{R}} + \lambda \, \chi(r) r^{\alpha} \sin(\alpha \theta)$$

$$\phi_e = \phi_{\mathcal{R}}^e + \lambda_e \, \chi(r) r^{\alpha} \sin(\alpha \theta)$$

where $\lambda = \int_{\Omega} \rho \, P_s \, dx$ and $\lambda_e = \int_{\Omega} \rho_e \, P_s \, dx$ with
 $-\Delta P_s = 0$ in $\Omega, \, P_s = 0$ on $\Gamma_1 \cup \Gamma_2$ and $\frac{\partial P_s}{\partial \nu} = 0$ on Γ_3 and
 $P_s = P_p + \tilde{P}$ where $P_p = \frac{1}{\pi} r^{-\alpha} \, \sin(\alpha \theta)$,

and $\tilde{P} \in H^1(\Omega)$ is the solution of

$$-\Delta \tilde{P} = \Delta P_p = 0, \qquad \tilde{P} = 0 \quad \text{on } \Gamma_1 \text{ and } \tilde{P} = P_p \quad \text{on } \Gamma_2 \cup \Gamma_3.$$

Let U be a bounded open subset of \mathbb{R}^+ , we denote by $L^p_{\tau}(U)$ the space

$$L^p_{\tau}(U) := \Big\{ u: \text{ measurable such that } ||u||^p_{L^p_{\tau}(U)} = \int_U |u(t)|^p t^{\tau} dt < \infty \Big\}.$$

We denote $W^{1,p}_{\tau}(U)$ the space of functions u of $L^p_{\tau}(U)$ such that $\frac{\partial u}{\partial t} \in L^p_{\tau}(U)$ where we denote the norm

$$||u||_{W^{1,p}_{\tau}(U)} = \left(\int_{U} |u(t)|^{p} t^{\tau} dt\right)^{1/p} + \left(\int_{U} |\frac{\partial u}{\partial t}|^{p} t^{\tau} dt\right)^{1/p}$$

Let \mathcal{H} and \mathcal{L} the linear operators defined by:

$$\mathcal{H}u(t) = \frac{1}{t} \int_0^t u(s) ds$$

$$\mathcal{L}u(t) = \frac{1}{t} \int_t^\infty u(s) ds$$

Thanks to [8] we have: If $\tau then$

$$||\mathcal{H}u||_{L^{p}_{\tau}(U)} \le C||u||_{L^{p}_{\tau}(U)}.$$
(6)

If $\tau > p-1$ then

$$||\mathcal{L}u||_{L^{p}_{\tau}(U)} \le C||u||_{L^{p}_{\tau}(U)}$$
(7)

where C is a constant which depends only on τ and p. Taking

$$w = \chi \phi_{\mathcal{R}} \text{ and } F = -\Delta w$$

where $F = -\chi \Delta \phi_{\mathcal{R}} - 2 \nabla \chi \nabla \phi_{\mathcal{R}} - \phi_{\mathcal{R}} \Delta \chi$. For all $p \in \left[\frac{2}{2-\alpha}, \frac{1}{1-\alpha}\right]$, we have $F \in L^{p}(\Omega)$
and $w \in W^{2,p}(\Omega) \cap W_{0}^{1,p}(\Omega)$, then $\nabla w \in [W^{1,p}(\Omega)]^{2}$ and $\frac{\partial w}{\partial \tau} = 0$, where $\tau(x)$ is a tangent
vector at a point $x \in \Gamma_{1}$.

Theorem 3.1 Let $p \in \left[\frac{2}{2-\alpha}, \frac{1}{1-\alpha}\right]$, ϕ be the solution to Problem (5) and $\phi_{\mathcal{R}} \in W^{2,p}(\Omega)$ the regular part, then there exists $g \in L^{\infty}(\Omega)$ such that

$$\chi(r)\phi_{\mathcal{R}}(r,\,\theta) = r\,\sin(\alpha\theta)g(r,\,\theta),\quad ||g(r,\,\theta)||_{L^{\infty}(\Omega)} \le C\,||\phi_{\mathcal{R}}||_{W^{2,p}(\Omega)}$$

where C is a constant which depends only on α , p and Ω .

Proof: Let $p \in \left[\frac{2}{2-\alpha}, \frac{1}{1-\alpha}\right]$, we have $w \in W^{2,p}(\Omega)$ then using the polar coordinates, it is not difficult to see that $\frac{\partial^2 w}{\partial r^2}, \frac{1}{r} \frac{\partial^2 w}{\partial r \partial \theta} - \frac{1}{r^2} \frac{\partial w}{\partial \theta}, \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} - \frac{1}{r} \frac{\partial w}{\partial r} \in L^p(\Omega)$. We have $\nabla w \in [W^{1,p}(\Omega)]^2$ with $p > \frac{2}{2-\alpha} \ge 2$, then $\nabla w \in [C(\overline{\Omega})]^2$. Since w = 0 in $\partial\Omega$, then $\frac{\partial w}{\partial \tau} = 0$ and the normal components of the gradient is perpendicular to line segment Γ_1 . By continuity, the gradient of w vanishes at r = 0 and $\frac{\partial w}{\partial r}(0,0) = 0$, then

$$r^{-1}\frac{\partial w(r,\,\theta)}{\partial r} = \frac{1}{r}\int_0^r \frac{\partial^2 w(\tilde{r},\,\theta)}{\partial r^2} d\,\tilde{r}.$$

Applying (6) with $\tau = 1$ and we integrate at θ , we obtain

$$\int_{0}^{\omega} \int_{0}^{R} \left| \frac{\partial w(r, \theta)}{\partial r} \right|^{p} r^{1-p} \, dr d\theta \le C \int_{0}^{\omega} \int_{0}^{R} \left| \frac{\partial^{2} w(r, \theta)}{\partial r^{2}} \right|^{p} r \, dr d\theta.$$
(8)

Similarly, we have $w(0, \theta) = 0$ and thanks to (6) we have

$$\int_{0}^{\omega} \int_{0}^{R} \left| \frac{w(r,\,\theta)}{r^{2}} \right|^{p} r \, d\, r d\, \theta \le C \int_{0}^{\omega} \int_{0}^{R} \left| \frac{\partial w(r,\,\theta)}{\partial r} \right|^{p} r^{1-p} \, d\, r d\, \theta < \infty.$$

$$\tag{9}$$

Thanks to (8) and (9), we have

$$\frac{\partial}{\partial r}(\frac{w}{r}) = \frac{1}{r}\frac{\partial w}{\partial r} - \frac{w}{r^2} \in L^p(\Omega) \quad \text{and} \quad \frac{w}{r} \in W_1^{1,p}\Big(0,R;L^p(]0,\omega[)\Big).$$

Now, we have

$$\Delta w = \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} \in L^p(\Omega)$$

then $\frac{\partial^2}{\partial \theta^2}(\frac{w}{r^2}) = \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} \in L^p(\Omega)$. Thanks to the Nirenberg inequality (see [9]), we have

$$||\frac{1}{r}\frac{\partial}{\partial\theta}(\frac{w}{r})||_{L^p(\Omega)} = ||\frac{\partial}{\partial\theta}(\frac{w}{r^2})||_{L^p(\Omega)} \le C(p,\Omega)||\frac{w}{r^2}||_{L^p(\Omega)}^{1/2}||\frac{\partial^2}{\partial\theta^2}(\frac{w}{r^2})||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^p(\Omega)}^{1/2}||_{L^$$

then $\frac{w}{r} \in W^{1,p}(\Omega) \subset C(\overline{\Omega})$ and therefore $\frac{w}{r}$ is bounded on $\overline{\Omega}$. Now, let us we prove that

$$\frac{1}{r}\frac{\partial w}{\partial \theta} \in W^{1,p}(\Omega)$$

Indeed, we have $\frac{1}{r}\frac{\partial}{\partial\theta}(\frac{1}{r}\frac{\partial w}{\partial\theta}) = \frac{1}{r^2}\frac{\partial^2 w}{\partial\theta^2} \in L^p(\Omega)$. Since $w \in W^{1,p}(\Omega)$, then $\frac{1}{r}\frac{\partial w}{\partial\theta} \in L^p(\Omega)$. On the other hand, we have $w \in W^{2,p}(\Omega)$ then

$$\frac{\partial}{\partial r}(\frac{1}{r}\frac{\partial w}{\partial \theta}) = \frac{1}{r}\frac{\partial^2 w}{\partial r \partial \theta} - \frac{1}{r^2}\frac{\partial w}{\partial \theta} \in L^p(\Omega)$$

hence $\frac{1}{r}\frac{\partial w}{\partial \theta} \in W^{1,p}(\Omega) \subset C(\overline{\Omega})$ and $||\frac{1}{r}\frac{\partial w}{\partial \theta}||_{L^{\infty}(\Omega)} \leq C||\frac{1}{r}\frac{\partial w}{\partial \theta}||_{W^{1,p}(\Omega)}$. We have $\frac{w}{r}(r, 0) = 0$, then

$$\frac{w(r,\theta)}{r} = \int_0^\theta \frac{\partial}{\partial \theta} (\frac{w(r,\tilde{\theta})}{r}) d\,\tilde{\theta},$$

and

$$\frac{w(r,\theta)}{r\,\theta} \leq ||\frac{\partial}{\partial\theta}(\frac{w}{r})||_{L^{\infty}(\Omega)} \leq C||\frac{1}{r}\frac{\partial w}{\partial\theta}||_{W^{1,p}(\Omega)}.$$

Near $\theta = 0$, we have $\theta \sim \frac{\sin(\alpha \theta)}{\alpha}$ and

$$\left|\frac{w(r,\theta)}{r\,\sin(\alpha\theta)}\right| \le \frac{C}{\alpha} \left|\left|\frac{1}{r}\frac{\partial w}{\partial \theta}\right|\right|_{W^{1,p}(\Omega)}.$$
(10)

In the same manner, we have $\frac{w}{r}(r, \frac{\pi}{\alpha}) = 0$, then $|\frac{w(r, \theta)}{r(\alpha \theta - \pi)}| \leq \frac{C}{\alpha} ||\frac{1}{r} \frac{\partial w}{\partial \theta}||_{W^{1,p}(\Omega)}$. Near $\theta = \frac{\pi}{\alpha}$, we have $\alpha \theta - \pi \sim \sin(\alpha \theta - \pi) = -\sin(\alpha \theta)$ and then (10) is satisfied. For $\theta \in [\theta_0, \frac{\pi}{\alpha} - \theta_0]$ with θ_0 small enough, there exists a constant c > 0 such that $|\sin(\alpha \theta)| \geq c |\alpha \theta|$ and the inequality (10) is satisfied. This completes the proof of Theorem.

Theorem 3.2 Let ϕ_1 and ϕ_2 be two solutions of problem (5) corresponding to M_1 and M_2 respectively.

If
$$M_1 \ge M_2$$
, then $\lambda_1 \ge \lambda_2$

where λ_1 and λ_2 are the singularity coefficients of ϕ_1 and ϕ_2 respectively.

Proof: Thanks to Theorem 3.1, we have

 $\phi_1 - \phi_2 \sim (\lambda_1 - \lambda_2) r^{\alpha} \sin(\alpha \theta)$ in a neighbourhood of the corner.

Thanks to Theorem 2.4, we have $\beta_1 \ge \beta_2$, then using the proposition 2.2 we obtain $\phi_1 \ge \phi_2$ and therefore $\lambda_1 \ge \lambda_2$.

Remark 3.3 1) Taking $\psi = \phi - \phi_e$ and let λ and λ_e the singularity coefficients of ϕ and ϕ_e respectively. If $\lambda \geq \lambda_e$ then $\psi^- \in H^2(\Omega)$. Indeed,

$$\phi - \phi_e \sim (\lambda - \lambda_e) r^\alpha \sin(\alpha \theta) \quad \forall (r, \theta) \in \mathcal{V},$$

where \mathcal{V} is the neighborhood of the corner. Since $\lambda \geq \lambda_e$, then $\psi^- = (\phi - \phi_e)^- = 0$ in \mathcal{V} . Consequently $\psi^- = (\phi_{\mathcal{R}} - \phi_{\mathcal{R}}^e)^- \in H^2(\Omega)$.

2) For $\rho_e \in L^{\infty}(\Omega)$ and $\beta \geq -G^{-1}(||\rho_e||_{\infty})$, we have $\phi \geq \phi_e$ in Ω . Subtracting the two equations verified by ϕ and ϕ_e , we have

$$-\Delta(\phi - \phi_e) - \left(G(\phi(x) - \phi_e(x) - \beta) - G(-\beta)\right) = G(-\beta) - \rho_e.$$

Taking $(\phi_e - \phi)^+$ as a test function in the weak formulation of this equation, using the same arguments as in the proposition 2.2, the monotonicity of the function γ and we find the desired result.

3.1 Behavior of Maxwell-Boltzmann Problem

In the case of Maxwellian velocity distribution e.g. $\gamma(s) = \exp(-s)$ the problem becomes:

$$\begin{aligned} & & -\Delta\phi = \kappa \exp(-\phi + \phi_e) := \rho, & \text{in } \Omega \\ & & \phi = 0 & & \text{on } \Gamma_1 \cup \Gamma_2 \\ & & & \frac{\partial\phi}{\partial\nu} = = 0 & & \text{on } \Gamma_3 \end{aligned}$$
(11)

and $\int \rho[f] = M$, where $\kappa = \exp(\beta)$.

Theorem 3.4 Let ϕ , ϕ_e , M and κ be given by the solution of the problem (11) and λ be the singularity coefficient of ϕ . As $M \to 0$, we have

$$\kappa \sim M \Big(\int_{\Omega} \exp(\phi_e) dx \Big)^{-1}$$
 and $\lambda \sim \kappa \int_{\Omega} \exp(\phi_e) P_s dx.$

Proof: We recall that $\rho = \kappa \exp(\phi_e - \phi)$ and $\kappa = M \Big(\int_{\Omega} \exp(\phi_e - \phi) dx \Big)^{-1}$, then $M \Big(\int_{\Omega} \exp(\phi_e) dx \Big)^{-1} \le \kappa \le M \exp(||\phi||_{\infty}) \Big(\int_{\Omega} \exp(\phi_e) dx \Big)^{-1}.$

We have

$$\Phi_2 \subset H^{1+\alpha-\varepsilon}(\Omega) \subset C(\overline{\Omega}) \subset L^{\infty}(\Omega),$$

then

$$||\phi||_{\infty} \leq C||\phi||_{\Phi_2} \leq C||\rho||_{L^2(\Omega)}.$$

On the other hand, we have

$$||\rho||_{L^{2}(\Omega)} \leq ||\rho||_{L^{1}(\Omega)}^{1/2} ||\rho||_{\infty}^{1/2} = M^{1/2} \cdot ||\rho||_{\infty}^{1/2} \text{ and } ||\rho||_{\infty} \leq \kappa \exp(||\phi_{e}||_{\infty}).$$

Then, $||\phi||_{\infty} \leq C(M \kappa \exp(||\phi_e||_{\infty}))^{1/2}$ and

$$M\Big(\int_{\Omega} \exp(\phi_e) dx\Big)^{-1} \le \kappa \le M \exp\left(C M \kappa \exp\left(||\phi_e||_{\infty}\right)^{1/2}\right) \Big(\int_{\Omega} \exp(\phi_e) dx\Big)^{-1} dx$$

Since κ is nondecreasing with respect to M, then for $M \to 0$, we have $\kappa \to \kappa_0 \ge 0$ and $C M \kappa \exp(||\phi_e||_{\infty}) \to 0$, therefore

$$\kappa \sim M \Big(\int_{\Omega} \exp(\phi_e) dx \Big)^{-1}.$$

Now, we prove that $\lambda \sim \kappa \int_{\Omega} \exp(\phi_e) P_s dx$. We have

$$\lambda = \int_{\Omega} \rho P_s dx = \kappa \int_{\Omega} \exp(\phi_e - \phi) P_s dx.$$

As $M \to 0$, we have $\kappa \to 0$, $||\rho||_{\infty} \to 0$ and $\phi \to 0$ a.e. in Ω , then

$$\exp(\phi_e - \phi) \to \exp(\phi_e).$$

Since $|\exp(\phi_e - \phi)| \leq \exp(\phi_e)$, the Lebesgue's Dominated Convergence Theorem give $\exp(\phi_e - \phi) \rightarrow \exp(\phi_e)$ in $L^2(\Omega)$ and therefore we obtain

$$\frac{\lambda}{\kappa} = \int \exp(\phi_e - \phi) P_s \to \int \exp(\phi_e) P_s \quad \text{as } M \to 0.$$

Which concludes the proof.

Now, We explicit the dependence of ϕ and ρ with respect to the parameter κ and taking

$$w_{\kappa} = \frac{\phi_{\kappa}}{\kappa}$$
 and $f_{\kappa} = \frac{\rho_{\kappa}}{\kappa}$.

Proposition 3.5 We have $\frac{1}{\sqrt{\kappa}} \frac{\nabla \rho_{\kappa}}{\rho_{\kappa}}$ and $\frac{1}{\sqrt{\kappa}} \nabla \phi_{\kappa}$ are uniformly bounded in $L^2(\Omega)$.

Proof: The problem (11) can be rewritten in f_{κ} as:

$$\begin{cases} f_{\kappa} - \frac{1}{\kappa} \Delta \log(f_{\kappa}) = \frac{1}{\kappa} \rho_{e}, & \text{in } \Omega \\ f_{\kappa} = 1 & \text{on } \Gamma_{1} \\ f_{\kappa} = \exp(\phi_{in}) & \text{on } \Gamma_{2} \\ \frac{\partial f_{\kappa}}{\partial \nu} = 0 & \text{on } \Gamma_{3} \end{cases}$$
(12)

Taking $\log(f_{\kappa})$ as a test function in the weak formulation of (12), then

$$\int_{\Omega} f_{\kappa} \log(f_{\kappa}) + \frac{1}{\kappa} \int_{\Omega} \left| \frac{\nabla f_{\kappa}}{f_{\kappa}} \right|^2 = \frac{1}{\kappa} \int_{\Omega} \rho_e \log(f_{\kappa}).$$

Using the Young inequality, we obtain

$$\int_{\Omega} f_{\kappa} \log(f_{\kappa}) + \frac{1}{\kappa} \int_{\Omega} \left| \frac{\nabla f_{\kappa}}{f_{\kappa}} \right|^2 \le \frac{C_p}{2\kappa} \int_{\Omega} \rho_e^2 + \frac{1}{2\kappa C_p} \int_{\Omega} |\log(f_{\kappa})|^2.$$

Thanks to the Poincaré inequality, we have $||\log(f_{\kappa})||_{L^{2}(\Omega)} \leq C \left\|\frac{\nabla f_{\kappa}}{f_{\kappa}}\right\|_{L^{2}(\Omega)}$ and then

$$\frac{1}{\kappa} \left(1 - \frac{C^2}{2C_p} \right) \int_{\Omega} \left| \frac{\nabla f_{\kappa}}{f_{\kappa}} \right|^2 \le \frac{C_p}{2\kappa} \int_{\Omega} \rho_e^2 - \int_{\Omega} \rho_{\kappa} \log(f_{\kappa}).$$

Taking $C_p = C^2$, we obtain

$$\frac{1}{\kappa} \int_{\Omega} \left| \frac{\nabla f_{\kappa}}{f_{\kappa}} \right|^2 \le \frac{C^2}{\kappa} \int_{\Omega} \rho_e^2 - 2 \int_{\Omega} \rho_{\kappa} \log(f_{\kappa}).$$

On the other hand, we have $\int_{\Omega} f_{\kappa} \log(f_{\kappa}) \ge |\Omega| \min_{x \in \mathbb{R}} (x \log(x))$ and consequently

$$\left\|\frac{1}{\sqrt{\kappa}}\frac{\nabla f_{\kappa}}{f_{\kappa}}\right\|_{L^{2}(\Omega)} \leq C_{1}$$

and this shows the first estimate. Now, we have $\phi_{\kappa} = \phi_e - \log(f_{\kappa})$, then

$$\left\|\frac{1}{\sqrt{\kappa}}\nabla\phi_{\kappa}\right\|_{L^{2}(\Omega)} \leq \left\|\frac{1}{\sqrt{\kappa}}\nabla\phi_{e}\right\|_{L^{2}(\Omega)} + \left\|\frac{1}{\sqrt{\kappa}}\frac{\nabla f_{\kappa}}{f_{\kappa}}\right\|_{L^{2}(\Omega)} \leq C_{2}.$$

and the second estimate follows immediately.

Theorem 3.6 As $\kappa \to \infty$, we have

$$\phi \to \infty$$
 a.e. in Ω , $\frac{\phi}{\kappa} \to 0$ in $H^1(\Omega) \cap C(\overline{\Omega})$ and
 $\frac{\rho}{\kappa} \to 0$ in $L^p(\Omega)$, $\forall p < \infty$ and uniformly in every compact of Ω .

Proof: We recall that

$$-\Delta w_{\kappa} = f_{\kappa}, \ f_{\kappa} = \exp(-\kappa w_{\kappa} + \phi_e), \ -\Delta \phi_e = \rho_e.$$

Thanks to proposition 2.3, for all x the sequence $f_{\kappa}(x)$ is a decreasing and bounded from below by 0, then there exists $f \geq 0$ such that

$$f_{\kappa} \to f$$
 a.e. in Ω .

We have $w_{\kappa} \geq 0$ and $\phi_e \in L^{\infty}(\Omega)$, then

$$|f_{\kappa}(x)| \le \exp(\phi_e(x))$$
 a.e. in Ω and $\exp(\phi_e) \in L^{\infty}(\Omega)$, (13)

using Lebesgue's Dominated Convergence Theorem, we have $f_{\kappa} \to f$ in $L^p(\Omega)$ for all $p < \infty$. On the other hand, we have $||w_{\kappa}||_{\Phi_p} = ||f_{\kappa}||_{L^p(\Omega)}$ then there exists $w \in \Phi_p$ such that

$$w_{\kappa} \to w \quad \text{in } \Phi_p.$$

We have

$$\Phi_p \subset H^{1+\alpha-\varepsilon}(\Omega) \Subset H^{\frac{3}{2}+\varepsilon}(\Omega) \subset C(\overline{\Omega}),$$

then

$$w_{\kappa} \to w \quad \text{in } C(\overline{\Omega}) \quad \text{and} \quad -\Delta w = f.$$

Now, we will show that f = 0 and w = 0. Thanks to proposition 3.5, we have

$$\left\|\nabla w_{\kappa}\right\|_{L^{2}(\Omega),} \leq \frac{C}{\sqrt{\kappa}},$$

then f = w = 0 in Ω and therefore

$$w_{\kappa} \to 0$$
 in Φ_p .

Consequently, we have $\phi_{\kappa} \to +\infty$ a.e. in Ω . Now, using Dini's Theorem [10], we have $f_{\kappa} \to 0$ uniformly in every compact of Ω .

References

- N. BEN ABDALLAH AND J. DOLBEAULT, Relative entropies for kinetic equations in bounded domain (irreversibility, stationary solution, uniqueness), Arch. Rational Mech. Anal. 168 (2003) 253-298
- [2] M.J. CÁCERES, J.A. CARRILLO, J. DOLBEAULT, Nonlinear stability in L^p for a confined system of charged particles, SIAM J. Math. Anal. 34 (2002) 478-494
- [3] J. DOLBEAULT, Free energy and solutions of the Vlasov-Poisson-Fokker-Planck system: External potential and confinement (large time behavior and steady states), J. Math. Pures Appl. (9), 78 (1999) 121157.

- [4] J.L. LIONS, Quelques méthodes de résolution des problèmes aux limites non linéaires, Dunod Paris, 1969
- [5] P. GRISVARD, Singularities in Boundary Value Problems, Masson, Paris 1992
- [6] P. GRISVARD, Espaces intermédiaires entre espaces de Sobolev avec poids, Annali della Scuola Normale Superiore di Pisa-Classe di Scienze, Sér. 3, 17 no. 3 (1963) 255-296
- [7] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics. 24 Pitman, Boston, 1985
- [8] G. H. HARDY, J. E. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge University Press, Cambridge1934
- [9] L. NIRENBERG, On elliptic partial differential equations, Annali della Scuola Normale Superiore di Pisa - Classe di Scienze, Sér. 3, 13 no. 2 (1959) 115-162
- [10] W. RUDIN, Principles of Mathematical Analysis, McGraw Hill, third edition, 1976