



HAL
open science

Growth and photosynthetic responses to copper stress of an invasive cordgrass,

Enrique Mateos-Naranjo, Susana Redondo-Gómez, Jesús Cambrollé, M.
Enrique Figueroa

► To cite this version:

Enrique Mateos-Naranjo, Susana Redondo-Gómez, Jesús Cambrollé, M. Enrique Figueroa. Growth and photosynthetic responses to copper stress of an invasive cordgrass,. *Marine Environmental Research*, 2008, 66 (4), pp.459. 10.1016/j.marenvres.2008.07.007 . hal-00501976

HAL Id: hal-00501976

<https://hal.science/hal-00501976>

Submitted on 13 Jul 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Growth and photosynthetic responses to copper stress of an invasive cordgrass,
Spartina densiflora

Enrique Mateos-Naranjo, Susana Redondo-Gómez, Jesús Cambrollé, M.
Enrique Figueroa

PII: S0141-1136(08)00194-3
DOI: [10.1016/j.marenvres.2008.07.007](https://doi.org/10.1016/j.marenvres.2008.07.007)
Reference: MERE 3276

To appear in: *Marine Environmental Research*

Received Date: 6 May 2008
Revised Date: 25 July 2008
Accepted Date: 28 July 2008

Please cite this article as: Mateos-Naranjo, E., Redondo-Gómez, S., Cambrollé, J., Figueroa, M.E., Growth and photosynthetic responses to copper stress of an invasive cordgrass, *Spartina densiflora*, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.07.007](https://doi.org/10.1016/j.marenvres.2008.07.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Growth and photosynthetic responses to copper stress of an invasive**
2 **cordgrass, *Spartina densiflora***

3
4 Enrique Mateos-Naranjo, Susana Redondo-Gómez, Jesús Cambrollé and M. Enrique
5 Figueroa

6 *Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de*
7 *Sevilla, Apartado 1095, 41080 - Sevilla, Spain*

8
9
10
11 Corresponding author:

12 Enrique Mateos-Naranjo

13 Dpto. Biología Vegetal y Ecología, Facultad de Biología, Universidad de Sevilla, Av

14 Reina Mercedes s/n, 41012 Sevilla, Spain.

15 E-mail: emana@us.es

16 Tel.: +34-95-4557165; fax: +34-95-4615780

1 **Abstract**

2 *Spartina densiflora* Brongn., is found in coastal marshes of south-west Spain, growing
3 in sediments with between 300 – 3000 mg Cu kg⁻¹ total soil DW (450-4500 mg Cu kg⁻¹
4 supposing that the soil porosity is 0.5). An experiment was designed to investigate the
5 effect of copper from 0 to 5000 mg kg⁻¹ (64 mmol l⁻¹) on the photosynthetic apparatus
6 and the growth of *S. densiflora*. We also determined total ash, copper, calcium,
7 magnesium and phosphorous concentrations, as well as C/N ratio. *S. densiflora* survived
8 to concentrations as high as 320 mg Cu kg⁻¹ DW in leaves, although excess of Cu
9 diminished water use efficiency and Ca-, Mg- and P-uptake. Also, quantum efficiency
10 of PSII, net photosynthetic rate, stomatal conductance and pigment concentrations
11 declined with increasing external Cu. Finally, the decline in the photosynthetic function
12 resulted in a biomass reduction of between 50 and 80% (for 600 and 5000 mg Cu kg⁻¹,
13 respectively).

14
15 **Keywords:** Copper; Growth rate; Invader; Photochemistry; Photosynthesis;
16 Photosynthetic pigments; Salt marshes; *Spartina densiflora*; Tissue analysis

1 **1. Introduction**

2

3 Copper is an essential micronutrient for plant growth, participating in important
4 biological reactions as an enzymatic cofactor and electron carrier in the photosynthetic
5 and respiratory processes (Andrade et al., 2004) but, when in excess, it becomes highly
6 toxic (Dewez et al., 2005). The excess of copper reduces plant growth as well as
7 photosynthetic and respiratory activities (Nalewajko and Olaveson, 1995). The
8 photosynthetic apparatus is particularly susceptible to this cation, resulting in a decrease
9 in the activity of photosystem II and electron transfer rates (Fernades and Henriques,
10 1991; Mallick and Mohn, 2003). Toxic concentrations of Cu may eventually build up in
11 certain estuarine and shallow coastal marine habitats near mining and industrial hot
12 spots (Nielsen and Nielsen, 2005).

13 The austral cordgrass, *Spartina densiflora* Brongn. (Poaceae), is invading salt
14 marshes as far apart as southern Europe (Tutin, 1980; Mateos-Naranjo et al., 2007),
15 North Africa (Fennane and Mathez, 1988) and North America (Kittelson and Boyd,
16 1997). This species has proven to be a vigorous invader and ecosystem engineer that
17 spreads by clonal growth and prolific seed production (Mateos-Naranjo et al., 2008a). In
18 the salt marshes of the joint estuary of the Tinto and Odiel rivers (SW Spain), one of the
19 most polluted areas by heavy metals in the world (Ruiz, 2001; Sáinz and Ruiz, 2006) on
20 account of the intense mining activity and waste from industrial zones, *S. densiflora*
21 grows over sediments with between 300 – 3000 ppm Cu per total soil DW (Nelson and
22 Lamothe, 1993; Sáinz et al., 2002), and with the greatest mobility and bioavailability
23 (Morillo et al., 2004). Other species of the *Spartina* genus have been shown to have a
24 great capacity to accumulate metals, such as copper, from the sediments. The

1 physiological impact of such bioaccumulation, nonetheless, has not been thoroughly
2 studied (Nybakken, 1988).

3 The main objective of the present study was to evaluate the effects of elevated
4 copper levels on *S. densiflora* growth and photosynthesis. The specific objectives were
5 to: (1) analyze the growth of plants in experimental copper treatments ranging from 0 to
6 64 mmol l⁻¹ Cu; (2) ascertain the extent to which the effects on the photosynthetic
7 apparatus (PSII chemistry), on gas exchange parameters and on photosynthetic
8 pigments determine plant performance; and (3) examine the possible role of
9 concentrations of mineral matter (ash), calcium, magnesium, phosphorous and nitrogen
10 in explaining the copper effects on growth.

1 **2. Material and methods**

2

3 *2.1. Plant material*

4

5 Seeds of *S. densiflora* were collected in December 2006 from the Odiel Marshes
6 (Huelva, SW Spain), and subsequently stored at 4°C (in darkness) for three months.
7 After the storing period, seeds were placed in a germinator (ASL Aparatos Científicos
8 M-92004, Madrid, Spain) and subjected to an alternating diurnal regime of 16 hours of
9 light (photon flux rate, 400-700 nm, 35 $\mu\text{mol m}^{-2} \text{s}^{-1}$) at 25°C and 8 hours of darkness at
10 12°C, for a month. The seedlings were planted in individual plastic pots (11 cm of
11 diameter) filled with perlite and placed in a glasshouse (during spring 2007) with
12 minimum-maximum temperatures of 21-25°C, 40-60% relative humidity and natural
13 daylight (minimum and maximum light flux: 200 and 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$). The pots were
14 carefully irrigated with 20% Hoagland's solution (Hoagland and Arnon, 1938) as
15 necessary. All the pots received the same irrigation.

16

17 *2.2. Stress treatments*

18

19 In April 2007, after a month of seedling cultures, the pots were allocated to five Cu
20 treatments in shallow trays (ten pots per tray, with one tray per Cu treatment): 0, 9, 15,
21 47 and 64 mmol l^{-1} Cu, in the same glasshouse. Cu treatments were established by
22 combining 20% Hoagland's solution and $\text{CuSO}_4 \cdot 7\text{H}_2\text{O}$ of the appropriate concentration.
23 The control, 0 mmol l^{-1} Cu treatment, had exactly 0.0005 mmol l^{-1} of Cu, since
24 Hoagland's solution contains a small amount of Cu as an essential trace nutrient. Cu
25 concentrations were chosen to cover variations recorded by Nelson and Lamothe (1993;

1 300-3000 mg Cu kg⁻¹ per total soil DW, i.e. 450-4500 mg Cu kg⁻¹ supposing that the
2 soil porosity is 0.5) and Sáinz et al. (2002) in the salt marshes of the joint estuary of the
3 Tinto and Odiel rivers.

4 At the beginning of the experiment, 3 L of the appropriate solution were placed in
5 each of the trays down to a depth of 1 cm. During the experiment, the levels in the trays
6 were monitored and they were topped up to the marked level with 20% Hoagland's
7 solution (without additional CuSO₄·7H₂O) as a way to limit the change in the Cu
8 concentration due to water evaporation of the nutritive solution. In addition, the entire
9 solution (including CuSO₄·7H₂O) was changed on a weekly basis (The total duration of
10 the experiment was 30 d)

11 The possibility of adding NaCl to the culture medium was disregarded, since the
12 absence of salt has been proven to affect neither the photosynthetic function of *S.*
13 *densiflora* nor its growth. In fact, Mateos-Naranjo et al. (2008b) found the highest rates
14 of net photosynthesis and leaf elongation in distilled water.

15

16 2.3. Growth

17

18 At the beginning and at the end of the experiment (i.e. after 30 d of treatment), four and
19 six entire plants (roots and leaves), from each treatment, respectively, were dried at
20 80°C for 48 h and then weighed. Dried, ground samples were ignited in lidded, ceramic
21 crucibles and ash weights were recorded; the furnace temperature was raised slowly
22 over 6 h to 550°C and this temperature was maintained for a further 8 h. Also, both
23 before and after the copper treatment, the number and height of all fully developed
24 tillers were measured.

25 The relative growth rate (RGR) in ash-free dry mass of whole plants was calculated

1 by using the formula:

2

$$3 \text{ RGR} = (\ln B_f - \ln B_i) \cdot D^{-1} \quad (\text{g g}^{-1} \text{day}^{-1})$$

4

5 where B_f = final dry mass (an average of the four plants from each treatment dried at
6 the beginning of the experiment), B_i = initial dry mass and D = duration of experiment
7 (days).

8 Leaf elongation rate (LER) was measured in random leaves ($n = 12$, per treatment;
9 two measurements per plant) by placing a marker of inert sealant at the base of the
10 youngest accessible leaf. The distance between the marker and the leaf base was
11 measured after 24 h (Ewing et al., 1995).

12

13 *2.4. Determination of copper, calcium, magnesium, phosphorous and nitrogen*

14

15 In accordance with protocols of Redondo-Gómez et al. (2007), at the end of the
16 experiment, leaf and root samples were dried at 80°C for 48 h and ground. Leaves and
17 roots were carefully washed with distilled water before any further analysis. Then 0.5 g
18 samples (in triplicate), taken from a mixture of the leaves or the roots belonging to the
19 six plants used for each treatment, digested with 6 ml HNO_3 , 0.5 ml HF and 1 ml H_2O_2 .
20 Ca, Mg, P and Cu were measured by inductively coupled plasma (ICP) spectroscopy
21 (ARL-Fison 3410, USA). Total N and C concentrations were determined for undigested
22 dry samples by means of an elemental analyzer (Leco CHNS-932, Spain).

23

24 *2.5. Chlorophyll fluorescence*

25

1 Chlorophyll fluorescence was measured in random fully developed penultimate leaves
2 ($n = 12$, two measurements per plant) using a portable modulated fluorimeter (FMS-2,
3 Hansatech Instrument Ltd., England) after 7 and 30 d of treatment. Light- and dark-
4 adapted fluorescence parameters were measured at dawn (stable, $50 \mu\text{mol m}^{-2} \text{s}^{-1}$
5 ambient light) and at midday ($1600 \mu\text{mol m}^{-2} \text{s}^{-1}$).

6 The plants were dark-adapted for 30 minutes, using leaf-clips exclusively designed
7 for this purpose. The minimal fluorescence level in the dark-adapted state (F_0) was
8 measured using a modulated pulse ($<0.05 \mu\text{mol m}^{-2} \text{s}^{-1}$ for $1.8 \mu\text{s}$) which was too small
9 to induce significant physiological changes in the plant. The data stored were an
10 average taken over a 1.6 second period. Maximal fluorescence in this state (F_m) was
11 measured after applying a saturating actinic light pulse of $15000 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 0.7s.
12 The values of the variable fluorescence ($F_v = F_m - F_0$) and maximum quantum efficiency
13 of PSII photochemistry (F_v/F_m) were calculated from F_0 and F_m . This ratio of variable
14 to maximal fluorescence correlates with the number of functional PSII reaction centres,
15 and dark adapted values of F_v/F_m can be used to quantify photoinhibition (Krivosheeva
16 et al., 1996).

17 The same leaf section of each plant was used to measure light-adapted parameters.
18 Steady state fluorescence yield (F_s) was recorded after adapting plants to ambient light
19 conditions for 30 minutes. A saturating actinic light pulse of $15000 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 0.7
20 s was then used to produce the maximum fluorescence yield (F_m') by temporarily
21 inhibiting PSII photochemistry.

22 Using fluorescence parameters determined in both light- and dark-adapted states, the
23 following were calculated: quantum efficiency of PSII ($\Phi_{\text{PSII}} = (F_m' - F_s)/F_m'$), which
24 measures the proportion of the light absorbed by chlorophyll associated with PSII that is
25 used in photochemistry (Maxwell and Johnson, 2000); and non-photochemical

1 quenching ($NPQ = (F_m - F_m') / F_m'$; Redondo-Gómez et al., 2006), which is linearly
2 related to heat dissipation (Maxwell and Johnson, 2000).

3

4 *2.6. Gas exchange*

5

6 Gas exchange measurements were taken on random fully expanded leaves ($n = 10$, a
7 measurement per plant and four extra measurements taken randomly) 7 and 30 days
8 after the treatment, by using an infrared gas analyzer in an open system (LI-6400, LI-
9 COR Inc., Neb., USA). Net photosynthetic rate (A), intercellular CO_2 concentration
10 (C_i) and stomatal conductance to CO_2 (G_s) were determined at ambient CO_2
11 concentration of $360 \mu\text{mol mol}^{-1}$, temperature of $20/25^\circ\text{C}$, $50 \pm 5\%$ relative humidity
12 and a photon flux density of $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$. A , C_i and G_s were calculated using
13 standard formulae of Von Caemmerer and Farquhar (1981). The water use efficiency
14 (WUE) was calculated as the ratio between A and transpiration rate [$\text{mmol } (CO_2$
15 assimilated) mol^{-1} (H_2O transpired)].

16

17 *2.7. Photosynthetic pigments*

18

19 At the end of the experiment period, photosynthetic pigments in fully expanded
20 penultimate leaves ($n = 10$, a measurement per plant and four extra measurements taken
21 randomly) from each treatment were extracted using 0.05 g of fresh material in 10 ml of
22 80% aqueous acetone. After filtering, 1 ml of the suspension was diluted with a further
23 2 ml of acetone and chlorophyll a (Chl a), chlorophyll b (Chl b) and carotenoids ($Cx+c$)
24 contents were determined with a Hitachi U-2001 spectrophotometer (Hitachi Ltd,
25 Japan), by making use of three wavelengths (663.2 , 646.8 and 470.0 nm). The

1 concentrations of pigments ($\mu\text{g g}^{-1}$ fwt) were obtained by calculation, using the method
2 of Lichtenthaler (1987).

3

4 2.8. *Statistical analysis*

5

6 Statistical analysis was carried out using Statistica v. 6.0 (Statsoft Inc.). Pearson
7 coefficients were calculated to assess the correlation between different variables. Data
8 were analyzed using a one- and two-way analysis of variance (*F*-test). Data were first
9 tested for normality with the Kolmogorov-Smirnov test and for homogeneity of
10 variance with the Brown-Forsythe test and arc-sine transformed (Fv/Fm at dawn and net
11 photosynthetic rate data) in order to normalize the error distribution for ANOVA.
12 Significant test results were followed by Tukey tests for identification of important
13 contrasts. Differences between measurements of fluorescence at dawn and midday were
14 compared by the Student test (*t*-test) to investigate whether Cu concentration affected
15 the sensitivity of plants to photoinhibition.

1 **3. Results**

2

3 *3.1. Determination of copper, calcium, magnesium, phosphorous and C/N ratio*

4

5 There was a marked decrease in the mineral (ash) content of leaves (but not of the roots)
6 with increasing external Cu concentration ($r = -0.76$, $p < 0.01$). The ash content of roots
7 dropped when exposed to 15 mmol l^{-1} Cu (Fig. 1A).

8 Tissue copper concentrations were greater in the roots than in leaves (two-way
9 ANOVA, $p < 0.001$), and increased with external Cu concentration ($r = 0.89$, $p <$
10 0.0001 ; $r = 0.98$, $p < 0.0001$, for root and leaf, respectively; Fig 1B). In contrast, leaf
11 and root Ca, Mg and P concentrations diminished with increasing Cu concentration
12 (Figs. 1C-E).

Fig. 1

Carbon/nitrogen (C/N) ratio was higher in the leaves than in the roots of *S.*
14 *densiflora* (Fig. 1F). Both root and leaf C/N ratios tended to increase with external Cu
15 concentration, although leaf ratio dropped sharply when exposed to the highest Cu
16 concentration.

17

18 *3.2. Chlorophyll fluorescence*

19

20 The fluorescence parameters presented in Figure 2 correspond to plants after 30 d of
21 treatment. Fluorescence parameters after 7 d treatment did not show any difference with
22 the control and are not presented.

23 Values of Fv/Fm diminished, either at dawn or midday, with increasing Cu
24 concentration (dawn: $r = -0.85$, $p < 0.001$; midday: $r = -0.68$, $p < 0.001$; Fig. 3A).

25 Furthermore, Fv/Fm values were significantly higher for the control at both dawn and

1 midday (ANOVA, $p < 0.0001$ for each) because of higher values of F_m (data not
 2 presented) and lower values of F_0 . F_v/F_m was lower at midday, except for 64 mmol l^{-1}
 3 Cu, and the reductions resulted again mainly from the fact that values of F_m were lower
 4 at midday than at dawn (t -test, $p < 0.05$; Fig. 2A). On the other hand, the minimal
 5 fluorescence (F_0) increased with increasing Cu concentration at both dawn and midday
 6 ($r = 0.80$, $p < 0.001$; $r = 0.82$, $p < 0.0001$, respectively; Fig. 2B), while F_m remained
 7 steady. F_0 was inversely correlated with F_v/F_m at both dawn and midday ($r = -0.95$, $p <$
 8 0.05 ; $r = -0.93$, $p < 0.05$, respectively).

9 Quantum efficiency of PSII (Φ_{PSII}) decreased with increasing Cu concentration,
 10 either at dawn or midday ($r = -0.88$, $p < 0.0001$; $r = -0.75$, $p < 0.0001$; Fig. 2C). Φ_{PSII}
 11 values were significantly higher at midday than at dawn.

12 Non-photochemical quenching (NPQ) did not vary with Cu concentration at dawn,
 13 whereas at midday it increased up to 47 mmol l^{-1} Cu, and then it declined with the
 14 further increases in Cu concentration (Fig. 2D).

Fig. 2

16 3.3. Gas exchange and pigment concentration

17

18 The gas exchange parameters presented in Figure 3 correspond to plants after 30 d of
 19 treatment. Net photosynthetic rate (A) after 7 d treatment did not show any differences
 20 with the control (c. $15 \mu\text{mol m}^{-2} \text{ s}^{-1}$; $p > 0.05$) and hence it is not presented here.

21 Figure 3A shows that A declined sharply with the copper concentration of 15
 22 mmol l^{-1} and then stabilized at the highest concentrations. There was a strong linear
 23 relationship between A and RGR ($r = 0.91$, $p < 0.05$), and Φ_{PSII} ($r = 0.89$, $p < 0.05$).

Fig. 3

24 Stomatal conductance (Gs) showed the same pattern as A (Fig. 3B). In contrast,
 25 intercellular CO_2 concentration (C_i) increased significantly with Cu (Fig. 3C). Water

1 use efficiency (WUE), on the other hand, decreased significantly with Cu concentration
2 (Fig. 3D).

3 Pigment concentrations decreased with Cu concentration, showing the same
4 pattern as A and Gs (Fig. 4).

Fig. 4

6 3.4. Growth

7

8 Relative growth rate (RGR) was inhibited by Cu excess, as proves the fact that the
9 maximum value was recorded at the control (ANOVA, $p < 0.001$; Fig. 5A). In addition,
10 Cu treated plants showed chlorosis, especially at the two highest Cu concentrations.
11 RGR decreased with increasing Cu concentration ($r = -0.68$, $p < 0.0001$), and was
12 directly correlated with leaf elongation rate (LER; $r = 0.89$, $p < 0.05$), number of tillers
13 ($r = 0.95$, $p < 0.05$) and mean height of tillers ($r = 0.99$, $p < 0.001$; Figs. 5A-D).

Fig. 5

1 4. Discussion and conclusions

2

3 *Spartina densiflora* demonstrated a high tolerance to copper stress, since all plants
4 survived even at concentrations as high as 64 mmol l⁻¹ of Cu during 30 d. Paschke and
5 Redente (2002) determined the lethal concentrations of copper (LC50, substrate copper
6 concentration that kills 50% of plants) for six grass species used in restoration activities
7 in concentrations close to 5 mmol l⁻¹ after 60 d of treatment. In our glasshouse
8 experiment, we found that Cu levels were much higher in *S. densiflora* below- (72-1480
9 ppm Cu DW) than above-ground (10-320 ppm Cu DW) parts. A similar trend was
10 recorded for *Spartina maritima* (Reboredo, 1985) and *Spartina anglica* (Otte et al.,
11 1991) under field conditions. Luque et al. (1999) found Cu concentrations c. 24 mg kg⁻¹
12 DW in young leaves of *S. densiflora* sampled from Odiel salt marshes. In this site, 30
13 and 200 mg Cu kg⁻¹ DW have been quantified in young and old leaves of this species,
14 respectively; and 150 and 600 mg Cu kg⁻¹ DW for young and old leaves, respectively,
15 from the Tinto Marshes (Huelva, SW Spain; unpublished data). The lower Cu
16 concentration in leaves of *S. densiflora*, compared to that present in roots, could be
17 accounted for by the development of such mechanisms as compartmentation, which
18 could control the ion transport into leaves, thereby improving its tolerance to heavy
19 metals. The sequestering of metals in tissues or cellular compartments, which are less
20 sensitive to metals, has been described as a tolerance mechanism (Weis and Weis,
21 2004), entailing a restriction of the upward movement into the shoots (an avoidance
22 mechanism) and the translocation of excessive metals into the leaves (Verkleij and
23 Schat, 1990). Nevertheless, with excessive levels of soil Cu, this element may be
24 translocated from the roots and accumulated in the tops of the plant (Kabata-Pendias

1 and Pendias, 2001), which could explain the increase of Cu concentration recorded in
2 leaves of *S. densiflora* with increasing external copper.

3 On the other hand, roots of *S. densiflora* could also act as a ‘barrier’ to the uptake
4 or transport of Cu, since leaf concentration hardly increased when the external
5 concentration exceeded 9 mmol l⁻¹ Cu. In this regard, Verkleij and Schat (1990) showed
6 that metal-chelating substances may be present in plant exudates, which act to decrease
7 metal uptake, and consequently toxicity. In other species of *Spartina*, some authors have
8 observed the formation of an iron plaque on the roots, with the ensuing accumulation of
9 metals in the rhizosphere (Doyle and Otte 1997), this plaque being present as well in *S.*
10 *densiflora* (Mateos-Naranjo et al., 2008b). On the other hand, the metal excess in the
11 root zone may inhibit water uptake, thus explaining the recorded decline of water use
12 efficiency.

13 Due to the significant functions of Cu in enzymes and its variable valence, ions
14 with an affinity to proteins and other compounds similar to that of Cu may have
15 antagonistic interrelationships, a fact which could be the explanation for the decrease in
16 ash content (Kabata-Pendias and Pendias 2001). However, a rise in ash concentration
17 was recorded in roots of *S. densiflora* at the highest Cu concentration, which could be
18 explained by the strong accumulation of this metal, as the results of internal metal
19 concentrations confirmed. In our experiment, the presence of Cu affected the
20 concentration of the macroelements, Ca, Mg, N and P. An excess of Cu inhibits the
21 activity of phosphatase, thereby diminishing the availability of P (Tyler, 1976). Lin and
22 Wu (1994) found that in *Lotus purshianus*, an excess of Cu reduced the concentration of
23 P in both root and leaf tissues. Cu-Ca interactions are highly complex and apparently
24 cross-linked with the range of pH in the growth media (Kabata-Pendias and Pendias,
25 2001), and Mg is replaced in the chlorophyll by Cu (Küpper et al., 2002). Llorens et al.

1 (2000) also reported an enhancement of Cu, in respect to N, due to changes in nitrogen
2 metabolism with a reduction of total nitrogen in plant tissues. However, a rise in N
3 concentration was recorded in leaves of *S. densiflora* at the highest Cu concentration. In
4 this regard, Kabata-Pendias and Pendias (2001) explained that the concentrations of the
5 two elements are relative to the formation of strong protein complexes with Cu.
6 Furthermore, it is possible that *S. densiflora* leaves became less rigid at high Cu levels,
7 since Fernandes and Henriques (1991) reported that the high sensitivity of diatoms to
8 Cu results from competition between Cu and Si(OH)_4 for the same transport site.

9 Excessive Cu accumulated in plants tissue can be toxic to plants, affecting several
10 physiological and biochemical processes (Balsberg Pahlsson, 1989). Ouzounidou
11 (1994) explained that the marked impact of copper could be attributed to an overall
12 inhibition of photosystem II (PSII), due to chloroplast membrane leakage; this may be
13 associated with a substantial stress response. In this manner, F_v/F_m decreases are more
14 permanent than increases of dynamic inhibition ' ΦPSII '; the latter usually relaxes as
15 light intensity is lowered. F_v/F_m did show a significant reduction at midday compared to
16 dawn values. At midday, the reduction in F_v/F_m values indicated that *S. densiflora*
17 experienced photodamage at the highest light flux. ΦPSII decreased as a consequence of
18 the decrease in photochemical quenching (qP) and the increase in NPQ (up to 47 mmol
19 l^{-1}), which indicates that the plants dissipated light as heat, in this way protecting the
20 leaf from light-induced damage (Maxwell and Johnson, 2000). Nonetheless, at the
21 highest Cu concentration, NPQ diminished, indicating that excess levels of Cu affected
22 this mechanism associated with the adaptation to high light irradiances (Demmig-
23 Adams, 1998).

24 The decline of F_v/F_m in the presence of copper resulted from an increase in F_0 ,
25 while F_m was maintained at a stable level. F_0 increased despite the decline of

1 chlorophyll concentrations, these results being similar to results obtained by Ralph and
2 Burchett (1998). The higher values of F_0 indicated an impact on the PSII reaction
3 centre, as well as to reduction in the transfer of energy from the collecting antennas to
4 reaction centres (Ralph and Burchett, 1998). Padinha et al. (2000) also found that
5 photosynthetic efficiency was lower in plants of *Spartina maritima* from polluted sites.

6 On the other hand, Cu had a negative effect on net photosynthetic rate (A) and
7 stomatal conductance (Gs), without direct relation between both parameters since there
8 was not a reduction in intercellular CO₂ concentration (Ci). Burzynski and Zurek (2007)
9 found that lower Gs in cotyledons of *Cucumis sativus* treated with copper did not
10 influence A, as Ci remained unchanged, and was high. The decrease of A by Cu was
11 attributed to the influence of this metal on the activities of photosynthetic carbon
12 reduction cycle enzymes (PGK and GAPDH were inhibited by Cu), as well as loss of
13 chlorophyll pigments (Ouzounidou, 1994). Burzynski and Klobus (2004) suggested that
14 Cu initially destroys the photosynthetic carbon reduction cycle and subsequently
15 influences the photosynthetic electron transport. In the case of *S. densiflora*, the
16 decrease in pigment concentrations or increase in its degradation, and consequent
17 negative effect on photosynthetic electron transport, could lead as well to a decline in
18 the photosynthetic function. The substitution the central magnesium ion in the
19 chlorophyll by Cu may also damage the chlorophyll synthesizing system (Küpper et al.,
20 2002). In this respect, we recorded a decrease of Mg, whereas copper content increased
21 in the leaf tissue.

22 The decline in the photosynthetic function resulted in a biomass reduction of *S.*
23 *densiflora* which was apparent in the RGR of ash-free dry mass, number of tillers, leaf
24 elongation rate and mean height of tillers. Thus, the growth of *S. densiflora* was
25 inhibited by leaf tissue concentrations above 0.03 mg g⁻¹ dry mass.

1 *Spartina densiflora* survived to concentrations up to 320 mg Cu kg⁻¹ DW in
2 leaves, when it was treated with 5000 mg Cu l⁻¹ (64 mmol l⁻¹), despite the fact that
3 concentrations of Cu between 20 and 100 mg Cu kg⁻¹ DW in mature leaf tissue are
4 considered excessive or toxic (Kabata-Pendias and Pendias, 2001). Copper inhibited
5 water use efficiency, Ca-, Mg- and P-uptake, quantum efficiency of PSII, net
6 photosynthesis, stomatal conductance, and pigment synthesis, this resulting in a decline
7 of growth.

8

9 **Acknowledgements**

10

11 We are grateful to Mr. F. Fernández-Muñoz for technical assistance. We also thank the
12 Spanish Science and Technology and Environmental Ministries for their support
13 (projects 04212007 and PCI2006-A7-0641, respectively), and Seville University
14 Glasshouse General Service for collaboration.

15

1 **References**

- 2
- 3 Andrade, L.R., Farina, M., Filho, A.M.G., 2004. Effects of copper on *Enteromorpha*
- 4 *flexuosa* (Chlorophyta) in vitro. *Ecotoxicology and Environmental Safety* 58, 117-
- 5 125.
- 6 Balsberg Pahlsson, A.M., 1989. Toxicity of heavy metal (Zn, Cu, Cd, Pb) to vascular
- 7 plants. *Water Air and Soil Pollution* 66, 163-171.
- 8 Burzynski, M., Klobus, G., 2004. Changes of photosynthetic parameters in cucumber
- 9 leaves under Cu, Cd, and Pb stress. *Photosynthetica* 42, 505-510.
- 10 Burzynski, M., Zurek, A., 2007. Effects of copper and cadmium on photosynthesis in
- 11 cucumber cotyledons. *Photosynthetica* 45, 239-244.
- 12 Demmig-Adams, B., 1998. Survey of thermal energy dissipation and pigment
- 13 composition in sun and shade leaves. *Plant Cell Physiology* 39, 474-482.
- 14 Dewez, D., Geoffroy, L., Vernet, G., Popovic, R., 2005. Determination of
- 15 photosynthetic and enzymatic biomarkers sensitivity used to evaluate toxic effects of
- 16 copper and fludioxonil in alga *Scenedesmus obliquus*. *Aquatic Toxicology* 74, 150-
- 17 159.
- 18 Doyle, M.O., Otte, M.L., 1997. Organism-induced accumulation of iron, zinc and
- 19 arsenic in wetlands soils. *Environmental Pollution* 96, 1-11.
- 20 Ewing, K., McKee, K., Mendelsohn, I., Hester, M., 1995. A comparison of indicators
- 21 of sublethal salinity stress in the salt marsh grass, *Spartina Patens* (Ait.) Muhl.
- 22 *Aquatic Botany* 52, 59-74.
- 23 Fennane, M., Mathez, J., 1988. Nouveaux materiaux pour la flore de Maroc. Fascicule
- 24 3. *Naturalia Montspeliensia* 52, 135-141.
- 25 Fernandes, J.C., Henriques, F.S., 1991. Biochemical, physiological and structural

- 1 effects of excess copper in plants. *Botanical Review* 57, 246-273.
- 2 Hoagland, D., Arnon, D.I., 1938. The water culture method for growing plants without
3 soil. *California Agricultural Experiment Station Bulletin* 347, 1-39.
- 4 Kabata-Pendias, A., Pendias, H., 2001. Trace elements in soils and plants, CRC Press,
5 Boca Ratón, Florida, pp. 106-119.
- 6 Kittelson, P.M., Boyd, M.J., 1997. Mechanisms of expansion for an introduced species
7 of cordgrass, *Spartina densiflora*, in Humboldt Bay, California. *Estuaries* 20, 770-
8 778.
- 9 Krivosheeva, A., Tao, D.L., Ottander, C., Wingsle, G., Dube, S.L., Öquist, G., 1996.
10 Cold acclimated and photoinhibition in Scots pine. *Planta* 200, 296-305.
- 11 Küpper, H., Setlik, I., Spiller, M., Küpper, F.C., Prásil, O., 2002. Heavy metal-induced
12 inhibition of photosynthesis: targets of in vivo heavy metal chlorophyll formation.
13 *Journal of Phycology* 38, 429-441.
- 14 Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic
15 biomembranes. *Methods in Enzymology* 148, 350-382.
- 16 Lin, S.L., Wu, L., 1994. Effect of copper concentration on mineral nutrient uptake and
17 copper accumulation in protein of copper-tolerant and non-tolerant *Lotus purshianus*
18 L. *Ecotoxicology and Environmental Safety* 29, 214-228.
- 19 Llorens, N., Arola, L., Blade, C., Mas, A., 2000. Effects of copper exposure upon
20 nitrogen metabolism in tissues cultured *Vitis vinifera*. *Plant Science* 160, 159-163.
- 21 Luque, C.J., Castellanos, E.M., Castillo, J.M., Gonzalez, M., Gonzalez-Vilches, M.C.,
22 Figueroa, M.E., 1999. Metals in halophytes of a contaminated estuary (Odiel
23 Saltmarshes, SW Spain). *Marine Pollution Bulletin* 38, 49-51.
- 24 Mallick, N., Mohn, F.H., 2003. Use of chlorophyll fluorescence in metal-stress
25 research: a case study with the green microalga *Scenedesmus*. *Ecotoxicology and*

- 1 Enriromental Safety 55, 64-69.
- 2 Mateos-Naranjo, E., Redondo-Gómez, S., Silva, J., Santos, R., Figueroa, M.E., 2007.
- 3 Effect of prolonged flooding on the invader *Spartina densiflora* Brong. Journal of
- 4 Aquatic Plant Management 45, 121-123.
- 5 Mateos-Naranjo, E., Redondo-Gómez, S., Luque, C.J., Castellanos, E.M., Davy, A.J.,
- 6 Figueroa, M.E. (2008a, in press). Environmental limitations on recruitment from
- 7 seed in invasive *Spartina densiflora* on a southern European salt marsh. Estuarine
- 8 Coastal and Shelf Science.
- 9 Mateos-Naranjo, E., Redondo-Gómez, S., Cambrollé, J., Luque, T., Figueroa, M.E.
- 10 (2008b, in press). Growth and photosynthetic responses to zinc stress of an
- 11 invasive cordgrass, *Spartina densiflora*. Plant Biology.
- 12 Maxwell, K., Johnson, G.N., 2000. Chlorophyll fluorescence- a practical guide. Journal
- 13 of Experimental Botany 51, 659-668.
- 14 Morillo, J., Usero, J., Gracia, I., 2004. Heavy metal distribution in marine sediments
- 15 from the southwest coast of Spain. Chemosphere 55, 431- 442.
- 16 Nalewajko, C., Olaveson, M.M., 1995. Differential responses of growth,
- 17 photosynthesis, respiration, and phosphate uptake to copper in copper-tolerant and
- 18 copper-intolereant strains of *Scenedesmus acutus* (Chlorophyceae). Canadian
- 19 Journal of Botany 73, 1295-1303.
- 20 Nelson, C.H., Lamothe, P.J., 1993. Heavy metal anomalies in the Tinto and Odiel River
- 21 and estuary system, Spain. Estuaries 16, 496-511.
- 22 Nielsen, H.D., Nielsen, S.L., 2005. Photosynthetic responses to Cu²⁺ exposure are
- 23 independent of light acclimation and uncoupled from growth inhibition in *Fucus*
- 24 *serratus* (Phaeophyceae). Marine Pollution Bulletin 51, 715-721.

- 1 Nybakken, J. W., 1988. *Marine Biology: an ecological approach* (2nd ed.). Harper and
2 Row Publishers, New York.
- 3 Ouzounidou, G., 1994. Copper-induced changes on growth, metal content and
4 photosynthetic function of *Alyssum montanum* L. plants. *Environmental and*
5 *Experimental Botany* 34, 165-172
- 6 Otte, M.L., Bestebroer, S.L., Van der Linden, J.M., Rozema, J., Broekman, R.A., 1991.
7 A survey of zinc, copper, and cadmium concentrations in salt marsh plants along
8 the Dutch coast. *Environmental Pollution* 72, 175-189.
- 9 Padinha, C., Santos, R., Brown, M.T., 2000. Evaluating environmental contamination in
10 Ria Formosa (Portugal) using stress indexes of *Spartina maritima*. *Marine*
11 *Environmental Research* 49, 67-78.
- 12 Paschke, M.W., Redente, E.F., 2002. Copper toxicity thresholds for important
13 restoration grass species of the western United States. *Environmental Toxicology*
14 *and Chemistry* 21, 2692-2697.
- 15 Ralph, P.J., Burchett, M.D., 1998. Photosynthesis response of *Halophila ovalis* to heavy
16 metals stress. *Environmental Pollution* 103, 91-101.
- 17 Reboredo, F.H.S., 1985. Heavy metal analysis of sediments and live tissues of *Spartina*
18 *maritima*, from salt marshes of Sado estuary Portugal. Proceedings of 5th
19 Conference on 'Heavy Metals in the Environment'. Heidelberg (CEP Consultants
20 Ltd, ed.), Edinburgh UK.
- 21 Redondo-Gómez, S., Wharmby, C., Castillo, J.M., Mateos-Naranjo, E., Luque, C.J., de
22 Cires, A., Luque, T., Davy, A.J., Figueroa, M.E., 2006. Growth and photosynthetic
23 responses to salinity in an extreme halophyte, *Sarcocornia fruticosa*. *Physiologia*
24 *Plantarum* 128, 116-124.
- 25 Redondo-Gómez, S., Mateos-Naranjo, E., Davy, A.J., Fernández-Muñoz, F.,

- 1 Castellanos, E., Luque, T., Figueroa, M.E., 2007. Growth and photosynthetic
2 responses to salinity of the salt-marsh shrub *Atriplex portulacoides*. *Annals of*
3 *Botany* 100, 555-563.
- 4 Ruiz, F., 2001. Trace metals in estuarine sediments of south-western Spain. *Marine*
5 *Pollution Bulletin* 42, 481-489.
- 6 Sáinz, A., Grande, J.A., De la Torre, M.L., Sánchez-Rodas, D., 2002. Characterisation
7 of sequential leachate discharge of mining waste rock dumps in the Tinto and
8 Odiel rivers. *Journal of Environmental Management* 64, 345-353.
- 9 Sáinz, A., Ruiz, F., 2006. Influence of the very polluted inputs of the Tinto-Odiel
10 system on the adjacent littoral sediments of southwestern Spain: A statistical
11 approach. *Chemosphere* 62, 1612-1622.
- 12 Tutin, T.G., 1980. *Spartina* Schreber, in: Tutin, T.G., Heywood, V.H., Burges, N.A.,
13 Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A. (Eds.), *Flora Europaea*,
14 Vol. 5. Cambridge University Press, Cambridge, pp. 259-260.
- 15 Tyler, G., 1976. Heavy metal pollution, phosphatase activity and mineralization of
16 organic phosphorous in forest soil. *Soil Biology & Biochemistry* 8, 327-332.
- 17 Verkleij, J.A., Schat, H., 1990. Mechanism of metal tolerance in higher plants, in:
18 Shaw, A.J., *Heavy metal tolerance in plants: evolutionary aspects*. CRC Press,
19 Boca Raton, FL, USA, pp. 179-194.
- 20 Von Caemmerer, S., Farquhar, G.D., 1981. Some relationships between the
21 biochemistry of photosynthesis and the gas exchange of leaves. *Planta* 153, 377-
22 387.
- 23 Weis, J.S., Weis, P., 2004. Metal uptake, transport and release by wetland plants:
24 implications for phytoremediation and restoration. *Environment international* 30,
25 687-700.


1 Fig 1. Ash (A), total copper (B), total calcium (C), total magnesium (D) and total
2 phosphorous (E) concentrations, and C/N ratio (F) for leaves (\circ) and roots (\bullet) of
3 *Spartina densiflora* in response to treatment with a range of Cu concentrations after a
4 month. Values represent mean \pm SE, n = 3.

5
6 Fig 2. Maximum quantum efficiency of PSII photochemistry, F_v/F_m (A), minimal
7 fluorescence, F_0 (B), quantum efficiency of PSII, Φ PSII (C), and non-photochemical
8 quenching, NPQ (D), at mid-day (\bullet) and at dawn (\circ) in randomly selected, fully
9 expanded penultimate leaves of *Spartina densiflora* in response to treatment with a
10 range of Cu concentrations for a month. Values represent mean \pm SE, n = 12.

11
12 Fig 3. Net photosynthetic rate, A (A), stomatal conductance, Gs (B), intercellular CO₂
13 concentration, Ci (C), and water use efficiency (D) in randomly selected, fully expanded
14 penultimate leaves of *Spartina densiflora* in response to treatment with a range of Cu
15 concentrations for a month. Values represent mean \pm SE, n = 10.

16
17 Fig 4. Chlorophyll a, chl a (A), Chlorophyll b, chl b (B), and carotenoid, Cx+c (C)
18 concentrations in randomly selected, fully expanded penultimate leaves of *Spartina*
19 *densiflora* in response to treatment with a range of Cu concentrations for a month.
20 Values represent mean \pm SE, n = 10.


21
22 Fig 5. Growth analysis of *Spartina densiflora* in response to treatment with a range of
23 Cu concentrations over a month. Relative growth rate (A), leaf elongation rate (B),
24 number of tillers (C), and mean height of tiller (D). Values represent mean \pm SE, n = 6.
25 The analysis was carried out on an ash-free dry mass basis.



1


2 Fig.1

1 Fig.2




A

1 Fig.3




1 Fig.4



A

1 Fig.5



2