



HAL
open science

Genetic diversity and fitness in small populations of partially asexual, self-incompatible plants.

Miguel Navascués, Solenn Stoeckel, Stephanie Mariette

► To cite this version:

Miguel Navascués, Solenn Stoeckel, Stephanie Mariette. Genetic diversity and fitness in small populations of partially asexual, self-incompatible plants.. *Heredity*, 2010, 104 (5), pp.482-92. 10.1038/hdy.2009.159 . hal-00505867v1

HAL Id: hal-00505867

<https://hal.science/hal-00505867v1>

Submitted on 31 Jan 2012 (v1), last revised 31 Jan 2012 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Genetic diversity and fitness in small populations of partially asexual, self-**
2 **incompatible plants**

3

4 Miguel Navascués^{1,2}, Solenn Stoeckel³ and Stéphanie Mariette⁴

5

6 ¹Unidad de Genética, Centro de Investigación Forestal, INIA, Carretera de La Coruña km 7.5,
7 28040 Madrid, Spain

8 ²Équipe Éco-évolution Mathématique, CNRS, UMR 7625 Écologie et Évolution, Université
9 Pierre et Marie Curie & École Normale Supérieure, 46 rue d'Ulm 75230 Paris Cedex 05,
10 France

11 ³UMR 1099 BIO3P (Biology of Organisms and Populations applied to Plant Protection),
12 INRA Agrocampus Rennes, F-35653 Le Rheu, France

13 ⁴Unité de Recherche sur les Espèces Fruitières, INRA, Domaine de la Grande Ferrade, 71
14 avenue Edouard Bourlaux, BP 81, 33883 Villenave d'Ornon Cedex, France

15 Author for correspondence: Stéphanie Mariette, Tel: (33) 557 122 383, Fax: (33) 557 122
16 439, E-mail: smariett@bordeaux.inra.fr

17

18 **Keywords**

19 Self-incompatibility, asexual reproduction, number of *S*-alleles, linkage disequilibrium,
20 inbreeding depression, mutation load

21 **Running title:** Asexuality and self-incompatibility in small populations

22 **Word count:** 5433

23

24 **Summary**

25 How self-incompatibility systems are maintained in plant populations is still a debated issue.
26 Theoretical models predict that self-incompatibility systems break down according to the
27 intensity of inbreeding depression and number of *S*-alleles. Other studies have explored the
28 role of asexual reproduction in the maintenance of self-incompatibility. However, the
29 population genetics of partially asexual, self-incompatible populations are poorly understood
30 and previous studies have failed to consider all possible effects of asexual reproduction or
31 could only speculate on those effects. In the present study, we investigated how partial
32 asexuality may affect genetic diversity at the *S*-locus and fitness in small self-incompatible
33 populations. A genetic model including an *S*-locus and a viability locus was developed to
34 perform forward simulations of the evolution of populations of various sizes. Drift combined
35 with partial asexuality produced a decrease in the number of alleles at the *S*-locus. In addition,
36 an excess of heterozygotes was present in the population, causing an increase in mutation
37 load. This heterozygote excess was enhanced by the self-incompatibility system in small
38 populations. In addition, in highly asexual populations, individuals produced asexually had
39 some fitness advantages over individuals produced sexually (due to the increased
40 heterozygosity, sex produces the homozygosis of deleterious alleles). Our results suggest that
41 future research on the role of asexuality for the maintenance of self-incompatibility will need
42 to (1) account for whole genome fitness (mutation load generated by asexuality, self-
43 incompatibility and drift) and (2) acknowledge that the maintenance of self-incompatibility
44 may not be independent of the maintenance of sex itself.

45

46 **Introduction**

47 Hermaphroditic plant species reproduce with variable rates of selfing, ranging from strict
48 selfing to strict outcrossing (Barrett, 2002). Self-incompatibility (SI) is a reproductive system
49 that prevents self-fertilization. In the case of heteromorphic self-incompatibility, distinct
50 morphologies result in distinct compatibility groups, whereas in the case of homomorphic
51 self-incompatibility, compatible individuals cannot be distinguished by their morphology (de
52 Nettancourt, 1977). Most SI systems depend on physiological mechanisms that prevent pollen
53 germination or pollen tube growth. In sporophytic self-incompatibility (SSI) systems, the
54 compatibility of a pollen grain depends on the diploid genotype of the plant that produced it.
55 In gametophytic self-incompatibility (GSI) systems, the compatibility of a pollen grain
56 depends on its haploid genotype. GSI is more widespread than SSI (Glémin *et al.*, 2001).
57 Fisher (1941) showed that self-fertilization should have a selective advantage because a
58 selfing genotype will transmit more copies of its genome than a non-selfing genotype (this has
59 been termed the automatic advantage of selfing). However, numerous studies have shown that
60 inbred offspring are less fit than outbred offspring. The relative decrease in the mean fitness
61 of selfed versus outcrossed individuals (inbreeding depression) is generally recognized as the
62 only main factor that counterbalances the selective advantage of selfing (Charlesworth and
63 Charlesworth, 1987).
64 Consequently, the level of inbreeding depression in populations should play a determining
65 role in the evolution of SI systems. Inbreeding depression decreases as population size
66 becomes smaller due to reduced polymorphism for selection to act on (Bataillon and
67 Kirkpatrick, 2000). Charlesworth and Charlesworth (1979) showed that the number of *S*-
68 alleles also plays a role in maintaining SI, since a low number of alleles will limit the number
69 of compatible crosses in the population. A decrease in population size can also cause a
70 reduction in the number of *S*-alleles (Brennan *et al.*, 2003), and a self-compatible mutant can

71 be positively selected for (Reinartz and Les, 1994). Thus, small populations may be
72 particularly prone to the breakdown of SI due to weak inbreeding depression and low
73 numbers of *S*-alleles. However, small self-incompatible populations may maintain high levels
74 of inbreeding depression due to a sheltered load of deleterious alleles linked to the *S*-locus
75 (Glémin *et al.*, 2001). The existence of a sheltered load has been demonstrated experimentally
76 in *Solanum carolinense* by Stone (2004) and Mena-Alí *et al.* (2009).

77 Overall, under a wide range of conditions, SI can evolve to self-compatibility. In effect, the
78 loss of SI systems is very frequent in plant evolution (Igic *et al.*, 2008). However, the reasons
79 for which some species maintain a SI system while other species lose it are not completely
80 understood. It has been suggested that asexual reproduction, “when an individual produces
81 new individuals that are genetically identical to the ancestor at all loci in the genome, except
82 at those sites that have experienced somatic mutations” (de Meeûs *et al.*, 2007), plays a role in
83 the maintenance or breakdown of SI. Two studies suggest that asexuality could relieve the
84 main selective pressures that favour the breakdown of SI. First, Chen *et al.* (1997) showed in
85 Australian Droseraceae that all self-incompatible taxa have effective forms of asexual
86 reproduction, whereas the obligatory sexual annual taxon, *Drosera glanduligera*, is self-
87 compatible. Their interpretation is that self-incompatible forms accumulate recessive lethal
88 polymorphisms, especially in association with biparental inbreeding generated by elevated
89 levels of asexual reproduction. The hypothetical high genetic load constitutes the selective
90 pressure that maintains the outbreeding mechanisms. Second, Vallejo-Marín and O'Brien
91 (2007) hypothesized that asexuality could provide reproductive assurance in cross-fertilizing
92 species subject to pollen limitation. They predicted that cross-fertilizing species subject to
93 pollen limitation would often have some means of asexual reproduction, and they found a
94 strong correlation between self-incompatibility and asexuality in *Solanum* (Solanaceae).
95 Conversely, Young *et al.* (2002) developed a contrasting hypothesis for the case of *Rutidosia*

96 *leiolepis* in which they found SSI and high rates of asexual reproduction. They suggested that
97 increased asexual reproduction causes mate limitation by reducing genotype diversity at the *S*-
98 locus, favouring a breakdown of self-incompatibility.

99 The net effect of asexual reproduction on the maintenance or breakdown of SI systems is
100 therefore still unknown. Moreover, it is unclear to what extent the effects of asexual
101 reproduction discussed by the various authors are actually present in partially asexual, self-
102 incompatible populations. Theoretical studies have described the effects of partial asexuality
103 on the inbreeding depression of self-compatible populations (Muirhead and Lande, 1997) and
104 the number of *S*-alleles (Vallejo-Marín and Uyenoyama, 2008). However, no model has been
105 developed so far to study the joint effect of asexuality, self-incompatibility and drift on the
106 evolution of diversity and fitness parameters for a hermaphroditic species. Using individual-
107 based simulations, we investigated the effect of (i) partial asexual reproduction and drift on
108 diversity at the *S*-locus, (ii) combined asexuality, self-incompatibility and drift on the
109 frequency of deleterious alleles. Thus, our main goal was to characterise the dynamics of two
110 key factors (number of *S*-alleles and inbreeding depression) for the maintenance of SI in
111 partially asexual, self-incompatible populations. It was not the aim of this work to estimate
112 the probability of invasion of self-compatible alleles or other modifiers of reproduction.

113

114 **Materials and Methods**

115 *Genetic model*

116 The model considered in this study was based on the model developed by Glémin *et al.*
117 (2001). This model consists of a population of N (four population sizes evaluated: 25, 50, 100
118 and 1000) diploid hermaphroditic individuals with a GSI system. In addition, our model also
119 included asexual reproduction at rate c (probability that an individual is generated by asexual
120 reproduction, seven values evaluated: 0, 0.5, 0.8, 0.9, 0.99, 0.999 and 1). We considered an

121 asexual reproduction event as the production of a new independent individual that is an exact
122 copy of the parental individual (or only different by somatic mutation) (e.g. de Meeûs *et al.*,
123 2007). As in Glémin *et al.* (2001), each individual genome possessed the *S*-locus, which
124 regulated SI, and a viability locus, whose state determined individual fitness. Two neutral loci
125 were also included for reference. The four loci were considered to be physically unlinked and
126 were inherited independently through sexual reproduction.

127 Sexual reproduction events were controlled by the *S*-locus: crosses between individuals were
128 only possible when the *S*-allele carried by the pollen grain was different from both *S*-alleles of
129 the pistil (i.e. at least three different *S*-alleles were necessary in the population for sexual
130 reproduction to occur). To focus on effects specifically attributable to asexuality on the
131 maintenance and breakdown of SI, we assumed unlimited pollen availability and that sexual
132 crosses were always fruitful. Thus, sexual reproduction was only limited when asexuality and
133 genetic drift reduced the number of *S*-alleles to less than three within a population. In this
134 case, independent of the original rate of asexual reproduction, all individuals reproduced
135 asexually ($c=1$) until mutation introduced a third *S*-allele. Then, c was reset to its original
136 value.

137 Individual fitness, f , was determined by the viability locus, which had two alleles A and a .
138 Strength of the dominance of allele A over deleterious allele a was regulated by the
139 dominance coefficient h , and strength of the selection against a was regulated by the selection
140 coefficient s . Thus, relative fitness of genotypes were: $f_{AA}=1$, $f_{Aa}=1-hs$ and $f_{aa}=1-s$ ($0 < s \leq 1$
141 and $0 < h \leq 1/2$). Three selection regimes at the viability locus were considered: (i) a neutral
142 case, to be able to study the effects of the interaction between SI and asexuality without the
143 interference of selection, where parameters s and h were set to zero; (ii) a case of mildly
144 deleterious and partially recessive mutations, for which values assigned to s and h were

145 respectively 0.1 and 0.2; and (iii) a case of highly recessive lethal mutations, values assigned
146 to s and h were respectively 1 and 0.02.

147 Mutations were allowed in all loci. Neutral loci mutated at rate $\mu_N=10^{-3}$, following a k -allele
148 model ($k=100$). Mutation at the S -locus followed an infinite-allele model (Kimura and Crow,
149 1964) with rate μ_S ; three values were considered for this rate ($\mu_S=10^{-3}$, 10^{-4} and 10^{-5} as in
150 Glémin *et al.*, 2001). Mutations at the viability locus occurred at rate $\mu_1=10^{-3}$ from A to a and
151 at rate $\mu_2=10^{-4}$ from a to A .

152

153 *Individual-based simulations*

154 Forward-time simulations of a population of N diploid individuals with no overlapping
155 generations were performed. Simulations started with all individuals carrying two unique
156 alleles at the S -locus ($2N$ different S -alleles in the whole population), alleles A or a randomly
157 assigned (with equal probability, 0.5) to the viability locus and any of the k alleles ($k=100$)
158 randomly assigned (with equal probability, $1/k$) to the two neutral loci.

159 At each generation, the number of individuals generated by asexual reproduction, x , was
160 drawn from a Poisson distribution with mean Nc , and the remaining $N-x$ individuals were
161 generated by sexual reproduction (parents contributing to these two types of offspring were
162 not determined in this step). Genotypes of the N individuals were generated one by one. If a
163 given individual originated by asexual reproduction, a genotype i was randomly drawn from
164 all individuals of the previous generation. That genotype was assigned to the individual with a
165 probability equal to the fitness of genotype i ; this will be called the selection step. If the
166 genotype was not successfully assigned, the genotype draft and the selection step was
167 repeated until a genotype passed the selection step. If the individual originated by sexual
168 reproduction, the simulation followed a different procedure. First, an individual from previous
169 generation was randomly drawn to act as a mother and the ovule haploid genotype was

170 randomly generated. Then, a second individual was drawn to act as a father and the pollen
 171 haploid genotype was randomly generated. If the *S*-allele at the pollen grain was not different
 172 from both *S*-alleles of the mother (i.e. incompatible pollen), the pollen was discarded. New
 173 fathers and pollen grain genotypes were generated until a compatible cross was obtained (i.e.
 174 pollen availability was unlimited). Then, a diploid genotype *i* was generated from the haploid
 175 genotypes of the ovule and the pollen. The selection step was carried out on genotype *i* as in
 176 the case of asexual reproduction; if the genotype was not assigned the process was repeated
 177 by drawing two new parents until a genotype was able to pass the selection step. Once all *N*
 178 genotypes were selected, mutations were applied to each individual. The numbers of mutants
 179 at the *S*-locus and at the two neutral loci were drawn from Poisson distributions with means
 180 $2N\mu_S$ and $2N\mu_N$ respectively. At the viability locus, a mutant was formed with a probability μ_1
 181 if the allele was *A* and a probability μ_2 if the allele was *a*. The rate of asexual reproduction
 182 was a fixed parameter of our model and did not change with the selection process.

183

184 *Monitored genetic indices*

185 Simulations were run for 50 000 generations before beginning to monitor the genetic indices,
 186 this allowed monitored indices to become stable and avoided the influence of initial
 187 conditions in their values (Glémin *et al.*, 2001). From generation 50 000, indices were
 188 calculated every 3 000 generations until 107 000 generations (i.e. 20 times). For the same
 189 combination of parameters, 50 replicates of the simulations were performed providing 1 000
 190 (indices calculated 20 times in each of the 50 replicates) evaluations of the monitored
 191 variables for each set of parameter values. Indices were calculated from all genotypes of the
 192 population (i.e. they were not estimates, but the true values of the population).

193 Genetic diversity indices: For each locus we calculated the number of alleles (n_a), allele

194 frequencies, effective number of alleles ($n_e = 1 / \sum_{i=1}^{n_a} p_i^2$, where p_i was the frequency of allele),

195 expected heterozygosity ($H_e = 1 - 1/n_e$), observed heterozygosity (H_o , proportion of
 196 heterozygous individuals) and the inbreeding coefficient ($F_{IS} = (H_e - H_o)/H_e$). Theoretical
 197 predictions for some of these indices were obtained from the work of several authors for some
 198 of the scenarios considered:

199 a) Vallejo-Marín and Uyenoyama (2008) studied the effects of partial asexuality in the
 200 number of alleles of the *S*-locus using a diffusion approximation and found that the number of
 201 common alleles (n_c , which is approximately equivalent to the effective number of *S*-alleles,
 202 n_e , Takahata, 1990) is determined by:

203 (1)
$$\frac{M}{\theta} = \frac{1 - F_{IS}}{2N\mu_s}$$

204 where $M = N(1-c)$ and $\theta = 2N\mu_s$. This equation was solved numerically to obtain the
 205 expected effective number of *S*-alleles in the simulations without selection on the viability
 206 locus.

207 b) Glémin *et al.* (2001) obtained the expected F_{IS} for a neutral locus linked to the *S*-locus as:

208 (2)
$$F_{IS} = \frac{f_S(1-c)}{1 + f_S(1-c)}$$

209 where f_S is the scaling factor of the genealogy of the *S*-locus and r' is the net recombination
 210 rate between the two loci. For the case of partial asexuality, the number of common alleles

211 was given by Equation 1 and the scaling factor was $f_S = \frac{r(1-c)}{2N\mu_s(1-c)}$

212 (Vallejo-Marín and Uyenoyama, 2008). To correct for the reduction of recombination due to
 213 asexuality, we used $r' = r(1-c)$, where r was the recombination rate during sexual
 214 reproduction (i.e. $r=0.5$ for unlinked loci). This approach ignored mutation on the neutral
 215 locus.

216 c) Finally, an expression of the expected F_{IS} of a neutral locus in a self-compatible, partially
 217 asexual population was derived following Balloux *et al.* (2003):

218 (3)
$$F_{IS} = \frac{\gamma(1-\nu)}{2N(1-\nu)s + \gamma}$$

219 where $\gamma = (1 - \mu_N)^2$ and s is the selfing rate. Selfing rate in Balloux *et al.* (2003) model is
 220 defined as the rate of self-fertilization of individuals; thus a fertilization event between two
 221 individuals from the same genet is considered an outcrossing event. Under random mating,
 222 $s = 1/N$ and for negligible mutation rates, $\gamma = 1$.

223 Linkage disequilibrium: The linkage disequilibrium between pairs of loci (S -locus-viability
 224 locus and the two neutral loci) was studied with the correlation coefficient between two loci,
 225 R_{GGD} , developed by Garnier-Géré and Dillmann (1992) which is particularly appropriate for
 226 partially asexual diploids (de Meeûs and Balloux, 2004).

227 Fitness: Mean fitness of the population (\bar{W}) was calculated from the genotype frequencies as
 228 $\bar{W} = \sum_{i,j} p_{ij} w_{ij}$, and the average fitness of individuals
 229 potentially produced by selfing (W_s), outcrossing (W_o) and asexuality (W_a) were calculated as

230
$$W_s = \sum_{i,j} p_{ij} w_{ij} s_{ij},$$

231 $W_o = \sum_{i,j} p_{ij} w_{ij} o_{ij}$ (where p_A and p_a are the frequencies of alleles
 232 A and a) and $W_a = \bar{W}$. Inbreeding depression (δ) and mutation load (L) were calculated as

233 $\delta = 1 - (W_s/W_o)$ and $L = 1 - \bar{W}$. In addition, we compared the fitness of individuals produced

234 by asexual reproduction (W_a) to the fitness of individuals produced sexually, using the ratio
 235 W_o/W_a . W_s , W_o and W_a were calculated as the fitness of potential offspring if all individuals
 236 were produced by selfing, by outcrossing and asexually, respectively. However, these modes
 237 of reproduction might not be actually present in the different scenarios considered (e.g. in

238 self-incompatible populations, no selfing occurs, but potential W_s and inbreeding depression
239 can be calculated).

240 Fixation probabilities: To interpret the results, we calculated the probabilities of four events:
241 fixation of alleles A and a in the viability locus, ‘fixation’ of viability locus heterozygote (Aa)
242 and fixation of a genotype in the S -locus (reduction of number of S -alleles to two, this was
243 only monitored in self-incompatible populations). These fixation probabilities were estimated
244 on 1 000 evaluations of the monitored variables as the proportion of times in which $p_a=0$,
245 $p_a=1$ and $H_o=1$ for the viability locus, or $n=2$ for the S -locus.

246

247 **Results**

248 *Effect of asexuality and drift on linkage disequilibrium between loci*

249 Probably the most obvious effect of partial asexual reproduction was the reduction of
250 recombination. Asexuality generated identical genotypes in offspring, so allele associations
251 were transmitted in the same way that they would be transmitted if they were physically
252 linked throughout the whole genome. In fully asexual populations, maximum linkage
253 disequilibrium was expected (equivalent to the linkage disequilibrium between fully linked
254 loci) and in partially asexual populations, recombination was reduced proportionally to the
255 rate of asexual reproduction. Figure 1 shows how non-random associations of alleles
256 increased with asexual reproduction rates. Linkage disequilibrium levels also depended on
257 population size because drift produced a departure from the expected values of frequencies of
258 allele combinations (Figure 1). Strong drift might have even caused some combinations of
259 alleles to be absent in the population.

260 In small populations, some generations with complete allelic association started to appear at
261 asexual reproduction rates of 0.8 and were predominant at rates of 0.99 and higher. In larger
262 populations, complete allelic association was only found in the fully asexual scenario. These

263 associations of alleles were observed between loci subject to selection and between neutral
264 loci, which showed that linkage disequilibrium was not a product of selective processes
265 (Figure 1a and Figure 1b).

266 Physical linkage between the *S*-locus and loci with deleterious alleles can decrease the
267 number of *S*-alleles (Uyenoyama, 2003). It can also dramatically increase inbreeding
268 depression in small populations (Glémin *et al.*, 2001). Therefore, for the present model, it was
269 important to discern whether the linkage disequilibrium generated by asexual reproduction is
270 sufficient for the selective forces of the *S*-locus to interact with the viability locus. The results
271 are presented below taking this perspective into account.

272

273 *Effect of asexuality and drift on diversity parameters: number of S-alleles and heterozygote*
274 *excess*

275 The effective number of *S*-alleles decreased with asexuality (Figure 2) as shown analytically
276 by Vallejo-Marín and Uyenoyama (2008). Balancing selection on the *S*-locus only occurred
277 during sexual reproduction events. Since these events were scarce at high rates of asexual
278 reproduction, the influence of balancing selection (which promotes high allelic diversity) on
279 the population diminished. This can temporarily stop sexual reproduction by reducing the
280 number of *S*-alleles to two (i.e. fixation of an *S*-locus genotype, Table 1).

281 No significant differences in the number of *S*-alleles were found among the three different
282 selection regimes for the viability locus (frequency distributions of the number of *S*-alleles
283 among different selection regimes were undistinguishable in a Kolmogorov-Smirnov test, *P*-
284 value>0.98). It must be noted that small populations (with high linkage disequilibrium) were
285 fixed for the wild type allele *A* most of the time (Table 2b) and this may explain the weak
286 influence of deleterious alleles on the *S*-locus.

287 Another general effect of partial asexual reproduction in diploids was the reduction of allele
288 segregation. This can increase heterozygosity by the independent accumulation of mutations
289 on the two alleles of an asexual lineage (Pamilo, 1987; Birky, 1996). To quantify this effect,
290 we measured the inbreeding coefficient F_{IS} , which compares observed and expected
291 heterozygosities.

292 Theoretical equilibrium values for F_{IS} at a neutral locus (unlinked to the S -locus) were very
293 similar in self-compatible and self-incompatible populations (Figure 3), and, for large
294 population sizes, the simulated populations followed the same pattern (Figure 3b). However,
295 for small population sizes, drift generated strong linkage disequilibrium between the S -locus
296 and other loci. Thus, balancing selection on the S -locus had a hitchhiking effect on other loci,
297 increasing their heterozygosity. This explained the contrasting F_{IS} values on simulated self-
298 compatible and self-incompatible populations (Figure 3a). Analytical predictions do not
299 reflect this dramatic difference because they do not account for linkage disequilibrium. F_{IS}
300 decreased with asexuality and this decrease was enhanced by self-incompatibility in small
301 populations, particularly noticeable at asexual reproduction rates higher than 0.8. This
302 decrease in F_{IS} observed in neutral loci also occurred at the viability locus (both for the mildly
303 deleterious and lethal recessive cases), which implied an increase in the frequency of the
304 deleterious allele (Figure 4).

305

306 *Effect of asexuality and drift on fitness*

307 For both self-compatible and self-incompatible populations, mutation load globally increased
308 with asexuality and population size, but a greater variance among generations and populations
309 was observed for very small populations (Figure 5). As mentioned above, the frequency of the
310 deleterious allele increased with asexuality in small self-incompatible populations, due to a
311 hitchhiking effect of the S -locus over the viability locus. In the self-incompatibility system,

312 when the population size was small enough, the mutation load increased with asexual
313 reproduction rates; this load resulted from the increase of the deleterious allele frequency
314 (Figure 5a and 5c). Under similar conditions in the self-compatible system, mutation loads
315 were lower than the mutation load observed in the self-incompatible system (Figure 5b and
316 5d). In contrast, inbreeding depression had similar values in self-incompatible and self-
317 compatible populations (Figure 6), showing some increases with asexual reproduction rate for
318 lethal recessive mutations that was more apparent in large populations.

319 The average fitness was higher in sexual populations than in asexual populations due to
320 Mendelian segregation (e.g. Figure 5; Chasnov, 2000; Kirkpatrick and Jenkins, 1989). The
321 relative fitness between individuals produced by sexual and asexual reproduction was
322 influenced very little by partial asexuality in a self-compatible population (Figure 7b and 7d).
323 However, in small self-incompatible populations, the asexual reproduction rate increased the
324 advantage of asexual compared to sexual reproduction (Figure 7a and 7c). These populations
325 consisted mainly of heterozygous individuals (Figure 3). Therefore, asexual reproduction
326 produced mainly Aa individuals, with only slightly lower fitness than the fittest AA haplotype.
327 However, sexual reproduction produced a high proportion of aa individuals, reducing the
328 average fitness of offspring, a reduction that was not compensated by the production of the
329 fittest AA individuals. This could potentially lead to a positive feedback effect on the asexual
330 reproduction rate, but this was not studied in our model where the asexual reproduction rate
331 was a fixed parameter.

332

333 **Discussion**

334 The number of S -alleles and inbreeding depression are considered to be the main factors that
335 influence the maintenance or breakdown of self-incompatibility systems (Charlesworth and
336 Charlesworth, 1979). A decrease in the number of S -alleles is expected to favour the

337 breakdown of self-incompatibility since self-compatible mutants can invade the population,
338 whereas the maintenance of inbreeding depression within populations prevents the breakdown
339 of self-incompatibility. In the present study, we examined how drift and partial asexuality
340 may modify the dynamics of these two key parameters in a self-incompatible population.

341

342 *Drift, combined with partial asexuality, dramatically reduced the number of S-alleles*

343 As already predicted by Yokoyama and Hetherington (1982), we showed that the number of
344 *S*-alleles decreases with the effective size of the population. Thus, in small populations,
345 repeated bottlenecks or founder events may lead to the breakdown of the self-incompatibility
346 system. However, Karron (1987) found no differences in mating systems between rare and
347 widespread congeners across several families (but see also a study on some Brassicaceae
348 species, Kunin and Shmida, 1997). As for the impact of asexuality, our simulation study
349 confirmed a recent result demonstrated analytically by Vallejo-Marín and Uyenoyama (2008).
350 The number of *S*-alleles decreased with asexuality due to the weakened influence of balancing
351 selection in a partially asexual population (Figure 2). In contrast with other effects of
352 asexuality that were only observed for very high rates of asexuality, intermediate asexual
353 reproduction rates were sufficient to significantly reduce the number of *S*-alleles. High rates
354 of asexual reproduction combined with drift produced extremely low numbers of *S*-alleles.
355 This effect was strong enough to even reduce the number of *S*-alleles to two (i.e. fixation of
356 an *S*-locus genotype, Table 1), which stopped all possibility of sexual reproduction until a
357 new *S*-allele arose from mutation or migration. Therefore, in such extreme cases, this drastic
358 effect of asexuality on the number of *S*-alleles should favour the breakdown of self-
359 incompatibility. However, paradoxically, the number of *S*-alleles may not be very relevant in
360 a system where asexuality serves to provide reproductive assurance.

361

362 *Drift, combined with partial asexuality and self-incompatibility, led to an increase in*
363 *mutation load*

364 Drift influenced the level of mutation load. As population size decreased, the frequency of the
365 deleterious allele decreased and mutation load decreased, due to more effective purging. We
366 also observed a much greater variance in mutation load in very small populations (Table 2
367 and Figure 5, $N=25$). In this case, selection was overwhelmed by drift and this caused a higher
368 probability of fixation of the deleterious allele, and an increase in mutation load. Similar
369 results have been described by Bataillon and Kirkpatrick (2000), Glémin (2003) and Haag and
370 Roze (2007).

371 Drift is also a key component of mutation load in asexual populations (Haag and Roze, 2007).
372 Indeed, large asexual and sexual populations showed comparable frequencies of the
373 deleterious allele and comparable mutation loads (Figure 5; see also Haag and Roze, 2007).
374 However, mutation load was greater in small asexual populations than in small sexual ones.
375 This could be explained by the absence of segregation in asexual populations, in which
376 heterozygous individuals are present in high frequencies (Table 2 and Figure 5; see also Haag
377 and Roze, 2007).

378 In this study, we additionally showed that self-incompatibility played a role in increasing
379 mutation load in partially asexual, self-incompatible populations. Partial asexual reproduction
380 and small population sizes produced strong associations (i.e. linkage disequilibrium) between
381 deleterious alleles (on any locus of the genome) and *S*-alleles. Glémin *et al.* (2001) have
382 already shown that deleterious mutations on loci linked to the *S*-locus may be sheltered by
383 balancing selection acting on rare *S*-alleles. Asexuality and self-incompatibility both favoured
384 the increase of the heterozygote genotype *Aa* within the population, leading to an increase of
385 mutation load in small self-incompatible, partially asexual populations.

386

387 *Inbreeding depression remained low in small, partially asexual, self-incompatible*
388 *populations*

389 Inbreeding depression followed different patterns than the ones observed for mutation load.
390 For very small populations, inbreeding depression is expected to decrease (Bataillon and
391 Kirkpatrick, 2000), due to the absence of polymorphism at the viability locus. This was
392 observed in small self-compatible and self-incompatible populations (Glémin *et al.*, 2001, and
393 our own simulations, Figure 6). As for the impact of asexuality, our results showed no large
394 differences between the self-compatible and the self-incompatible cases for small populations.
395 The effect of strong drift, in the presence of strong asexuality, was to reduce genotype
396 diversity. In most cases, populations were fixed at one allele, but sometimes the population
397 was fixed at the heterozygote genotype *Aa*. This mainly occurred in self-incompatible
398 populations (see Table 2). With no or very low genotype diversity, inbreeding depression had
399 very low values, even if deleterious alleles were present at high frequencies in the population.
400 Chen *et al.* (1997) proposed that asexuality would favour the maintenance of SI. Their
401 reasoning was that asexual reproduction would increase a population's genetic load due to
402 lethal recessive mutations, which was confirmed by our simulation results. However,
403 mutation load and inbreeding depression did not show parallel trends, and the high level of
404 mutation load due to the combination of drift, asexuality and SI was accompanied by low
405 levels of inbreeding depression. Nevertheless, inbreeding depression generally tended to
406 increase with asexuality due to the accumulation of deleterious alleles and this should favour
407 the maintenance of SI.

408

409 *Evolution of partially asexual, self-incompatible populations: a third possible outcome*

410 Until now, the evolutionary interest of partially asexual, self-incompatible populations has
411 been focused on two possible outcomes: (1) the system is stable and SI is maintained or (2)

412 the system is unstable and SI disappears (Vallejo-Marín, 2007; Vallejo-Marín and O'Brien,
413 2007; Chen *et al.*, 1997; Young *et al.*, 2002). However, a third possible outcome should be
414 considered: being the unstable system, sexual reproduction completely disappears. This type
415 of event was observed in our simulations due to a reduction in the number of *S*-alleles to two
416 (see above). However, we also studied how selection on the viability locus could contribute to
417 the potential loss of sexuality. Our results showed that the fitness of individuals produced by
418 asexual reproduction (W_a) was greater than that of individuals produced by random sexual
419 mating (W_o) in small populations (Figure 7). Fitness of individuals (potentially) produced by
420 selfing (W_s) was also lower than W_a in small self-incompatible populations, suggesting that
421 asexuality may have an immediate advantage over selfing under such circumstances.

422 Therefore, we speculate that asexual reproduction could take over sexual reproduction more
423 easily than self-compatibility over self-incompatibility when low numbers of *S*-alleles reduce
424 mate availability. Despite the effect of asexuality on inbreeding depression and *S*-locus
425 diversity, asexual reproduction may offer an alternative means of reproduction under these
426 adverse conditions and self-incompatibility may be temporarily dormant instead of breaking
427 down. Asexual populations could subsequently recover a sexual self-incompatible
428 reproductive system when new *S*-alleles are introduced in the population (for instance, by
429 migration). However, prolonged periods of asexual reproduction may facilitate the
430 accumulation of mutations affecting sexual traits, including self-incompatibility, since
431 selective pressures are ineffective for these traits (Eckert, 2001). Thus, alternative periods of
432 sexual and asexual reproduction may be necessary for such dynamics, which is probably the
433 case for plants combining asexual and sexual reproduction (Bengtsson and Cephitis, 2000).
434 Nevertheless, the scenarios described here need to be confirmed by further and more complex
435 simulations and other factors may favour self-compatible sexual reproduction, such as the
436 more effective purging of deleterious mutations.

437

438 *Perspectives*

439 In this study, we used a model with physically unlinked loci and a single viability locus.
440 However, because the linkage disequilibrium produced by asexuality affected the whole
441 genome, population genomic models may be necessary to fully study the effect of deleterious
442 alleles on the *S*-locus for small, partially asexual populations. In addition, physically linked
443 loci with deleterious alleles (which were not studied here) may reduce the number of *S*-alleles
444 (Uyenoyama, 2003) and the strength of that effect may be influenced by the rate of asexuality
445 as discussed by Vallejo-Marín and Uyenoyama (2008). Moreover, in the present study, it was
446 not possible to investigate the evolutionary dynamics of mating system and asexuality
447 modifiers since these were assumed to be fixed parameters. It would be then interesting to
448 develop a model to test the invasion propensity of self-compatible or asexual mutants. Finally,
449 models incorporating migration or population structure are necessary to study more realistic
450 dynamics of partially asexual, self-incompatible populations. Population structure is expected
451 to affect the genetic diversity of viability and the *S*-locus, for instance by introducing a third
452 allele via migration within subpopulations fixed at two *S*-alleles. These models should also
453 take into account the specificities of the different types of asexual reproduction where, for
454 instance, the asexual propagules may have a lower dispersal capacity (e.g. vegetative
455 reproduction) and may create spatial clusters of individuals with the same genotype.

456 Recent theoretical work (Vallejo-Marín and Uyenoyama, 2008) and our own model lead to
457 several predictions about *S*-locus diversity, mutation load and inbreeding depression within
458 partially asexual self-incompatible populations. However, we need more experimental
459 observations to test and validate these theoretical advances. First, as outlined by Vallejo-
460 Marín and O'Brien (2007), if asexuality relieves selective pressures favouring the breakdown
461 of SI, the co-occurrence of asexuality and self-incompatibility should be frequent. Association

462 of self-incompatibility and asexuality has been studied in only a few monospecific studies
463 (*Rutidosia leiolepis*, Young *et al.*, 2002; *Eucalyptus morrisbyi*, Jones *et al.*, 2005; *Prunus*
464 *avium*, Stoeckel *et al.*, 2006). The association was found at a multispecific level by Chen *et*
465 *al.* (1997) in Australian Droseraceae and by Vallejo-Marín and O'Brien (2007) in Solanaceae.
466 More reports on species combining self-incompatibility and asexuality would enhance our
467 understanding of pressures explaining their maintenance. Second, experimental measurements
468 of asexual reproduction rates and number of *S*-alleles in species showing various levels of
469 asexual reproduction would be useful to test predictions from our simulations. Although
470 experimental measurements of inbreeding depression in self-incompatible species are
471 available (Mena-Alí *et al.*, 2008), studies comparing inbreeding depression from different
472 populations with contrasting sizes and rates of asexual reproduction are lacking.

473

474 *Conclusions*

475 This study investigated for the first time the effect of partial asexual reproduction on the
476 fitness of self-incompatible populations. For loci unlinked to the *S*-locus and large
477 populations, fitness values were similar to those of a self-compatible population. However, in
478 small populations, the combination of drift, asexuality and SI increased mutation load due to
479 the accumulation of deleterious alleles. This increase in the mutation load was accompanied
480 by a selective advantage of asexually produced offspring compared to sexually produced
481 offspring. Therefore, future studies addressing the maintenance of SI in partially asexual
482 populations (by studying the invasibility of self-compatible genotypes) will need to (1)
483 account for whole genome fitness and (2) acknowledge that the process may not be
484 independent of the maintenance of sex itself (i.e. will need to establish whether modifiers of
485 asexual reproduction rates invade more easily than self-compatible alleles).

486

487 **Acknowledgements**

488 Collaboration between MN and SM was promoted through the implementation of the
489 REPROFOR project, financed by the Spanish Ministry for Education and Science. We would
490 like to thank Santiago C. González-Martínez and Ricardo Alía for helping to develop this
491 collaboration. E. Porcher is acknowledged for helpful discussions on self-incompatibility
492 evolution. We are also very thankful to P. Garnier-Géré for a detailed discussion on linkage
493 disequilibrium estimators.

494

495 **References**

- 496 Balloux F, Lehmann L, de Meeûs T (2003). The population genetics of clonal and partially
497 clonal diploids. *Genetics* **164**: 1635-1644.
- 498 Barrett SCH (2002). The evolution of plant sexual diversity. *Nat Rev Genet* **3**: 274-284.
- 499 Bataillon T, Kirkpatrick M (2000). Inbreeding depression due to mildly deleterious mutations
500 in finite populations: size does matter. *Genet Res* **75**: 75-81.
- 501 Bengtsson BO, Ceplitis A (2000). The balance between sexual and asexual reproduction in
502 plants living in variable environments. *J Evolution Biol* **13**: 415-422.
- 503 Birky CW (1996). Heterozygosity, Heteromorphy, and Phylogenetic Trees in Asexual
504 Eukaryotes. *Genetics* **144**, 427-437.
- 505 Brennan AC, Harris SA, Hiscock SJ (2003). The population genetics of sporophytic self-
506 incompatibility in *Senecio squalidus* L. (Asteraceae): avoidance of mating constraints
507 imposed by low S-allele number. *Philos T R Soc B* **358**: 1047-1050.
- 508 Charlesworth D, Charlesworth B (1987). Inbreeding depression and its evolutionary
509 consequences. *Annu Rev EcolEvol S* **18**: 237-268.
- 510 Charlesworth D, Charlesworth B (1979). The evolution and breakdown of S-allele systems.
511 *Heredity* **43**: 41-55.
- 512 Chasnov JR (2000). Mutation-selection balance, dominance and the maintenance of sex.
513 *Genetics* **156**: 1419-1425.
- 514 Chen L, Stace HM, James SH (1997). Self-incompatibility, seed abortion and clonality in the
515 breeding systems of several Western Australian *Drosera* species (Droseraceae) . *Aust J*
516 *Bot* **45**: 191-201.
- 517 Eckert CG (2001). The loss of sex in clonal plants. *Evol Ecol* **15**: 501-520.
- 518 Fisher RA (1941). Average excess and average effect of a gene substitution. *Ann Eugen* **11**:
519 53-63.

520 Garnier-Géré P, Dillmann C (1992). A computer program for testing pairwise linkage
521 disequilibria in subdivided populations. *J Hered* **83**: 239.

522 Glémin S (2003). How are deleterious mutations purged? Drift versus nonrandom mating.
523 *Evolution* **57**: 2678—2687.

524 Glémin S, Bataillon T, Ronfort J, Mignot A, Olivieri I (2001). Inbreeding depression in small
525 populations of self-incompatible plants. *Genetics* **159**: 1217-1229.

526 Haag CR, Roze D (2007). Genetic load in sexual and asexual diploids: segregation,
527 dominance and genetic drift. *Genetics* **176**: 1663-1678.

528 Iqbal B, Lande R, Kohn JR (2008). Loss of self-incompatibility and its evolutionary
529 consequences. *Int J Plant Sci* **169**: 93-104.

530 Jones RC, McKinnon GE, Potts BM, Vaillancourt RE (2005). Genetic diversity and mating
531 system of an endangered tree *Eucalyptus morrisbyi*. *Aust J Bot* **53**: 367-377.

532 Karron JD (1987). A comparison of levels of genetic polymorphism and self-compatibility in
533 geographically restricted and widespread plant congeners. *Evol Ecol* **1**: 47-58.

534 Kimura M, Crow JF (1964). The number of alleles that can be maintained in a finite
535 population. *Genetics* **49**: 725-738.

536 Kirkpatrick M, Jenkins CD (1989). Genetic segregation and the maintenance of sexual
537 reproduction. *Nature*, **339**: 300-301.

538 Kunin WE, Shmida A (1997). Plant reproductive traits as a function of local, regional, and
539 global abundance. *Conserv Biol* **11**: 183-192.

540 de Meeûs T, Balloux F (2004). Clonal reproduction and linkage disequilibrium in diploids: a
541 simulation study. *Infect Genet Evol* **4**: 345—351.

542 de Meeûs T, Prugnolle F, Agnew P (2007). Asexual reproduction: genetics and evolutionary
543 aspects. *Cell Mol Life Sci* **64**: 1355-1372.

544 Mena-Alí J, Keser L, Stephenson A (2008). Inbreeding depression in *Solanum carolinense*
545 (Solanaceae), a species with a plastic self-incompatibility response. *BMC Evol Biol* **8**:
546 10.

547 Mena-Alí J, Keser L, Stephenson A (2009). The effect of sheltered load on reproduction in
548 *Solanum carolinense*, a species with variable self-incompatibility. *Sex Plant Reprod*
549 **22**: 63-71.

550 Muirhead CA, Lande R (1997). Inbreeding depression under joint selfing, outcrossing, and
551 asexuality. *Evolution* **51**: 1409-1415.

552 de Nettancourt D (1977). *Incompatibility in Angiosperms*, New York: Springer, Berlin
553 Heidelberg .

554 Pamilo P (1987). Heterozygosity in apomictic organisms. *Hereditas* **107**: 95-101.

555 Reinartz JA, Les DH (1994). Bottleneck-induced dissolution of self-incompatibility and
556 breeding system consequences in *Aster furcatus* (Asteraceae). *Am J Bot* **81**: 446-455.

557 Stoeckel S, Grange J, Fernandez-Manjarres JF, Bilger I, Frascaria-Lacoste N, Mariette S
558 (2006). Heterozygote excess in a self-incompatible and partially clonal forest tree
559 species - *Prunus avium* L. *Mol Ecol* **15**: 2109-2118.

560 Stone JL (2004). Sheltered load associated with S-alleles in *Solanum carolinense*. *Heredity*
561 **92**: 335-342.

562 Takahata N (1990). A simple genealogical structure of strongly balanced allelic lines and
563 trans-species evolution of polymorphism. *PNAS* **87**: 2419-2423.

564 Uyenoyama MK (2003). Genealogy-dependent variation in viability among self-
565 incompatibility genotypes. *Theor Popul Biol* **63**: 281-293.

566 Vallejo-Marín M (2007). The paradox of clonality and the evolution of self-incompatibility.
567 *Plant Signal Behav* **2**: 265-266.

- 568 Vallejo-Marín M, O'Brien HE (2007). Correlated evolution of self-incompatibility and clonal
569 reproduction in *Solanum* (Solanaceae). *New Phytol* **173**: 415-421.
- 570 Vallejo-Marín M, Uyenoyama MK (2008). On the evolutionary modification of self-
571 incompatibility: implications of partial clonality for allelic diversity and genealogical
572 structure. In *Self-Incompatibility in Flowering Plants. Evolution, Diversity, and*
573 *Mechanisms* . pp. 53-71.
- 574 Yokoyama S, Hetherington LE (1982). The expected number of self-incompatibility alleles in
575 finite plant populations. *Heredity* **48**: 299-303.
- 576 Young AG, Hill JH, Murray BG, Peakall R (2002). Breeding system, genetic diversity and
577 clonal structure in the sub-alpine forb *Rutidosia leiolepis* F. Muell. (Asteraceae). *Biol*
578 *Conserv* **106**: 71-78.
- 579

580 **Figure legends**

581 **Figure 1** Linkage disequilibrium (R_{GDD} index, Garnier-Géré & Dillmann 1992) at increasing
582 rates of asexual reproduction. Box-plots represent median (black line), first and third quartiles
583 (box) and 5% and 95% percentiles (whiskers) of 1 000 observations from simulations
584 performed at seven rates of asexual reproduction ($c=0, 0.5, 0.8, 0.9, 0.99, 0.999$ and 1) and
585 four population sizes ($N=25, 50, 100$ and 1000). (a) R_{GDD} index between S -locus and viability
586 locus ($s=0.1, h=0.2$); (b) R_{GDD} index between two neutral loci.

587 **Figure 2** Effective number of S -alleles at increasing rates of asexual reproduction. Box-plots
588 represent median (black line), first and third quartiles (grey box) and 5% and 95% percentiles
589 (whiskers) of 1 000 observations from the simulations performed at six rates of asexuality
590 ($c=0, 0.5, 0.8, 0.9, 0.99$ and 1), population size $N=25$ (a) and $N=100$ (b), mutation rate $\mu_S=10^{-5}$,
591 $s=0$ and $h=0$ for the viability locus. Grey line represents the theoretical expected number of
592 common S -alleles obtained numerically from Equation 1 (Equation 3.7 in Vallejo-Marín &
593 Uyenoyama, 2008).

594 **Figure 3** Inbreeding coefficient F_{IS} at a neutral locus at increasing rates of asexual
595 reproduction. Box-plots represent median (black line), first and third quartiles (white and grey
596 boxes, for self-compatible and self-incompatible populations respectively) and 5% and 95%
597 percentiles (whiskers) of 1 000 observations from the simulations performed at six rates of
598 asexuality ($c=0, 0.5, 0.8, 0.9, 0.99$ and 1), mutation rates $\mu_S=10^{-5}$ and $\mu_N=10^{-3}$, and population
599 size (a) $N=25$ and (b) $N=100$. Theoretical equilibrium values for F_{IS} at a neutral locus in a
600 self-compatible, random-mating population (dotted black line, from Equation 3, neglecting
601 mutation) and in a self-incompatible population (continuous grey line, from Equation 2) are
602 also represented.

603 **Figure 4** Frequency of deleterious alleles at increasing rates of asexual reproduction under
604 two selection regimes. Box-plots represent median (black line), first and third quartiles (white

605 and grey boxes, for self-compatible and self-incompatible populations respectively) and 5%
606 and 95% percentiles (whiskers) of 1000 observations from the simulations performed at seven
607 rates of asexuality ($c=0, 0.5, 0.8, 0.9, 0.99, 0.999$ and 1), mutation rates $\mu_S=10^{-5}$ and $\mu_N=10^{-3}$.
608 Population size and viability locus coefficients: (a) $N=25, s=0.1, h=0.2$, (b) $N=25, s=1,$
609 $h=0.02$, (c) $N=100, s=0.1, h=0.2$, and (d) $N=100, s=1, h=0.02$.

610 **Figure 5** Mutation load at increasing rates of asexual reproduction under two different
611 selection regimes according to reproductive system. Box-plots represent median (black line),
612 first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1 000 observations
613 from simulations performed at seven rates of asexual reproduction ($c=0, 0.5, 0.8, 0.9, 0.99,$
614 0.999 and 1) and four population sizes ($N=25, 50, 100$ and 1000). (a) Self-incompatible
615 population with a viability locus with mildly deleterious allele ($s=0.1, h=0.2$), (b) self-
616 compatible population with a viability locus with mildly deleterious allele ($s=0.1, h=0.2$), (c)
617 self-incompatible population with a viability locus with highly recessive lethal allele ($s=1,$
618 $h=0.02$), and (d) self-compatible population with a viability locus with highly recessive lethal
619 allele ($s=1, h=0.02$).

620 **Figure 6** Inbreeding depression at increasing rates of asexual reproduction and under two
621 different selection regimes according to reproductive system. Box-plots represent median
622 (black line), first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1000
623 observations from simulations performed at seven rates of asexual reproduction ($c=0, 0.5, 0.8,$
624 $0.9, 0.99, 0.999$ and 1) and four population sizes ($N=25, 50, 100$ and 1000). (a) Self-
625 incompatible population with a viability locus with mildly deleterious allele ($s=0.1, h=0.2$),
626 (b) self-compatible population with a viability locus with mildly deleterious allele ($s=0.1,$
627 $h=0.2$), (c) self-incompatible population with a viability locus with highly recessive lethal
628 allele ($s=1, h=0.02$), and (d) self-compatible population with a viability locus with highly
629 recessive lethal allele ($s=1, h=0.02$).

630 **Figure 7** Fitness ratio between sexually and asexually produced individuals at increasing rates
631 of asexual reproduction under two different selection regimes according to reproductive
632 system. Box-plots represent median (black line), first and third quartiles (box) and 5% and
633 95% percentiles (whiskers) of 1 000 observations from simulations performed at seven rates
634 of asexual reproduction ($c=0, 0.5, 0.8, 0.9, 0.99, 0.999$ and 1) and four population sizes
635 ($N=25, 50, 100$ and 1000). (a) Self-incompatible population with a viability locus with mildly
636 deleterious allele ($s=0.1, h=0.2$), (b) self-compatible population with a viability locus with
637 mildly deleterious allele ($s=0.1, h=0.2$), (c) self-incompatible population with a viability locus
638 with highly recessive lethal allele ($s=1, h=0.02$), and (d) self-compatible population with a
639 viability locus with highly recessive lethal allele ($s=1, h=0.02$).
640

641 **Table 1** Genotype fixation probability at the *S*-locus under two different selection regimes at
 642 the viability locus

Population size	Selection regime	Rate of asexual reproduction						
		<i>c</i> =0	<i>c</i> =0.5	<i>c</i> =0.8	<i>c</i> =0.9	<i>c</i> =0.99	<i>c</i> =0.999	<i>c</i> =1
<i>N</i> =25	Mildly deleterious ¹	0.000	0.000	0.036	0.325	0.784	0.807	0.997
<i>N</i> =25	Lethal recessive ²	0.000	0.000	0.018	0.315	0.790	0.805	0.994
<i>N</i> =50	Mildly deleterious	0.000	0.000	0.000	0.002	0.072	0.117	0.990
<i>N</i> =50	Lethal recessive	0.000	0.000	0.000	0.001	0.075	0.090	0.989
<i>N</i> =100	Mildly deleterious	0.000	0.000	0.000	0.000	0.001	0.001	0.983

643 The probability of genotype fixation at the *S*-locus (only two alleles in the population) was
 644 estimated as the proportion of observations in which this event was recorded. Results in this
 645 table are for populations with a mutation rate of $\mu_S=10^{-5}$ at the *S*-locus.

646 ¹ Viability locus selection and dominance coefficients: $s=0.1$ and $h=0.2$

647 ² Viability locus selection and dominance coefficients: $s=1$ and $h=0.02$

648

649 **Table 2** Fixation probabilities at the viability locus in a self-incompatible population under
 650 two different selection regimes.

Population size	Selection regime	Rate of asexual reproduction						
		$c=0$	$c=0.5$	$c=0.8$	$c=0.9$	$c=0.99$	$c=0.999$	$c=1$
(a) Fixation of deleterious allele a								
$N=25$	Mildly deleterious ¹	0.066	0.049	0.056	0.050	0.085	0.081	0.067
$N=50$	Mildly deleterious	0.000	0.000	0.001	0.000	0.001	0.001	0.001
$N=100$	Mildly deleterious	0.000	0.000	0.000	0.000	0.000	0.000	0.000
(b) Fixation of allele A								
$N=25$	Mildly deleterious	0.670	0.714	0.663	0.519	0.264	0.279	0.082
$N=25$	Lethal recessive ²	0.831	0.783	0.767	0.581	0.293	0.317	0.085
$N=50$	Mildly deleterious	0.583	0.541	0.574	0.507	0.474	0.466	0.139
$N=50$	Lethal recessive	0.661	0.635	0.596	0.564	0.518	0.493	0.150
$N=100$	Mildly deleterious	0.314	0.306	0.320	0.294	0.274	0.290	0.187
$N=100$	Lethal recessive	0.437	0.361	0.318	0.313	0.269	0.296	0.189
(c) Fixation of genotype Aa								
$N=25$	Mildly deleterious	0.000	0.000	0.023	0.184	0.472	0.467	0.723
$N=25$	Lethal recessive	0.000	0.000	0.013	0.212	0.551	0.569	0.879
$N=50$	Mildly deleterious	0.000	0.000	0.000	0.000	0.026	0.050	0.585
$N=50$	Lethal recessive	0.000	0.000	0.000	0.000	0.032	0.041	0.721
$N=100$	Mildly deleterious	0.000	0.000	0.000	0.000	0.000	0.000	0.230
$N=100$	Lethal recessive	0.000	0.000	0.000	0.000	0.000	0.000	0.317


651 The probability of fixation at the viability locus (fixation of genotypes, AA , aa or Aa) was
 652 estimated as the proportion of observations in which fixation was recorded.

653 ¹ Viability locus selection and dominance coefficients: $s=0.1$ and $h=0.2$

654 ² Viability locus selection and dominance coefficients: $s=1$ and $h=0.02$

655


656 **Figure 1**



657

658


659 **Figure 2**



660

661


662 **Figure 3**



663

664


665 **Figure 4**



666

667


668 **Figure 5**



669

670


671 **Figure 6**



672

673

674 **Figure 7**



675

676