



HAL
open science

TGF- β and fibrosis in different organs - molecular pathway imprints

Dirk Pohlers, Julia Brenmoehl, Ivonne Löffler, Cornelia K. Müller, Carola Leipner, Stefan Schultze-Mosgau, Andreas Stallmach, Raimund W. Kinne, Gunter Wolf

► **To cite this version:**

Dirk Pohlers, Julia Brenmoehl, Ivonne Löffler, Cornelia K. Müller, Carola Leipner, et al.. TGF- β and fibrosis in different organs - molecular pathway imprints. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, 2009, 1792 (8), pp.746. 10.1016/j.bbadis.2009.06.004 . hal-00506515

HAL Id: hal-00506515

<https://hal.science/hal-00506515>

Submitted on 28 Jul 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript


TGF- β and fibrosis in different organs - molecular pathway imprints

Dirk Pohlers, Julia Brenmoehl, Ivonne Löffler, Cornelia K. Müller, Carola Leipner, Stefan Schultze-Mosgau, Andreas Stallmach, Raimund W. Kinne, Gunter Wolf

PII: S0925-4439(09)00135-5
DOI: doi:[10.1016/j.bbadis.2009.06.004](https://doi.org/10.1016/j.bbadis.2009.06.004)
Reference: BBADIS 62965

To appear in: *BBA - Molecular Basis of Disease*

Received date: 2 March 2009
Revised date: 11 June 2009
Accepted date: 12 June 2009



Please cite this article as: Dirk Pohlers, Julia Brenmoehl, Ivonne Löffler, Cornelia K. Müller, Carola Leipner, Stefan Schultze-Mosgau, Andreas Stallmach, Raimund W. Kinne, Gunter Wolf, TGF- β and fibrosis in different organs - molecular pathway imprints, *BBA - Molecular Basis of Disease* (2009), doi:[10.1016/j.bbadis.2009.06.004](https://doi.org/10.1016/j.bbadis.2009.06.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **TGF- β and fibrosis in different organs – molecular pathway imprints**

2

3 Dirk Pohlers¹, Julia Brenmoehl², Ivonne Löffler³, Cornelia K. Müller⁴, Carola Leipner⁵,

4 Stefan Schultze-Mosgau⁴, Andreas Stallmach², Raimund W. Kinne¹, Gunter Wolf³

5

6 1 Experimental Rheumatology Group, Department of Orthopedics, Waldkrankenhaus “Rudolf

7 Elle” Eisenberg, University Hospital, Friedrich Schiller University, Jena

8 2 Department of Gastroenterology, Hepatology and Infectious Diseases, Clinic of Internal

9 Medicine II, University Hospital, Friedrich Schiller University, Jena

10 3 Clinic of Internal Medicine III, University Hospital, Friedrich Schiller University, Jena

11 4 Clinic of Oromaxillofacial Surgery/Plastic Surgery, University Hospital, Friedrich Schiller

12 University, Jena

13 5 Institute of Animal Research, University Hospital, Friedrich Schiller University, Jena

14

15 Corresponding author:

16 Raimund W. Kinne

17 Experimental Rheumatology Unit, University Hospital Jena

18 Department of Orthopedics, Waldkrankenhaus “Rudolf Elle”

19 Klosterlausnitzer Str. 81

20 07607 Eisenberg

21 e-mail: Raimund.W.Kinne@med.uni-jena.de

22 Tel./Fax +49 36691 81228/6

23

24 Key words: TGF- β , fibrosis, arthritis, nephritis, inflammatory bowel disease, Crohn’s disease,

25 wound-healing, myocarditis

26 **ABSTRACT**

27 The action of transforming-growth-factor (TGF)- β following inflammatory responses is
28 characterized by increased production of extracellular matrix (ECM) components, as well as
29 mesenchymal cell proliferation, migration, and accumulation. Thus, TGF- β is important for
30 the induction of fibrosis often associated with chronic phases of inflammatory diseases. This
31 common feature of TGF-related pathologies is observed in many different organs. Therefore,
32 in addition to the description of the common TGF- β -pathway, this review focuses on TGF- β -
33 related pathogenetic effects in different pathologies/organs, i. e., arthritis, diabetic
34 nephropathy, colitis/Crohn's disease, radiation-induced fibrosis, and myocarditis (including
35 their similarities and dissimilarities). However, TGF- β exhibits both exacerbating and
36 ameliorating features, depending on the phase of disease and the site of action. Due to its
37 central role in severe fibrotic diseases, TGF- β nevertheless remains an attractive therapeutic
38 target, if targeted locally and during the fibrotic phase of disease.

39

40 1. INTRODUCTION

41 The fibrotic reaction of the connective tissue following an inflammatory response is mainly
42 characterized by an increased production of extracellular matrix (ECM) components and
43 mesenchymal cell proliferation, migration and accumulation. Despite the existence of
44 numerous distinct causes of chronic inflammatory diseases in different organs and tissues,
45 these diseases are generally characterized by: i) severe and intermittent progression with
46 phases of acute exacerbation and remission; ii) immigration of inflammatory cells
47 (macrophages, granulocytes and T-cells); and iii) increased expression of pro-inflammatory
48 mediators (Figure 1). These processes result in the proliferation of local fibroblasts and, by
49 interaction with epithelial cells, in their differentiation into myofibroblasts and can be
50 regarded as a misguided wound healing. Finally, the inflammatory process comes to rest, but
51 the massive fibrosis prevents a rebuilding of functionally intact tissue and organs.

52 Transforming growth factor (TGF)- β is an ubiquitously expressed cytokine belonging
53 to a large superfamily of activins/bone morphogenetic proteins [1]. This mediator plays an
54 active role in the processes discussed above, such as proliferation, wound healing [2], and
55 synthesis of ECM molecules [3]. TGF- β , therefore, strongly contributes to fibrotic disorders
56 such as diabetic nephropathy, Crohn's disease, rheumatoid arthritis, radiation-induced
57 fibrosis, and myocarditis. However, TGF- β is clearly a bi- (or multi-) functional molecule
58 with strong effects on the immune system [4,5].

59

60 2. TGF- β SIGNALING PATHWAY

61 TGF- β is synthesized as one part of a large molecule, the pro-TGF- β containing the latency-
62 associated proteins (LAP; Figure 2). The latter is cleaved from TGF- β in the Golgi apparatus,
63 but remains non-covalently associated with the growth factor. The disulfide-bound latent-

64 TGF- β -binding proteins 1/2 (LTBP 1/2) connect the whole complex to the ECM (details
65 described in [6]). Release of TGF- β from the pro-TGF- β complex can be achieved through
66 proteolytic activity by plasmin [7] or matrix-metalloproteinase (MMP)-2 and MMP-9 [8],
67 through integrins [9,10], recently reviewed in [11], treatment with mild acids [12], or through
68 the action of thrombospondin (THBS; [13]) by disrupting the non-covalent interactions
69 between LAP and TGF- β 1. Once released, TGF- β mediates signals through pairs of type I
70 and type II receptors [14]. The type III receptor (betaglycan) acts - probably in conjunction
71 with other heparan sulfate glycans like syndecan [15]- as a co-receptor for binding/presenting
72 TGF and a regulator of TGF- β signaling [16]. The result of ligand binding is the activation of
73 the type II receptor (TGFBR2), which then phosphorylates the type I receptor (TGFBR1). The
74 active receptor complex then phosphorylates the so-called R (receptor)-Smad-2 or Smad-3
75 that propagates the signal [17]. The phosphorylation of Smad2/3 decreases the affinity for the
76 Smad-anchor for receptor activation (SARA), which in non-stimulated cells mediates the
77 retention of Smad2/3 in the cytoplasm by interaction, and increases the affinity of Smad2/3
78 for Smad-4 (a so-called co-Smad). This complex is now able to enter the nucleus, to bind
79 transcriptional co-activators like p300 and Creb-binding-protein (CBP) or repressors like SkiL
80 or TGIF [18], and to regulate the transcriptional activity of various genes.

81 Mitogen-activated protein kinases (MAPKs) and protein kinase C can also interfere
82 with either the nuclear translocation or binding of Smad3/4 complexes to DNA and regulate
83 TGF- β 1 signalling [19,20]. Moreover, the serine-threonine protein kinase B can directly
84 interact with Smad3, thereby preventing its phosphorylation and nuclear translocation [21].
85 Other pathways can be directly activated by TGF- β 1. These include components of the
86 MAPK pathway, such as ras, raf, ERK, p38 and JNK, the phosphatidylinositol-3 kinase
87 cascade, as well as the regulators of cadherin junctions, RhoA and Rac ([19,22,23]; Fig. 1).
88 Recently, an involvement of the focal adhesion kinase (FAK) was established in

89 myofibroblast differentiation and in remodeling of the connective tissue following stimulation
90 with TGF [24,25]. In line with these results, TGF-induced FAK-signaling is required for the
91 activation of TAK [26] or MEKK1 and, subsequently, of JNK [27], all factors shown to be
92 essential for the transcription of pro-fibrotic genes.

93 Notably, that signal transduction pathways have their own intracellular regulators. An
94 inhibitory Smad (Smad7) blocks TGF- β 1 signaling by physical interaction with the activated
95 TGFBR1 receptor and prevents the docking and phosphorylation of Smad2/3 [28,29].

96 This complex TGF- β signaling pathway (Figure 2) contains numerous ligands,
97 receptors, and signaling molecules which, as potential targets of dysregulation via increased
98 or decreased expression, activation or interaction, may be partially involved in fibrotic
99 reactions in the diseases discussed below.

100

101 **3. TGF- β -RELATED MOLECULES IN RHEUMATOID ARTHRITIS**

102 The importance of TGF- β 1 in rheumatoid arthritis (RA) ranges from an association with
103 certain vascularization patterns in the synovial membrane (SM) [30], and an association of
104 TGF- β polymorphisms with the radiological signs of joint destruction [31] to an induction of
105 pro-inflammatory cytokines, MMP [32], aggrecanase [33] and urokinase-type plasminogen
106 activator [34]. In addition, TGF- β plays an important role for the function of regulatory T-
107 cells [5] in the suppression of autoimmunity. Indeed, increased levels of TGF- β 1 have been
108 found in the synovial membrane of patients with RA by northern blot [35],
109 immunohistochemistry [36,37], western blot [38] and in synovial effusions [39]. Also
110 TGFBR2 was detected at higher levels than in normal synovial tissue [35].

111 Using “pathway-directed” software following genome-wide comparison between
112 synovial fibroblasts (SFB) from patients with RA and osteoarthritis (OA) with Affymetrix

113 arrays, gene expression of TGF- β 1, TGF- β 3, LTBP1/2, THBS1, TGFBR1, SARA, CBP, SkiL
114 – belonging to the TGF- β -pathway (Figure 2) - was elevated in RA [40]. After validating
115 array data at the mRNA and protein levels using quantitative PCR and western
116 blot/immunohistochemistry, we confirmed an upregulated TGF- β pathway in RA-SFB. The
117 presence of TGF- β 1, in conjunction with increased amounts of TGF- β -releasing THBS1 and a
118 higher expression of TGFBR1, may thus lead to an amplified response of RA-SFB to TGF- β .
119 This RA-specific response has been confirmed by increased expression of MMP-11 following
120 TGF- β stimulation, implying a pathogenetic relevance of the TGF- β -pathway for MMP-
121 induced degradation or remodeling processes in RA [40].

122 The importance of TGF- β for the pathogenesis of arthritis is emphasized by a number
123 of animal models. The abundant expression of TGF- β 1, 2, and 3 as well as the TGFBR1 and -
124 2 in rat synovium was increased after onset of the collagen-induced arthritis (CIA) [41], and
125 direct intra-articular injection of TGF- β 1 or TGF- β 2 induced synovial erythema, swelling,
126 and cellular infiltration resulting in synovial inflammation and hyperplasia [42]. Conversely,
127 neutralization of TGF- β inhibited acute and chronic arthritis induced by streptococcal cell
128 wall (SCW) [43]. In line with these observations, an adenovirus-mediated overexpression of
129 TGF- β 1 in rabbit knees led to an increased glycosaminoglycan release, nitric oxide
130 production and, most notably, to fibrosis and muscle edema [44]. The prevention of CIA by
131 administration of a TGFBR1 inhibitor (HTS466284), which concomitantly reduced the
132 expression of vascular endothelial growth factor (VEGF), platelet-derived growth factor
133 (PDGF)-AA, TNF- α , and cellular proliferation [45], further underlines the pathogenetic role
134 of TGF- β in arthritis.

135 In contrast to the above-mentioned studies, numerous observations show beneficial
136 effects of TGF- β in arthritis. If administered systemically, TGF- β 1 suppressed SCW-induced

137 arthritis, as measured by cellular infiltration and joint erosion [46]. In addition, investigation
138 of the cytokine expression during CIA demonstrated a strong upregulation of TGF- β 1/2 in the
139 remission state of disease, possibly reflecting the anti-inflammatory regulation of T-cells by
140 TGF- β in arthritis. If TGF- β signaling was inhibited by expression of dominant-negative
141 TGFBR2 in T-cells [47] the susceptibility and the clinical severity of CIA was strongly
142 increased. Likewise, if TGF- β 1 was retrovirally overexpressed in arthritogenic splenocytes,
143 CIA could not be transferred to SCID mice and established disease was ameliorated [48].
144 Therefore, it may be important to inhibit TGF- β only at the site of inflammation without
145 targeting regulatory lymphocytes at extra-articular sites.

146

147 **4. TGF- β -RELATED MOLECULES IN DIABETIC NEPHROPATHY**

148 TGF- β and its signal transduction play a major role in diabetic nephropathy and have
149 therefore been thoroughly studied [49-53]. Among the features of the diabetic milieu,
150 hyperglycemia, increased non-enzymatic glycation of proteins, *de novo* synthesis of
151 diacylglycerol and subsequent activation of protein kinase C, increased intracellular
152 glucosamine production, and enhanced renal production of vasoactive agents (angiotensin II,
153 endothelins, thromboxane) have all been shown to increase the expression of TGF- β in
154 cultured renal cells and animal models of diabetic nephropathy [50,52,54].

155 The TGF- β level is elevated in the kidneys of insulin-dependent diabetic animals
156 during both early and late stages of disease [53,54]. Treatment of the streptozotocin (STZ)-
157 diabetic rat with sufficient insulin to reduce hyperglycemia suppressed the enhanced
158 expression of TGF- β and matrix components in the glomeruli. In the STZ-diabetic rat and
159 mouse, increased TGF- β 1 expression in the renal cortex and glomeruli as well as up-
160 regulation of the TGFBR2 mRNA and protein was noted early after the onset of diabetes

161 [53].. The *db/db* mouse, a model of type 2 diabetes, characterized by hyperglycemia, obesity,
162 and insulin resistance, develops increased amounts of TGF- β 1 localized in the glomerular
163 compartments [55]. In contrast, the mRNA and protein levels of the TGFBR2 are
164 significantly up-regulated in both the glomerular and the tubulointerstitial compartments [55].

165 The development of diabetic renal hypertrophy and glomerulosclerosis is likely caused
166 by heightened activity of the TGF- β system [56-63]. Short-term treatment of the STZ-diabetic
167 mouse with a neutralizing monoclonal antibody against all three isoforms of TGF- β prevented
168 glomerular hypertrophy, reduced the increment in kidney weight by 50%, and significantly
169 attenuated the increase in TGF- β 1, α 1(IV) collagen, and fibronectin mRNAs without
170 affecting glycemic control [56]. The results of this study suggested a cause-and-effect
171 relationship between the renal TGF- β system and the development of early structural changes
172 in diabetic nephropathy. Systemic anti-TGF- β therapy for eight weeks prevented the
173 mesangial matrix expansion of diabetic glomerulosclerosis and, most importantly, preserved
174 kidney function, showing for the first time that neutralization of TGF- β activity prevents the
175 progression of renal failure in diabetes [57]. However, the anti-TGF- β antibody did not reduce
176 albuminuria, which itself may promote the progression of renal insufficiency [57]. The
177 paradox of preserved renal function in view of persistent albuminuria may be explained by
178 postulating that the deleterious effects of proteinuria are themselves mediated by the TGF- β
179 system [63-65].

180 In mouse mesangial and tubular cells, high glucose stimulates the transcription of
181 fibronectin and, in addition, potentiates the transcriptional activation of fibronectin by TGF-
182 β 1 [58,59]. This particular effect of TGF- β 1 appears to be mediated by Smad3, because over-
183 expression of Smad3 alone was able to induce fibronectin promoter activity [66]. In
184 conjunction with exogenous TGF- β 1, Smad3 over-expression synergistically increased

185 fibronectin expression, as if the extra Smad3 had increased the efficiency of TGF- β signaling
186 [66]. Finally, transfection of a Smad3-dominant-negative construct inhibited TGF- β 1
187 stimulated fibronectin promoter activity [67,68]. However, part of the TGF- β 1-induced
188 fibronectin expression may be mediated in parallel by the p38MAPK pathway [68,69].
189 Finally, there is evidence that Smad3 is a central mediator in the TGF- β 1-induced increase of
190 mRNA expression for α 1(I) collagen [68,70]. TGF- β -induced MAPK activation also leads to
191 N-terminal phosphorylation of p53 that enables its interaction with TGF- β -activated Smads
192 [64], an indication for cross-talk between the various TGF- β signaling pathways.

193 TGF- β 1 may have additional effects besides the stimulation of extracellular matrix
194 production. In podocytes, TGF- β 1 induces apoptosis through Smad7 by inhibiting nuclear
195 translocation of the cell survival factor NF- κ B [70]. TGF- β 1-mediated activation of Smad7 is
196 specific for podocytes and is not found in mesangial cells in a limited series of biopsies from
197 patients with diabetic nephropathy [64]. Since podocytes are terminally differentiated cells
198 unable to undergo cell division due to the up-regulation of the cell cycle inhibitory proteins
199 p57 and p27^{Kip1}, apoptotic loss of cells was not replaced.

200 Studies performed in diabetic patients with various degrees of nephropathy also
201 underline the importance of the renal TGF- β system in disease development [71-73]. All three
202 isoforms of TGF- β are elevated in both the glomerular and the tubulointerstitial compartments
203 of patients with established diabetic nephropathy [71,72]. Furthermore, glomerular TGF- β 1
204 mRNA is markedly increased in biopsy specimens from patients with proven diabetic kidney
205 disease. These investigations suggests that increased renal TGF- β levels closely correlate with
206 the degree of mesangial matrix expansion, interstitial fibrosis, and renal insufficiency.

207 Another study, designed to assess renal production of TGF- β [73] measured aortic,
208 renal vein, and urinary levels of TGF- β in 14 type 2 diabetic and 11 non-diabetic control
209 patients undergoing elective coronary artery catheterization [73]. Both groups were roughly
210 matched with regard to the range of renal function and the presence of hypertension and
211 proteinuria. Renal blood flow was measured to calculate the net mass balance across the
212 kidney. The gradient of TGF- β 1 concentration across the renal vascular bed was negative in
213 the non-diabetic patients indicating net renal extraction of TGF- β 1, whereas the gradient was
214 positive in the diabetic patients indicating net renal production of TGF- β 1. When the renal
215 TGF- β 1 mass balance was calculated, a similar pattern was observed with the non-diabetic
216 kidney removing approximately 3500 ng/min of TGF- β 1 from the circulation, and the diabetic
217 kidney adding approximately 1000 ng/min of TGF- β 1 [73]. In addition, the level of
218 bioassayable TGF- β was increased four-fold in the urine of diabetic *versus* non-diabetic
219 patients. This was not simply a function of enhanced glomerular permeability to protein since
220 diabetic patients both with and without microalbuminuria displayed similarly high rates of
221 urinary TGF- β excretion [73]. These results demonstrated that the kidneys of diabetic patients
222 overproduce TGF- β 1 protein; further details, e. g., the exact contribution of the different renal
223 cell types, need to be investigated.

224 An interesting *post-hoc* study assessed whether treatment with the angiotensin
225 converting enzyme inhibitor captopril would lower serum TGF- β 1 levels in a small subset of
226 patients with diabetic nephropathy who had been enrolled in the Collaborative Study Group
227 [74,75]. After six months, the serum TGF- β 1 level decreased significantly by 21% in the
228 captopril-treated group, whereas it increased slightly by 11% in the placebo-treated group
229 [75]. Interestingly, the captopril-treated patients with decreased serum TGF- β 1 levels tended
230 to have better preserved renal function over the ensuing two-year period [75]. This association

231 was even more pronounced in the subset of patients with an initial glomerular filtration rate of
232 less than 75 ml/min. These results suggest that TGF- β 1 plays a pivotal role in the progression
233 of diabetic nephropathy and that angiotensin converting enzyme inhibitor therapy may protect
234 the kidney by lowering TGF- β 1 production.

235

236 **5. TGF- β -RELATED MOLECULES IN INTESTINAL INFLAMMATION – A DOUBLE-** 237 **SIDED SWORD**

238 TGF- β is constitutively expressed by epithelial cells, fibroblasts, and mononuclear cells in the
239 gastrointestinal tract [76]. Its critical role in intestinal homeostasis as a negative master
240 regulator of inflammation is well-established [77,78] and indisputable. However, translation
241 of elegant mouse experiments into therapeutic interventions in humans requires a clearer
242 understanding of TGF- β activity in the human gut. Mice with global TGF- β defects, such as
243 TGF- β -null mice or transgenic mice expressing a dominant negative TGFBR2 chain, are
244 unresponsive to TGF- β 1 signaling. The former die soon after birth due to systemic
245 inflammation, and the latter develop severe colonic and pulmonary inflammation [79,80].
246 These manifestations of global TGF- β 1 defects are mirrored in mouse models of colitis in that
247 the secretion of TGF- β 1 is consistently associated with either the protection from colitis or a
248 greatly diminished severity of colitis. This is seen both in the T_H1 model of colitis induced by
249 the haptening reagent trinitrobenzene sulfonic acid (TNBS), which mimics Crohn's disease
250 (CD), or the T_H2 model of colitis induced by the haptening agent oxazolone, which mimics
251 ulcerative colitis (UC) [81,82]. In addition, it is seen in the colitis of SCID or RAG2-deficient
252 mice receiving CD45RB^{high} (naive) T cells in which the protective effect of the cotransfer of
253 CD45RB^{low} T (memory) cells is abolished by concomitant administration of a neutralizing
254 TGF- β 1 antibody [83]. These and other studies quite conclusively establish that TGF- β 1 plays

255 an essential, regulatory role in the control of colitis. On the other hand, in mice TGF- β 1 in
256 combination with IL-6 induces a strong pro-inflammatory T_H17 cell differentiation [84-86].
257 Notably, in humans TGF- β 1 does not have a direct effect on the development of human T_H17
258 cells, but it can indirectly favour the development of these cells by suppressing the expression
259 of T-bet and selectively inhibiting the expansion of IFN- γ -producing T cells [87-89].

260 The important role of Smad3 as an essential mediator of TGF- β 1-induced anti-
261 inflammatory and suppressive activities at the mucosal level emerges from studies in mice
262 with targeted deletion of the Smad3 gene. The animals are viable, but die from defects in
263 mucosal immunity at 1–6 months of age. Mutant mice show diminished cell responsiveness to
264 TGF- β 1, massive infiltration of T cells, and multiple pyogenic abscesses in the stomach and
265 intestine [87,90]. When Smad signaling was studied in normal human gut mucosa, whole
266 biopsies or isolated lamina propria mononuclear cells, a basal level of phosphorylated
267 (phospho-) Smad3 was observed which was rapidly upregulated by the addition of exogenous
268 TGF- β 1 [91].

269 TGF- β has been also implicated as a key inducer of epithelial-mesenchymal transition
270 (EMT) [92-97]. Amongst others, EMT is an essential component of tissue remodeling and
271 wound repair (reviewed in [98,99]) and fibrosis (reviewed in [100]). During this transition,
272 the epithelial phenotype, characterized by strong cell-cell junctions and polarity, is replaced
273 by a mesenchymal phenotype, with reduced cell-cell interactions, a fibroblastic morphology
274 and increased motility. TGF- β stimulates the proliferation of many cell types, particularly
275 those of mesenchymal origin, and it is also a potent inhibitor of epithelial cell proliferation.
276 EMT in response to TGF- β 1 and in fibrosis is mediated predominantly via Smad-dependent
277 (mainly Smad3) pathways [101,102]. A loss of Smad3 in mice blocked both morphological

278 changes of lens epithelium to a mesenchymal phenotype and expression of EMT markers in
279 response to injury *in vivo* or to exposure to exogenous TGF- β in organ culture [101].

280 CD is a chronic, progressive disease of the gastrointestinal tract with an unknown
281 etiology. It is characterized by transmural inflammation of all layers of the bowel wall. The
282 formation of stenoses and strictures is common in this disease, which causes abdominal pain,
283 anorexia, and weight loss. Approximately 50% of CD patients undergo surgery for this type
284 of complication during a 10-year course; however, the recurrence rate after surgery is high. In
285 contrast, UC rarely causes intestinal stenosis. Cytokines released from inflammatory cells
286 have long been implicated in the pathogenesis of intestinal fibrosis. TGF- β /Smad signaling
287 plays an important role in CD [103-105]. The transmural infiltrate of CD is responsible for
288 initiating and maintaining a series of connective tissue changes not only involving the
289 mucosa, but also the submucosa and *muscularis mucosae* and *muscularis propria*, where a
290 marked increase of collagen type I, III, and V mRNA is observed [106,107]. TGF- β has been
291 identified as one of the central growth factors/cytokines that specifically induces a fibrotic
292 response after inflammatory injury in the intestinal tract. In CD, there is a marked
293 overexpression of TGF- β 1 and TGFBRs in the colonic mucosa [76,108]. Fibrosis in CD can
294 therefore be interpreted as an aberrant healing response to mucosal injury [109]. In addition,
295 TGF- β appears to be involved in intestinal fibrosis in other enteropathies, such as radiation
296 enteritis, collagenous colitis, and intestinal graft-versus host disease [110-112]. Both TGF- β
297 and its receptors are overexpressed in the intestine of patients with CD [113]. Intestinal
298 fibroblast expression of TGF- β isoforms varies according to the nature of tissue. Fibroblasts
299 from normal and inflamed mucosa both express the TGF- β 1 and TGF- β 3 isoforms, while
300 those from fibrotic tissue show reduced expression of TGF- β 3, but enhanced expression of
301 TGF- β 2 and TGF- β 1 [114]. This is remarkable, since the TGF- β 1 and TGF- β 2 isoforms have

302 been specifically implicated in pathogenic fibrosis, while TGF- β 3 appears to have antifibrotic
303 properties [115]. Monteleone and colleagues have provided insight into the failure of TGF- β
304 down-regulation in CD. Despite the abundant expression of TGF- β in the mucosa of patients
305 with CD, phospho-Smad3 is diminished in the mucosa compared to control mucosal samples,
306 as is the complex of Smad3 with Smad4. This may be due to the induction and overexpression
307 of Smad7 in the mucosa of patients with CD and UCs [116]. However, upregulation of Smad7
308 is not specific for inflammatory bowel disease (IBD), but also occurs in *Helicobacter pylori*-
309 induced gastritis [117]. In addition, mucosal T cells in both whole tissue and isolated cells
310 show defective TGF- β 1 signaling as measured by reduced immuno-reactivity against
311 phospho-Smad3 [117]. Specific anti-sense oligonucleotides for Smad7 reduce expression of
312 Smad7 in cells isolated from IBD patients which then become responsive to exogenous TGF-
313 β 1. TGF- β 1 cannot inhibit pro-inflammatory cytokine production in isolated lamina propria
314 mononuclear cells from CD patients, but inhibition of Smad7 with anti-sense oligonucleotides
315 restores TGF- β 1 signaling and allows TGF- β 1 to inhibit cytokine production. In inflamed
316 mucosal tissue explants from CD patients, inhibition of Smad7 also restores phospho-Smad3
317 and decreases pro-inflammatory cytokine production, an effect which is partially blocked by
318 anti-TGF- β 1. The extension of these studies examined the interactions between Smad
319 signaling and NF- κ B activation in inflamed gut: While TGF- β 1 is a potent inhibitor of TNF-
320 α -induced NF- κ B activation in normal gut, it has no activity in inflamed gut. This can be
321 attributed to over-expression of Smad7, since treatment of cells from inflamed gut with anti-
322 sense to Smad7 allows TGF- β 1 to rapidly down-regulate NF- κ B activation [117].

323 If it becomes possible to specifically inhibit Smad7, endogenous TGF- β 1 in the
324 inflamed gut may negatively regulate pro-inflammatory cytokine production and NF- κ B
325 activation, the major components of the immune overactivity which drives tissue injury in

326 IBD. At the same time, however, it is important to discover the factors which control Smad7
327 expression in the inflamed gut. Furthermore, cell-specific expression of Smad7 will be
328 important because TGF- β 1 has different effects on different cell types. Thus, while blocking
329 Smad7 will allow TGF- β 1 to reduce pro-inflammatory cytokine production by T cells and
330 macrophages, it may allow TGF- β 1 to increase collagen production in myofibroblasts,
331 resulting in fibrosis. However, at the moment the relative importance of the known inducers
332 of Smad7 in the gut, such as IFN- γ or TNF- α , or even whether TGF- β 1 itself induces Smad7
333 in a negative regulatory loop, is still unclear.

334

335 **6. TGF- β -RELATED MOLECULES IN RADIATION-INDUCED FIBROSIS**

336 TGF- β 1 levels are increased in irradiated mouse skin [118,119] and decrease slowly after
337 irradiation in both pig and human skin [120,121]. Following microvascular hard or soft tissue
338 transfer, TGF- β 1 is again upregulated in a biphasic manner. The first expression peak on day
339 3 post operation is due to enhanced activation of latent TGF- β 1 by extracellular enzymes
340 while the second between day 14 and 28 after surgery is a result of *de novo* synthesis [122].
341 Its most important signaling receptor TGFBR2 is upregulated in irradiated graft beds as well
342 [123]. Signaling leads to increased nucleoplasmatic shuttling of active Smad2/3 and induction
343 of TGF- β 1 target genes in fibrotic healing, which is mainly due to decrease in cytoplasmatic
344 levels of the inhibitory Smad7 [124].

345 As a consequence, the extracellular matrix is qualitatively and quantitatively altered
346 [125]. Prolyl-hydroxyprolinase- β overexpression [126] promotes synthesis of collagen I, III
347 and IV [124], while repression of degrading enzymes, such as MMP-1 and induction of tissue
348 inhibitors [127] suppresses the degrading pathways. Moreover, integrin surface receptors,
349 such as α 2 β 1 integrin are up-regulated as well and modulate transmission of tensile forces

350 [128]. In the presence of such forces fibroblasts differentiate into myofibroblasts resulting in
351 constrictive fibrosis [129].

352 Irradiation-induced fibrosis and damage to the microvasculature lead to wound healing
353 disorders following surgery in previously irradiated areas. Such disorders are dependent, in
354 part, on the radiation dosage and the timing of surgery after irradiation [130,131] and reduce
355 the success rate of free flaps to 90% compared with 94% in non-irradiated graft beds [132].

356 Taking the central role of TGF- β 1 in radiation-impaired wound healing into account,
357 the question of whether TGF- β 1-levels differ in the healthy tissue between different patients
358 is of utmost importance. It has been demonstrated that patients with an increased TGF- β 1
359 plasma level exhibit an increased risk of developing skin fibrosis following irradiation [133].
360 To find a particular predictor which sufficiently corresponds with the frequency to develop
361 wound healing complications following surgery in previously irradiated graft beds would give
362 a large clinical impact on planning individual treatment protocols. The availability of reliable
363 markers may eventually allow the prediction of outcome prior to commencement of treatment,
364 and thus allow modification of combined protocols to minimize late adverse effects without
365 compromising tumor control.

366

367 **7. TGF- β -RELATED MOLECULES IN MYOCARDITIS**

368 Heart fibrosis is a hallmark feature of the chronic stage of viral myocarditis [134,135]. Human
369 pathogenic coxsackievirus B3 (CVB3) is considered the most frequent viral cause of chronic
370 myocarditis in men [136]. Clinical manifestations of acute myocarditis vary from flu-like
371 symptoms to the fulminant fatal forms. Frequently, acute myocarditis, with distinct onset,
372 follows a monophasic clinical course, and the majority of patients recover spontaneously after
373 several days of congestive heart failure. Some patients progress into subacute or chronic

374 forms, which ultimately lead to death. The molecular mechanisms underlying fibrosis
375 development in chronic myocarditis are currently not well understood.

376 Like humans, mice develop a marked age-related susceptibility to CVB infections
377 [137]. The myocardial lesions in mice closely resemble those seen in human disease [138];
378 therefore, experimental murine models of coxsackievirus-induced myocarditis have been
379 developed to investigate the pathogenesis of this disease. Although chronic inflammation is
380 characteristic for the human disease as well as for our mouse model, fibrosis is more prevalent
381 in the latter.

382 Excessive fibrosis, as it occurs under conditions of chronic myocarditis, can be
383 classified as either replacement fibrosis, when functional tissue is replaced by connective
384 tissue, or as reactive fibrosis, which is part of an adaptive process [139]. In addition to
385 collagen, tenascin C, and fibronectin, splice variants of these molecules are often part of the
386 fibrotic tissue [140,141]. Various cytokines and growth factors are believed to contribute to
387 the induction of fibrosis, including TGF- β [142], IL-1, [143], TNF- α [144,145], and PDGFs
388 [146,147]. For example, persistent expression of cytokines in the chronic stage of CVB3-
389 induced myocarditis has been described for various mouse models [148-150], as well as for
390 human dilatative cardiomyopathy [151]. New observations suggest that sustained pro-
391 inflammatory signaling is associated with a pro-fibrotic phenotype based on TGF- β -mediated
392 signaling [152].

393 For fibrogenesis in the heart, members of the PDGF family are apparently important
394 mediators. Transgenic overexpression of PDGF-C and PDGF-D, two more recently
395 discovered PDGF isoforms [153], in the heart leads to massive cardiac fibrosis [146,147].
396 Therefore, the importance of PDGF for the pathology of chronic myocarditis was investigated
397 in mice with CVB3-induced myocarditis. Interestingly, all analysed isoforms of PDGF-A, -B,
398 and -C were upregulated in close correlation with the inflammatory process. High levels of

399 the growth factors persisted only in MHC class II knockout mice, which develop a chronic
400 myocarditis upon CVB3 infection, whereas immunocompetent C57BL/6 wild-type mice
401 exhibit only an acute, completely reversible myocarditis [154,155].

402 Furthermore, it has been shown that the PDGF-receptor (PDGFR) blocker Imatinib
403 inhibits activation of resident PDGFR, and attenuates fibrosis in this mouse model
404 significantly [156]. These data strongly suggest that elevation of PDGF levels and subsequent
405 activation of PDGFR causally contribute to the type of cardiac fibrosis which occurs in this
406 model. Efficacy of Imatinib for attenuation of fibrosis has recently been reported also for
407 other organs, i.e., liver, joints, and kidney [157-159]. It can therefore be assumed that
408 Imatinib-sensitive tyrosine kinases play a more general role in fibrogenesis. In addition to the
409 PDGFR [157,160], the Abelson tyrosine kinase, a mediator of TGF- β -signaling (c-Abl) [161],
410 has also been proposed as a relevant target for Imatinib in the inhibition of fibrogenesis
411 [159,162]. While our data [156], and data of others [146,147] strongly suggest a causal
412 involvement of the PDGFR in fibrogenesis, they do not exclude that c-Abl activity is also
413 involved in fibrogenetic signalling. It could function downstream of TGFBRs, but also
414 partially mediate signalling of the activated PDGFR [163]. It is known that TGF- β can drive
415 cardiac fibrosis when overexpressed in the mouse heart [164] and the use of genetic mouse
416 models to understand the role of TGF- β signalling in the heart is reviewed in [165]. But the
417 relative contribution of the different TGF- β -mediated signalling mechanisms to fibrosis in our
418 model of CVB3-induced chronic myocarditis remains to be fully elucidated.

419

420 **8. DIFFERENCES AND SIMILARITIES OF TGF- β IN DIFFERENT ORGANS**

421 As presented in Table 1, numerous molecules have similar expression patterns in the diseases
422 mentioned in this review. Due to limited information from the human system, some data were
423 replaced by results derived from the respective animal model (fibrosis following irradiation).

424 TGF- β 1 is elevated in most pathologies in its active or latent form and shows
425 similarities with the isoform TGF- β 2 in most cases. In contrast, the expression of TGF- β 3
426 varies from down-regulation during wound healing and CD, abundance without regulation in
427 rheumatoid arthritis and UC, to up-regulation during diabetic nephropathy. If determined, the
428 expression of TGFBR2 is often enhanced during disease. The widespread appearance of the
429 term “n.d.” points out the urgent need for further investigation of the expression and the
430 effects of TGF- β -related molecules in human diseases associated with fibrosis, particularly at
431 the stage of signaling and transcriptional regulation.

432

433 **9. THERAPEUTIC STRATEGIES**

434 As either increased or decreased activity of the TGF- β pathway has been implicated in the
435 pathogenesis of different human diseases, methods for increasing or decreasing signaling
436 through these pathways are required [166]. There are some tools for enhancing TGF- β
437 signals, e. g., direct administration of the ligand, usage of agonists, and increased expression
438 of receptors or decreased expression of signaling antagonists.

439 On the other hand, there are different ways to inhibit TGF- β signaling, e. g.,
440 administration of neutralizing antibodies, application of soluble receptors, usage of antisense
441 nucleotides, and chemically synthesized inhibitors of the receptor serine/threonine kinases.

442 Regarding TGF- β inhibition, some clinical trials have been performed with
443 neutralizing antibodies, especially in fibrotic diseases, including a TGF- β 2 neutralizing
444 antibody (lerdelimumab) which effectively decreased the amount of scarring after glaucoma
445 surgery [167]. A TGF- β 1 neutralizing antibody (CAT-192, metelimumab) has been
446 administered intravenously to patients with systemic sclerosis, which causes scarring in skin
447 and internal organs, however, without any evidence of efficacy and with more adverse events

448 and serious adverse events in *verum* than in placebo patients [168]. This raises the question of
449 whether broader or even complete blockade of the TGF- β axis will be safe or rather
450 associated with substantial toxicity [168]. Despite these concerns, a pan-TGF- β (GC-1008)
451 neutralizing antibody has recently been administered to patients with idiopathic pulmonary
452 fibrosis in a phase I clinical trial for safety evaluation [169] and is being administered to
453 patients with focal segmental glomerulosclerosis [170], renal cell carcinoma, and malignant
454 melanoma [171]; however, safety and efficacy data from these studies are not yet available.
455 Similar to neutralizing antibodies, soluble TGF- β superfamily receptors abrogate signaling at
456 the ligand level by binding the ligand and preventing it from binding to its cell surface
457 receptors. Due to the lack of studies in humans, only data from animal models are available.
458 For example, soluble TGFBR3 has demonstrated efficacy against renal damage progression
459 associated with diabetes in mice [172]. Similarly, soluble TGFBR2 has anti-cancer effects in
460 mice, as it suppresses the growth and metastasis of pancreatic cancer cells [173] and also
461 inhibits breast cancer cell growth, migration, invasion, and metastasis [174].
462 Using anti-sense oligonucleotides to reduce the expression of TGF- β superfamily members is
463 a relatively new therapeutic tool that has been already successfully applied in clinical trials for
464 cancer treatment [175,176]. Also, small molecule inhibitors have been introduced to block
465 TGF- β signaling. The TGFBR1-inhibitor Ly573636, which blocks intrinsic receptor-kinase
466 activity, is currently being assessed in patients with certain cancers (see [176]), whereas other
467 TGFBR1-inhibitors have shown to be efficient in mouse tumor models [177].

468

469 **10. CONCLUSIONS**

470 Due to the central role of TGF- β in severe fibrotic diseases (with similarities and
471 dissimilarities among organs/diseases) it is an attractive therapeutic target, as already shown
472 in several clinical trials (see above).

473 The strongly bifunctional role of TGF- β , however (pro-fibrotic, but anti-inflammatory),
474 requires great care for the application of TGF- β -directed treatments. Future strategies will
475 therefore have to focus on developing suitable tools to address this bifunctionality such as:
476 1) locally or regionally restricted administration of agonists or antagonists (in particular,
477 broad TGF- β blockade by agents such as TGF- β antibodies, TGF- β receptors (soluble/anti-
478 sense), latency-associated peptide or Smad7; 2) phase-dependent application in those periods
479 of disease dominated by the critical pathogenetic features of TGF- β ; and 3) development of
480 local or systemic biomarkers indicative of a future favorable response. The systemic
481 applicability of broad TGF- β blockade will have to await the safety data from ongoing trials.

482

483 **ACKNOWLEDGEMENTS**

484 The authors thank Ulrike Körner, Simone Schönfelder, Soo-Yeon Lee, Christin Thiele and
485 Nora Moll for excellent technical assistance. Monique Nürnberger and E. Palombo-Kinne are
486 gratefully acknowledged for critical reading and /or editing of the manuscript. The work of R.
487 W. Kinne was supported by the Deutsche Forschungsgemeinschaft (Ki439/6, Ki439/7), the
488 Interdisciplinary Center of Clinical Research Jena (FKZ 010405), and the Jena Center of
489 Bioinformatics (FKZ0313652B). G. Wolf received grants from the Deutsche
490 Forschungsgemeinschaft (Wo 460/14-1, 14-2) and the Interdisciplinary Center of Clinical
491 Research Jena.

492

493

494 **REFERENCES**

495

496 [1] Y. Shi, J. Massague, Mechanisms of TGF-beta signaling from cell membrane to the
497 nucleus, *Cell* 113 (2003) 685-700.

498 [2] B.J. Falser, R.A. Macsata, D. Plummer, L. Mishra, A.N. Sidawy, Transforming growth
499 factor-beta and wound healing, *Perspect. Vasc. Surg. Endovasc. Ther.* 18 (2006) 55-
500 62.

501 [3] B.A. Hocevar, P.H. Howe, Analysis of TGF-beta-mediated synthesis of extracellular
502 matrix components, *Methods Mol. Biol.* 142 (2000) 55-65.

503 [4] M.O. Li, R.A. Flavell, TGF-beta: a master of all T cell trades, *Cell* 134 (2008) 392-
504 404.

505 [5] S.M. Wahl, W. Chen, Transforming growth factor-beta-induced regulatory T cells
506 referee inflammatory and autoimmune diseases, *Arthritis Res. Ther.* 7 (2005) 62-68.

507 [6] J.P. Annes, J.S. Munger, D.B. Rifkin, Making sense of latent TGFbeta activation, *J.*
508 *Cell Sci.* 116 (2003) 217-224.

509 [7] Y. Sato, D.B. Rifkin, Inhibition of endothelial cell movement by pericytes and smooth
510 muscle cells: activation of a latent transforming growth factor-beta 1-like molecule by
511 plasmin during co-culture, *J. Cell Biol.* 109 (1989) 309-315.

512 [8] Q. Yu, I. Stamenkovic, Cell surface-localized matrix metalloproteinase-9
513 proteolytically activates TGF-beta and promotes tumor invasion and angiogenesis,
514 *Genes Dev.* 14 (2000) 163-176.

515 [9] J.S. Munger, X. Huang, H. Kawakatsu, M.J. Griffiths, S.L. Dalton, J. Wu, J.F. Pittet,
516 N. Kaminski, C. Garat, M.A. Matthay, D.B. Rifkin, D. Sheppard, The integrin alpha v
517 beta 6 binds and activates latent TGF beta 1: a mechanism for regulating pulmonary
518 inflammation and fibrosis, *Cell* 96 (1999) 319-328.

519 [10] D. Mu, S. Cambier, L. Fjellbirkeland, J.L. Baron, J.S. Munger, H. Kawakatsu, D.
520 Sheppard, V.C. Broaddus, S.L. Nishimura, The integrin alpha(v)beta8 mediates
521 epithelial homeostasis through MT1-MMP-dependent activation of TGF-beta1, *J. Cell*
522 *Biol.* 157 (2002) 493-507.

523 [11] P.J. Wipff, B. Hinz, Integrins and the activation of latent transforming growth factor
524 beta1 - an intimate relationship, *Eur. J. Cell Biol.* 87 (2008) 601-615.

- 525 [12] R.M. Lyons, J. Keski-Oja, H.L. Moses, Proteolytic activation of latent transforming
526 growth factor-beta from fibroblast-conditioned medium, *J. Cell Biol.* 106 (1988) 1659-
527 1665.
- 528 [13] J.E. Murphy-Ullrich, M. Poczatek, Activation of latent TGF-beta by thrombospondin-
529 1: mechanisms and physiology, *Cytokine Growth Factor Rev.* 11 (2000) 59-69.
- 530 [14] R. Derynck, Y.E. Zhang, Smad-dependent and Smad-independent pathways in TGF-
531 beta family signalling, *Nature* 425 (2003) 577-584.
- 532 [15] L. Chen, C. Klass, A. Woods, Syndecan-2 regulates transforming growth factor-beta
533 signaling, *J. Biol. Chem.* 279 (2004) 15715-15718.
- 534 [16] H.J. You, M.W. Bruinsma, T. How, J.H. Ostrander, G.C. Blobe, The type III TGF-
535 beta receptor signals through both Smad3 and the p38 MAP kinase pathways to
536 contribute to inhibition of cell proliferation, *Carcinogenesis* 28 (2007) 2491-2500.
- 537 [17] J. Massague, How cells read TGF-beta signals, *Nat. Rev. Mol. Cell Biol.* 1 (2000)
538 169-178.
- 539 [18] D. Wotton, J. Massague, Smad transcriptional corepressors in TGF beta family
540 signaling, *Curr. Top. Microbiol. Immunol.* 254 (2001) 145-164.
- 541 [19] J. Massague, Integration of Smad and MAPK pathways: a link and a linker revisited,
542 *Genes Dev.* 17 (2003) 2993-2997.
- 543 [20] I. Yakymovych, D.P. Ten, C.H. Heldin, S. Souchelnytskyi, Regulation of Smad
544 signaling by protein kinase C, *FASEB J.* 15 (2001) 553-555.
- 545 [21] A.R. Conery, Y. Cao, E.A. Thompson, C.M. Townsend, Jr., T.C. Ko, K. Luo, Akt
546 interacts directly with Smad3 to regulate the sensitivity to TGF-beta induced
547 apoptosis, *Nat. Cell Biol.* 6 (2004) 366-372.
- 548 [22] A. Masszi, C.C. Di, G. Sirokmany, W.T. Arthur, O.D. Rotstein, J. Wang, C.A.
549 McCulloch, L. Rosivall, I. Mucsi, A. Kapus, Central role for Rho in TGF-beta1-
550 induced alpha-smooth muscle actin expression during epithelial-mesenchymal
551 transition, *Am. J. Physiol Renal Physiol* 284 (2003) F911-F924.
- 552 [23] N. Muller, A. Reinacher-Schick, S. Baldus, H.J. van, G. Berx, A. Baar, R.F. van, W.
553 Schmiegel, I. Schwarte-Waldhoff, Smad4 induces the tumor suppressor E-cadherin
554 and P-cadherin in colon carcinoma cells, *Oncogene* 21 (2002) 6049-6058.
- 555 [24] H. Wang, V. Radjendirane, K.K. Wary, S. Chakrabarty, Transforming growth factor
556 beta regulates cell-cell adhesion through extracellular matrix remodeling and

- 557 activation of focal adhesion kinase in human colon carcinoma Moser cells, *Oncogene*
558 23 (2004) 5558-5561.
- 559 [25] V.J. Thannickal, D.Y. Lee, E.S. White, Z. Cui, J.M. Larios, R. Chacon, J.C. Horowitz,
560 R.M. Day, P.E. Thomas, Myofibroblast differentiation by transforming growth factor-
561 beta1 is dependent on cell adhesion and integrin signaling via focal adhesion kinase, *J.*
562 *Biol. Chem.* 278 (2003) 12384-12389.
- 563 [26] X. Shi-Wen, S.K. Parapuram, D. Pala, Y. Chen, D.E. Carter, M. Eastwood, C.P.
564 Denton, D.J. Abraham, A. Leask, Requirement of transforming growth factor beta-
565 activated kinase 1 for transforming growth factor beta-induced alpha-smooth muscle
566 actin expression and extracellular matrix contraction in fibroblasts, *Arthritis Rheum.*
567 60 (2009) 234-241.
- 568 [27] S. Liu, S.W. Xu, L. Kennedy, D. Pala, Y. Chen, M. Eastwood, D.E. Carter, C.M.
569 Black, D.J. Abraham, A. Leask, FAK is required for TGFbeta-induced JNK
570 phosphorylation in fibroblasts: implications for acquisition of a matrix-remodeling
571 phenotype, *Mol. Biol. Cell* 18 (2007) 2169-2178.
- 572 [28] H. Hayashi, S. Abdollah, Y. Qiu, J. Cai, Y.Y. Xu, B.W. Grinnell, M.A. Richardson,
573 J.N. Topper, M.A. Gimbrone, Jr., J.L. Wrana, D. Falb, The MAD-related protein
574 Smad7 associates with the TGFbeta receptor and functions as an antagonist of
575 TGFbeta signaling, *Cell* 89 (1997) 1165-1173.
- 576 [29] A. Nakao, M. Afrakhte, A. Moren, T. Nakayama, J.L. Christian, R. Heuchel, S. Itoh,
577 M. Kawabata, N.E. Heldin, C.H. Heldin, D.P. Ten, Identification of Smad7, a
578 TGFbeta-inducible antagonist of TGF-beta signalling, *Nature* 389 (1997) 631-635.
- 579 [30] G. Salvador, R. Sanmarti, B. Gil-Torregrosa, A. Garcia-Peiro, J.R. Rodriguez-Cros,
580 J.D. Canete, Synovial vascular patterns and angiogenic factors expression in synovial
581 tissue and serum of patients with rheumatoid arthritis, *Rheumatology. (Oxford)* 45
582 (2006) 966-971.
- 583 [31] S.Y. Kim, S.W. Han, G.W. Kim, J.M. Lee, Y.M. Kang, TGF-beta1 polymorphism
584 determines the progression of joint damage in rheumatoid arthritis, *Scand. J.*
585 *Rheumatol.* 33 (2004) 389-394.
- 586 [32] H. Cheon, S.J. Yu, D.H. Yoo, I.J. Chae, G.G. Song, J. Sohn, Increased expression of
587 pro-inflammatory cytokines and metalloproteinase-1 by TGF-beta1 in synovial
588 fibroblasts from rheumatoid arthritis and normal individuals, *Clin. Exp. Immunol.* 127
589 (2002) 547-552.

- 590 [33] Y. Yamanishi, D.L. Boyle, M. Clark, R.A. Maki, M.D. Tortorella, E.C. Arner, G.S.
591 Firestein, Expression and regulation of aggrecanase in arthritis: the role of TGF-beta,
592 J. Immunol. 168 (2002) 1405-1412.
- 593 [34] J.A. Hamilton, D.S. Piccoli, T. Leizer, D.M. Butler, M. Croatto, A.K. Royston,
594 Transforming growth factor beta stimulates urokinase-type plasminogen activator and
595 DNA synthesis, but not prostaglandin E2 production, in human synovial fibroblasts,
596 Proc. Natl. Acad. Sci. U. S. A 88 (1991) 7180-7184.
- 597 [35] F. Taketazu, M. Kato, A. Gobl, H. Ichijo, P. ten Dijke, J. Itoh, M. Kyogoku, J.
598 Ronnelid, K. Miyazono, C.H. Heldin, ., Enhanced expression of transforming growth
599 factor-beta s and transforming growth factor-beta type II receptor in the synovial
600 tissues of patients with rheumatoid arthritis, Lab. Invest. 70 (1994) 620-630.
- 601 [36] C.Q. Chu, M. Field, E. Abney, R.Q. Zheng, S. Allard, M. Feldmann, R.N. Maini,
602 Transforming growth factor-beta 1 in rheumatoid synovial membrane and
603 cartilage/pannus junction, Clin. Exp. Immunol. 86 (1991) 380-386.
- 604 [37] Z. Szekanecz, G.K. Haines, L.A. Harlow, M.R. Shah, T.W. Fong, R. Fu, S.J. Lin, G.
605 Rayan, A.E. Koch, Increased synovial expression of transforming growth factor
606 (TGF)-beta receptor endoglin and TGF-beta 1 in rheumatoid arthritis: possible
607 interactions in the pathogenesis of the disease, Clin. Immunol. Immunopathol. 76
608 (1995) 187-194.
- 609 [38] D.H. Goddard, S.L. Grossman, W.V. Williams, D.B. Weiner, J.L. Gross, K. Eidsvoog,
610 J.R. Dasch, Regulation of synovial cell growth. Coexpression of transforming growth
611 factor beta and basic fibroblast growth factor by cultured synovial cells, Arthritis
612 Rheum. 35 (1992) 1296-1303.
- 613 [39] R. Fava, N. Olsen, J. Keski-Oja, H. Moses, T. Pincus, Active and latent forms of
614 transforming growth factor beta activity in synovial effusions, J. Exp. Med. 169 (1989)
615 291-296.
- 616 [40] D. Pohlars, A. Beyer, D. Koczan, T. Wilhelm, H.J. Thiesen, R.W. Kinne, Constitutive
617 upregulation of the transforming growth factor-beta pathway in rheumatoid arthritis
618 synovial fibroblasts, Arthritis Res. Ther. 9 (2007) R59.
- 619 [41] A. Müssener, K. Funo, S. Kleinau, L. Klareskog, Dynamic expression of transforming
620 growth factor-betas (TGF-beta) and their type I and type II receptors in the synovial
621 tissue of arthritic rats, Clin. Exp. Immunol. 107 (1997) 112-119.

- 622 [42] J.B. Allen, C.L. Manthey, A.R. Hand, K. Ohura, L. Ellingsworth, S.M. Wahl, Rapid
623 onset synovial inflammation and hyperplasia induced by transforming growth factor
624 beta, *J. Exp. Med.* 171 (1990) 231-247.
- 625 [43] S.M. Wahl, J.B. Allen, G.L. Costa, H.L. Wong, J.R. Dasch, Reversal of acute and
626 chronic synovial inflammation by anti-transforming growth factor beta, *J. Exp. Med.*
627 177 (1993) 225-230.
- 628 [44] Z. Mi, S.C. Ghivizzani, E. Lechman, J.C. Glorioso, C.H. Evans, P.D. Robbins,
629 Adverse effects of adenovirus-mediated gene transfer of human transforming growth
630 factor beta 1 into rabbit knees, *Arthritis Res. Ther.* 5 (2003) R132-R139.
- 631 [45] M. Sakuma, K. Hatsushika, K. Koyama, R. Katoh, T. Ando, Y. Watanabe, M. Wako,
632 M. Kanzaki, S. Takano, H. Sugiyama, Y. Hamada, H. Ogawa, K. Okumura, A. Nakao,
633 TGF-beta type I receptor kinase inhibitor down-regulates rheumatoid synoviocytes
634 and prevents the arthritis induced by type II collagen antibody, *Int. Immunol.* 19
635 (2007) 117-126.
- 636 [46] M.E. Brandes, J.B. Allen, Y. Ogawa, S.M. Wahl, Transforming growth factor beta 1
637 suppresses acute and chronic arthritis in experimental animals, *J. Clin. Invest.* 87
638 (1991) 1108-1113.
- 639 [47] C. Schramm, J. Kriegsmann, M. Protschka, S. Huber, T. Hansen, E. Schmitt, P.R.
640 Galle, M. Blessing, Susceptibility to collagen-induced arthritis is modulated by
641 TGFbeta responsiveness of T cells, *Arthritis Res. Ther.* 6 (2004) R114-R119.
- 642 [48] Y. Chernajovsky, G. Adams, K. Triantaphyllopoulos, M.F. Ledda, O.L. Podhajcer,
643 Pathogenic lymphoid cells engineered to express TGF beta 1 ameliorate disease in a
644 collagen-induced arthritis model, *Gene Ther.* 4 (1997) 553-559.
- 645 [49] F.N. Ziyadeh, K. Sharma, M. Ericksen, G. Wolf, Stimulation of collagen gene
646 expression and protein synthesis in murine mesangial cells by high glucose is
647 mediated by autocrine activation of transforming growth factor-beta, *J. Clin. Invest.* 93
648 (1994) 536-542.
- 649 [50] F.N. Ziyadeh, Evidence for the involvement of transforming growth factor- in the
650 pathogenesis of diabetic kidney disease: Are Koch's postulates fulfilled?, *Curr. Pract.*
651 *Med.* 1 (1998) 87-89.
- 652 [51] G. Wolf, K. Sharma, Y. Chen, M. Ericksen, F.N. Ziyadeh, High glucose-induced
653 proliferation in mesangial cells is reversed by autocrine TGF-beta, *Kidney Int.* 42
654 (1992) 647-656.

- 655 [52] G. Wolf, E. Mueller, R.A. Stahl, F.N. Ziyadeh, Angiotensin II-induced hypertrophy of
656 cultured murine proximal tubular cells is mediated by endogenous transforming
657 growth factor-beta, *J. Clin. Invest.* 92 (1993) 1366-1372.
- 658 [53] T. Yamamoto, T. Nakamura, N.A. Noble, E. Ruoslahti, W.A. Border, Expression of
659 transforming growth factor beta is elevated in human and experimental diabetic
660 nephropathy, *Proc. Natl. Acad Sci. U. S. A* 90 (1993) 1814-1818.
- 661 [54] K. Sharma, F.N. Ziyadeh, Renal hypertrophy is associated with upregulation of TGF-
662 beta 1 gene expression in diabetic BB rat and NOD mouse, *Am. J. Physiol.* 267 (1994)
663 F1094-01.
- 664 [55] S.W. Hong, M. Isono, S. Chen, Iglesias-de la Cruz MC, D.C. Han, F.N. Ziyadeh,
665 Increased glomerular and tubular expression of transforming growth factor-beta1, its
666 type II receptor, and activation of the Smad signaling pathway in the db/db mouse,
667 *Am. J. Pathol.* 158 (2001) 1653-1663.
- 668 [56] K. Sharma, Y. Jin, J. Guo, F.N. Ziyadeh, Neutralization of TGF-beta by anti-TGF-beta
669 antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene
670 expression in STZ-induced diabetic mice, *Diabetes* 45 (1996) 522-530.
- 671 [57] F.N. Ziyadeh, B.B. Hoffman, D.C. Han, Iglesias-de la Cruz MC, S.W. Hong, M.
672 Isono, S. Chen, T.A. McGowan, K. Sharma, Long-term prevention of renal
673 insufficiency, excess matrix gene expression, and glomerular mesangial matrix
674 expansion by treatment with monoclonal antitransforming growth factor-beta antibody
675 in db/db diabetic mice, *Proc. Natl. Acad Sci. U. S. A* 97 (2000) 8015-8020.
- 676 [58] M. Isono, A. Mogyrosi, D.C. Han, B.B. Hoffman, F.N. Ziyadeh, Stimulation of TGF-
677 beta type II receptor by high glucose in mouse mesangial cells and in diabetic kidney,
678 *Am. J. Physiol. Renal Physiol.* 278 (2000) F830-F838.
- 679 [59] G. Wolf, G. Zahner, R. Schroeder, R.A. Stahl, Transforming growth factor beta
680 mediates the angiotensin-II-induced stimulation of collagen type IV synthesis in
681 cultured murine proximal tubular cells, *Nephrol. Dial. Transplant.* 11 (1996) 263-269.
- 682 [60] D.C. Han, B.B. Hoffman, S.W. Hong, J. Guo, F.N. Ziyadeh, Therapy with antisense
683 TGF-beta1 oligodeoxynucleotides reduces kidney weight and matrix mRNAs in
684 diabetic mice, *Am. J. Physiol. Renal Physiol.* 278 (2000) F628-F634.
- 685 [61] M. Abbate, C. Zoja, M. Morigi, D. Rottoli, S. Angioletti, S. Tomasoni, C. Zanchi, L.
686 Longaretti, R. Donadelli, G. Remuzzi, Transforming growth factor-beta1 is up-
687 regulated by podocytes in response to excess intraglomerular passage of proteins: a

- 688 central pathway in progressive glomerulosclerosis, *Am. J. Pathol.* 161 (2002) 2179-
689 2193.
- 690 [62] R.E. Gilbert, A. Cox, L.L. Wu, T.J. Allen, U.L. Hulthen, G. Jerums, M.E. Cooper,
691 Expression of transforming growth factor-beta1 and type IV collagen in the renal
692 tubulointerstitium in experimental diabetes: effects of ACE inhibition, *Diabetes* 47
693 (1998) 414-422.
- 694 [63] G. Wolf, S. Chen, F.N. Ziyadeh, From the periphery of the glomerular capillary wall
695 toward the center of disease: podocyte injury comes of age in diabetic nephropathy,
696 *Diabetes* 54 (2005) 1626-1634.
- 697 [64] D.T. Wu, M. Bitzer, W. Ju, P. Mundel, E.P. Bottinger, TGF-beta concentration
698 specifies differential signaling profiles of growth arrest/differentiation and apoptosis in
699 podocytes, *J. Am. Soc. Nephrol.* 16 (2005) 3211-3221.
- 700 [65] S.L. Dallas, K. Miyazono, T.M. Skerry, G.R. Mundy, L.F. Bonewald, Dual role for the
701 latent transforming growth factor-beta binding protein in storage of latent TGF-beta in
702 the extracellular matrix and as a structural matrix protein, *J. Cell Biol.* 131 (1995) 539-
703 549.
- 704 [66] M. Isono, S. Chen, S.W. Hong, Iglesias-de la Cruz MC, F.N. Ziyadeh, Smad pathway
705 is activated in the diabetic mouse kidney and Smad3 mediates TGF-beta-induced
706 fibronectin in mesangial cells, *Biochem. Biophys. Res Commun.* 296 (2002) 1356-
707 1365.
- 708 [67] Y. Furuse, N. Hashimoto, M. Maekawa, Y. Toyama, A. Nakao, I. Iwamoto, K.
709 Sakurai, Y. Suzuki, K. Yagui, S. Yuasa, K. Toshimori, Y. Saito, Activation of the
710 Smad pathway in glomeruli from a spontaneously diabetic rat model, OLETF rats,
711 *Nephron Exp. Nephrol.* 98 (2004) e100-e108.
- 712 [68] M. Schiffer, L.E. Schiffer, A. Gupta, A.S. Shaw, I.S. Roberts, P. Mundel, E.P.
713 Bottinger, Inhibitory smads and tgf-Beta signaling in glomerular cells, *J. Am. Soc.*
714 *Nephrol.* 13 (2002) 2657-2666.
- 715 [69] M. Cordenonsi, M. Montagner, M. Adorno, L. Zacchigna, G. Martello, A. Mamidi, S.
716 Soligo, S. Dupont, S. Piccolo, Integration of TGF-beta and Ras/MAPK signaling
717 through p53 phosphorylation, *Science* 315 (2007) 840-843.
- 718 [70] M. Schiffer, P. Mundel, A.S. Shaw, E.P. Bottinger, A novel role for the adaptor
719 molecule CD2-associated protein in transforming growth factor-beta-induced
720 apoptosis, *J. Biol. Chem.* 279 (2004) 37004-37012.

- 721 [71] K. Yoshioka, T. Takemura, K. Murakami, M. Okada, S. Hino, H. Miyamoto, S. Maki,
722 Transforming growth factor-beta protein and mRNA in glomeruli in normal and
723 diseased human kidneys, *Lab. Invest.* 68 (1993) 154-163.
- 724 [72] T. Yamamoto, N.A. Noble, A.H. Cohen, C.C. Nast, A. Hishida, L.I. Gold, W.A.
725 Border, Expression of transforming growth factor-beta isoforms in human glomerular
726 diseases, *Kidney Int.* 49 (1996) 461-469.
- 727 [73] K. Sharma, F.N. Ziyadeh, B. Alzahabi, T.A. McGowan, S. Kapoor, B.R. Kurnik, P.B.
728 Kurnik, L.S. Weisberg, Increased renal production of transforming growth factor-
729 beta1 in patients with type II diabetes, *Diabetes* 46 (1997) 854-859.
- 730 [74] S. Tsakas, D.S. Goumenos, Accurate measurement and clinical significance of urinary
731 transforming growth factor-beta1, *Am. J. Nephrol.* 26 (2006) 186-193.
- 732 [75] K. Sharma, B.O. Eltayeb, T.A. McGowan, S.R. Dunn, B. Alzahabi, R. Rohde, F.N.
733 Ziyadeh, E.J. Lewis, Captopril-induced reduction of serum levels of transforming
734 growth factor-beta1 correlates with long-term renoprotection in insulin-dependent
735 diabetic patients, *Am. J. Kidney Dis.* 34 (1999) 818-823.
- 736 [76] M.W. Babyatsky, G. Rossiter, D.K. Podolsky, Expression of transforming growth
737 factors alpha and beta in colonic mucosa in inflammatory bowel disease,
738 *Gastroenterology* 110 (1996) 975-984.
- 739 [77] G.C. Blobe, W.P. Schiemann, H.F. Lodish, Role of transforming growth factor beta in
740 human disease, *N. Engl. J. Med.* 342 (2000) 1350-1358.
- 741 [78] A. Izcue, J.L. Coombes, F. Powrie, Regulatory T cells suppress systemic and mucosal
742 immune activation to control intestinal inflammation, *Immunol. Rev.* 212 (2006) 256-
743 271.
- 744 [79] L. Gorelik, S. Constant, R.A. Flavell, Mechanism of transforming growth factor beta-
745 induced inhibition of T helper type 1 differentiation, *J. Exp. Med.* 195 (2002) 1499-
746 1505.
- 747 [80] A.B. Kulkarni, C.G. Huh, D. Becker, A. Geiser, M. Lyght, K.C. Flanders, A.B.
748 Roberts, M.B. Sporn, J.M. Ward, S. Karlsson, Transforming growth factor beta 1 null
749 mutation in mice causes excessive inflammatory response and early death, *Proc. Natl.*
750 *Acad. Sci. U. S. A* 90 (1993) 770-774.
- 751 [81] M. Boirivant, I.J. Fuss, A. Chu, W. Strober, Oxazolone colitis: A murine model of T
752 helper cell type 2 colitis treatable with antibodies to interleukin 4, *J. Exp. Med.* 188
753 (1998) 1929-1939.

- 754 [82] M.F. Neurath, S. Pettersson, K.H. Meyer zum Buschenfelde, W. Strober, Local
755 administration of antisense phosphorothioate oligonucleotides to the p65 subunit of
756 NF-kappa B abrogates established experimental colitis in mice, *Nat. Med.* 2 (1996)
757 998-1004.
- 758 [83] F. Powrie, J. Carlino, M.W. Leach, S. Mauze, R.L. Coffman, A critical role for
759 transforming growth factor-beta but not interleukin 4 in the suppression of T helper
760 type 1-mediated colitis by CD45RB(low) CD4+ T cells, *J. Exp. Med.* 183 (1996)
761 2669-2674.
- 762 [84] E. Bettelli, Y. Carrier, W. Gao, T. Korn, T.B. Strom, M. Oukka, H.L. Weiner, V.K.
763 Kuchroo, Reciprocal developmental pathways for the generation of pathogenic
764 effector TH17 and regulatory T cells, *Nature* 441 (2006) 235-238.
- 765 [85] P.R. Mangan, L.E. Harrington, D.B. O'Quinn, W.S. Helms, D.C. Bullard, C.O. Elson,
766 R.D. Hatton, S.M. Wahl, T.R. Schoeb, C.T. Weaver, Transforming growth factor-beta
767 induces development of the T(H)17 lineage, *Nature* 441 (2006) 231-234.
- 768 [86] M. Veldhoen, R.J. Hocking, C.J. Atkins, R.M. Locksley, B. Stockinger, TGFbeta in
769 the context of an inflammatory cytokine milieu supports de novo differentiation of IL-
770 17-producing T cells, *Immunity* 24 (2006) 179-189.
- 771 [87] V. Santarlasci, L. Maggi, M. Capone, F. Frosali, V. Querci, P.R. De, F. Liotta, L.
772 Cosmi, E. Maggi, S. Romagnani, F. Annunziato, TGF-beta indirectly favors the
773 development of human Th17 cells by inhibiting Th1 cells, *Eur. J. Immunol.* 39 (2009)
774 207-215.
- 775 [88] E. Volpe, N. Servant, R. Zollinger, S.I. Bogiatzi, P. Hupe, E. Barillot, V. Soumelis, A
776 critical function for transforming growth factor-beta, interleukin 23 and
777 proinflammatory cytokines in driving and modulating human T(H)-17 responses, *Nat.*
778 *Immunol.* 9 (2008) 650-657.
- 779 [89] L. Yang, D.E. Anderson, C. Baecher-Allan, W.D. Hastings, E. Bettelli, M. Oukka,
780 V.K. Kuchroo, D.A. Hafler, IL-21 and TGF-beta are required for differentiation of
781 human T(H)17 cells, *Nature* 454 (2008) 350-352.
- 782 [90] X. Yang, J.J. Letterio, R.J. Lechleider, L. Chen, R. Hayman, H. Gu, A.B. Roberts, C.
783 Deng, Targeted disruption of SMAD3 results in impaired mucosal immunity and
784 diminished T cell responsiveness to TGF-beta, *EMBO J.* 18 (1999) 1280-1291.

- 785 [91] G. Monteleone, A. Kumberova, N.M. Croft, C. McKenzie, H.W. Steer, T.T.
786 MacDonald, Blocking Smad7 restores TGF-beta1 signaling in chronic inflammatory
787 bowel disease, *J. Clin. Invest* 108 (2001) 601-609.
- 788 [92] N.A. Bhowmick, M. Ghiassi, A. Bakin, M. Aakre, C.A. Lundquist, M.E. Engel, C.L.
789 Arteaga, H.L. Moses, Transforming growth factor-beta1 mediates epithelial to
790 mesenchymal transdifferentiation through a RhoA-dependent mechanism, *Mol. Biol.*
791 *Cell* 12 (2001) 27-36.
- 792 [93] V. Ellenrieder, S.F. Hendler, W. Boeck, T. Seufferlein, A. Menke, C. Ruhland, G.
793 Adler, T.M. Gress, Transforming growth factor beta1 treatment leads to an epithelial-
794 mesenchymal transdifferentiation of pancreatic cancer cells requiring extracellular
795 signal-regulated kinase 2 activation, *Cancer Res.* 61 (2001) 4222-4228.
- 796 [94] K. Fujimoto, H. Sheng, J. Shao, R.D. Beauchamp, Transforming growth factor-beta1
797 promotes invasiveness after cellular transformation with activated Ras in intestinal
798 epithelial cells, *Exp. Cell Res.* 266 (2001) 239-249.
- 799 [95] K. Lehmann, E. Janda, C.E. Pierreux, M. Rytomaa, A. Schulze, M. McMahon, C.S.
800 Hill, H. Beug, J. Downward, Raf induces TGFbeta production while blocking its
801 apoptotic but not invasive responses: a mechanism leading to increased malignancy in
802 epithelial cells, *Genes Dev.* 14 (2000) 2610-2622.
- 803 [96] M. Oft, K.H. Heider, H. Beug, TGFbeta signaling is necessary for carcinoma cell
804 invasiveness and metastasis, *Curr. Biol.* 8 (1998) 1243-1252.
- 805 [97] G. Portella, S.A. Cumming, J. Liddell, W. Cui, H. Ireland, R.J. Akhurst, A. Balmain,
806 Transforming growth factor beta is essential for spindle cell conversion of mouse skin
807 carcinoma in vivo: implications for tumor invasion, *Cell Growth Differ.* 9 (1998) 393-
808 404.
- 809 [98] A.M. Arias, Epithelial mesenchymal interactions in cancer and development, *Cell* 105
810 (2001) 425-431.
- 811 [99] J.P. Thiery, D. Chopin, Epithelial cell plasticity in development and tumor
812 progression, *Cancer Metastasis Rev.* 18 (1999) 31-42.
- 813 [100] R. Kalluri, E.G. Neilson, Epithelial-mesenchymal transition and its implications for
814 fibrosis, *J. Clin. Invest* 112 (2003) 1776-1784.
- 815 [101] S. Saika, S. Kono-Saika, Y. Ohnishi, M. Sato, Y. Muragaki, A. Ooshima, K.C.
816 Flanders, J. Yoo, M. Anzano, C.Y. Liu, W.W. Kao, A.B. Roberts, Smad3 signaling is

- 817 required for epithelial-mesenchymal transition of lens epithelium after injury, *Am. J.*
818 *Pathol.* 164 (2004) 651-663.
- 819 [102] M. Sato, Y. Muragaki, S. Saika, A.B. Roberts, A. Ooshima, Targeted disruption of
820 TGF-beta1/Smad3 signaling protects against renal tubulointerstitial fibrosis induced
821 by unilateral ureteral obstruction, *J. Clin. Invest* 112 (2003) 1486-1494.
- 822 [103] P.L. Beck, D.K. Podolsky, Growth factors in inflammatory bowel disease, *Inflamm.*
823 *Bowel. Dis.* 5 (1999) 44-60.
- 824 [104] C. Fiocchi, TGF-beta/Smad signaling defects in inflammatory bowel disease:
825 mechanisms and possible novel therapies for chronic inflammation, *J. Clin. Invest* 108
826 (2001) 523-526.
- 827 [105] K.B. Hahm, Y.H. Im, T.W. Parks, S.H. Park, S. Markowitz, H.Y. Jung, J. Green, S.J.
828 Kim, Loss of transforming growth factor beta signalling in the intestine contributes to
829 tissue injury in inflammatory bowel disease, *Gut* 49 (2001) 190-198.
- 830 [106] H. Matthes, H. Herbst, D. Schuppan, A. Stallmach, S. Milani, H. Stein, E.O. Riecken,
831 Cellular localization of procollagen gene transcripts in inflammatory bowel diseases,
832 *Gastroenterology* 102 (1992) 431-442.
- 833 [107] A. Stallmach, D. Schuppan, H.H. Riese, H. Matthes, E.O. Riecken, Increased collagen
834 type III synthesis by fibroblasts isolated from strictures of patients with Crohn's
835 disease, *Gastroenterology* 102 (1992) 1920-1929.
- 836 [108] I.C. Lawrance, L. Maxwell, W. Doe, Inflammation location, but not type, determines
837 the increase in TGF-beta1 and IGF-1 expression and collagen deposition in IBD
838 intestine, *Inflamm. Bowel. Dis.* 7 (2001) 16-26.
- 839 [109] J.B. Pucilowska, K.L. Williams, P.K. Lund, Fibrogenesis. IV. Fibrosis and
840 inflammatory bowel disease: cellular mediators and animal models, *Am. J. Physiol*
841 *Gastrointest. Liver Physiol* 279 (2000) G653-G659.
- 842 [110] L.M. Liem, W.E. Fibbe, H.C. van Houwelingen, E. Goulmy, Serum transforming
843 growth factor-beta1 levels in bone marrow transplant recipients correlate with blood
844 cell counts and chronic graft-versus-host disease, *Transplantation* 67 (1999) 59-65.
- 845 [111] M. Stahle-Backdahl, J. Maim, B. Veress, C. Benoni, K. Bruce, A. Egesten, Increased
846 presence of eosinophilic granulocytes expressing transforming growth factor-beta1 in
847 collagenous colitis, *Scand. J. Gastroenterol.* 35 (2000) 742-746.

- 848 [112] J. Wang, H. Zheng, C.C. Sung, K.K. Richter, M. Hauer-Jensen, Cellular sources of
849 transforming growth factor-beta isoforms in early and chronic radiation enteropathy,
850 *Am. J. Pathol.* 153 (1998) 1531-1540.
- 851 [113] F.F. di Mola, H. Friess, A. Scheuren, S.P. Di, H. Graber, B. Egger, A. Zimmermann,
852 M. Korc, M.W. Buchler, Transforming growth factor-betas and their signaling
853 receptors are coexpressed in Crohn's disease, *Ann. Surg.* 229 (1999) 67-75.
- 854 [114] B.C. McKaig, K. Hughes, P.J. Tighe, Y.R. Mahida, Differential expression of TGF-
855 beta isoforms by normal and inflammatory bowel disease intestinal myofibroblasts,
856 *Am. J. Physiol Cell Physiol* 282 (2002) C172-C182.
- 857 [115] A. Leask, D.J. Abraham, TGF-beta signaling and the fibrotic response, *FASEB J.* 18
858 (2004) 816-827.
- 859 [116] G. Monteleone, B.G. Del Vecchio, I. Monteleone, D. Fina, R. Caruso, V. Gioia, S.
860 Ballerini, G. Federici, S. Bernardini, F. Pallone, T.T. MacDonald, Post-transcriptional
861 regulation of Smad7 in the gut of patients with inflammatory bowel disease,
862 *Gastroenterology* 129 (2005) 1420-1429.
- 863 [117] G. Monteleone, B.G. Del Vecchio, G. Palmieri, P. Vavassori, I. Monteleone, A.
864 Colantoni, S. Battista, L.G. Spagnoli, M. Romano, M. Borrelli, T.T. MacDonald, F.
865 Pallone, Induction and regulation of Smad7 in the gastric mucosa of patients with
866 *Helicobacter pylori* infection, *Gastroenterology* 126 (2004) 674-682.
- 867 [118] K. Randall, J.E. Coggle, Expression of transforming growth factor-beta 1 in mouse
868 skin during the acute phase of radiation damage, *Int. J. Radiat. Biol.* 68 (1995) 301-
869 309.
- 870 [119] K. Randall, J.E. Coggle, Long-term expression of transforming growth factor TGF
871 beta 1 in mouse skin after localized beta-irradiation, *Int. J. Radiat. Biol.* 70 (1996)
872 351-360.
- 873 [120] M. Martin, J.L. Lefaix, P. Pinton, F. Crechet, F. Daburon, Temporal modulation of
874 TGF-beta 1 and beta-actin gene expression in pig skin and muscular fibrosis after
875 ionizing radiation, *Radiat. Res.* 134 (1993) 63-70.
- 876 [121] M. Martin, J. Lefaix, S. Delanian, TGF-beta1 and radiation fibrosis: a master switch
877 and a specific therapeutic target?, *Int. J. Radiat. Oncol. Biol. Phys.* 47 (2000) 277-290.
- 878 [122] S. Schultze-Mosgau, F. Wehrhan, G. Grabenbauer, K. Amann, M. Radespiel-Troger,
879 F.W. Neukam, F. Rodel, Transforming growth factor beta1 and beta2 (TGFbeta2 /

- 880 TGFbeta2) profile changes in previously irradiated free flap beds, *Head Neck* 24
881 (2002) 33-41.
- 882 [123] S. Schultze-Mosgau, F. Wehrhan, F. Rodel, K. Amann, M. Radespiel-Troger, G.G.
883 Grabenbauer, Transforming growth factor-beta receptor-II up-regulation during wound
884 healing in previously irradiated graft beds in vivo, *Wound Repair Regen.* 11 (2003)
885 297-305.
- 886 [124] S. Schultze-Mosgau, M.A. Blaese, G. Grabenbauer, F. Wehrhan, J. Kopp, K. Amann,
887 H.P. Rodemann, F. Rodel, Smad-3 and Smad-7 expression following anti-
888 transforming growth factor beta 1 (TGFbeta1)-treatment in irradiated rat tissue,
889 *Radiother. Oncol.* 70 (2004) 249-259.
- 890 [125] W.A. Border, N.A. Noble, Transforming growth factor beta in tissue fibrosis, *N. Engl.*
891 *J. Med.* 331 (1994) 1286-1292.
- 892 [126] S. Schultze-Mosgau, J. Kopp, M. Thorwarth, F. Rodel, I. Melnychenko, G.G.
893 Grabenbauer, K. Amann, F. Wehrhan, Plasminogen activator inhibitor-I-related
894 regulation of procollagen I (alpha1 and alpha2) by antitransforming growth factor-
895 beta1 treatment during radiation-impaired wound healing, *Int. J. Radiat. Oncol. Biol.*
896 *Phys.* 64 (2006) 280-288.
- 897 [127] D. Ulrich, F. Lichtenegger, M. Eblenkamp, D. Repper, N. Pallua, Matrix
898 metalloproteinases, tissue inhibitors of metalloproteinases, aminoterminal propeptide
899 of procollagen type III, and hyaluronan in sera and tissue of patients with capsular
900 contracture after augmentation with Trilucent breast implants, *Plast Reconstr Surg* 114
901 (2004) 229-236.
- 902 [128] J.A. Schiro, B.M. Chan, W.T. Roswit, P.D. Kassner, A.P. Pentland, M.E. Hemler,
903 A.Z. Eisen, T.S. Kupper, Integrin alpha 2 beta 1 (VLA-2) mediates reorganization and
904 contraction of collagen matrices by human cells, *Cell* 67 (1991) 403-410.
- 905 [129] P.D. Arora, N. Narani, C.A. McCulloch, The compliance of collagen gels regulates
906 transforming growth factor-beta induction of alpha-smooth muscle actin in fibroblasts,
907 *Am. J. Pathol.* 154 (1999) 871-882.
- 908 [130] M.K. Tibbs, Wound healing following radiation therapy: a review, *Radiother. Oncol.*
909 42 (1997) 99-106.
- 910 [131] R. Tokarek, E.F. Bernstein, F. Sullivan, J. Uitto, J.B. Mitchell, Effect of therapeutic
911 radiation on wound healing, *Clin. Dermatol.* 12 (1994) 57-70.

- 912 [132] S. Schultze-Mosgau, G.G. Grabenbauer, M. Radespiel-Troger, J. Wiltfang, J. Ries,
913 F.W. Neukam, F. Rodel, Vascularization in the transition area between free grafted
914 soft tissues and pre-irradiated graft bed tissues following preoperative radiotherapy in
915 the head and neck region, *Head Neck* 24 (2002) 42-51.
- 916 [133] C. Li, P.B. Wilson, E. Levine, J. Barber, A.L. Stewart, S. Kumar, TGF-beta1 levels in
917 pre-treatment plasma identify breast cancer patients at risk of developing post-
918 radiotherapy fibrosis, *Int. J. Cancer* 84 (1999) 155-159.
- 919 [134] A.M. Feldman, D. McNamara, Myocarditis, *N. Engl. J. Med.* 343 (2000) 1388-1398.
- 920 [135] B. Maisch, A. Richter, A. Sandmoller, I. Portig, S. Pankuweit, Inflammatory dilated
921 cardiomyopathy (DCMI), *Herz* 30 (2005) 535-544.
- 922 [136] L. Rey, V. Lambert, P. Watre, L. Andreoletti, Detection of enteroviruses ribonucleic
923 acid sequences in endomyocardial tissue from adult patients with chronic dilated
924 cardiomyopathy by a rapid RT-PCR and hybridization assay, *J. Med. Virol.* 64 (2001)
925 133-140.
- 926 [137] J.L. Melnik, Enteroviruses: polioviruses, coxsackieviruses, echoviruses, and newer
927 enteroviruses, in: Fields BN, Knipe DM, Howley PM (Eds.), *Virology*, Lippincott-
928 Raven, Philadelphia, 1996, pp. 655-712.
- 929 [138] N.R. Rose, D.A. Neumann, A. Herskovitz, Coxsackievirus myocarditis, in: G.H.
930 Stollerman, J. T. LaMont, J. J. Leonard, M. D. Siperstein (Eds.), *Advances in internal*
931 *medicine*, Mosby-Year Book, Inc., St.Louis, 1992, pp. 411-429.
- 932 [139] B. Maisch, H. Rupp, [Myocardial fibrosis: a cardiopathophysiologic Janus head], *Herz*
933 31 (2006) 260-268.
- 934 [140] U. Eismann, M. Sommer, H. Kosmehl, D. Appenroth, C. Fleck, G. Stein, Fibronectin
935 splice variants--prognostic markers for the stage of renal interstitial fibrosis in the rat,
936 *Nephron* 92 (2002) 379-388.
- 937 [141] A. Tamura, S. Kusachi, K. Nogami, A. Yamanishi, Y. Kajikawa, S. Hirohata, T. Tsuji,
938 Tenascin expression in endomyocardial biopsy specimens in patients with dilated
939 cardiomyopathy: distribution along margin of fibrotic lesions, *Heart* 75 (1996) 291-
940 294.
- 941 [142] S. Rosenkranz, M. Fleisch, K. Amann, C. Haeuseler, H. Kilter, U. Seeland, K.D.
942 Schluter, M. Bohm, Alterations of beta-adrenergic signaling and cardiac hypertrophy
943 in transgenic mice overexpressing TGF-beta(1), *Am. J. Physiol. Heart. Circ. Physiol.*
944 283 (2002) H1253-H1262.

- 945 [143] K. Isoda, Y. Kamezawa, N. Tada, M. Sato, F. Ohsuzu, Myocardial hypertrophy in
946 transgenic mice overexpressing human interleukin 1alpha, *J. Card. Fail.* 7 (2001) 355-
947 364.
- 948 [144] D. Bryant, L. Becker, J. Richardson, J. Shelton, F. Franco, R. Peshock, M. Thompson,
949 B. Giroir, Cardiac failure in transgenic mice with myocardial expression of tumor
950 necrosis factor-alpha, *Circulation* 97 (1998) 1375-1381.
- 951 [145] T. Kubota, C.F. McTiernan, C.S. Frye, A.J. Demetris, A.M. Feldman, Cardiac-specific
952 overexpression of tumor necrosis factor-alpha causes lethal myocarditis in transgenic
953 mice, *J. Card. Fail.* 3 (1997) 117-124.
- 954 [146] A. Ponten, X. Li, P. Thoren, K. Aase, T. Sjoblom, A. Ostman, U. Eriksson, Transgenic
955 overexpression of platelet-derived growth factor-C in the mouse heart induces cardiac
956 fibrosis, hypertrophy, and dilated cardiomyopathy, *Am. J. Pathol.* 163 (2003) 673-682.
- 957 [147] A. Ponten, E.B. Folestad, K. Pietras, U. Eriksson, Platelet-derived growth factor D
958 induces cardiac fibrosis and proliferation of vascular smooth muscle cells in heart-
959 specific transgenic mice, *Circ. Res.* 97 (2005) 1036-1045.
- 960 [148] B. Gluck, M. Schmidtke, I. Merkle, A. Stelzner, D. Gemsa, Persistent expression of
961 cytokines in the chronic stage of CVB3-induced myocarditis in NMRI mice, *J. Mol.*
962 *Cell Cardiol.* 33 (2001) 1615-1626.
- 963 [149] M. Schmidtke, B. Gluck, I. Merkle, P. Hofmann, A. Stelzner, D. Gemsa, Cytokine
964 profiles in heart, spleen, and thymus during the acute stage of experimental
965 coxsackievirus B3-induced chronic myocarditis, *J. Med. Virol.* 61 (2000) 518-526.
- 966 [150] T. Shioi, A. Matsumori, S. Sasayama, Persistent expression of cytokine in the chronic
967 stage of viral myocarditis in mice, *Circulation* 94 (1996) 2930-2937.
- 968 [151] M. Satoh, G. Tamura, I. Segawa, A. Tashiro, K. Hiramori, R. Satodate, Expression of
969 cytokine genes and presence of enteroviral genomic RNA in endomyocardial biopsy
970 tissues of myocarditis and dilated cardiomyopathy, *Virchows Arch.* 427 (1996) 503-
971 509.
- 972 [152] Y. Sakata, A.L. Chancey, V.G. Divakaran, K. Sekiguchi, N. Sivasubramanian, D.L.
973 Mann, Transforming growth factor-beta receptor antagonism attenuates myocardial
974 fibrosis in mice with cardiac-restricted overexpression of tumor necrosis factor, *Basic*
975 *Res. Cardiol.* 103 (2008) 60-68.
- 976 [153] C.H. Heldin, U. Eriksson, A. Ostman, New members of the platelet-derived growth
977 factor family of mitogens, *Arch. Biochem. Biophys.* 398 (2002) 284-290.

- 978 [154] K. Grun, B. Markova, F.D. Bohmer, A. Berndt, H. Kosmehl, C. Leipner, Elevated
979 expression of PDGF-C in coxsackievirus B3-induced chronic myocarditis, *Eur. Heart*
980 *J.* 26 (2005) 728-739.
- 981 [155] R. Wessely, K. Klingel, K.U. Knowlton, R. Kandolf, Cardioslective infection with
982 coxsackievirus B3 requires intact type I interferon signaling: implications for mortality
983 and early viral replication, *Circulation* 103 (2001) 756-761.
- 984 [156] C. Leipner, K. Grun, A. Muller, E. Buchdunger, L. Borsi, H. Kosmehl, A. Berndt, T.
985 Janik, A. Uecker, M. Kiehntopf, F.D. Bohmer, Imatinib mesylate attenuates fibrosis in
986 coxsackievirus b3-induced chronic myocarditis, *Cardiovasc. Res.* 79 (2008) 118-126.
- 987 [157] M. Neef, M. Ledermann, H. Saegesser, V. Schneider, N. Widmer, L.A. Decosterd, B.
988 Rochat, J. Reichen, Oral imatinib treatment reduces early fibrogenesis but does not
989 prevent progression in the long term, *J. Hepatol.* 44 (2006) 167-175.
- 990 [158] R.T. Paniagua, O. Sharpe, P.P. Ho, S.M. Chan, A. Chang, J.P. Higgins, B.H.
991 Tomooka, F.M. Thomas, J.J. Song, S.B. Goodman, D.M. Lee, M.C. Genovese, P.J.
992 Utz, L. Steinman, W.H. Robinson, Selective tyrosine kinase inhibition by imatinib
993 mesylate for the treatment of autoimmune arthritis, *J. Clin. Invest.* 116 (2006) 2633-
994 2642.
- 995 [159] S. Wang, M.C. Wilkes, E.B. Leof, R. Hirschberg, Imatinib mesylate blocks a non-
996 Smad TGF-beta pathway and reduces renal fibrogenesis in vivo, *FASEB J.* 19 (2005)
997 1-11.
- 998 [160] Y. Aono, Y. Nishioka, M. Inayama, M. Ugai, J. Kishi, H. Uehara, K. Izumi, S. Sone,
999 Imatinib as a novel antifibrotic agent in bleomycin-induced pulmonary fibrosis in
1000 mice, *Am. J. Respir. Crit. Care Med.* 171 (2005) 1279-1285.
- 1001 [161] M.C. Wilkes, H. Mitchell, S.G. Penheiter, J.J. Dore, K. Suzuki, M. Edens, D.K.
1002 Sharma, R.E. Pagano, E.B. Leof, Transforming growth factor-beta activation of
1003 phosphatidylinositol 3-kinase is independent of Smad2 and Smad3 and regulates
1004 fibroblast responses via p21-activated kinase-2, *Cancer Res.* 65 (2005) 10431-10440.
- 1005 [162] C.E. Daniels, M.C. Wilkes, M. Edens, T.J. Kottom, S.J. Murphy, A.H. Limper, E.B.
1006 Leof, Imatinib mesylate inhibits the profibrogenic activity of TGF-beta and prevents
1007 bleomycin-mediated lung fibrosis, *J. Clin. Invest.* 114 (2004) 1308-1316.
- 1008 [163] R. Plattner, L. Kadlec, K.A. DeMali, A. Kazlauskas, A.M. Pendergast, c-Abl is
1009 activated by growth factors and Src family kinases and has a role in the cellular
1010 response to PDGF, *Genes Dev.* 13 (1999) 2400-2411.

- 1011 [164] U. Seeland, C. Haeuseler, R. Hinrichs, S. Rosenkranz, T. Pfitzner, K. Scharffetter-
1012 Kochanek, M. Bohm, Myocardial fibrosis in transforming growth factor-beta(1) (TGF-
1013 beta(1)) transgenic mice is associated with inhibition of interstitial collagenase, *Eur. J.*
1014 *Clin. Invest.* 32 (2002) 295-303.
- 1015 [165] H. Xiao, Y.Y. Zhang, Understanding the role of transforming growth factor-beta
1016 signalling in the heart: overview of studies using genetic mouse models, *Clin. Exp.*
1017 *Pharmacol. Physiol.* 35 (2008) 335-341.
- 1018 [166] K.J. Gordon, G.C. Blobe, Role of transforming growth factor-beta superfamily
1019 signaling pathways in human disease, *Biochim. Biophys Acta* 1782 (2008) 197-228.
- 1020 [167] A.L. Mead, T.T. Wong, M.F. Cordeiro, I.K. Anderson, P.T. Khaw, Evaluation of anti-
1021 TGF-beta2 antibody as a new postoperative anti-scarring agent in glaucoma surgery,
1022 *Invest. Ophthalmol. Vis. Sci.* 44 (2003) 3394-3401.
- 1023 [168] C.P. Denton, P.A. Merkel, D.E. Furst, D. Khanna, P. Emery, V.M. Hsu, N. Silliman, J.
1024 Streisand, J. Powell, A. Akesson, J. Coppock, F. Hoogen, A. Herrick, M.D. Mayes, D.
1025 Veale, J. Haas, S. Ledbetter, J.H. Korn, C.M. Black, J.R. Seibold, Recombinant human
1026 anti-transforming growth factor beta1 antibody therapy in systemic sclerosis: a
1027 multicenter, randomized, placebo-controlled phase I/II trial of CAT-192, *Arthritis*
1028 *Rheum.* 56 (2007) 323-333.
- 1029 [169] [<http://clinicaltrials.gov/ct2/show/NCT00125385>]
- 1030 [170] [<http://clinicaltrials.gov/ct2/show/NCT00464321>]
- 1031 [171] [<http://clinicaltrials.gov/ct2/show/NCT00356460>]
- 1032 [172] P. Juarez, M.M. Vilchis-Landeros, J. Ponce-Coria, V. Mendoza, R. Hernandez-Pando,
1033 N.A. Bobadilla, F. Lopez-Casillas, Soluble betaglycan reduces renal damage
1034 progression in db/db mice, *Am. J. Physiol. Renal. Physiol.* 292 (2007) F321-F329.
- 1035 [173] M.A. Rowland-Goldsmith, H. Maruyama, K. Matsuda, T. Idezawa, M. Ralli, S. Ralli,
1036 M. Korc, Soluble type II transforming growth factor-beta receptor attenuates
1037 expression of metastasis-associated genes and suppresses pancreatic cancer cell
1038 metastasis, *Mol. Cancer Ther.* 1 (2002) 161-167.
- 1039 [174] R.S. Muraoka, N. Dumont, C.A. Ritter, T.C. Dugger, D.M. Brantley, J. Chen, E.
1040 Easterly, L.R. Roebuck, S. Ryan, P.J. Gotwals, V. Koteliansky, C.L. Arteaga,
1041 Blockade of TGF-beta inhibits mammary tumor cell viability, migration, and
1042 metastases, *J. Clin. Invest.* 109 (2002) 1551-1559.

- 1043 [175] P. Hau, P. Jachimczak, R. Schlingensiepen, F. Schulmeyer, T. Jauch, A. Steinbrecher,
1044 A. Brawanski, M. Proescholdt, J. Schlaier, J. Buchroithner, J. Pichler, G. Wurm, M.
1045 Mehdorn, R. Strege, G. Schuierer, V. Villarrubia, F. Fellner, O. Jansen, T. Straube, V.
1046 Nohria, M. Goldbrunner, M. Kunst, S. Schmaus, G. Stauder, U. Bogdahn, K.H.
1047 Schlingensiepen, Inhibition of TGF-beta2 with AP 12009 in recurrent malignant
1048 gliomas: from preclinical to phase I/II studies, *Oligonucleotides* 17 (2007) 201-212.
- 1049 [176] J. Nemunaitis, R.O. Dillman, P.O. Schwarzenberger, N. Senzer, C. Cunningham, J.
1050 Cutler, A. Tong, P. Kumar, B. Pappen, C. Hamilton, E. DeVol, P.B. Maples, L. Liu, T.
1051 Chamberlin, D.L. Shawler, H. Fakhrai, Phase II study of belagenpumatucel-L, a
1052 transforming growth factor beta-2 antisense gene-modified allogeneic tumor cell
1053 vaccine in non-small-cell lung cancer, *J. Clin. Oncol.* 24 (2006) 4721-4730.
- 1054 [177] E. Suzuki, S. Kim, H.K. Cheung, M.J. Corbley, X. Zhang, L. Sun, F. Shan, J. Singh,
1055 W.C. Lee, S.M. Albelda, L.E. Ling, A novel small-molecule inhibitor of transforming
1056 growth factor beta type I receptor kinase (SM16) inhibits murine mesothelioma tumor
1057 growth in vivo and prevents tumor recurrence after surgical resection, *Cancer Res* 67
1058 (2007) 2351-2359.
- 1059 [178] N.G. Frangogiannis, G. Ren, O. Dewald, P. Zymek, S. Haudek, A. Koerting, K.
1060 Winkelmann, L.H. Michael, J. Lawler, M.L. Entman, Critical role of endogenous
1061 thrombospondin-1 in preventing expansion of healing myocardial infarcts, *Circulation*
1062 111 (2005) 2935-2942.
- 1063 [179] Chuva de Sousa Lopes SM, A. Feijen, J. Korving, O. Korchynskyi, J. Larsson, S.
1064 Karlsson, P. ten Dijke, K.M. Lyons, R. Goldschmeding, P. Doevendans, C.L.
1065 Mummery, Connective tissue growth factor expression and Smad signaling during
1066 mouse heart development and myocardial infarction, *Dev. Dyn.* 231 (2004) 542-550.
- 1067 [180] I. Gotis-Graham, P.J. Hogg, H.P. McNeil, Significant correlation between
1068 thrombospondin 1 and serine proteinase expression in rheumatoid synovium, *Arthritis*
1069 *Rheum.* 40 (1997) 1780-1787.
- 1070 [181] F. Wu, T. Dassopoulos, L. Cope, A. Maitra, S.R. Brant, M.L. Harris, T.M. Bayless, G.
1071 Parmigiani, S. Chakravarti, Genome-wide gene expression differences in Crohn's
1072 disease and ulcerative colitis from endoscopic pinch biopsies: insights into distinctive
1073 pathogenesis, *Inflamm. Bowel. Dis.* 13 (2007) 807-821.
- 1074 [182] N.A. Wahab, L. Schaefer, B.S. Weston, O. Yiannikouris, A. Wright, A. Babelova, R.
1075 Schaefer, R.M. Mason, Glomerular expression of thrombospondin-1, transforming

- 1076 growth factor beta and connective tissue growth factor at different stages of diabetic
1077 nephropathy and their interdependent roles in mesangial response to diabetic stimuli,
1078 *Diabetologia* 48 (2005) 2650-2660.
- 1079 [183] F.M. Brennan, D. Chantry, M. Turner, B. Foxwell, R. Maini, M. Feldmann, Detection
1080 of transforming growth factor-beta in rheumatoid arthritis synovial tissue: lack of
1081 effect on spontaneous cytokine production in joint cell cultures, *Clin. Exp. Immunol.*
1082 81 (1990) 278-285.
- 1083 [184] J. Hao, H. Ju, S. Zhao, A. Junaid, T. Scammell-La Fleur, I.M. Dixon, Elevation of
1084 expression of Smads 2, 3, and 4, decorin and TGF-beta in the chronic phase of
1085 myocardial infarct scar healing, *J Mol. Cell Cardiol.* 31 (1999) 667-678.
- 1086 [185] J. Waltenberger, L. Lundin, K. Oberg, E. Wilander, K. Miyazono, C.H. Heldin, K.
1087 Funai, Involvement of transforming growth factor-beta in the formation of fibrotic
1088 lesions in carcinoid heart disease, *Am J Pathol* 142 (1993) 71-78.
- 1089 [186] S. Klahr, Mechanisms of progression of chronic renal damage, *J. Nephrol.* 12 Suppl 2
1090 (1999) S53-S62.
- 1091 [187] S. Schultze-Mosgau, F. Wehrhan, K. Amann, M. Radespiel-Troger, F. Rodel, G.G.
1092 Grabenbauer, In Vivo TGF-beta 3 expression during wound healing in irradiated
1093 tissue. An experimental study, *Strahlenther. Onkol.* 179 (2003) 410-416.
- 1094 [188] S. Schultze-Mosgau, F. Wehrhan, F. Rodel, K. Amann, M. Radespiel-Troger, G.G.
1095 Grabenbauer, Transforming growth factor-beta receptor-II up-regulation during wound
1096 healing in previously irradiated graft beds in vivo, *Wound Repair Regen.* 11 (2003)
1097 297-305.
- 1098
- 1099

Table 1. Expression of TGF- β related proteins in human disease compared to controls (normal or inflammatory)

Organ	Irradiated skin	Heart	Synovial membrane		Gut		Kidney
Disease	Wound healing	Fibrotic heart diseases*	RA	CD inflamed	CD fibrotic	UC	Diabetic nephropathy
Molecule							
THBS1	n.d.	↑[178,179]	↑[40,180]	↓[181]	n.d.	↔[181]	↑[182]
Latent-TGF- β	↑[122]	n.d.	↔[183], ↑[39]	n.d.	n.d.	n.d.	↑[182]
TGF- β 1	↑[122]	↑[184,185]	↔[35,36,40,183], ↑[37,38]	↑[76,108,114]	↑[108,114]	↑[76,108,114]	↔[186], ↑[53,72]
TGF- β 2	↑[122]	↔[185]	↔[35,183]	↑[114]	↑[114]	↔[114]	↑[72]
TGF- β 3	↓[187]	↑[185]	↔[35,183]	↓[114]	↓[114]	↔[114]	↔[186], ↑[72]
TGFBR1	n.d.	↓[184]	↑[40]	↔[113]	n.d.	n.d.	n.d.
TGFBR2	↑[188]	↔[184]	↑[35,183]	↑[113]	n.d.	n.d.	n.d.

↑ increased, ↓ decreased, ↔ no change, **n.d.** not determined; *mostly following myocardial infarction

FIGURE LEGENDS

Figure 1: TGF- β 1 induces different, but overlapping response in different organ systems.

Figure 2: Molecules of the TGF- β -pathway ranging from the release of TGF- β , binding of receptors, signaling through Smads, FAK and MAPK to its entry into the nucleus and the regulation of the transcriptional activity (adapted from [40]).




Figure 1




Figure 2