



Nucleon transfer in heavy ion collisions with the time-dependent Hartree-Fock theory using a particle number projection technique

Cédric Simenel

► To cite this version:

Cédric Simenel. Nucleon transfer in heavy ion collisions with the time-dependent Hartree-Fock theory using a particle number projection technique. 2010. hal-00510315v1

HAL Id: hal-00510315

<https://hal.science/hal-00510315v1>

Preprint submitted on 18 Aug 2010 (v1), last revised 6 Nov 2010 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nucleon transfer in heavy ion collisions with the time-dependent Hartree-Fock theory using a particle number projection technique

Cédric Simenel^{1, 2, *}

¹*Department of Nuclear Physics, Research School of Physics and Engineering,
Australian National University, Canberra, Australian Capital Territory 0200, Australia*

²*CEA, Centre de Saclay, IRFU/Service de Physique Nucléaire, F-91191 Gif-sur-Yvette, France.*

(Dated: August 18, 2010)

We study nucleon transfer in the $^{16}\text{O}+^{208}\text{Pb}$ reaction below the fusion barrier with the time-dependent Hartree-Fock (TDHF) mean-field theory. A particle-number projection technique is used to calculate the proton and neutron distributions of the fragments after the collision. The agreement with experimental data for the sum of the proton-transfer channels is good, considering that TDHF has no parameter adjusted on reaction mechanism. Some perspectives for extensions beyond TDHF to include cluster-transfers are discussed.

Binary collisions of many-body systems are of fundamental interest to test dynamical approaches of the quantum many-body problem. During the collision, the systems may retain their entities [(in)elastic scattering] or new ones may be produced, for instance, if they fuse or transfer some of their constituents. Examples of transfer reactions include electron transfer in ion or cluster collisions [1], and nucleon transfer in collisions of atomic nuclei [2]. The prediction of the outcome of such reactions is one of the main challenges of modern quantum many-body dynamics theories. In particular, the transfer products may be in a coherent superposition of fragments with different constituent numbers, and quantities such as transfer probabilities should be computed to allow comparison with experiments.

The coupled channel framework, where the relative motion of the collision partners is coupled to their internal degrees of freedom, is surely amongst the most popular approaches to study transfer reactions [3, 4]. It usually allows a detailed reproduction of experimental data, providing the fact that the structure of the collision partners (ground and excited states) as well as their interaction potential are well known. For numerical tractability, however, only few states are usually included in the one hand, and, in the other hand, all information on the structure of the reactants is not always available, as, e.g., for exotic nuclei. It is then important to develop other approaches with less parameters, to enhance their predictive power. In this spirit, recent works have pushed the envelope of describing binary collisions of many-body systems both quantum mechanically and microscopically, with no parameter adjusted on reaction mechanisms. For instance, the dynamics of the valence electrons in collisions of atoms, molecules, or atomic clusters, is usually given by the time-dependent density functional theory (TDDFT) (see, e.g., [5] and references therein). In nuclear physics, these approaches usually consider independent particles evolving in a mean-field as a starting point, as in the time-dependent Hartree-Fock (TDHF) theory [6, 7]. Although they have been mostly applied to fusion reactions, several recent attempts of describ-

ing nucleon transfer in heavy ion collisions within TDHF have been made [8–13].

Here, a step forward in TDHF studies of nucleon transfer in heavy-ion collisions is made by computing the probabilities for different transfer products in the exit channel. This is done using a particle number projection technique applied on the fragments of the total independent particle state. This technique is standard in beyond-mean-field models for nuclear structure when the number of particles is only given in average [14]. Note that this technique could be generalized to determine the particle number distribution in fragments of any many-body system, for instance, following electron transfer or ionization in atomic clusters, nuclear fission...

As a first application, we investigate sequential transfer of nucleons in $^{16}\text{O}+^{208}\text{Pb}$ collisions. In such a collision, nucleon transfer may occur when the projectile has enough energy to overcome the Coulomb repulsion and reach the vicinity of its collision partner, that is, at energies around and above the so-called fusion barrier. Here, we focus on sub-barrier central collisions and compare our calculations with experimental data. Transfer probabilities are expected to be sensitive to nucleon clusters. In particular, pair-transfers are interesting as they may be used to populate giant pairing vibrations and to study the Josephson effect in nuclei [3]. Various perspectives of this work in terms of beyond-TDHF improvements to treat properly correlations responsible for transfer of nucleon clusters are then discussed.

The time-dependent Hartree-Fock (TDHF) theory has been introduced by Dirac [18]. In nuclear physics, it is usually used with a Skyrme energy density functional (EDF) [19] to generate the nuclear mean-field [9, 20, 21]. The EDF is the only phenomenological ingredient which is adjusted only on few nuclear structure properties [22]. The same EDF is used to compute the initial Hartree-Fock ground state of the nuclei and the time evolution.

In TDHF, the state of the N particles is constrained to be an anti-symmetrized independent particle state (Slater determinant) at any time. The state vector reads




FIG. 1: (color online) Density evolution for the central collision of a ^{16}O (initially on the right side) with a ^{208}Pb (left) at $E_{c.m.} = 74.44$ MeV. The snapshots run from $t = 7.5$ to 37.5 zs by steps of 7.5 zs.

$|\phi\rangle = \prod_{i=1}^N \hat{a}_i^\dagger |-\rangle$ where \hat{a}_i^\dagger creates a particle in the state $|i\rangle$ when applied on the particle vacuum $|-\rangle$. The one-body density matrix of such a state reads $\rho(\mathbf{r}sq, \mathbf{r}'s'q') = \sum_{i=1}^N n_i \varphi_i^{sq}(\mathbf{r}) \varphi_i^{s'q'*}(\mathbf{r}')$, where $\varphi_i^{sq}(\mathbf{r}) = \langle \mathbf{r}sq | i \rangle$ is a single-particle wave function, \mathbf{r} , s and q denote the nucleon position, spin, and isospin, respectively, and $n_i = 1$ for occupied states and 0 otherwise. It evolves according to the TDHF equation $i\hbar \frac{\partial}{\partial t} \rho = [h[\rho], \rho]$. The single particle Hamiltonian $h[\rho]$ is related to the Skyrme EDF, noted $E[\rho]$, which depends on local densities [23] by $h[\rho](\mathbf{r}sq, \mathbf{r}'s'q') = \frac{\delta E[\rho]}{\delta \rho(\mathbf{r}'s'q', \mathbf{r}sq)}$.

Realistic TDHF calculations in 3 dimensions are now possible with modern Skyrme functionals including spin-orbit term [6, 7, 24, 25]. Here, the TDHF equation is solved iteratively in time using the TDHF3D code with the SLy4d parameterization of the Skyrme EDF [6]. A time step $\Delta t = 1.5 \times 10^{-24}$ s is used. The spatial grid has $N_x \times N_y \times N_z/2 = 84 \times 28 \times 14$ points with a plane of symmetry (the collision plane $z = 0$) and a lattice spacing $\Delta x = 0.8$ fm. The initial distance between the nuclei is 44.8 fm.

The density evolution of the $^{16}\text{O} + ^{208}\text{Pb}$ central collision at a center of mass energy $E_{c.m.} = 74.44$ MeV (just below the fusion barrier) plotted in Fig. 1 shows that the two nuclei form a di-nuclear system with a neck and then re-separate. There is *a priori* no reason that these two fragments conserve the same average neutron and proton numbers as in the entrance channel [11] (except for symmetric reactions). Indeed, between the touching and re-separation, nucleons can be exchanged. In TDHF calculations, this exchange is treated through the time-dependent distortion of single-particle wave-functions which can eventually be partially transferred from one partner to the other.

The following operator written in r-space counts the number of particles with isospin q in the right side of the

separation plane (defined arbitrarily as $x > 0$):

$$\hat{N}_R^q = \sum_s \int d\mathbf{r} \hat{a}^\dagger(\mathbf{r}sq) \hat{a}(\mathbf{r}sq) \mathcal{H}(x) \quad (1)$$

where $\mathcal{H}(x) = 1$ if $x > 0$ and 0 elsewhere, and $\hat{a}(\mathbf{r}sq) = \sum_i \varphi_i^{sq}(\mathbf{r}) \hat{a}_i$. Let us write

$$\langle i|j\rangle_R^q = \sum_s \int d\mathbf{r} \varphi_i^{sq*}(\mathbf{r}) \varphi_j^{sq}(\mathbf{r}) \mathcal{H}(x) \quad (2)$$

the overlap in the $x > 0$ region between two single-particle states with isospin q . Using $\langle \hat{a}_i^\dagger \hat{a}_j \rangle = n_i \delta_{ij}$, we obtain the average number of particles in the $x > 0$ region as $\langle \hat{N}_R^q \rangle = \sum_i \langle i|i \rangle_R^q n_i$. Applied to the average proton and neutron numbers of the small fragment after a central collision at $E_{c.m.} = 74.44$ MeV (see Fig. 1), we get ~ 6.1 protons and ~ 8.1 neutrons, respectively. This indicates that the proton transfer probability from the light to the heavy fragment is so high at the barrier that two protons, in average, have been sequentially transferred. Decreasing the energy induces a rapid convergence of the average proton and neutron numbers towards the ^{16}O ones. Indeed, $\langle \hat{N}_R^p \rangle \simeq \langle \hat{N}_R^n \rangle \simeq 8.0$ at $E_{c.m.} = 70$ MeV, i.e., 6% below the barrier.

Well below the barrier, where transfer is prohibited, the variance of \hat{N}_R is strictly zero: $\sigma_R = \sqrt{\langle \hat{N}_R^2 \rangle - \langle \hat{N}_R \rangle^2} = 0$. (Here and in the following, we omit the isospin q for simplicity.) This property is lost at higher energies where transfer occurs. Then, the system in the exit channel is not an eigenstate of \hat{N}_R , and each fragment is not described by an eigenstate of the particle number operator (e.g., a Slater determinant) anymore. Note that the upper limit of the variance obeys $\sigma_R^2 \leq \langle \hat{N}_R \rangle \left(1 - \frac{\langle \hat{N}_R \rangle}{N_t}\right)$ for a Slater determinant [26], where N_t is the total number of protons or neutrons. This is an intrinsic limitation of independent particle systems. In case of violent collisions such as deep-inelastic reactions, experimental variances may exceed this limit [26], and inclusion of correlations is then needed. However, for less violent collisions such as sub-barrier transfer reactions studied here, smaller experimental variances are expected, and a mean-field approach like TDHF might give reasonable estimates of the variances.

Let us calculate the variance σ_R after the reaction. Using anti-commutation relations for fermions and $\langle \hat{a}_i^\dagger \hat{a}_j^\dagger \hat{a}_k \hat{a}_l \rangle = n_i n_j (\delta_{il} \delta_{jk} - \delta_{ik} \delta_{jl})$ for a Slater determinant, we get [26] $\sigma_R^2 = \sum_{i=1}^N \langle i|i \rangle_R - \sum_{i,j=1}^N |\langle i|j \rangle_R|^2$. Applying this formula to the small fragment in the exit channel of the reaction at $E_{c.m.} = 74.44$ MeV shown in Fig. 1, we get $\sigma_R^p \simeq 0.5$ for protons and $\sigma_R^n \simeq 0.3$ for neutrons. At $E_{c.m.} = 70$ MeV, we get $\sigma_R^p \simeq \sigma_R^n \simeq 0.2$, showing that transfer occurs at this energy, although it does not change the average number of protons and neutrons as discussed before. These finite variances clearly indicate

that the many-body systems on each side of the separation plane are not eigenstates of the particle number operator anymore.

To get a deeper insight into these TDHF predictions of transfer, we go one step further by computing the transfer probabilities. It is possible to extract the component of the wave function associated to a specific transfer channel using a particle number projector onto N protons or neutrons in the $x > 0$ region. Such a projector is written $\hat{P}_R(N) = \frac{1}{2\pi} \int_0^{2\pi} d\theta e^{i\theta(\hat{N}_R - N)}$ [14]. It can be used to compute the probability to find N nucleons in $x > 0$ in the state $|\phi\rangle$,

$$|\hat{P}_R(N)|\phi\rangle|^2 = \frac{1}{2\pi} \int_0^{2\pi} d\theta e^{-i\theta N} \langle\phi|\phi_R(\theta)\rangle, \quad (3)$$

where $|\phi_R(\theta)\rangle = e^{i\theta\hat{N}_R}|\phi\rangle$. Note that $|\phi_R(\theta)\rangle$ is an independent particle state. The last term in Eq. (3) is then the determinant of the matrix of the single particle overlaps: $\langle\phi|\phi_R(\theta)\rangle = \det(F)$ with

$$F_{ij} = \sum_s \int d\mathbf{r} \varphi_i^{s*}(\mathbf{r}) \varphi_j^s(\mathbf{r}) e^{i\theta\mathcal{H}(x)} = \delta_{ij} + \langle i|j\rangle_R (e^{i\theta} - 1).$$

The integral in Eq. (3) is discretized using $\theta_n = 2\pi n/M$ with the integer $n = 1 \dots M$. Choosing $M = 300$ ensures convergence. The resulting probabilities are shown in Fig. 2 for central collisions at $E_{c.m.} = 74.44$ MeV and 65 MeV ($\sim 13\%$ below the barrier). At the barrier, the most probable channel is a two-proton transfer leading to a ^{14}C nucleus in the exit channel. At the lower energy, the transfer probabilities are typically one or several orders of magnitude lower than at the barrier and the (in)elastic channels are by far the dominant ones. Note that the probability for proton stripping (transfer from the light to the heavy nucleus) is higher than for proton pickup (transfer from the heavy to the light nucleus), while neutron pickup is more probable than neutron stripping.

In transfer experiments, one usually measures angular differential cross-sections for multi-nucleon transfer channels. It is numerically heavy and time consuming to compute such cross-sections. A standard alternative is to translate the experimental angular cross-sections at sub-barrier energies into transfer probabilities as a function of the distance of closest approach R_{min} between the collision partners assuming a Rutherford trajectory [28]: $R_{min} = Z_1 Z_2 e^2 [1 + \text{cosec}(\alpha/2)] / 2E_{c.m.}$ where α is the center of mass scattering angle, and $Z_{1,2}$ the proton numbers of the colliding nuclei. Experimental transfer probabilities can then be calculated from the ratio of sub-barrier transfer to Rutherford cross-sections [28] for a given distance of closest approach.

The evolutions of the main proton-transfer channels with the distance of closest approach predicted by TDHF are shown in Fig. 3 in solid, dashed and dotted lines for zero, one and two-proton stripping, respectively. In




FIG. 2: Neutron (circles) and proton (squares) number probability distributions of the lightest fragment in exit channel of a head-on $^{16}\text{O}+^{208}\text{Pb}$ collision at $E_{c.m.} = 74.44$ MeV (solid lines) and 65 MeV (dotted lines).

fact, the TDHF probability for two-proton transfer behaves roughly as the square of the one-proton transfer probability (if the latter is small compared to one), which is a signature for sequential transfer [2]. The two-proton transfer in TDHF is, then, much smaller than the one-proton one (except at the barrier, corresponding to $R_{min} \simeq 12.7$ fm, not shown in Fig. 3 but in Fig. 2).

Multi-proton transfer has been measured for $^{16}\text{O}+^{208}\text{Pb}$ at $E_{c.m.} = 74.3$ MeV by Videbæk *et al.* [27]. One and two-proton stripping has been observed at this energy, and no proton-pickup, in qualitative agreement with TDHF calculations. However, it is well known that the two-proton stripping in this reaction occurs mainly as a cluster transfer, i.e., as a pair or alpha-transfer [27, 29]. The treatment of such nucleon-clusters involves correlations beyond TDHF. As a consequence, TDHF is not expected to reproduce the ratio between the one and two-proton transfer probabilities, but only their sums which should be less affected by such cluster structures. The experimental sum of the one and two-proton transfer probabilities are shown in Fig. 3 (squares). They can be compared with the one-proton transfer in TDHF, as the two-proton sequential transfer is negligible in this energy range. The overall agreement is good, considering the fact that TDHF has no parameter adjusted on reaction mechanism.

To investigate the transfer of particle-cluster, one should consider theories beyond the standard independent particle approximation. Pair transfer could be investigated with the TDHF-Bogolyubov theory which accounts for pairing [30, 31], or with the time-dependent density-matrix theory which treats explicitly two-body correlations [32]. Stochastic techniques can also be used to account for zero point motion [13]. The Balian-




FIG. 3: Proton number probability distribution as function of the distance of closest approach obtained with TDHF (lines). Experimental data (squares) are adapted from Ref. [27] using $E_{c.m.} = 74.3$ MeV data and show the sum of the one and two-proton transfer channels.

Vénérioni variational approach also contains fluctuations beyond TDHF [35]. One limitation of TDHF is that all exit channels follow the same trajectory. Several TDHF trajectories with different external potentials "forcing" transfer could be used to build a more general state using the time-dependent Generator Coordinate Method [36]. Finally, investigations of the excitation energies of the transfer products should be studied with, e.g., the density-constrained TDHF approach [37].

The calculations have been performed on the Centre de Calcul Recherche et Technologie of the Commissariat à l'Énergie Atomique, France. The author is thankful to D. Lacroix and B. Avez for their collaboration on the subject. M. Dasgupta, M. Evers and D. J. Hinde are thanked for stimulating discussions and their continuous support during this work, and a careful reading of the paper. In particular, the author is grateful to M. Evers for his illuminating help on the extraction of experimental probabilities.

* cedric.simenel@cea.fr

- [1] E. E. B. Campbell and F. Rohmund, Rep. Prog. Phys. **63**, 1061 (2000).
- [2] L. Corradi, G. Pollaro, and S. Szilner, J. Phys. G **36**, 113101 (2009).
- [3] W. von Oertzen and A. Vitturi, Rep. Prog. Phys. **64**, 1247 (2001).
- [4] I. J. Thompson and F. M. Nunes, *Nuclear Reactions for Astrophysics: Principles, Calculation and Applications of Low-Energy Reactions* (Cambridge University Press, 2009), ISBN 0521856353.
- [5] Z. Wang, P. Dinh, P.-G. Reinhard, E. Suraud, G. Bruny, C. Montano, S. Feil, S. Eden, H. Abdoul-Carime, B. Fari-zon, et al., Int. J. Mass Spectrom. **285**, 143 (2009), ISSN 1387-3806.
- [6] K.-H. Kim, T. Otsuka, and P. Bonche, J. Phys. G **23**, 1267 (1997).
- [7] A. S. Umar and V. E. Oberacker, Phys. Rev. C **73**, 054607 (2006).
- [8] D. J. Kedziora and C. Simenel, Phys. Rev. C **81**, 044613 (2010).
- [9] C. Simenel, D. Lacroix, and B. Avez, *Quantum Many-Body Dynamics: Applications to Nuclear Reactions* (VDM Verlag, 2010).
- [10] C. Simenel and B. Avez, Int. J. Mod. Phys. E **17**, 31 (2008).
- [11] A. Umar, V. Oberacker, and J. Maruhn, Eur. Phys. J. A **37**, 245 (2008).
- [12] C. Golabek and C. Simenel, Phys. Rev. Lett. **103**, 042701 (2009).
- [13] K. Washiyama, S. Ayik, and D. Lacroix, Phys. Rev. C **80**, 031602 (2009).
- [14] M. Bender, P.-H. Heenen, and P.-G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).
- [15] M. Dasgupta, D. J. Hinde, N. Rowley, and A. M. Stefanini, Ann. Rev. Nucl. Part. Sci. **48**, 401 (1998).
- [16] A. Chatterjee, A. Navin, A. Shrivastava, S. Bhattacharyya, M. Rejmund, N. Keeley, V. Nanal, J. Nyberg, R. G. Pillay, K. Ramachandran, et al., Phys. Rev. Lett. **101**, 032701 (2008).
- [17] A. Astier, P. Petkov, M.-G. Porquet, D. S. Delion, and P. Schuck, Phys. Rev. Lett. **104**, 042701 (2010).
- [18] P. A. M. Dirac, Proc. Camb. Phil. Soc. **26**, 376 (1930).
- [19] T. Skyrme, Phil. Mag. **1**, 1043 (1956).
- [20] P. Bonche, S. Koonin, and J. W. Negele, Phys. Rev. C **13**, 1226 (1976).
- [21] J. W. Negele, Rev. Mod. Phys. **54**, 913 (1982).
- [22] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, Nuclear Physics A **635**, 231 (1998).
- [23] Y. M. Engel, D. M. Brink, K. Goeke, S. J. Krieger, and D. Vautherin, Nuc. Phys. A **249**, 215 (1975).
- [24] J. A. Maruhn, P.-G. Reinhard, P. D. Stevenson, and M. R. Strayer, Phys. Rev. C **74**, 027601 (2006).
- [25] T. Nakatsukasa and K. Yabana, Phys. Rev. C **71**, 024301 (2005).
- [26] C. H. Dasso, T. Døssing, and H. C. Pauli, Z. Phys. A **289**, 395 (1979).
- [27] F. Videbæk, R. B. Goldstein, L. Grodzins, S. G. Steadman, T. A. Belote, and J. D. Garrett, Phys. Rev. C **15**, 954 (1977).
- [28] R. A. Broglia and A. Winther, *Heavy Ion Reactions: Lecture Notes the Elementary Processes (Frontiers in Physics)* (Addison Wesley Publishing Company, 1991), ISBN 0201513927.
- [29] I. J. Thompson, M. A. Nagarajan, J. S. Lilley, and M. J. Smithson, Nucl. Phys. A **505**, 84 (1989), ISSN 0375-9474.
- [30] B. Avez, C. Simenel, and P. Chomaz, Phys. Rev. C **78**, 044318 (2008).
- [31] S. Ebata, T. Nakatsukasa, T. Inakura, K. Yoshida, Y. Hashimoto, and K. Yabana (2010), arXiv:1007.0785 [nucl-th].
- [32] M. Assié and D. Lacroix, Phys. Rev. Lett. **102**, 202501 (2009).
- [33] M. Tohyama and A. S. Umar, Phys. Rev. C **65**, 037601 (2002).
- [34] S. Ayik, K. Washiyama, and D. Lacroix, Phys. Rev. C

- 79**, 054606 (2009).
- [35] J. M. A. Broomfield and P. D. Stevenson, J. Phys. G **35**, 095102 (2008).
- [36] H. Goutte, J. F. Berger, P. Casoli, and D. Gogny, Phys. Rev. C **71**, 024316 (2005).
- [37] A. S. Umar, V. E. Oberacker, J. A. Maruhn, and P. G. Reinhard, Phys. Rev. C **80**, 041601 (2009).