



HAL
open science

Lipschitz stability in an inverse problem for the Kuramoto-Sivashinsky equation

Lucie Baudouin, Eduardo Cerpa, Emmanuelle Crépeau, Alberto Mercado

► **To cite this version:**

Lucie Baudouin, Eduardo Cerpa, Emmanuelle Crépeau, Alberto Mercado. Lipschitz stability in an inverse problem for the Kuramoto-Sivashinsky equation. *Applicable Analysis*, 2013, 92 (10), pp. 2084-2102. hal-00510506v3

HAL Id: hal-00510506

<https://hal.science/hal-00510506v3>

Submitted on 19 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lipschitz stability in an inverse problem for the Kuramoto-Sivashinsky equation

Lucie Baudouin*, Eduardo Cerpa†, Emmanuelle Crépeau‡ and Alberto Mercado†

Abstract

In this article, we present an inverse problem for the nonlinear 1-d Kuramoto-Sivashinsky (K-S) equation. More precisely, we study the nonlinear inverse problem of retrieving the anti-diffusion coefficient from the measurements of the solution on a part of the boundary and also at some positive time in the whole space domain. The Lipschitz stability for this inverse problem is our main result and it relies on the Bukhgeim-Klibanov method. The proof is indeed based on a global Carleman estimate for the linearized K-S equation.

Keywords: Inverse problem, Kuramoto-Sivashinsky equation, Carleman estimate

AMS subject classifications: 35R30, 35K55.

1 Introduction

We focus in this paper on an inverse problem that consists in the determination of a coefficient in a partial differential equation (pde) from the partial knowledge of a given single solution of the equation. This class of problems (single-measurement coefficient inverse problems) was investigated using Carleman estimates for the first time in [7] by Bukhgeim and Klibanov. See [20], [21] and the recent book [3] for details about the so-called Bukhgeim-Klibanov method. This method was initially used to prove uniqueness for inverse problems (i.e. that each measurement corresponds to only one coefficient) from local Carleman estimates (estimates valid for solutions with compact support in the interior of the domain), as in [7]. Regarding the continuity of the inverse problem of recovering the source term, the first Lipschitz stability result for a multidimensional wave equation was obtained by Puel and Yamamoto [25] using the uniqueness result and a compactness-uniqueness argument.

*CNRS, LAAS, 7 avenue du colonel Roche, F-31400 Toulouse, France.

Univ de Toulouse, LAAS, F-31400 Toulouse, France.

E-mail: lucie.baudouin@laas.fr

†Departamento de Matemática, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile.

E-mail: eduardo.cerpa@usm.cl, alberto.mercado@usm.cl

‡Laboratoire de Mathématiques, Université de Versailles Saint-Quentin en Yvelines, 78035 Versailles, France.

E-mail: emmanuelle.crepeau@math.uvsq.fr

Global Carleman estimates (valid for solutions considered in the whole domain and satisfying boundary conditions) were applied to parabolic equations by first time in [18], where Lipschitz stability of an inverse problem is established. Since then, this type of inverse problems for parabolic equations has received a large amount of attention. The primary difference with respect to hyperbolic inverse problems is that parabolic problems are not time-reversible: therefore, an additional measurement must be added if that method is applied. As one can read in the discussion of the introduction of [18], the knowledge of the full-state of the solution for some positive time is required. Proving the Lipschitz stability without this assumption, which is usually needed when global Carleman inequalities are used, is still an open problem. Nevertheless, there are some uniqueness results with less assumptions on the measurements, that can be found in the literature, such as [27] or some other inversion method in [20, 22].

Recent results regarding linear parabolic problems can be found in [5] (discontinuous coefficient), [13] (systems), [19] (network) and the references therein. In [6, 15, 27], nonlinear parabolic equations were even considered.

Among others pde's coefficient/source inverse problems where Carleman estimates have been used we can mention, without being exhaustive, logarithmic stability [4], Calderón problem [14] or Schrödinger equation [2].

In this paper, we consider a 1D nonlinear fourth-order parabolic equation called Kuramoto-Sivashinsky (K-S) equation. This equation was proposed independently by Kuramoto and Tsuzuki [23] as a model for the phase turbulence in reaction diffusion systems, and by Sivashinsky [28], as a model the physical phenomena of plane flame propagation, where the combined influence of diffusion and thermal conduction of a gas is described.

The K-S equation with non-constant coefficients describing the diffusion $\sigma = \sigma(x)$, and the anti-diffusion $\gamma = \gamma(x)$, is given as

$$\left\{ \begin{array}{ll} y_t + (\sigma(x)y_{xx})_{xx} + \gamma(x)y_{xx} + yy_x = g, & \forall (t, x) \in Q, \\ y(t, 0) = h_1(t), \quad y(t, 1) = h_2(t), & \forall t \in (0, T), \\ y_x(t, 0) = h_3(t), \quad y_x(t, 1) = h_4(t), & \forall t \in (0, T), \\ y(0, x) = y_0(x), & \forall x \in (0, 1), \end{array} \right. \quad (1)$$

where $Q := (0, T) \times (0, 1)$, $\sigma : [0, 1] \rightarrow \mathbb{R}_+^*$, and the functions y_0, g, h_j are the initial condition, the source term and the boundary data respectively. All these terms are assumed to be known and compatible.

In this nonlinear pde, the fourth-order term models the diffusion, and the second-order term models the incipient instabilities. We consider the inverse problem of retrieving the anti-diffusion coefficient γ from boundary measurements of the solution. This corresponds for instance to getting information on the instability of a reaction-diffusion media by measuring a single solution, which could represent a flame propagating on the domain. Concerning the boundary measurements we will make, it is worth to mention that in a fourth-order

parabolic problem like KS, boundary data u_{xx} and u_{xxx} are referred to as Neumann data, which in fact represent heat flux [24] in this kind of models.

To the knowledge of the authors there are no results in the literature concerning the determination of coefficients for this nonlinear equation. However, a Carleman estimate has been used to obtain the null-controllability of the K-S equation in reference [10] for the constant coefficient case. Other results on the control of the KS equation can be found in [17, 1, 12, 24, 9].

Since the linearized equation is parabolic, we know that boundary measurements will not be sufficient to prove stability and we must consider an additional measurement of the full solution for a given time T_0 (as in [5, 18] among others).

Our first result involves the local well-posedness of the nonlinear equation (1). A less regular framework can be used for this equation but the method applied in this paper requires the solution and its time-derivative to be at least in $L^2(0, T; H^4(0, 1))$. Therefore, let us introduce the following notations for the functional spaces appearing in this paper:

$$\begin{aligned}\mathcal{Y}_k &:= C([0, T]; H^k(0, 1)) \cap L^2(0, T; H^{k+2}(0, 1)), \quad \text{for } k \in \mathbb{N}; \\ \mathcal{F} &:= \{f \in L^2(0, T; H^4(0, 1)) / f_t \in L^2(0, T; L^2(0, 1))\}; \\ \mathcal{Z} &:= \{z \in \mathcal{Y}_6 / z_t \in \mathcal{Y}_2\}.\end{aligned}\tag{2}$$

Theorem 1.1 *Let $\gamma \in H^4(0, 1)$ and $\sigma \in H^4(0, 1)$ be such that*

$$\sigma(x) \geq \sigma_0 > 0, \quad \forall x \in (0, 1).\tag{3}$$

There exists $\varepsilon > 0$ such that if $y_0 \in H^6(0, 1)$, $g \in \mathcal{F}$, and $h_j \in H^2(0, T)$ for $j = 1, \dots, 4$ satisfy the compatibility conditions

$$y_0(0) = h_1(0), \quad y_{0,x}(0) = h_3(0), \quad y_0(1) = h_2(0), \quad y_{0,x}(1) = h_4(0),\tag{4}$$

and

$$\|y_0\|_{H^6(0,1)} \leq \varepsilon, \quad \|g\|_{\mathcal{F}} \leq \varepsilon, \quad \|h_j\|_{H^2(0,T)} \leq \varepsilon \quad \text{for } j = 1, \dots, 4,\tag{5}$$

then the K-S equation (1) has a unique solution $y \in \mathcal{Z}$.

Once the existence of solutions to the K-S equation has been established (see Section 2), the following inverse problem is addressed:

Is it possible to retrieve the anti-diffusion coefficient $\gamma = \gamma(x)$ from the measurement of $y_{xx}(t, 0)$ and $y_{xxx}(t, 0)$ on $(0, T)$ and from the measurement of $y(T_0, x)$ on $(0, 1)$, where y is the solution to Equation (1) and $T_0 \in (0, T)$?

A local answer for this nonlinear inverse problem is given (see section 4). To be more specific, let $\tilde{\gamma}$ be given and fixed. We denote by \tilde{y} the solution to Equation (1) with γ replaced by $\tilde{\gamma}$. This paper focuses on the following question concerning the unknown γ and y .

Stability: Is it possible to estimate $\|\tilde{\gamma} - \gamma\|_{L^2(0,1)}$ by suitable norms $\|\tilde{y}(T_0, x) - y(T_0, x)\|$ in space and $\|\tilde{y}_{xx}(t, 0) - y_{xx}(t, 0)\|$, $\|\tilde{y}_{xxx}(t, 0) - y_{xxx}(t, 0)\|$ in time?

Of course, a positive answer implies the usual uniqueness result.

Uniqueness: Do the equalities of the measurements $\tilde{y}_{xx}(t, 0) = y_{xx}(t, 0)$ and $\tilde{y}_{xxx}(t, 0) = y_{xxx}(t, 0)$ for $t \in (0, T)$ and $\tilde{y}(T_0, x) = y(T_0, x)$ for $x \in (0, 1)$ imply $\tilde{\gamma} = \gamma$ on $(0, 1)$?

In order to answer these questions, we use the Bukhgeim-Klibanov method. First, a global Carleman estimate for the linearized K-S equation with non-constant coefficients is obtained. It is then used to prove the main result which can be stated as follows.

To precisely state the results we prove in this article, we introduce, for $m > 0$, the set

$$L_{\leq m}^{\infty}(0, 1) = \{\gamma \in L^{\infty}(0, 1) \text{ s.t. } \|\gamma\|_{L^{\infty}(0,1)} \leq m\}.$$

Theorem 1.2 *Let us consider $\sigma \in H^4(0, 1)$ satisfying (3), $\gamma \in H^4(0, 1)$, $g \in \mathcal{F}$ and the data $y_0 \in H^6(0, 1)$ and $h_j \in H^2(0, T)$ for $j = 1, \dots, 4$ under the compatibility conditions (4). Let $y \in \mathcal{Z}$ be the solution of (1), and $\tilde{y} \in \mathcal{Z}$ the solution corresponding to a given $\tilde{\gamma} \in H^4(0, 1)$ instead of γ . We assume that there exists $\eta > 0$ and $T_0 \in (0, T)$ such that*

$$\inf \{|\tilde{y}_{xx}(T_0, x)|, x \in (0, 1)\} \geq \eta, \quad (6)$$

Then, given $M > 0$, there exists a positive constant C depending on the parameters (T, m, M, η) , such that for every $\gamma \in L_{\leq m}^{\infty}(0, 1)$,

$$\begin{aligned} \|\gamma - \tilde{\gamma}\|_{L^2(0,1)}^2 &\leq C \|y_{xx}(\cdot, 0) - \tilde{y}_{xx}(\cdot, 0)\|_{H^1(0,T)}^2 + C \|y_{xxx}(\cdot, 0) - \tilde{y}_{xxx}(\cdot, 0)\|_{H^1(0,T)}^2 \\ &\quad + C \|y(T_0, \cdot) - \tilde{y}(T_0, \cdot)\|_{H^4(0,1)}^2 + C \|y(T_0, \cdot) - \tilde{y}(T_0, \cdot)\|_{H^1(0,1)}^4 \end{aligned} \quad (7)$$

for all y satisfying

$$\|y\|_{\mathcal{Z}} \leq M.$$

This inequality states the stability of the inverse problem. Before giving the outline of our paper and the proofs of the different steps, we want to give several comments on this result.

Remark 1.3 *For numerical purposes it would be interesting to know explicitly how the constant C in (7) depends on the diffusion σ or on the time T . This kind of question has been addressed in [11, 16] for observability constant in the framework of second-order parabolic equations. In those papers the authors got an exponential dependence on both the constant diffusion and the time.*

Remark 1.4 *One can show that there exist solutions satisfying assumption (6). We present two different arguments:*

1. *We take $\varepsilon > 0$ given by Theorem 1.1, and some $y^0 \in H^6(0, 1)$ such that $\inf_{x \in (0,1)} |y_{xx}^0| \geq \varepsilon/2$. For arbitrary boundary data and source term belonging to the corresponding spaces, by Theorem 1.1 there exists a solution $\tilde{y} \in C([0, 1]; H^6(0, 1))$ with $\tilde{y}(0, \cdot) = y^0$. Using*

Sobolev injection and continuity, we obtain the existence of a time $T_0 > 0$ such that (6) is fulfilled with $\eta = \varepsilon/4$.

2. We can also prove that there exist solutions satisfying (6) without asking T_0 to be small, but instead, constraining the source term and boundary data as follows: Let y_0 be the initial data and T_0 belong to $(0, T)$. Let us pick up a state $y_1 = y_1(x)$ strictly convex. We consider the trajectory $\tilde{y}(t, x) = \frac{T_0-t}{T_0}y_0(x) + \frac{t}{T_0}y_1(x)$, which is the solution of equation (1) with source term given by $g = \tilde{y}_t + (\sigma(x)\tilde{y}_{xx})_{xx} + \gamma(x)\tilde{y}_{xx} + \tilde{y}\tilde{y}_x$ and the boundary data given by the traces of \tilde{y} . Thus, $\tilde{y}(T_0, x) = y_1(x)$ and hence the trajectory \tilde{y} satisfies (6).

Therefore, the set of data and solutions where our stability result is valid is not empty.

Remark 1.5 We obtain the same result if \tilde{y} has a different initial condition than y . See in Section 4, that the term $v(x, 0)$ of system (46) does not play any role in the result.

Remark 1.6 We can complete inequality (7) by the following:

$$\begin{aligned} & \|y_{xx}(\cdot, 0) - \tilde{y}_{xx}(\cdot, 0)\|_{H^1(0, T)}^2 + \|y_{xxx}(\cdot, 0) - \tilde{y}_{xxx}(\cdot, 0)\|_{H^1(0, T)}^2 \\ & \quad + \|y(T_0, \cdot) - \tilde{y}(T_0, \cdot)\|_{H^4(0, 1)}^2 + \|y(T_0, \cdot) - \tilde{y}(T_0, \cdot)\|_{H^1(0, 1)}^4 \\ & \leq C \left(\|y - \tilde{y}\|_{H^1(0, T; H^4(0, 1))}^2 + \|y - \tilde{y}\|_{C([0, T]; H^1(0, 1))}^4 \right). \end{aligned}$$

This inequality follows directly from standard Sobolev injections. It indicates that the required measurements are finite if y and \tilde{y} belong to the space $H^1(0, T; H^4(0, 1))$ and this is true if y and \tilde{y} are solutions in \mathcal{Z} provided by Theorem 1.1.

Remark 1.7 As stated in the introduction, an internal measurement at $t = T_0$ is required if this method, using Carleman estimates, is used to prove the stability for this type of inverse problem for parabolic equations. Nevertheless, this is probably a technical point since there is no counter-example that demonstrates whether this assumption is required for stability. In [27], uniqueness (but not stability) is proven using a very different technique in an inverse problem for a parabolic equation and without any internal measurements in the whole space domain. One can also mention a method in [22] that can deliver uniqueness from hyperbolic equations to parabolic ones.

Remark 1.8 In this paper, the boundary measurements are located at $x = 0$, but the result would be the same if we measure at $x = 1$ instead. Indeed, the choice of a suitable weight function in the proof of the Carleman estimate in Section 3 is critical to impose the side of measurement.

This article is organized as follows. The well-posedness result stated in Theorem 1.1 is proved in Section 2. A global Carleman estimate for a general K-S equation is given and proved in Section 3. Finally, Section 4 contains the use of the Bukhgeim-Klibanov method to prove the Lipschitz stability of the inverse problem stated in Theorem 1.2.

2 On the Cauchy problem for KS equation

This section presents a proof of Theorem 1.1 in a more general case including time dependent lower-order coefficients. We consider the following K-S system

$$\begin{cases} y_t + (\sigma(x)y_{xx})_{xx} + \gamma(x)y_{xx} + G_1y_x + G_2y + yy_x = g, & \forall (t, x) \in Q, \\ y(t, 0) = h_1(t), \quad y(t, 1) = h_2(t), & \forall t \in (0, T), \\ y_x(t, 0) = h_3(t), \quad y_x(t, 1) = h_4(t), & \forall t \in (0, T), \\ y(0, x) = y_0(x), & \forall x \in (0, 1), \end{cases} \quad (8)$$

where G_1, G_2 belong to $H^1(0, T; H^4(0, 1))$, $g \in \mathcal{F}$ and $y_0 \in H^6(0, 1)$ is compatible with $h_j \in H^2(0, T)$ for $j = 1, \dots, 4$. Recall that the coefficients satisfy $\gamma \in H^4(0, 1)$, $\sigma \in H^4(0, 1)$ and hypothesis (3).

First, we only consider the main part of the linear differential operator in the next proposition.

Proposition 2.1 *Let $z_0 \in H^6 \cap H_0^2(0, 1)$ and $f \in \mathcal{F}$. Then, the following equation*

$$\begin{cases} z_t + (\sigma(x)z_{xx})_{xx} = f, & \forall (t, x) \in Q, \\ z(t, 0) = 0, \quad z(t, 1) = 0, & \forall t \in (0, T), \\ z_x(t, 0) = 0, \quad z_x(t, 1) = 0, & \forall t \in (0, T), \\ z(0, x) = z_0(x), & \forall x \in (0, 1), \end{cases} \quad (9)$$

has a unique solution $z \in \mathcal{Z}$ and there exists $C > 0$ such that

$$\|z\|_{\mathcal{Z}} \leq C (\|f\|_{\mathcal{F}} + \|z_0\|_{H^6}).$$

Proof. The operator

$$\begin{aligned} H^4 \cap H_0^2(0, 1) \subset L^2(0, 1) &\longrightarrow L^2(0, 1) \\ z &\longmapsto (\sigma(x)z''(x))'', \end{aligned}$$

is simultaneously positive, coercive and self-adjoint. By the Hille-Yosida-Phillips Theorem (see [8]), it generates a strongly continuous semigroup in $L^2(0, 1)$. Therefore, for each $z_0 \in H^4 \cap H_0^2(0, 1)$ and $f \in C^1([0, T]; L^2(0, 1))$, Equation (9) has a unique solution $z \in C([0, T]; H^4 \cap H_0^2(0, 1)) \cap C^1([0, T]; L^2(0, 1))$.

We will demonstrate that the solutions $z \in \mathcal{Z}$ (refer to the notation introduced in (2)), can be obtained by taking z_0 and f sufficiently regular.

We now search for some energy estimates that indicate the space where the solutions lie on depending on the regularity of the data. Suppose that there are solutions sufficiently regular to perform the following computations. Equation (9) is multiplied by z and integrated over $(0, 1)$ in space. Some integrations by parts give

$$\frac{d}{dt} \left(\int_0^1 |z(t, x)|^2 dx \right) + \int_0^1 |z_{xx}(t, x)|^2 dx \leq C \left(\int_0^1 |f(t, x)|^2 dx + \int_0^1 |z(t, x)|^2 dx \right). \quad (10)$$

Throughout this paper, C denotes a positive constant that may vary from line to line. To make the reading easier, we denote for any function u of x and t ,

$$\iint_Q u = \int_0^T \int_0^1 u(t, x) dx dt$$

Using Gronwall's lemma, we first obtain that for all $t > 0$,

$$\int_0^1 |z(t, x)|^2 dx \leq C \left(\iint_Q |f|^2 + \int_0^1 |z_0|^2 dx \right). \quad (11)$$

Then, (10) is integrated over $[0, T]$ and (11) is used to get

$$\iint_Q |z_{xx}|^2 \leq C \left(\iint_Q |f|^2 + \int_0^1 |z_0|^2 dx \right). \quad (12)$$

Inequalities (11) and (12) finally imply that

$$\|z\|_{\mathcal{Y}_0}^2 \leq C \iint_Q |f|^2 + C \int_0^1 |z_0|^2 dx. \quad (13)$$

Now, equation (9) is multiplied by $(\sigma z_{xx})_{xx}$ and integrated over $(0, 1)$ in space. Some integrations by parts give also

$$\frac{1}{2} \frac{d}{dt} \left(\int_0^1 \sigma |z_{xx}(t, x)|^2 dx \right) + \int_0^1 |(\sigma z_{xx}(t, x))_{xx}|^2 dx = \int_0^1 f(t, x) (\sigma z_{xx}(t, x))_{xx} dx.$$

Using the inequality $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$, we get

$$\frac{d}{dt} \left(\int_0^1 \sigma |z_{xx}(t, x)|^2 dx \right) + \int_0^1 |(\sigma z_{xx}(t, x))_{xx}|^2 dx \leq \int_0^1 |f(t, x)|^2 dx. \quad (14)$$

Using Gronwall's lemma, from(14) and (3) we obtain that for all $t > 0$,

$$\int_0^1 |z_{xx}(t, x)|^2 dx \leq C \left(\iint_Q |f|^2 + \int_0^1 |z_0''|^2 dx \right). \quad (15)$$

Then, (14) is integrated over $[0, T]$ and (15) is used to get

$$\iint_Q |(\sigma z_{xx})_{xx}|^2 \leq C \left(\iint_Q |f|^2 + \int_0^1 |z_0''|^2 dx \right), \quad (16)$$

and then, taking into account that $\sigma \in H^4$, we get

$$\iint_Q |z_{xxxx}|^2 \leq C \left(\iint_Q |f|^2 + \int_0^1 |z_0''|^2 dx \right) + C \|z\|_{L^2(0, T; H^3(0, 1))}. \quad (17)$$

For any $\varepsilon > 0$, from Ehrling's Lemma (see Theorem 7.30 in [26]) and (11), we have that

$$\begin{aligned} \|z\|_{L^2(0, T; H^3(0, 1))} &\leq \varepsilon \|z\|_{L^2(0, T; H^4(0, 1))} + C \|z\|_{L^2(0, T; L^2(0, 1))} \\ &\leq \varepsilon \|z\|_{L^2(0, T; H^4(0, 1))} + C \left(\iint_Q |f|^2 + \int_0^1 |z_0|^2 dx \right). \end{aligned} \quad (18)$$

Taking $\varepsilon > 0$ small enough, inequalities (15), (17) and (18) imply that

$$\|z\|_{\mathcal{Y}_2}^2 \leq C \iint_Q |f|^2 + C \|z_0\|_{L^2(0, T; H^2(0, 1))}^2. \quad (19)$$

On the other hand, Equation (9) is derived with respect to time. Thus $q := z_t$ satisfies

$$\begin{cases} q_t + (\sigma(x)q_{xx})_{xx} = f_t, & \forall (t, x) \in Q, \\ q(t, 0) = 0, \quad q(t, 1) = 0, & \forall t \in (0, T), \\ q_x(t, 0) = 0, \quad q_x(t, 1) = 0, & \forall t \in (0, T), \\ q(0, x) = f(0, x) - (\sigma z_0''(x))'', & \forall x \in (0, 1). \end{cases} \quad (20)$$

Using estimate (19), we obtain $q \in \mathcal{Y}_2$ if $(f(0, x) - (\sigma z_0''(x))'') \in H^2(0, 1)$ and $f_t \in L^2(0, T; L^2(0, 1))$. These hypotheses are fulfilled if $z_0 \in H^6 \cap H_0^2(0, 1)$ and $f \in \mathcal{F}$. Note that $\mathcal{F} \subset C([0, T]; H^2(0, 1))$. From the equation satisfied by z and the fact that $f \in \mathcal{F}$ and $z_t \in \mathcal{Y}_2$, we determine that $z \in \mathcal{Y}_6$, which concludes the proof of Proposition 2.1. \square

Then, we focus on the linear problem with non-homogenous boundary conditions and low-order coefficients that depend on time.

Proposition 2.2 *Let $z_0 \in H^6(0, 1)$, $\hat{f} \in \mathcal{F}$, $G_1, G_2 \in H^1(0, T; H^4(0, 1))$ and $h_j \in H^2(0, T)$ for $j = 1, \dots, 4$ satisfying the compatibility conditions with z_0 . Then, the equation*

$$\begin{cases} z_t + (\sigma(x)z_{xx})_{xx} + \gamma(x)z_{xx} + G_1z_x + G_2z = \hat{f}, & \forall (t, x) \in \mathcal{Q}, \\ z(t, 0) = h_1(t), \quad z(t, 1) = h_2(t), & \forall t \in (0, T), \\ z_x(t, 0) = h_3(t), \quad z_x(t, 1) = h_4(t), & \forall t \in (0, T), \\ z(0, x) = z_0(x), & \forall x \in (0, 1), \end{cases} \quad (21)$$

has a unique solution $z \in \mathcal{Z}$ and there exists $C > 0$ such that

$$\|z\|_{\mathcal{Z}} \leq C \left(\|\hat{f}\|_{\mathcal{F}} + \|z_0\|_{H^6} + \sum_{j=1}^4 \|h_j\|_{H^2} \right).$$

Proof. We first prove this result for null boundary data (i.e. for $h_j = 0$ for $j = 1, \dots, 4$ and therefore $z_0 \in H^6 \cap H_0^2(0, 1)$).

For any $\hat{w} \in \mathcal{Z}$, $\Pi(\hat{w})$ is defined as the solution of (9) with $f = (\hat{f} - \gamma(x)\hat{w}_{xx} - G_1\hat{w}_x - G_2\hat{w})$. Note that $f \in \mathcal{F}$ and therefore $\Pi(\hat{w}) \in \mathcal{Z}$ is well defined.

If T is small enough, then Π is a contraction. Indeed, for any $w, \hat{w} \in \mathcal{Z}$, we have

$$\begin{aligned} \|\Pi(\hat{w}) - \Pi(w)\|_{\mathcal{Z}} &\leq C \|\gamma(x)(w_{xx} - \hat{w}_{xx}) + G_1(w_x - \hat{w}_x) + G_2(w - \hat{w})\|_{\mathcal{F}} \\ &\leq C \|w - \hat{w}\|_{L^2(H^6)} + C \|w_t - \hat{w}_t\|_{L^2(H^2)} \end{aligned} \quad (22)$$

$$\begin{aligned} &\leq CT^{\frac{1}{4}} \|w - \hat{w}\|_{L^4(H^6)} + CT^{\frac{1}{4}} \|w_t - \hat{w}_t\|_{L^4(H^2)} \\ &\leq CT^{\frac{1}{4}} \|w - \hat{w}\|_{\mathcal{Y}_6} + CT^{\frac{1}{4}} \|w_t - \hat{w}_t\|_{\mathcal{Y}_2} \\ &\leq CT^{\frac{1}{4}} \|w - \hat{w}\|_{\mathcal{Z}}, \end{aligned} \quad (23)$$

where the space $L^m(0, T; H^n(0, 1))$ is denoted as $L^m(H^n)$.

Hence, the operator Π has a unique fixed point in \mathcal{Z} , which is the solution of (21) with $h_j = 0$ for $j = 1, \dots, 4$. Using standard arguments and the linearity of this equation, the solution can be extended to a larger time interval.

In order to prove the general case, take $h_j \in H^2(0, T)$, $j = 1, \dots, 4$ compatible with z_0 . It is not difficult to find a function $\psi \in H^2(0, T; C^\infty([0, 1]))$ satisfying the boundary conditions of (21). For instance take $\psi(x, t) = \sum_{j=1}^4 p_j(x)h_j(t)$ where $p_1(x) = 2x^3 - 3x^2 + 1$, $p_2(x) = -2x^3 + 3x^2$, $p_3(x) = x^3 - 2x^2 + x$ and $p_4(x) = x^3 - x^2$. In particular we have $L\psi := \psi_t + (\sigma(x)\psi_{xx})_{xx} + \gamma(x)\psi_{xx} + G_1\psi_x + G_2\psi \in \mathcal{F}$. Then, if w is the solution of equation (21) with null boundary data, initial condition $w_0 - \psi(\cdot, 0)$, and right-hand side equal to $\hat{f} - L\psi$, let us define $z = w + \psi$. It is not difficult to see that z is the required solution. \square

Remark 2.3 The third-order term z_{xxx} can be added to Equation (21). Indeed, in that case (22) becomes $C\|w - \hat{w}\|_{L^2(H^7)} + C\|w_t - \hat{w}_t\|_{L^2(H^3)}$, which is bounded by

$$CT^{\frac{1}{4}}\|w - \hat{w}\|_{L^\infty(H^6)}^{1/2}\|w - \hat{w}\|_{L^2(H^8)}^{1/2} + CT^{\frac{1}{4}}\|w_t - \hat{w}_t\|_{L^\infty(H^2)}^{1/2}\|w_t - \hat{w}_t\|_{L^2(H^4)}^{1/2}.$$

This last expression is bounded by (23). The remainder of the proof is the same.

Again, by using a fixed point theorem, we can prove Theorem 1.1 for equation (8). Let $y_0 \in H^6(0, 1)$, $h_j \in H^2(0, 1)$ compatible with y_0 , and $g \in \mathcal{F}$. For any $v \in \mathcal{Z}$, we define $\Lambda(v)$ as the solution of (21) with $\hat{f} = (g - vv_x)$ and $z_0 = y_0$. Note that $\hat{f} \in \mathcal{F}$ and therefore $\Lambda(v) \in \mathcal{Z}$ is well defined. Indeed, if $v \in \mathcal{Y}_3$ and $v_t \in \mathcal{Y}_0$, then we have

$$(vv_x)_{xxxx} = (10v_{xx}v_{xxx} + 5v_xv_{xxxx} + vv_{xxxx}) \in L^2(0, T; L^2(0, 1))$$

and

$$(vv_x)_t = v_tv_x + vv_{xt} \in L^2(0, T; L^2(0, 1)).$$

Furthermore, we can prove

$$\begin{aligned} \|\Lambda(v)\|_{\mathcal{Z}} &\leq C \left(\|g\|_{\mathcal{F}} + \|vv_x\|_{\mathcal{F}} + \|y_0\|_{H^6} + \sum_{j=1}^4 \|h_j\|_{H^2} \right) \\ &\leq C \left(\|g\|_{\mathcal{F}} + \|v\|_{\mathcal{Z}}^2 + \|y_0\|_{H^6} + \sum_{j=1}^4 \|h_j\|_{H^2} \right). \end{aligned} \quad (24)$$

Let $\varepsilon > 0$ and suppose that y_0 , h_j and g satisfy (5). Consider v such that $\|v\|_{\mathcal{Z}} \leq R$ with $R > 0$ satisfying $C(6\varepsilon + R^2) < R$. From (24), we obtain $\|\Lambda(v)\|_{\mathcal{Z}} < R$. Thus, the application Λ maps the ball $B_R := \{v \in \mathcal{Z} / \|v\|_{\mathcal{Z}} \leq R\}$ into itself.

We will now prove that $\Lambda : B_R \rightarrow B_R$ is a contraction. For any $z, v \in B_R$, $\Lambda(z) - \Lambda(v)$ is the solution of (21) with $z_0 = 0$, $h_j = 0$ for $j = 1, \dots, 4$ and $\hat{f} = vv_x - zz_x$. We obtain the estimate

$$\|\Lambda(z) - \Lambda(v)\|_{\mathcal{Z}} \leq C\|vv_x - zz_x\|_{\mathcal{F}} \leq C(\|(v - z)v_x\|_{\mathcal{F}} + \|z(v_x - z_x)\|_{\mathcal{F}}).$$

Using the definition (2) of the space \mathcal{F} , $v, z \in C([0, 1]; H^6(0, 1)) \hookrightarrow L^\infty(0, T; W^{5, \infty}(0, 1))$ and $v_t, z_t \in C([0, 1]; H^6(0, 1)) \hookrightarrow L^\infty(0, T; W^{1, \infty}(0, 1))$, we obtain

$$\|\Lambda(z) - \Lambda(v)\|_{\mathcal{Z}} \leq C(\|v\|_{\mathcal{Z}} + \|z\|_{\mathcal{Z}})\|v - z\|_{\mathcal{Z}} \leq 2CR\|v - z\|_{\mathcal{Z}},$$

which implies that Λ is a contraction if R is chosen small enough. More precisely, we can choose R, ε such that $2CR < 1$ and $C(6\varepsilon + R^2) < R$. Hence, the map Λ has a unique fixed point $y \in \mathcal{Z}$, which is the unique solution of (8). Thus, we have proven Theorem 1.1. \square

3 Global Carleman inequality

In this section, a global Carleman inequality will be proved for the linearized K-S equation. We define the space

$$\mathcal{V} = \{v \in L^2(0, T; H^4 \cap H_0^2(0, 1)) / Lv \in L^2((0, T) \times (0, 1))\} \quad (25)$$

where

$$Lv = v_t + (\sigma v_{xx})_{xx} + q_2 v_{xx} + q_1 v_x + q_0 v$$

with $q_j \in L^\infty(\Omega)$ for $j = 0, 1, 2$.

Consider $\beta \in C^4([0, 1])$ such that for some $r > 0$ we have, for all $x \in (0, 1)$:

$$\begin{aligned} 0 < r &\leq \beta(x), \\ 0 < r &\leq \beta'(x), \\ \beta''(x) &\leq -r < 0, \\ |\sigma'(x)\beta'(x)| &\leq \frac{r}{4} \min_{z \in [0, 1]} \{\sigma(z)\}. \end{aligned} \tag{26}$$

For instance, if σ is constant, we can consider $\beta(x) = \sqrt{1+x}$.

On the other hand, given $T_0 \in (0, T)$ we can choose $\phi_0 \in C^1([0, T])$ such that

$$\begin{aligned} \phi_0(0) = \phi_0(T) &= 0, \quad \text{and} \\ 0 < \phi_0(t) &\leq \phi_0(T_0) \quad \text{for each } t \in (0, T). \end{aligned} \tag{27}$$

For example, if $T_0 = T/2$, we can use $\phi_0(t) = t(T-t)$.

We finally define the function

$$\phi(t, x) = \frac{\beta(x)}{\phi_0(t)}, \tag{28}$$

for $(t, x) \in (0, T) \times [0, 1]$, which is the weight function of the Carleman estimate. From (26) and (27) it is not difficult to see that ϕ satisfies the following properties:

$$\begin{aligned} \exists C > 0 \text{ such that } \phi &\leq C\phi_x \text{ and} \\ \phi^n &\leq C\phi^m \text{ for each positive integers } n < m. \end{aligned} \tag{29}$$

Theorem 3.1 *Let ϕ be a function defined by (28) and $m > 0$. Then there exists $\lambda_0 > 0$ and a constant $C = C(T, \lambda_0, r, m) > 0$ such that if $\|q_i\|_{L^\infty((0, T) \times (0, 1))} \leq m$ for $i = 0, 1, 2$ then we have*

$$\begin{aligned} \int_0^T \int_0^1 e^{-2\lambda\phi} \left(\frac{|v_t|^2 + |(\sigma v_{xx})_{xx}|^2}{\lambda\phi} + \lambda^7 \phi^7 |v|^2 + \lambda^5 \phi^5 |v_x|^2 + \lambda^3 \phi^3 |v_{xx}|^2 + \lambda\phi |v_{xxx}|^2 \right) dx dt \\ \leq C \int_0^T \int_0^1 e^{-2\lambda\phi} |Lv|^2 dx dt \\ + C \int_0^T e^{-2\lambda\phi(t, 0)} \left(\lambda^3 \phi_x^3(t, 0) \sigma(0)^2 |v_{xx}(t, 0)|^2 + \lambda\phi_x(t, 0) \sigma^2(0) |v_{xxx}(t, 0)|^2 \right) dt \end{aligned} \tag{30}$$

for all $v \in \mathcal{V}$, for all $\lambda \geq \lambda_0$.

As we pointed out in the Introduction, a Carleman estimate for the K-S equation with constant coefficients σ and γ was previously obtained in [10]. The final goal in that work was to prove null-controllability with boundary controls. Thus, (30) is a generalization to the case of non-constant coefficients.

Proof. Consider the following operator P defined in $\mathcal{W}_\lambda := \{e^{-\lambda\phi}v : v \in \mathcal{V}\}$ by

$$Pw = e^{-\lambda\phi}L(e^{\lambda\phi}w).$$

We then obtain the decomposition $Pw = P_1w + P_2w + Rw$, where

$$P_1w = 6\lambda^2\phi_x^2\sigma w_{xx} + \lambda^4\phi_x^4\sigma w + (\sigma w_{xx})_{xx} + 6\lambda^2(\phi_x^2\sigma)_x w_x \quad (31)$$

$$P_2w = w_t + 4\lambda^3\phi_x^3\sigma w_x + 4\lambda\phi_x\sigma w_{xxx} + 4\lambda^3\phi_x(\phi_x^2\sigma)_x w \quad (32)$$

$$\begin{aligned} Rw = & \lambda\phi_t w + 2\lambda\phi_x\sigma_{xx}w_x + \lambda^2\phi_x^2\sigma_{xx}w + \lambda\phi_{xx}\sigma_{xx}w \\ & + 6\lambda\phi_x\sigma_x w_{xx} + 6\lambda^2\phi_x\phi_{xx}\sigma_x w + 6\lambda\phi_{xx}\sigma_x w + 2\lambda\phi_{xxx}\sigma_x w \\ & + 4\lambda^2\phi_x\phi_{xxx}\sigma w + 6\lambda\phi_{xx}\sigma w_{xx} + 3\lambda^2\phi_{xx}^2\sigma w + 4\lambda\phi_{xxx}\sigma w_x \\ & + \lambda\phi_{xxxx}\sigma w + q_0 w + q_1 w_x + q_1 \lambda\phi_x w \\ & + q_2 w_{xx} + 2\lambda q_2 \phi_x w_x + \lambda^2 q_2 \phi_x^2 w + \lambda\phi_{xx} q_2 w \\ & - 2\lambda^3\phi_x^2\phi_{xx}\sigma w - 2\lambda^3\phi_x^3\sigma_x w. \end{aligned} \quad (33)$$

Thus,

$$\|Pw - Rw\|_{L^2(Q)}^2 = \|P_1w\|_{L^2(Q)}^2 + 2\langle P_1w, P_2w \rangle + \|P_2w\|_{L^2(Q)}^2$$

where $\langle \cdot, \cdot \rangle$ is the $L^2(Q)$ scalar product.

For any $v \in \mathcal{V}$ we obtain $v_t \in L^2(0, T; L^2(0, 1))$ and then $v \in C([0, T]; L^2(0, 1))$. From the construction of ϕ (see (27)), we obtain $w \in C([0, T]; L^2(0, 1))$ and $w(x, 0) = w(x, T) = 0$ for any $w \in \mathcal{W}_\lambda$.

Let us define the notations

$$\begin{aligned} I(w) &= -6\lambda^7 \int_0^T \int_0^1 \phi_x^6 \phi_{xx} \sigma^2 |w|^2 dx dt, \\ I(w_x) &= -\lambda^5 \int_0^T \int_0^1 \phi_x^4 \sigma (30\phi_{xx}\sigma + 12\phi_x\sigma_x) |w_x|^2 dx dt, \\ I(w_{2x}) &= -\lambda^3 \int_0^T \int_0^1 \phi_x^2 \sigma (58\phi_{xx}\sigma + 40\phi_x\sigma_x) |w_{xx}|^2 dx dt, \\ I(w_{3x}) &= -\lambda \int_0^T \int_0^1 \sigma (2\phi_{xx}\sigma - 4\phi_x\sigma_x) |w_{xxx}|^2 dx dt, \end{aligned}$$

and

$$I_x = \int_0^T (10\lambda^3\phi_x^3\sigma^2 |w_{xx}|^2 + 2\lambda\phi_x\sigma\sigma_{xx} |w_{xx}|^2 + 2\lambda\phi_x\sigma^2 |w_{xxx}|^2) \Big|_{x=0}^1 dt$$

The following weighted norm is defined, for any $w \in \mathcal{W}_\lambda$, as

$$\|w\|_{\lambda, \phi}^2 = \int_0^T \int_0^1 (\lambda^7\phi^7 |w|^2 + \lambda^5\phi^5 |w_x|^2 + \lambda^3\phi^3 |w_{xx}|^2 + \lambda\phi |w_{xxx}|^2) dx dt.$$

We first require the following

Lemma 3.2 *Under the hypothesis of Theorem 3.1, there exists $\delta > 0$ such that*

$$\langle P_1w, P_2w \rangle_{L^2(Q)} \geq \delta \|w\|_{\lambda, \phi}^2 + I_x \quad (34)$$

for λ large enough and for all $w \in \mathcal{W}_\lambda$.

Proof. It is sufficient to prove that

$$\langle P_1 w, P_2 w \rangle_{L^2} = \sum_{k=0}^3 I(w_{kx}) + R_0(w) + I_x \quad (35)$$

for a large enough λ , for all $w \in \mathcal{W}_\lambda$, where $|R_0(w)| \leq \lambda^{-1} \|w\|_{\lambda, \phi}^2$.

Indeed, let us first assume that we have (35). From the hypotheses in (26) we easily check that there exists $\varepsilon > 0$ such that ϕ satisfies for all $x \in (0, 1)$,

$$\begin{aligned} \phi_{xx}(x) &\leq -\varepsilon\phi < 0, \\ 30\phi_{xx}(x)\sigma(x) + 12\phi_x(x)\sigma_x(x) &\leq -\varepsilon\phi < 0, \\ 58\phi_{xx}(x)\sigma(x) + 40\phi_x(x)\sigma_x(x) &\leq -\varepsilon\phi < 0, \text{ and} \\ 2\phi_{xx}(x)\sigma(x) - 4\phi_x(x)\sigma_x(x) &\leq -\varepsilon\phi < 0. \end{aligned} \quad (36)$$

Then from (29) and assuming (35) we obtain, for λ large enough,

$$\begin{aligned} \langle P_1 w, P_2 w \rangle_{L^2} &= \sum_{k=0}^3 I(w_{kx}) + R_0(w) + I_x \\ &\geq 2\delta \|w\|_{\lambda, \phi}^2 - |R_0(w)| + I_x \\ &\geq \delta \|w\|_{\lambda, \phi}^2 + I_x. \end{aligned} \quad (37)$$

Let us now prove (35): we write $\langle P_1 w, P_2 w \rangle_{L^2(Q)} = \sum_{i,j=1}^4 I_{i,j}$ where $I_{i,j}$ denotes the L^2 -product between the i -th term of $P_1 w$ in (31) and the j -th term of $P_2 w$ in (32).

Integrations by parts in time or space are performed on each expression $I_{i,j}$. Each resulting expression will be included in one of the terms of the right-hand side of (35). The results are listed below, and we indicate for each term where it will be included.

- $I_{1,1} = -I_{4,1} + \underbrace{3\lambda^2 \iint_Q (\phi_x^2 \sigma)_t |w_x|^2}_{R_0(w)}$
- $I_{1,2} = -\underbrace{12\lambda^5 \iint_Q (\phi_x^5 \sigma^2)_x |w_x|^2}_{I(w_x)}$
- $I_{1,3} = -\underbrace{12\lambda^3 \iint_Q (\phi_x^3 \sigma^2)_x |w_{xx}|^2}_{I(w_{2x})} + \underbrace{12\lambda^3 \int_0^T \left[\phi_x^3 \sigma^2 |w_{xx}|^2 \right]_0^1 dt}_{I_x}$
- $I_{1,4} = \underbrace{12\lambda^5 \iint_Q [\phi_x^3 \sigma (\phi_x^2 \sigma)_x]_{xx} |w|^2}_{R_0(w)} - \underbrace{24\lambda^5 \iint_Q \phi_x^3 \sigma (\phi_x^2 \sigma)_x |w_x|^2}_{I(w_x)}$
- $I_{2,1} = -\underbrace{\frac{\lambda^4}{2} \iint_Q (\phi_x^4 \sigma)_t |w|^2}_{R_0(w)}$
- $I_{2,2} = -\underbrace{2\lambda^7 \iint_Q (\phi_x^7 \sigma^2)_x |w|^2}_{I(w)}$
- $I_{2,3} = -\underbrace{2\lambda^5 \iint_Q (\phi_x^5 \sigma^2)_{xxx} |w|^2}_{R_0(w)} + \underbrace{6\lambda^5 \iint_Q (\phi_x^5 \sigma^2)_x |w_x|^2}_{I(w_x)}$

- $I_{2,4} = \underbrace{4\lambda^7 \iint_Q \phi_x^5 \sigma (\phi_x^2 \sigma)_x |w|^2}_{I(w)}.$
- $I_{3,1} = \frac{1}{2} \int_0^1 \left[\sigma |w_{xx}|^2 \right]_0^T dx = 0.$
- $I_{3,2} = \underbrace{-2\lambda^3 \iint_Q [(\phi_x^3 \sigma)_{xx} \sigma]_x |w_x|^2}_{R_0(w)} + \underbrace{4\lambda^3 \iint_Q (\phi_x^3 \sigma)_x \sigma |w_{xx}|^2}_{I(w_{2x})}$
 $+ \underbrace{2\lambda^3 \iint_Q (\phi_x^3)_x \sigma^2 |w_{xx}|^2}_{I(w_{2x})} - \underbrace{2\lambda^3 \int_0^T \left[\phi_x^3 \sigma^2 |w_{xx}|^2 \right]_0^1}_{I_x} dt.$
- $I_{3,3} = \underbrace{2\lambda \int_0^T \left[\phi_x \sigma \sigma_{xx} |w_{xx}|^2 \right]_0^1}_{I_x} dt - \underbrace{2\lambda \iint_Q (\phi_x \sigma \sigma_{xx})_x |w_{xx}|^2}_{R_0(w)} + \underbrace{8\lambda \iint_Q \phi_x \sigma \sigma_x |w_{3x}|^2}_{I(w_{xxx})}$
 $+ \underbrace{2\lambda \int_0^T \left[\phi_x \sigma^2 |w_{xxx}|^2 \right]_0^1}_{I_x} dt - \underbrace{2\lambda \iint_Q (\phi_x \sigma^2)_x |w_{xxx}|^2}_{I(w_{xxx})}.$
- $I_{3,4} = \underbrace{4\lambda^3 \iint_Q (\phi_x (\phi_x^2 \sigma)_{xx}) \sigma w_{xx}}_{R_0(w)} - \underbrace{4\lambda^3 \iint_Q (\phi_x (\phi_x^2 \sigma)_x)_x \sigma |w_x|^2}_{R_0(w)}$
 $+ \underbrace{4\lambda^3 \iint_Q \phi_x (\phi_x^2 \sigma)_x \sigma |w_{2x}|^2}_{I(w_{2x})}.$
- $I_{4,1} = 6\lambda^2 \iint_Q (\phi_x^2 \sigma)_x w_x w_t$, which is canceled when adding with $I_{1,1}$.
- $I_{4,2} = \underbrace{24\lambda^5 \iint_Q (\phi_x^2 \sigma)_x \phi_x^3 \sigma |w_x|^2}_{I(w_x)}.$
- $I_{4,3} = \underbrace{12\lambda^3 \iint_Q [(\phi_x^2 \sigma)_x \phi_x \sigma]_{xx} |w_x|^2}_{R_0(w)} - \underbrace{24\lambda^3 \iint_Q (\phi_x^2 \sigma)_x \phi_x \sigma |w_{xx}|^2}_{I(w_{2x})}.$
- $I_{4,4} = \underbrace{-12\lambda^5 \iint_Q (\phi_x^2 \sigma)_x (\phi_x^3 \sigma)_x |w|^2}_{R_0(w)}.$

Summing up all the terms, we obtain (35). \square

Then, we will prove a Carleman inequality for the conjugated operator P .

Lemma 3.3 *There exists $\lambda_0 > 0$ such that for all $\lambda \geq \lambda_0$ we have, for all $w \in \mathcal{W}_\lambda$,*

$$\int_0^T \int_0^1 (\lambda^7 \phi^7 |w|^2 + \lambda^5 \phi^5 |w_x|^2 + \lambda^3 \phi^3 |w_{xx}|^2 + \lambda \phi |w_{xxx}|^2) dx dt$$

$$+ \|P_1 w\|_{L^2(Q)}^2 + \|P_2 w\|_{L^2(Q)}^2 \leq C \|Pw\|_{L^2(Q)}^2 - I_x.$$

Proof.

From hypothesis (26) and the inequalities listed in (36), we know that there exists $\delta > 0$ such that

$$\sum_{k=0}^3 I(w_{kx}) \geq \delta \|w\|_{\lambda, \phi}^2 \quad (38)$$

for a parameter λ large enough.

Besides, from the definition (33), the fact $\|q_i\|_{L^\infty((0,T) \times (0,1))} \leq m$ for $i = 0, 1, 2$, and (29), it is trivial to check that

$$\begin{aligned} \|Rw\|_{L^2((0,T) \times (0,1))}^2 &\leq C \left(\lambda^6 \iint_Q \phi^6 |w|^2 + \lambda^2 \iint_Q \phi^2 |w_x|^2 + \lambda^2 \iint_Q \phi^2 |w_{xx}|^2 \right) \\ &\leq C \lambda^{-1} \|w\|_{\lambda, \phi}^2. \end{aligned} \quad (39)$$

Thus, for λ large enough, we have

$$\begin{aligned} \|P_1 w\|_{L^2}^2 + 2 \langle P_1 w, P_2 w \rangle + \|P_2 w\|_{L^2}^2 &= \|Pw - Rw\|_{L^2}^2 \\ &\leq 2 \|Pw\|_{L^2}^2 + 2 \|Rw\|_{L^2}^2 \\ &\leq 2 \|Pw\|_{L^2}^2 + C \lambda^{-1} \|w\|_{\lambda, \phi}^2. \end{aligned} \quad (40)$$

From Lemma 3.2 and estimates (40) and (38), we conclude the proof of Lemma 3.3. \square

To complete the proof of Theorem 3.1, we have to deal with the norms for $P_1 w$ and $P_2 w$ appearing in Lemma 3.3. From the definition of $P_2 w$, and because (26) holds, we have

$$\frac{1}{\lambda \phi} |w_t|^2 \leq \frac{2}{\lambda \phi} |P_2 w|^2 + C (\lambda^5 \phi^5 |w|^2 + \lambda^5 \phi^5 |w_x|^2 + \lambda \phi |w_{xxx}|^2)$$

and

$$\iint_Q \frac{1}{\lambda \phi} |w_t|^2 \leq C \iint_Q |P_2 w|^2 + C \|w\|_{\lambda, \phi}^2$$

for λ large enough. A similar result is proven for $(\sigma w_{xx})_{xx}$ and $P_1 w$, and we then have

$$\iint_Q \frac{1}{\lambda \phi} (|w_t|^2 + |(\sigma w_{xx})_{xx}|^2) \leq C \iint_Q (|P_1 w|^2 + |P_2 w|^2) + C \|w\|_{\lambda, \phi}^2. \quad (41)$$

From (41) and Lemma 3.3 we obtain

$$\begin{aligned} \iint_Q \frac{1}{\lambda \phi} (|w_t|^2 + |(\sigma w_{xx})_{xx}|^2) + \lambda^7 \phi^7 |w|^2 + \lambda^5 \phi^5 |w_x|^2 + \lambda^3 \phi^3 |w_{xx}|^2 + \lambda \phi |w_{xxx}|^2 \\ \leq C \iint_Q |Pw|^2 - CI_x. \end{aligned} \quad (42)$$

To handle the terms in I_x , we note that for any $x \in (0, 1)$ and λ large enough,

$$-C \lambda \int_0^T \phi_x(x, t) \sigma(x) \sigma_{xx}(x) |w_{xx}(x, t)|^2 dt \leq C \lambda^3 \int_0^T \phi_x(x, t)^3 \sigma(x)^2 |w_{xx}(x, t)|^2 dt.$$

Then

$$-CI_x \leq C \lambda^3 \int_0^T \phi_x(0, t)^3 \sigma(0)^2 |w_{xx}(0, t)|^2 dt + C \lambda \int_0^T \phi_x(0, t) \sigma(0)^2 |w_{xxx}(0, t)|^2 dt \quad (43)$$

and from (42) and (43) we obtain

$$\begin{aligned} \iint_Q \frac{1}{\lambda \phi} (|w_t|^2 + |(\sigma w_{xx})_{xx}|^2) + \|w\|_{\lambda, \phi}^2 &\leq C \iint_Q |Pw|^2 \\ &+ C \lambda^3 \int_0^T \phi_x(0, t)^3 \sigma(0)^2 |w_{xx}(0, t)|^2 dt + C \lambda \int_0^T \phi_x(0, t) \sigma(0)^2 |w_{xxx}(0, t)|^2 dt. \end{aligned} \quad (44)$$

Computing the derivatives of $e^{\lambda\phi}w$ it is trivial to prove that

$$\left| \partial_x^k v \right|^2 = \left| \partial_x^k (e^{\lambda\phi}w) \right|^2 \leq C \sum_{j=0}^k \left| \lambda^{k-j} \phi^{k-j} \partial_x^j w \right|^2$$

for each $k = 0, \dots, 3$. Therefore

$$\begin{aligned} \int_0^T \int_0^1 e^{-2\lambda\phi} \left(\lambda^7 \phi^7 |e^{\lambda\phi}w|^2 + \lambda^5 \phi^5 |(e^{\lambda\phi}w)_x|^2 + \lambda^3 \phi^3 |(e^{\lambda\phi}w)_{xx}|^2 + \lambda \phi |(e^{\lambda\phi}w)_{xxx}|^2 \right) dx dt \\ \leq C \|w\|_{\lambda, \phi}. \end{aligned}$$

Considering finally that $Pw = e^{-\lambda\varphi}Lv$, we obtain Carleman estimate (30). \square

Remark 3.4 *We considered the function β to be increasing. This allows the Carleman inequality to be obtained with boundary terms at $x = 0$. If a decreasing function β was used instead, then an inequality with boundary terms at $x = 1$ would have been obtained. As discussed in the following section, the boundary terms in the Carleman inequality are related to the location of the observations in the inverse problem.*

4 Inverse Problem

In this section, the local stability of the nonlinear inverse problem stated in Theorem 1.2 will be proved following the ideas of [7] and [21]. The proof is splitted in several steps.

Step 1. Local study of the inverse problem

Let γ , $\tilde{\gamma}$, y and \tilde{y} be defined as in Theorem 1.2. If we set $u = y - \tilde{y}$ and $f = \tilde{\gamma} - \gamma$, then u solves the following K-S equation:

$$\begin{cases} u_t + (\sigma(x)u_{xx})_{xx} + \gamma u_{xx} + \tilde{y}u_x + \tilde{y}_x u + uu_x = f(x)\tilde{y}_{xx}(x, t), & \forall (t, x) \in Q, \\ u(t, 0) = u(t, 1) = 0, & \forall t \in (0, T), \\ u_x(t, 0) = u_x(t, 1) = 0, & \forall t \in (0, T), \\ u(0, x) = 0, & \forall x \in (0, 1). \end{cases} \quad (45)$$

Then, in order to prove the stability of the inverse problem mentioned in the introduction, it is sufficient to obtain an estimate of f in terms of $u_{xx}(\cdot, 0)$, $u_{xxx}(\cdot, 0)$ and $u(T_0, \cdot)$, where $\tilde{\gamma}$ and \tilde{y} are given, $\gamma \in H^4(0, 1)$ and u is the solution of Equation (45).

We begin by deriving Equation (45) with respect to time. Thus, $v = u_t$ satisfies the following equation:

$$\begin{cases} v_t + (\sigma v_{xx})_{xx} + \gamma v_{xx} + \tilde{y}v_x + \tilde{y}_x v = f\tilde{y}_{xxt} - g, & \forall (t, x) \in Q, \\ v(t, 0) = v(t, 1) = 0, & \forall t \in (0, T), \\ v_x(t, 0) = v_x(t, 1) = 0, & \forall t \in (0, T), \\ v(0, x) = fR(x, 0), & \forall x \in (0, 1), \end{cases} \quad (46)$$

where $g(x, t) = u(x, t)y_{xt}(x, t) + u_x(x, t)y_t(x, t)$.

The proof of Theorem 1.2 relies on the use of the Carleman estimate given in Theorem 3.1. This result will be used twice. First, Equation (46) allows to estimate v in terms of f , \tilde{y}_{xx} and g . Then, Equation (45) will be used to handle the terms u and u_x , which appear in the expression of the source term g . The details are given in the next step below.

Step 2. First use of the Carleman estimate

Similarly to the proof of the Carleman estimate, we set $w = e^{-\lambda\phi}v$. Then, we work on the term

$$I = 2 \int_0^1 \int_0^{T_0} w(t, x) w_t(t, x) dt dx.$$

On the one hand, we can calculate I and bound it from below. Indeed, using $w(0, x) = e^{-\lambda\phi(0, x)}v(0, x) = 0$ for all $x \in (0, 1)$ and Equation (45), we can easily obtain

$$\begin{aligned} I &= \int_0^1 |w(T_0, x)|^2 dx \\ &= \int_0^1 e^{-2\lambda\phi(T_0, x)} |(f\tilde{y}_{xx} - (\sigma u_{xx})_{xx} - \gamma u_{xx} - \tilde{y}u_x - \tilde{y}_x u - uu_x)(T_0, x)|^2 dx \\ &\geq \int_0^1 e^{-2\lambda\phi(T_0, x)} |f(x)|^2 |\tilde{y}_{xx}(T_0, x)|^2 dx - C \|u(T_0)\|_{H^4(0,1)}^2 - C \|u(T_0)\|_{H^1(0,1)}^4 \end{aligned}$$

where C depends on $\|\gamma\|_{L^\infty(0,1)}$, $\|\tilde{y}(T_0)\|_{W^{1,\infty}(0,1)}$ and $\|\sigma\|_{W^{2,\infty}(0,1)}$.

On the other hand, in order to estimate I from above we apply the Carleman estimate (44) to Equation (46) using $q_0 = \tilde{y}_x$ and $q_1 = \tilde{y}$, which are uniformly bounded in $L^\infty((0, T) \times (0, 1))$ by the hypothesis in Theorem 1.2. We obtain that

$$\begin{aligned} I &= 2 \int_0^1 \int_0^{T_0} w(t, x) w_t(t, x) dt dx \\ &\leq \left(\int_0^1 \int_0^{T_0} \lambda\phi(t, x) |w(t, x)|^2 dt dx \right)^{\frac{1}{2}} \left(\int_0^1 \int_0^{T_0} \frac{|w_t(t, x)|^2}{\lambda\phi(t, x)} dt dx \right)^{\frac{1}{2}} \\ &\leq C\lambda^{-3} \int_0^1 \int_0^T e^{-2\lambda\phi} |f(x)\tilde{y}_{xxt}(x, t)|^2 dx dt + C\lambda^{-3} \int_0^1 \int_0^T e^{-2\lambda\phi} |g(x, t)|^2 dx dt \\ &\quad + C\lambda^{-3} \int_0^T e^{-2\lambda\phi(0, t)} (\lambda^3 \phi_x^3(0, t) \sigma^2(0) |v_{xx}(0, t)|^2 + \lambda\phi_x(0, t) \sigma^2(0) |v_{xxx}(0, t)|^2) dt. \end{aligned}$$

Step 3. Second use of the Carleman estimate

Considering that $g = uy_{xt} + u_x y_t$, we will now use a Carleman estimate for the solution of Equation (45) in order to manage the term in g of the previous inequality. The unknown trajectory y is nevertheless such that y_{xt} and y_t belong to $L^\infty(0, T; L^\infty(0, 1))$ since $y \in \mathcal{Z}$. Thus, we have

$$\begin{aligned} \iint_Q e^{-2\lambda\phi} |g|^2 &\leq 2 \iint_Q e^{-2\lambda\phi} |u|^2 |y_{xt}|^2 + 2 \iint_Q e^{-2\lambda\phi} |u_x|^2 |y_t|^2 \\ &\leq C \iint_Q e^{-2\lambda\phi} (|u|^2 + |u_x|^2). \end{aligned}$$

Then we can apply Carleman estimate (44) to equation (45), using the identity $\tilde{y}_x u + uu_x = uy_x$, and taking $q_0 = y_x$ and $q_1 = \tilde{y}$, which are bounded in $L^\infty((0, T) \times (0, 1))$. We

can choose λ_0 as large as possible in Theorem 3.1: we then obtain

$$\begin{aligned}
& \iint_Q e^{-2\lambda\phi} |g|^2 \leq C\lambda^{-5} \iint_Q e^{-2\lambda\phi} (\lambda^7 |u|^2 + \lambda^5 |u_x|^2) \\
& \leq C\lambda^{-5} \iint_Q e^{-2\lambda\phi} |f\tilde{y}_{xx}|^2 \\
& \quad + C\lambda^{-5} \int_0^T e^{-2\lambda\phi(0,t)} (\lambda^3 \phi_x^3(0,t) \sigma^2(0) |u_{xx}(0,t)|^2 + \lambda \phi_x(0,t) \sigma^2(0) |u_{xxx}(0,t)|^2) dt.
\end{aligned}$$

Gathering all the estimates of I and g that were obtained above, we have

$$\begin{aligned}
& \int_0^1 e^{-2\lambda\phi(T_0,x)} |f(x)|^2 |\tilde{y}_{xx}(T_0,x)|^2 dx - C \|u(T_0)\|_{H^4(0,1)}^2 - C \|u(T_0)\|_{H^1(0,1)}^4 \\
& \leq C\lambda^{-3} \iint_Q e^{-2\lambda\phi} |f\tilde{y}_{xxt}|^2 + C\lambda^{-8} \iint_Q e^{-2\lambda\phi} |f\tilde{y}_{xx}|^2 \\
& \quad + C\lambda^{-8} \int_0^T e^{-2\lambda\phi(0,t)} (\lambda^3 \phi_x^3(0,t) \sigma^2(0) |u_{xx}(0,t)|^2 + \lambda \phi_x(0,t) \sigma^2(0) |u_{xxx}(0,t)|^2) dt \\
& \quad + C\lambda^{-3} \int_0^T e^{-2\lambda\phi(0,t)} (\lambda^3 \phi_x^3(0,t) \sigma^2(0) |v_{xx}(0,t)|^2 + \lambda \phi_x(0,t) \sigma^2(0) |v_{xxx}(0,t)|^2) dt.
\end{aligned}$$

From the hypothesis of the theorem, we have $\tilde{y} \in C([0, T]; H^6(0, 1))$, $\tilde{y}_t \in C([0, T]; H^2(0, 1))$, and $|\tilde{y}_{xx}(T_0, \cdot)| > \eta > 0$ in $(0, 1)$. Using also that the Carleman weight function satisfies (27) thus $e^{-2\lambda\phi(t,x)} \leq e^{-2\lambda\phi(T_0,x)}$ in $(0, T) \times (0, 1)$, we obtain

$$\begin{aligned}
& \int_0^1 e^{-2\lambda\phi(T_0,x)} |f(x)|^2 dx \\
& \leq C \left(\lambda^{-3} \int_0^1 e^{-2\lambda\phi(T_0,x)} |f(x)|^2 dx + \|u(T_0)\|_{H^4(0,1)}^2 + \|u(T_0)\|_{H^1(0,1)}^4 \right. \\
& \quad + \lambda^{-8} \int_0^T e^{-2\lambda\phi(0,t)} (\lambda^3 \phi_x^3(0,t) \sigma^2(0) |u_{xx}(0,t)|^2 + \lambda \phi_x(0,t) \sigma^2(0) |u_{xxx}(0,t)|^2) dt \\
& \quad \left. + \lambda^{-3} \int_0^T e^{-2\lambda\phi(0,t)} (\lambda^3 \phi_x^3(0,t) \sigma^2(0) |v_{xx}(0,t)|^2 + \lambda \phi_x(0,t) \sigma^2(0) |v_{xxx}(0,t)|^2) dt \right).
\end{aligned}$$

Therefore, the regularity of ϕ (that come from the assumptions on β and ϕ_0) allows to prove that choosing λ_0 large enough, we obtain the existence of a constant C that depends on r, K, T, λ_0, m such that $\forall \lambda > \lambda_0$,

$$\begin{aligned}
\|f(x)\|_{L^2(0,1)}^2 & \leq C \left(\|u(T_0, \cdot)\|_{H^4(0,1)}^2 + \|u(T_0, \cdot)\|_{H^1(0,1)}^4 \right. \\
& \quad \left. + \|u_{xx}(\cdot, 0)\|_{H^1(0,T)}^2 + \|u_{xxx}(\cdot, 0)\|_{H^1(0,T)}^2 \right).
\end{aligned}$$

This estimate leads to the local stability of the initial inverse problem since $f = \tilde{\gamma} - \gamma$ and $u = y - \tilde{y}$ and we have proved Theorem 1.2.

Acknowledgments: This work began while L. Baudouin and E. Crépeau were visiting the Universidad Técnica Federico Santa María on the framework of the MathAmsud project CIP-PDE. This study was partially supported by Fondecyt #11080130, Fondecyt #11090161, ANR C-QUID and CISIFS, and CMM-Basal grants.

References

- [1] A. Armaou and P. D. Christofides, *Feedback control of the Kuramoto-Sivashinsky equation*, Phys. D, **137** (2000), 49–61.
- [2] L. Baudouin and J.-P. Puel, *Uniqueness and stability in an inverse problem for the Schrödinger equation*, Inverse Problems **18** (2002), no. 6, 1537–1554.
- [3] L. Beilina and M. Klibanov, *Approximate Global Convergence and Adaptivity for Coefficient Inverse Problems*. Springer, 2012.
- [4] M. Bellassoued and M. Yamamoto, *Logarithmic stability in determination of a coefficient in an acoustic equation by arbitrary boundary observation*, J. Math. Pures Appl. (9) **85** (2006), no. 2, 193–224.
- [5] A. Benabdallah, P. Gaitan, and J. Le Rousseau, *Stability of discontinuous diffusion coefficients and initial conditions in an inverse problem for the heat equation*, SIAM J. Control Optim. **46** (2007), no. 5, 1849–1881.
- [6] M. Boulakia, C. Grandmont, and A. Osses, *Some inverse stability results for the bistable reaction-diffusion equation using Carleman inequalities*, C. R. Math. Acad. Sci. Paris **347** (2009), no. 11-12, 619–622.
- [7] A. L. Bukhgeim and M. V. Klibanov, *Uniqueness in the large of a class of multidimensional inverse problems*, Dokl. Akad. Nauk SSSR **260** (1981), no. 2, 269–272.
- [8] T. Cazenave and A. Haraux, *An Introduction to Semilinear Evolution Equations*, (Oxford Science Publications), (1998) Ed Oxford University Press.
- [9] E. Cerpa, *Null controllability and stabilization of a linear Kuramoto-Sivashinsky equation*, Commun. Pure Appl. Anal. **9** (2010), pp. 91–102.
- [10] E. Cerpa and A. Mercado, *Local exact controllability to the trajectories of the 1-D Kuramoto-Sivashinsky equation*, J. Differential Equations **250** (2011), no. 4, 2024–2044.
- [11] J.-M. Coron and S. Guerrero, *Singular optimal control: A linear 1-D parabolic-hyperbolic example*, Asymptotic Analysis. **44** (2005), 237–257.
- [12] P. D. Christofides and A. Armaou, *Global stabilization of the Kuramoto-Sivashinsky equation via distributed output feedback control*, Systems Control Lett., **39** (2000), 283–294.
- [13] M. Cristofol, P. Gaitan, and H. Ramoul, *Inverse problems for a 2×2 reaction-diffusion system using a Carleman estimate with one observation*, Inverse Problems **22** (2006), no. 5, 1561–1573.
- [14] D. Dos Santos Ferreira, C. E. Kenig, M. Salo, and G. Uhlmann. Limiting Carleman weights and anisotropic inverse problems. *Invent. Math.*, 178(1):119–171, 2009.
- [15] H. Egger, H. W. Engl, and M. V. Klibanov, *Global uniqueness and Hölder stability for recovering a nonlinear source term in a parabolic equation*, Inverse Problems **21** (2005), no. 1, 271–290.

- [16] E. Fernández-Cara and S. Guerrero, *Global Carleman inequalities for parabolic systems and applications to controllability*, SIAM J. Control Optim. 45 (2006), no. 4, pp. 1395–1446.
- [17] C. Hu and R. Temam, *Robust control of the Kuramoto-Sivashinsky equation*, Dyn. Contin. Discrete Impuls. Syst. Ser. B Appl. Algorithms, **8** (2001), 315–338.
- [18] O. Yu. Imanuvilov and M. Yamamoto, *Lipschitz stability in inverse parabolic problems by the Carleman estimate*, Inverse Problems **14** (1998), no. 5, 1229–1245.
- [19] L. I. Ignat, A. F. Pazoto and L. Rosier, *Inverse Problem for the heat equation and the Schrodinger equation on a tree*, 2011, <http://hal.archives-ouvertes.fr/hal-00595147/fr/>
- [20] M. V. Klibanov, *Inverse problems and Carleman estimates*, Inverse Problems **8** (1992), no. 4, 575–596.
- [21] M. V. Klibanov and J. Malinsky, *Newton-Kantorovich method for three-dimensional potential inverse scattering problem and stability of the hyperbolic Cauchy problem with time-dependent data*, Inverse Problems **7** (1991), no. 4, 577–596.
- [22] M. V. Klibanov and A. Timonov. *Carleman estimates for coefficient inverse problems and numerical applications*. Inverse and Ill-posed Problems Series. VSP, Utrecht, 2004.
- [23] Y. Kuramoto and T. Tsuzuki. *On the formation of dissipative structures in reaction-diffusion systems*, Prog. Theor. Phys (1975). 54 687-99.
- [24] W.-J. Liu and M. Krstić, *Stability enhancement by boundary control in the Kuramoto-Sivashinsky equation*, Nonlinear Anal. Ser. A: Theory Methods, **43** (2001), 485–507.
- [25] J.-P. Puel and M. Yamamoto, *On a global estimate in a linear inverse hyperbolic problem*, Inverse Problems **12** (1996), no. 6, 995–1002.
- [26] M. Renardy and R. Rogers, *An Introduction to Partial Differential Equations*, (Texts in Applied Mathematics), ed Springer-Verlag New York, Inc. 2004.
- [27] L. Roques and M. Cristofol, *On the determination of the nonlinearity from localized measurements in a reaction-diffusion equation*, Nonlinearity **23** (2010), no. 3, 675–686.
- [28] G. I. Sivashinsky, *Nonlinear analysis of hydrodynamic instability in laminar flames I: Derivation of basic equations*, Acta Astronaut. (1977), 4 1177-1206.