



# LTBP2 NULL MUTATIONS IN AN AUTOSOMAL RECESSIVE OCULAR SYNDROME WITH MEGALOCORNEA, SPHEROPHAKIA, AND SECONDARY GLAUCOMA

Julie Desir, Yves Sznajer, Fanny Depasse, Françoise Roulez, Marc Schrooyen,  
Françoise Meire, Marc J Abramowicz

## ► To cite this version:

Julie Desir, Yves Sznajer, Fanny Depasse, Françoise Roulez, Marc Schrooyen, et al.. LTBP2 NULL MUTATIONS IN AN AUTOSOMAL RECESSIVE OCULAR SYNDROME WITH MEGALOCORNEA, SPHEROPHAKIA, AND SECONDARY GLAUCOMA. European Journal of Human Genetics, 2010, n/a (n/a), pp.n/a-n/a. <10.1038/ejhg.2010.11>. <hal-00511180>

**HAL Id: hal-00511180**

**<https://hal.science/hal-00511180v1>**

Submitted on 24 Aug 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

***LTBP2* NULL MUTATIONS IN AN AUTOSOMAL RECESSIVE OCULAR SYNDROME  
WITH MEGALOCORNEA, SPHEROPHAKIA, AND SECONDARY GLAUCOMA**

Julie Désir (1,2), Yves Sznajer (2,3), Fanny Depasse (4), Françoise Roulez (5), Marc Schrooyen  
(4), Françoise Meire (4,5,6), Marc Abramowicz (1,2)

1. IRIBHM, Université Libre de Bruxelles (ULB), Brussels, Belgium
2. Department of Medical Genetics, Hôpital Erasme-ULB, Brussels, Belgium
3. Clinical Genetics Unit, HUDERF-ULB, Brussels, Belgium
4. Ophthalmology Department, Hôpital Erasme-ULB, Brussels, Belgium
5. Ophthalmology Department, HUDERF-ULB, Brussels, Belgium
6. Ophthalmology Department, University Hospital, Gent, Belgium

Correspondence to:

Julie Désir, MD, PhD  
Medical Genetics Department  
Hôpital Erasme-ULB  
808 Route de Lennik  
B-1070 Brussels  
Belgium  
[jdesir@ulb.ac.be](mailto:jdesir@ulb.ac.be)  
Tel: +3225557076  
Fax: +3225554212

## **ABSTRACT**

The latent TGF $\beta$  binding proteins (LTBPs) and fibrillins are a superfamily of large, multidomain proteins with structural and TGF $\beta$ -signalling roles in the extracellular matrix. Their importance is underscored by fibrillin-1 mutations responsible for Marfan syndrome, but their respective roles are still incompletely understood. We report two families where children from healthy, consanguineous parents, presented with megalocornea and impaired vision associated with small, round, dislocated lenses (microspherophakia and ectopia lentis) and myopia, as well as a high-arched palate, and, in older children, tall stature with an abnormally high arm span over body height ratio, i.e., associated features of Marfan syndrome. Glaucoma was not present at birth, but was diagnosed in older children. Whole genome homozygosity mapping followed by candidate gene analysis identified homozygous truncating mutations of *LTBP2* gene in patients from both families. Fibroblast mRNA analysis was consistent with nonsense-mediated mRNA decay, with no evidence of mutated exon skipping. We conclude that biallelic null *LTBP2* mutations cause the ocular phenotype in both families and could lead to Marfan-like features in older children. We suggest that intraocular pressures should be followed-up in young children with an ocular phenotype consisting of megalocornea, spherophakia and/or lens dislocation, and recommend *LTBP2* gene analysis in these patients.

**KEYWORDS:** microspherophakia, megalocornea, Marfan

## INTRODUCTION

Marfan syndrome (MFS, MIM 154700) is a connective tissue disorder with autosomal dominant inheritance <sup>1</sup>. MFS typically affects the skeletal system with a disproportionate increase in the linear growth of long bones (tall stature with increased arm span and decreased upper-to-lower body ratio, arachnodactyly, dolichostenomelia, pectus deformity and scoliosis) and joint hyperlaxity; the ocular system with ectopia lentis in 55% of affected <18 years <sup>2</sup>, as well as microspherophakia (small round lens) and megalocornea (large cornea) in severe congenital forms <sup>3,4</sup>; and the cardiovascular system with thick, elongated cardiac valves and progressive aortic root enlargement. Additional features of MFS include striae distensae of the skin, a decrease in muscle and fat mass, and a predisposition to spontaneous pneumothorax. Together with progressive aortic enlargement or rupture, these features indicate an increased elastolysis and defective tissue repair <sup>5</sup>. MFS is caused by mutations in *FBNI*, whose protein product fibrillin-1 (FBN1) is a major structural component of the microfibrils <sup>6</sup>. Beyond the structural defect of microfibrils however, abnormal signaling by growth factors belonging to the TGF $\beta$  family play an important role in MFS <sup>7,8</sup>. An altered cellular behavior during development, associated with an increased TGF $\beta$  activity, is further supported by mouse models of MFS <sup>9</sup>. Aortic aneurysm in a mouse model of MFS can be prevented by TGF $\beta$  antagonists such as a TGF $\beta$ -neutralizing antibody or the angiotensin II type 1 receptor blocker, losartan <sup>10</sup>.

The TGF $\beta$ s are synthesized as prepropolypeptides, and processed in the Golgi apparatus into mature TGF $\beta$  tightly associated with its propeptide, the latency associated peptide (LAP). In the extracellular matrix, the LAP.TGF $\beta$  complex is further covalently bound to a latent TGF $\beta$ -binding protein (LTBP) <sup>11</sup>. The LTBPs are a family of proteins that show important structural homologies with the fibrillins <sup>7</sup>. They are composed of a variety of repeated cystein-rich modules, including an 8-cysteine modular structure found only in fibrillins and LTBPs <sup>12</sup>. Some, but not all, of the 8-cystein modules can mediate binding to the LAP.TGF $\beta$ 1 complex <sup>13,14</sup>. While

LTBP1, LTBP3, and to a lesser extent LTBP4 were reported to form a covalent complex with LAP.TGF $\beta$ 1, LTBP2 and the fibrillins do not<sup>15</sup>. In addition, the LTBPs and fibrillins contain at least one hybrid cysteine-rich domain, which can mediate intermolecular disulfide bonding and may hence play an important role in microfibril assembly<sup>16</sup>. LTBP1, and presumably other LTBP molecules, is transglutaminase cross-linked to the extracellular matrix via its amino-terminal region<sup>7</sup>. This model allows for fine temporal and spatial regulation of TGF $\beta$  sequestering or release in the extracellular matrix. LTBP1, LTBP2 and LTBP4, interact through their carboxy-terminal region with FBN1<sup>7,17</sup>. LTBP2 also interacts through its amino-terminal domain with fibulin-5 suggesting a role for LTBP2 in the regulation of elastic fiber assembly<sup>18</sup>. In spite of recent advances, the biology of fibrillins and related molecules, e.g., the extent of the redundancy between molecules of the LTBP-fibrillin superfamily, remains incompletely understood. Recently LTBP2 null mutations were reported in primary congenital glaucoma (PCG)<sup>19,20</sup>. We here report biallelic null LTBP2 mutations in patients with eye features consisting primarily of microspherophakia, ectopia lentis, megalocornea and myopia, and some features of MFS in older children.

## **PATIENTS**

### **Family 1**

The proband (VI.4), a boy, was first examined in the ophthalmology clinic at 18 months of age because of impaired vision. Spherophakia with infero-nasal displacement of both lenses was observed. The cornea was abnormally large (14mm diameter in both eyes), with flat irides and iridodonesis. There was no nystagmus, and the eye pressure was normal (16mm Hg in both eyes). There was axial myopia (Right Eye (RE) 26.1mm, Left Eye (LE) 27.2mm at 6 years). Bilateral phakophagia was performed at age 2 years for the lens luxation (Figure 1, D). Retinal detachment presented in both eyes 4 to 5 years later and ocular hypertension developed (up to 30mm Hg). A proband's sister (VI.7) was referred at age 10 months because of megalocornea. She had good fixation and no nystagmus, but megalocornea (14.5mm diameter), spherophakia, iridodonesis and axial myopia were present. She had very deep anterior chamber, and miotic oval pupil with no well defined border. Eye pressure was normal. Biometry performed at age 5 years showed an increased axial length of the globe (RE 27.0mm, LE 28.1mm), a deep anterior chamber (RE 3.9mm, LE 4.4mm), and increased lens diameter (RE 5.5mm, LE 5.6mm). Posterior lens dislocation was complete at age 6 years (Figure 1, B). Ocular hypertony was noticed at age 8 years. The third affected child in this family (VI.8), another sister of the proband, had good fixation and no nystagmus, but presented at age 18 months with axial myopia, megalocornea (14.5mm diameter) (Figure 1, C), iridodonesis, iris anomalies (anterior synechiae and iris mamillations), deep anterior chambers and spherophakia with no displacement of the lenses. Eye pressure was normal. At 5 years of age intermittent anterior dislocation of the lens in the left eye was observed, causing acute glaucoma crises and the left lens was extracted in emergency. Left retinal detachment developed a few weeks after. Progressive posterior luxation of the right lens, and bilateral ocular hypertony was observed.

The family was referred to our Genetics department when the proband was 14 years of age. The parents were first cousins, of Moroccan descent. The proband's height was 184cm, with an increased arm span of 204cm, and a decreased upper to lower body ratio (83cm/101cm) (Figure 1, A). The hands were normal, with relatively long fingers (Figure 1, E-F), albeit no frank arachnodactyly as the Walker sign (wrist sign) and Steinberg sign (thumb sign) were not present. The face was elongated. The palate was high-arched. The upper-to-lower body ratio was normal in both sisters with similar ocular abnormalities. The skin was normal on physical examination, with no striae distensae. The teeth and gums were unremarkable. The parents presented no features of MFS or Weill-Marchesani syndrome. Ophthalmologic evaluation was normal in both parents and in the other six siblings. Of note, the proband and elder affected sister were in a special education school, and their IQ were just at or below normal (75 and 68 respectively). The younger affected sister had no systematic evaluation of mental capacities because psychomotor development and education were normal. Intelligence was normal in all other members of the family. A heart ultrasound showed a normal heart with normal valves and a normal aortic root, in the proband and both affected sisters. Respiratory function tests were normal in the proband. No increased level of homocystine was found in the urine. A skin biopsy was performed for fibroblast culture. Marfan syndrome was considered and direct analysis of the *FBN1* gene was performed, but no mutation was identified. The clinical parameters are shown in Table 1.

## **Family 2**

We examined a second family of Macedonian descent with two children, one boy in good health and one affected girl. She was referred at age 26 months because of megalocornea. On ophthalmic examination the diameter of her corneas was 14mm bilaterally. Slit lamp examination revealed a few patches of iris atrophy with iris transillumination and a very flat anterior chamber. The latter was associated with an anterior and superior dislocation of the lens. The lenses were abnormally round (spherophakia). The fundus and the eye pressure were normal. Biometry

showed normal axial length of 21.91mm (RE) and 22.5mm (LE) (normal values for this age 21,63mm +/- 0,77). Keratometry was within normal range (mean of 42.5 D RE and 43.37 D LE). Her refraction was hypermetropic with astigmatism in the right +16 (+7 135°) and in the left eye +12 (+7 45°). An iridectomy was performed but the anterior chamber remained very shallow. A clear lens was extracted in order to prevent acute angle closure glaucoma and to improve vision. At age 3 years, she didn't so far develop glaucoma. At 2 ½ years, her arm span was 87cm, with a height of 88cm and a sitting height of 51cm, giving normal arm span/height ratio and sitting height/height ratio for age. The hands were unremarkable (Figure 1, G). The face was normal. The palate was high-arched. The skin was normal, with no striae distensae. The teeth and gums were unremarkable. Psychomotor development was normal (sitting at 5 months, walking at 1 year, speaking of two languages). A heart ultrasound showed a normal heart with normal valves and a normal aortic root. The clinical parameters are shown in Table 1.



## METHODS

A peripheral blood sample was obtained for genetic analysis with informed consent from the parents, and from the patients and unaffected siblings when old enough to consent, according to our study protocol as approved by the ethical committee of ULB – Hôpital Erasme. Control DNA subjects from Moroccan and Caucasian origin were recruited from our Genetic Centre and from a cohort of volunteers recruited in Occupational Medicine Department of our institution. Leukocyte DNA was extracted by the standard phenol-chloroform method<sup>21</sup>. DNA samples were purified using the QiaAmp DNA microkit and resuspended in Tris-HCl buffer. A genome-wide search for homozygosity was performed in the affected siblings of family 1 using an Affymetrix 10K GeneChip® microarrays of 10,000 autosomal single nucleotide polymorphisms (Affymetrix UK Ltd, High Wycombe, United Kingdom) following the instructions of the manufacturer. Regions of homozygosity were delineated using the Excludear algorithm<sup>22</sup>. In chromosomal regions with apparent homozygosity by descent, additional microsatellites were genotyped in individual subjects. Marker order was obtained from the Cooperative Human Linkage Center genetic map (<http://gai.nci.nih.gov/CHLC/>) and from the University of California at Santa Cruz (UCSC) physical map (<http://genome.ucsc.edu/cgi-bin/hgGateway>). DNA was amplified by PCR using 15 ng of DNA from each individual in a final volume of 15 µl, followed by polyacrylamide gel electrophoresis and silver staining. A multipoint LOD score was computed using the MAPMAKER/HOMOZ software postulating a gene frequency of 0.005 and marker allele frequencies as observed in a series of control subjects, with a minimal minor allele frequency of 0.10. The 36 coding exons and intron/exon junctions of LTBP2 were amplified using primers designed by the ExonPrimer Perl script engine ® (Munich, Germany). Polymerase chain reaction products were purified using the Exosap-IT kit ® (USB, Cleveland, OH) and directly sequenced by the Big Dye Terminator cycle sequencing kit v2 ® (Applied Biosystems, Foster City, CA). Sequencing products were purified on Egde Biosystems Performa DTR Ultra 96-Well

Plate Kit columns ® (Sopachem, Eke, Belgium) and analyzed on a 3130 Genetic Analyzer sequencing machine (Applied Biosystems). Sequences were inspected *in silico* for mutations using the SeqScape software version 2.0 ® (Applied Biosystems).

Skin fibroblasts of the proband were primary cultured using standard procedures. The cells were then lysed in TRIzol reagent (Invitrogen) and the cellular RNA was extracted by a conventional phenol-chloroform method. The RNA was reverse-transcribed in single-stranded cDNA using the SuperScript First Strand Synthesis System kit for RT-PCR (Invitrogen). The second cDNA strand was synthesized and a PCR was performed using Taq DNA polymerase (Invitrogen), in order to amplify a 625 bp fragment of *LTBP2* cDNA with direct primer situated in exon 7 and reverse primer in exon 12 (Direct primer sequence: ACAGCAACAACATCCCTGC; Reverse primer sequence: GCGTAGGTGTAGCCGTGG) . The amplified products were detected by running a 1% agarose gel with Lambda DNA/Hind III and Phi X 174 DNA/Hae III Mix molecular weight marker (NEB). The gels were stained in ethidium bromide, observed under UV transilluminator and then photographed .

## RESULTS

### **Family 1**

The initial homozygosity mapping performed in the three patients of the first family using the 10K GeneChip identified two large homozygous segments, on 9q31.3-q33.2 and on 14q23.3-24.3. We re-analyzed these regions with microsatellite markers in all family members. This showed that the 9q31.3-q33.2 region was also homozygous in an asymptomatic brother, with normal habitus and in whom ophthalmologic examination was normal. Assuming complete penetrance of the trait, we hence rejected linkage to chromosome 9 and considered the 14q region only (Figure 2). A maximal multipoint LOD score of 2.87 at D14S1002 was calculated using the MAPMAKER/HOMOZ software. The 14q linkage segment contained 199 annotated genes. Computer-assisted in silico inspection using the Endeavour software program <sup>23</sup>, indicated *LTBP2* gene as a strong candidate. Direct sequencing of *LTBP2*, identified a homozygous mutation in the proband and both affected sisters, consisting of a one-base pair insertion in exon 9, causing a frameshift responsible for a premature stop codon (c.1796dupC; p.Val600GlyfsX2). Both parents were heterozygous for the mutation, and no unaffected sibling was homozygous. The mutation was not found in 100 Moroccan control subjects (Figure 3). A fragment of the *LTBP2* mRNA encompassing exons 7 through 12, i.e., surrounding the frameshift mutation and premature stop codon in exon 9, was amplified from the proband's cultured skin fibroblasts by RT-PCR. A band of the expected size (625 bp) was observed in unrelated controls. This band was absent in the patient, and no other band was detected (Figure 3). This result is consistent with nonsense-mediated mRNA decay.

### **Family 2**

In the second family, direct sequencing of the *LTBP2* gene identified a homozygous non-sense mutation in exon 4 (c.895C>T; p.Arg299X). Both parents were heterozygous carriers of the mutation, which was absent from 100 Caucasian controls (Figure 3).

## DISCUSSION

We found biallelic *LTBP2* null mutations in two unrelated families ascertained via the ophthalmology clinic. Megalocornea and microspherophakia with early ectopia lentis probably secondary to the small round lens appears to be the most specific findings in these families. Megalocornea and microspherophakia may be encountered in congenital Marfan syndrome<sup>3,4</sup>, but the incidence in the total Marfan population is probably very low. Megalocornea was present in all *LTBP2* mutated patients reported to date<sup>19,20</sup>. However, the patients in this report presented no corneal clouding, contrary to the patients previously reported<sup>19,20</sup>. Megalocornea with microspherophakia is a condition that must be considered in the differential diagnosis of PCG<sup>24</sup>. Glaucoma may be congenital or may develop during childhood. The pathogenesis of the glaucoma is incompletely understood and normal gonioscopy can be found with megalocornea<sup>25</sup>. Recurrent attacks of pupillary block may lead to the development of glaucoma at very young age and may therefore be clinically misdiagnosed as PCG. Glaucoma in our patients developed later than previously reported<sup>19,20</sup>: after the age of 5 years in Family 1, not present at 3 years in Family 2. The intraocular pressures were normal in all patients at initial evaluation. Glaucoma was thus secondary, not primary nor congenital. Microspherophakia was identified by byometry and by biomicroscopy (after pupil dilatation the edge (equator) of each crystalline lens was easily visible with slit lamp). Spherophakia and iridodonesis were present already at the age of 10 months, and dislocation developed in all 8 eyes beginning as early as the age of 2 years, with inferonasal and backward displacement, producing an image of broken zonule, as in homocystinuria in Family 1. The displacement was anterosuperior in Family 2. Complete posterior lens dislocation occurred in 3/8 eyes.

Some signs of increased growth of the long bones were seen in older patients. All had a high-arched palate. The proband in Family 1 had a height of 184cm at age 14 years which contrasted with the heights of his mother and father (165cm and 172cm), a slender habitus, and an arm span

of 204cm, giving an arm span/height ratio of 1.11 (Normal <1.05). An arm span greater than height was present in his 8 years-old sister, perhaps indicating a progression of increased linear growth over time. Thus, some MFS features could be part of the LTBP2 defect-associated phenotypic spectrum. While ectopia lentis and marfanoid habitus were found in some of the recently reported patients, including tall stature, an arm span greater than height, and arachnodactyly, no detailed data were provided <sup>19,20</sup>. Interestingly, none of the patients had cardiovascular symptoms, and all had a normal cardiac ultrasound. The skin did not show striae distensae. Therefore, we found no evidence of intense elastolysis and defective tissue repair as reported in MFS <sup>9</sup>. This is in line with in vitro data showing that LTBP2 is not able to form a covalent complex with LAP.TGFβ1 <sup>15</sup>.

LTBP2 is expressed in the trabecular meshwork and ciliary processes, with a high expression in the lens capsule/lens epithelium layer in the adult mouse and cow eye <sup>19</sup> and in the human eye <sup>20</sup>. Isolated, non-syndromic familial ectopia lentis has been reported as an autosomal recessive trait associated with ADAMTSL4 mutations <sup>26</sup>, or as an autosomal dominant trait associated with *FBN1* mutations <sup>27</sup>. Ectopia lentis is reported in MFS, and in Weill-Marchesani syndrome (MIM277600), a rare connective tissue disorder characterized by short stature, brachydactyly, joint stiffness, and lens abnormalities, associated with ADAMTSL10 gene mutations <sup>28</sup>. It is also a feature of homocystinuria (MIM236200) and of sulfocysteinuria (MIM272300). The two latter disorders suggest that ectopia lentis may result from a chemical alteration of disulfide bonds in the microfibrillar network of the ocular zonule that are essential for holding the lens in a dynamic equilibrium. Taken together, we speculate that LTBP2 and FBN1 might have partially redundant structural functions in the zonule, with additional modelling by ADAMTSL10 and possibly other proteases.

The zonule and the periodontal ligament share the feature of being rich in microfibrils and devoid of elastin <sup>29</sup>. Therefore we thoroughly examined the teeth, periodontal sulci and gingivae in our patients, but found no abnormality.

The two elder affected sibs in family 1 had an IQ just at or below the normal range. It is presently unclear whether mental or developmental delay co-segregates with the *LTBP2* defect, as the third affected child in family 1 as well as the patient in family 2 had normal milestones .

Mutation c.1796dupC (p.Val600GlyfsX2) found in Moroccan Family 1 affects the hybrid domain and has not been reported before. No Moroccan patients with *LTBP2* mutations have been reported to date <sup>19,20</sup>. Mutation c.895C>T (p.Arg299X) found in family 2 was recently described in PCG patients from Pakistan and in European Gypsies<sup>19</sup>. The parents in Family 2 are Macedonian Gypsies, not known to be related. The region where they originated from (Skopje and Chtip) encompasses gypsy minorities ([http://en.wikipedia.org/wiki/Roma\\_in\\_the\\_Republic\\_of\\_Macedonia](http://en.wikipedia.org/wiki/Roma_in_the_Republic_of_Macedonia)). This finding suggests that the p.R299X *LTBP2* defect may represent an ancient founder mutation and may present either as PCG or as megalocornea, spherophakia, and secondary glaucoma.

Both mutations are predicted to produce truncated proteins. To date, 5 of the 6 *LTBP2* mutations reported, and the mutation p.Val600GlyfsX2 affecting the hybrid domain, are located in the amino-terminal part of the protein, interfering with both fibrillin-1 and fibulin-5 binding <sup>19,20</sup>. So far, only one mutation has been described affecting only its carboxy-terminal domain, suggesting that interference with fibulin-5 alone can be deleterious <sup>20</sup>. The relatively mild phenotype of these null mutations in man contrasts with in utero lethality of *ltbp2* <sup>-/-</sup> mice, that die between embryonic day 3.5 (E3.5) and E6.5 <sup>29</sup>. *LTBP2* is a very large, highly repetitive molecule, and many of its modules are encoded by single, in-frame exons <sup>11</sup>. No evidence of exon skipping was found in family 1 (Figure 3).

In summary, we here report homozygous null mutations of *LTBP2* in two unrelated families from different ethnic backgrounds, associated with an autosomal recessive syndrome of microspherophakia and progressive ectopia lentis, myopia and no primary congenital glaucoma, as well as some Marfan features in the older patient. Glaucoma developed after the age of 5 years in the first family. We suggest that intraocular pressure should be followed-up in young children with megalocornea, spherophakia and/or lens dislocation. *LTBP2* gene analysis should be offered to these patients.

## **ACKNOWLEDGMENTS**

We thank JF. Laes, Biovallee, Gosselies, Belgium, for initial analysis of the 10K Affymetrix GeneChips, Z. Ajarchouh for expert technical work, and G. Vassart for support. We are indebted to P. Coucke and B. Loeys, Gent, for FBN1 gene sequencing and useful discussion. We thank V. Cormier-Daire, Paris, for interest and discussion. JD is a fellow of the Belgian Kids' Fund and of the Erasme Fund for Scientific Research. MJA is supported by grant # 3.4593.07 of the Fonds de la Recherche Scientifique Médicale (FRSM), Belgium.



## REFERENCES

- 1 De Paepe A, Devereux RB, Dietz HC, Hennekam RC, Pyeritz RE: Revised diagnostic criteria for the Marfan syndrome. *Am J Med Genet* 1996; **62**: 417-426.
- 2 Faivre L, Masurel-Paulet A, Collod-Beroud G, et al.: Clinical and molecular study of 320 children with Marfan syndrome and related type I fibrillinopathies in a series of 1009 probands with pathogenic FBN1 mutations. *Pediatrics* 2009; **123**: 391-398.
- 3 Meire FM, Delleman WJ, Bleeker-Wagemakers EM: Ocular manifestations of congenital Marfan syndrome with contractures (CMC syndrome). *Ophthalmic Paediatr Genet* 1991; **12**: 1-9.
- 4 Meire FM, Van EJ, Hanssens M: Congenital Marfan syndrome with contractures. A clinicopathological report. *Bull Soc Belge Ophtalmol* 1992; **245**:91-97.
- 5 Ramirez F and Dietz HC: Fibrillin-rich microfibrils: Structural determinants of morphogenetic and homeostatic events. *J Cell Physiol* 2007; **213**: 326-330.
- 6 Dietz HC, Cutting GR, Pyeritz RE, et al.: Marfan syndrome caused by a recurrent de novo missense mutation in the fibrillin gene. *Nature* 1991; **352**: 337-339.
- 7 Isogai Z, Ono RN, Ushiro S, et al.: Latent transforming growth factor beta-binding protein 1 interacts with fibrillin and is a microfibril-associated protein. *J Biol Chem* 2003; **278**: 2750-2757.
- 8 Gregory KE, Ono RN, Charbonneau NL, Kuo CL, Keene DR, Bachinger HP, Sakai LY: The prodomain of BMP-7 targets the BMP-7 complex to the extracellular matrix. *J Biol Chem* 2005; **280**: 27970-27980.

- 9 Neptune ER, Frischmeyer PA, Arking DE, Myers L, Bunton TE, Gayraud B, Ramirez F, Sakai LY, Dietz HC: Dysregulation of TGF-beta activation contributes to pathogenesis in Marfan syndrome. *Nat Genet* 2003; **33**: 407-411.
- 10 Habashi JP, Judge DP, Holm TM, et al.: Losartan, an AT1 antagonist, prevents aortic aneurysm in a mouse model of Marfan syndrome. *Science* 2006; **312**: 117-121.
- 11 Massague J: The TGF-beta family of growth and differentiation factors. *Cell* 1987; **49**: 437-438.
- 12 Robinson PN, Arteaga-Solis E, Baldock C, et al.: The molecular genetics of Marfan syndrome and related disorders. *J Med Genet* 2006; **43**: 769-787.
- 13 Gleizes PE, Beavis RC, Mazzieri R, Shen B, Rifkin DB: Identification and characterization of an eight-cysteine repeat of the latent transforming growth factor-beta binding protein-1 that mediates bonding to the latent transforming growth factor-beta1. *J Biol Chem* 1996; **271**: 29891-29896.
- 14 Saharinen J, Taipale J, Keski-Oja J: Association of the small latent transforming growth factor-beta with an eight cysteine repeat of its binding protein LTBP-1. *EMBO J* 1996; **15**: 245-253.
- 15 Saharinen J and Keski-Oja J: Specific sequence motif of 8-Cys repeats of TGF-beta binding proteins, LTBP1s, creates a hydrophobic interaction surface for binding of small latent TGF-beta. *Mol Biol Cell* 2000; **11**: 2691-2704.
- 16 Reinhardt DP, Gambie JE, Ono RN, Bachinger HP, Sakai LY: Initial steps in assembly of microfibrils. Formation of disulfide-cross-linked multimers containing fibrillin-1. *J Biol Chem* 2000; **275**: 2205-2210.

- 17 Hirani R, Hanssen E, Gibson MA: LTBP-2 specifically interacts with the amino-terminal region of fibrillin-1 and competes with LTBP-1 for binding to this microfibrillar protein. *Matrix Biol* 2007; **26**: 213-223.
- 18 Hirai M, Horiguchi M, Ohbayashi T, Kita T, Chien KR, Nakamura T: Latent TGF-beta-binding protein 2 binds to DANCE/fibulin-5 and regulates elastic fiber assembly. *EMBO J* 2007; **26**: 3283-3295.
- 19 Ali M, McKibbin M, Booth A, et al.: Null mutations in LTBP2 cause primary congenital glaucoma. *Am J Hum Genet* 2009; **84**: 664-671.
- 20 Narooie-Nejad M, Paylakhi SH, Shojae S, et al.: Loss of function mutations in the gene encoding latent transforming growth factor beta binding protein 2, LTBP2, cause primary congenital glaucoma. *Hum Mol Genet* 2009; **18**: 3969-3977.
- 21 Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual. 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory, 1989
- 22 Woods CG, Valente EM, Bond J, Roberts E: A new method for autozygosity mapping using single nucleotide polymorphisms (SNPs) and EXCLUDEAR. *J Med Genet* 2004; **41**: e101
- 23 Aerts S, Lambrechts D, Maity S, et al.: Gene prioritization through genomic data fusion. *Nat Biotechnol* 2006; **24**: 537-544.
- 24 Roche O, Dureau P, Uteza Y, Dufier JL: Congenital megalocornea. *J Fr Ophtalmol* 2002; **25**: 312-318.
- 25 Meire FM: Megalocornea. Clinical and genetic aspects. *Doc Ophthalmol* 1994; **87**: 1-121.

- 26 Ahram D, Sato TS, Kohilan A, et al.: A homozygous mutation in ADAMTSL4 causes autosomal-recessive isolated ectopia lentis. *Am J Hum Genet* 2009; **84**: 274-278.
- 27 Ades LC, Holman KJ, Brett MS, Edwards MJ, Bennetts B: Ectopia lentis phenotypes and the FBN1 gene. *Am J Med Genet A* 2004; **126A**: 284-289.
- 28 Dagoneau N, Benoist-Lasselin C, Huber C, et al.: ADAMTS10 mutations in autosomal recessive Weill-Marchesani syndrome. *Am J Hum Genet* 2004; **75**: 801-806.
- 29 Shipley JM, Mecham RP, Maus E, et al.: Developmental expression of latent transforming growth factor beta binding protein 2 and its requirement early in mouse development. *Mol Cell Biol* 2000; **20**: 4879-4887.

## FIGURE LEGENDS

### 1. Clinical findings

A. Proband (VI.4) of the first family, whose height was 184cm and arm span 204cm at age 14 years. B. Ultrasound scan of the eye of patient VI.7 showing complete posterior luxation of the lens, as well as increased sphericity and short size of the lens. C. Eye of younger affected daughter (VI.8) of family 1 showing megalocornea. D. Eye of proband (V4) showing inferonasal dislocation of the lens. E,F,G. Hands of patients VI.7 and VI.8 from the first family and of proband from second family.

### 2. Linkage analysis in family 1

The parents were second cousins. The 14q linkage region was initially identified by a genome-wide 10K GeneChip microarray of SNPs in the three affected sibling, then further analyzed by microsatellite polymorphisms in affected and unaffected siblings and parents, as shown. A single linkage region was found where all patients and no unaffected siblings were homozygous, yielding a maximal multipoint LOD score of 2.87 at D14S1002.

### 3. LTBP2 gene sequencing and mRNA analysis



**A.** Family 1. Top to bottom, direct sequencing of genomic DNA from father, proband, and an ethnically matched unrelated control, showing c.1796\_1797insC (p.Val600GlyfsX2) mutation in exon 9 of *LTBP2*, heterozygous in father and homozygous in proband. **B.** Family 2. Top to bottom, direct sequencing of genomic DNA from father, proband, and an ethnically matched unrelated control, showing c.895C>T (p.Arg299X) mutation in exon 4 of *LTBP2*, heterozygous in father and homozygous in proband. **C.** Analysis of *LTBP2* transcript in cultured fibroblasts using primers situated in exons 7 and 12. 1-3: RT-PCR of three aliquots of mRNA extracted from fibroblasts culture of Family 1's proband; 4-5: RT-PCR of two unrelated control fibroblast cultures, showing a band of expected size (625 bp); 6: Negative control with H<sub>2</sub>O used in place

of mRNA template; 7: Lambda DNA/Hind III and Phi X 174 DNA/Hae III Mix molecular weight marker.


**TABLE 1. Clinical parameters**

<b>Clinical features</b>	<b>Family I ; VI.4</b>	<b>Family I ; VI.7</b>	<b>Family I ; VI.8</b>	<b>Family II</b>
<b>Megalocornea (N&lt;13mm)</b>	14/14	15/15	14.5/14.5	14/14
<b>Lens dislocation</b>	Nasal/nasal	Inferonasal/ inferonasal	Inferonasal/ inferonasal	Anterosuperior/ anterosuperior
<b>Microspherophakia</b>	+/+	+/+	+/+	+/+
<b>Axial length of eye (mm)</b>	26.1/27.2	27.0/28.1	22.0/24.0	21.91/22.5
<b>Retinal detachment</b>	+/+	-/-	-/+	-/-
<b>Glaucoma (age at onset)</b>	6y	8y	5y	No glaucoma at 3y
<b>Visual acuity</b>	5/10/1/20	4/10/4/10	1/20/2/10	Not evaluated <sup>1</sup>
<b>Height (cm) (age)</b>	184 (14y)	138 (9y)	117.5 (5y)	88 (2y 4/12)
<b>U/L or SH/H <sup>2</sup></b>	-2SD	+0.5SD	-1SD	-1SD
<b>High-arched palate</b>	+	+	+	+
<b>Aortic root ectasia</b>	-	-	-	-
<b>Cardiac valvular anomalies</b>	-	-	-	-
<b>Cognitive defect (IQ)</b>	At limit (75)	Mild (68)	Not evaluated, normal development	Not evaluated, normal development

1. The child was not compliant to test visual acuity, but she looked objects at 10 cms of her eyes before lens was extracted
2. U/L: Upper/Lower body ratio; SH/H: Sitting Height/Height







D14S592 11  
 D14S981 11  
 D14S1069 12  
 D14S588 12  
 D14S251 12  
 D14S1002 11  
 D14S1025 12  
 D14S53 12

12  
 11  
 23  
 12  
 12  
 11  
 23  
 13

cM  
 D14S592  
 3.01 D14S981  
 3.21 D14S1069  
 2.58 D14S588  
 1.34 D14S251  
 1.45 D14S1002  
 4.16 D14S1025  
 3.73 D14S53

12	12	11	12	12	11	12
11	11	11	11	11	11	11
13	23	22	23	13	22	22
11	12	22	12	11	22	22
11	12	22	12	11	22	22
11	11	11	11	11	11	11
13	23	22	23	13	22	22
23	11	11	13	13	13	11

