



**HAL**  
open science

## Excited-state non-adiabatic dynamics simulations of pyrrole

Hans Lischka, Mario Barbatti, Mario Vazdar, Mirjana Eckert-Maksic

► **To cite this version:**

Hans Lischka, Mario Barbatti, Mario Vazdar, Mirjana Eckert-Maksic. Excited-state non-adiabatic dynamics simulations of pyrrole. *Molecular Physics*, 2009, 107 (08-12), pp.845-854. 10.1080/00268970802665639 . hal-00513244

**HAL Id: hal-00513244**

**<https://hal.science/hal-00513244v1>**

Submitted on 1 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Excited-state non-adiabatic dynamics simulations of pyrrole**

Journal:	<i>Molecular Physics</i>
Manuscript ID:	TMPH-2008-0381.R1
Manuscript Type:	Special Issue Paper - Fritz Schaefer
Date Submitted by the Author:	27-Nov-2008
Complete List of Authors:	Lischka, Hans; University of Vienna, Institute for theoretical Chemistry Barbatti, Mario; University of Vienna, Institute for Theoretical Chemistry Vazdar, Mario; Rudjer Bošković Institute Eckert-Maksic, Mirjana; Rudjer Bošković Institute
Keywords:	non-adiabatic dynamics, conical intersection, photochemistry



# Excited-state non-adiabatic dynamics simulations of pyrrole

Mario Vazdar,<sup>a</sup> Mirjana Eckert-Maksić,<sup>a\*</sup> Mario Barbatti,<sup>b\*</sup> Hans Lischka<sup>b\*</sup>

<sup>a</sup> *Laboratory for Physical-Organic Chemistry – Division of Organic Chemistry and Biochemistry. Rudjer Bošković Institute, 10002 Zagreb, Croatia;* <sup>b</sup> *Institute for Theoretical Chemistry – University of Vienna, Waehringerstrasse 17, A 1090 Vienna, Austria.*

## Abstract

Non-adiabatic on-the-fly-dynamics simulations of the photodynamics of pyrrole were performed at multireference configuration interaction level involving five electronic states with a simulation time of 200 fs. The analysis of the time dependence of the average state occupations shows that the deactivation of pyrrole to the electronic ground state takes place in about 140 fs. This deactivation time agrees very well with the experimentally measured time constant of 110 fs for the formation of fast hydrogen atoms. After excitation into the S<sub>4</sub> state, 80% of the trajectories followed the NH-stretching mechanism giving rise to a population of fast H atoms. The computed average kinetic energy is in good accord with the experimentally observed average kinetic energy of the fast hydrogen atoms. It is found that 10% of trajectories followed the ring-puckering mechanism and 3% followed the ring-opening mechanism. This latter mechanism was characterized in pyrrole for the first time and involves the conical intersection of lowest energy of this molecule.

**Keywords:** non-adiabatic dynamics; conical intersection; photochemistry; pyrrole

\* Corresponding authors: H. Lischka (hans.lischka@univie.ac.at), M. Barbatti (mario.barbatti@univie.ac.at) and M. Eckert-Maksić (mmaksic@emma.irb.hr)

## 1. Introduction

Pyrrole is one of the simplest biologically relevant heteroaromatic compounds. For this reason, its electronic states have been intensively studied both experimentally and theoretically during the last decades with the emphasis on its UV spectrum [1-10] and its photodynamics [11-24]. In particular, it is known that the deactivation of UV-excited pyrrole to the ground state occurs at a very short (femtosecond) time scale [19] with low luminescence quantum yields [25], indicating the dominance of internal conversion processes. Sobolewski and coworkers [13] have proposed that the deactivation of pyrrole and related heteroatomic compounds occurs via the NH-stretching mechanism along a  ${}^1\pi\sigma_{\text{NH}}^*$  repulsive state. This mechanism, which has been examined in detail [15, 16, 22, 23] by means of wave packet dynamics simulations, can fully explain the presence of fast H atoms in the photofragmentation spectra [26]. Nevertheless, the mechanism responsible for the formation of slow H atoms and of other experimentally observed fragments such as HCN and  $\text{CNH}_2$  [11, 14, 17, 19, 20] is still subject of considerable debate [15, 19, 26-28].

Recently, we have suggested [27] that non-adiabatic deactivation of pyrrole may also proceed via a ring-puckering mechanism. This second kind of mechanism could not only be the source of heavy fragments, but also partially explain the slow H atoms [29]. Also recently, a third deactivation mechanism that can be relevant for pyrrole was identified in thiophene [30], furan [31], imidazole [32], and in the imidazole group of adenine [33]. In this mechanism the deactivation of five-membered rings proceeds by a planar ring-opening deformation. This process was observed to occur in a minor fraction of trajectories during dynamics simulations of adenine [34]. Based on these findings, we have currently attempted and succeeded to locate this type of mechanism in pyrrole, too.

1  
2  
3 Profant *et al.* [35] and Poterya *et al.* [28] have experimentally investigated the  
4 photolysis of pyrrole clusters. In addition, they have also performed theoretical calculations  
5 on the isolated pyrrole and on pyrrole complexes [28]. They have found that in presence of  
6 solvent molecules the NH stretching mechanism is inhibited, which results in a strong  
7 reduction of the fast H atom elimination process while keeping the slow H atom elimination.  
8 These are important results that on one hand once more confirm the role of the NH-stretching  
9 mechanism for the fast H atoms formation and on the other hand indicate that ring  
10 deformation mechanisms should be involved in the slow H atoms formation.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

22 The strong dependence of different fragment yields on the excitation energy [26]  
23 indicates that the individual mechanisms are in mutual competition and can play different  
24 roles depending on the initial conditions. Excited-state energy surfaces have been investigated  
25 in detail under special consideration of crossings between different energy surfaces as already  
26 mentioned above [1, 18, 23, 27, 28, 36] and reaction paths have been constructed  
27 subsequently. This information led to substantial progress in the understanding of the  
28 photochemical processes in pyrrole. However, it turned out to be very difficult to estimate the  
29 importance of individual intersections and related reaction pathways. In order to better  
30 understand how these mechanisms are activated, it is desirable to perform dynamics  
31 simulations. Such simulations exhibit a substantial complexity. For instance, as for selecting  
32 the proper quantum chemical methods, it needs to be taken into account that: first, the non-  
33 adiabatic dynamics of pyrrole involves multiple excited states showing often multireference  
34 character and, secondly, that it is essentially impossible to identify just a few important  
35 internal degrees of freedom by which the photochemical reaction mechanism can be  
36 described. Therefore, an essential condition is the usage of the full set of nuclear coordinates.  
37 These are usual requirements to be met e.g. in simulations of organic chromophores  
38 exhibiting high density of excited states [34, 37, 38]. One convenient way to satisfy especially  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 the second condition is to use mixed-quantum classical dynamics methods [39-44]. In this  
4  
5 work surface hopping dynamics is performed using the fewest-switches algorithm of Tully  
6  
7 [45]. The advantage of this approach is that it allows the application of an “on-the-fly”  
8  
9 strategy [42, 43, 46] where a pre-selection of certain internal degrees of freedom and any  
10  
11 fitting of pre-computed potential energy points is avoided by computing at each time step the  
12  
13 energies, the complete energy gradient and non-adiabatic coupling terms required for the  
14  
15 integration of Newton’s equations of motion and the time-dependent Schrödinger equation.  
16  
17 This on-the-fly strategy is computationally very expensive and requires analytical energy  
18  
19 gradients and non-adiabatic coupling vectors for computational efficiency. Due to the  
20  
21 stringent computational requirements most of the photodynamical simulations have been  
22  
23 performed so far at the relatively cost-effective complete active space self consistent field  
24  
25 level (CASSCF). Since in this case dynamical electron correlation effects are mostly  
26  
27 neglected, the relative balance of electronic states of different character can be strongly  
28  
29 violated. It should be stressed that the non-adiabatic dynamics simulations presented here  
30  
31 were carried out at a significantly higher level using the MR-CISD method including five  
32  
33 electronic states. This represents the state-of-the-art approach for this kind of simulations,  
34  
35 which has not been documented before for molecules of the size of pyrrole to the best of our  
36  
37 knowledge. The present calculations have been made possible by use of the analytic gradient  
38  
39 features of the program package COLUMBUS [47-49] as it will be described below.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 **2. Computational details**

51  
52 Multireference configuration interaction (MRCI) and complete active space self-consistent  
53  
54 field (CASSCF) calculations were performed for pyrrole. The CAS space was comprised of  
55  
56 four  $\pi$  electrons in five orbitals (two  $\pi$  orbitals, two  $\pi^*$  orbitals and one Rydberg 3s orbital).  
57  
58 This space will be conventionally designated as CAS(4,5) in the text. State averaging was  
59  
60

1  
2  
3 performed over five singlet states with equal weights (ground state, two valence  $\pi\pi^*$  states  
4 and two Rydberg  $\pi 3s$  states), which will be denoted as SA-5. MRCI calculations were  
5 performed based on the orbitals computed by the SA-5-CASSCF(4,5) wave function. The  
6 reference configurations for the MRCI were constructed within the CAS(4,5) by allowing  
7 single and double excitations from the two  $\pi$  orbitals into the two  $\pi^*$  orbitals and the Rydberg  
8 3s orbital. The final configuration space was constructed by allowing all single and double  
9 excitations from the reference configurations into the virtual orbital space (MR-CISD). All  
10 core electrons and the lowest eight additional orbitals were frozen in the MRCI calculations  
11 and the interacting space restriction [50] was applied. The basis set was composed of aug'-cc-  
12 pVDZ type [51] on the nitrogen and carbon atoms (the prime indicates that d-aug functions  
13 were removed). On the hydrogen atom connected to nitrogen, the cc-pVDZ basis set was  
14 used, whereas for the remaining hydrogen atoms the cc'-pVDZ basis set was used (the prime  
15 signifies that p-functions were deleted). This hybrid basis set will be denoted as BS.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35 The MRCI approach and the basis set were selected by balancing the accuracy  
36 requirements of the calculations of four excited states of different character (see Table 1) and  
37 the need for computational efficiency, since an on-the-fly approach requires several tens of  
38 thousands of individual MRCI calculations to be carried out. Therefore, before starting the  
39 dynamics simulations an extensive set of calculations had been performed, including the  
40 Franck-Condon region, the seam of conical intersections, and reaction pathways. For the  
41 determination of minima on the crossing seam (MXS), starting geometries were selected from  
42 our previous MRCI calculations on pyrrole [27] and were reoptimized with the above-  
43 described MRCI method. Reaction paths for the two ring-deformation processes were  
44 constructed by the method of linear interpolation of internal coordinates (LIIC) between the  
45 ground-state geometry and the corresponding ring-deformed conical intersections. The  
46 reaction path for the NH-stretching process was constructed by rigidly stretching the NH  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 distance in steps of 0.2 Å starting from the ground state equilibrium structure up to a NH  
4  
5 distance of 2.6 Å.  
6  
7

8  
9 All energy calculations and MXS optimizations were performed by using analytical  
10 gradient and non-adiabatic coupling procedures described in References [52-56]. For vertical  
11 excitation energy calculations, the Davidson correction (+Q) [53, 57, 58] was used in order to  
12 describe higher order excitation effects. For the  $C_{2v}$  labeling of the states, the  $x$  axis was  
13 assumed to be oriented perpendicular to the ring plane.  
14  
15  
16  
17  
18  
19

20  
21 Mixed quantum-classical dynamics calculations were performed for pyrrole by using  
22 an on-the-fly approach [42, 43, 46, 59, 60]. Energies, gradients, and non-adiabatic coupling  
23 vectors were computed at each time step at the MR-CISD/SA-5-CASSCF(4,5)/BS level of  
24 theory. The nuclear motion was represented by classical trajectories computed by numerical  
25 integration of Newton's equations by the velocity-Verlet algorithm [61]. Non-adiabatic effects  
26 were taken into account by means of the surface hopping approach [45]. Time-dependent  
27 adiabatic populations were corrected for decoherence effects [62] ( $\alpha = 0.1$  hartree) and used to  
28 calculate surface hopping probabilities in accordance to the Tully's fewest switches approach  
29 [45]. In order to alleviate the computational costs, no coupling vectors were calculated  
30 between non-consecutive states [44]. In total, 90 trajectories were computed. The initial  
31 Cartesian coordinates and momenta were selected from a quantum harmonic oscillator  
32 (Wigner) distribution in the ground state. The trajectories were started in the  $S_4$  state at these  
33 geometries. This procedure gave rise to a composition of 60% of trajectories initially in the  
34  $\pi\pi^*$  states and 40% in the  $\pi 3s/\pi\sigma_{NH}^*$  states. The minimum excitation energy was 6.36 eV  
35 while the average was 6.76 eV with a standard deviation of 0.26 eV. The trajectories were  
36 then propagated for a maximum time of 200 fs with a time step of 0.5 fs.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The structures of the puckered geometries were described in terms of the Cremer-Pople parameters  $Q$  and  $\phi$  [63]. While the parameter  $Q$  measures the extent of puckering ( $Q =$



0 Å indicates a planar structure), the parameter  $\phi$  describes the kind of puckering. For 5-membered rings, there are only few kinds of puckered conformations available: envelope conformations with atom  $k$  above ( ${}^kE$ ) or below ( $E_k$ ) the ring plane and twisted conformations with atom  $k$  above the ring plane and atom  $k-1$  below the ring plane ( ${}^kT_{k-1}$ ). Because of the pyrrole symmetry,  $\phi$  can be reduced to the  $0^\circ - 90^\circ$  range by projecting all values on this quadrant.

All CASSCF and MR-CISD+Q calculations were performed with the COLUMBUS [47-49] program package. The atomic orbital (AO) integrals and AO gradient integrals have been calculated with program modules taken from DALTON [64]. The dynamic simulations were carried out using the NEWTON-X program [42, 65] with an interface to the COLUMBUS program package.

Table 1. (around here)

### 3. Analysis of the energy surfaces

In order to investigate the reliability of the MRCI method used in the dynamics study, we have performed a series of tests and comparisons with other previously published results. Specifically, we have compared vertical excitation energies, reaction paths, and MXS structures with results obtained with methods of higher level of theory.

#### 3.1 Vertical excitation energies

The theoretical computation of vertical excitation energies of pyrrole and the assignment of the experimental UV spectrum have been a matter of discussion for a long period of time [1, 3, 4, 8, 9]. The currently calculated values are compared to other available theoretical and experimental results in Table 1. The comparison reveals that vertical excitation energies computed by the MR-CISD/SA-5-CASSCF(4,5)/BS method are in good accordance with

1  
2  
3 results obtained previously by the MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ  
4 method [27] where the auxiliary (AUX) orbital represents the 3s Rydberg orbital into which  
5 single excitations from the valence CAS(6,5) are allowed. Most of the calculated vertical  
6 excitation energies differ by ca. 0.1-0.2 eV, except in the case of the  $^1B_1$  state where this  
7 difference is 0.35 eV. Furthermore, the present results for the  $^1A_2$  and  $^1B_1$  Rydberg states are  
8 in excellent agreement with experimental values assigned in Ref. [9]. The current energies of  
9 the  $\pi\pi^*$  valence states are higher than in most of the other methods with the deviation being  
10 particularly large in comparison to the CASPT2 results. Nevertheless, a series of different  
11 methods, like MRCI [2, 4], EOM-CCSD [8], CC3 [3], and TDDFT [4] indicates that CASPT2  
12 might be underestimating these transition energies. Therefore, we conclude that the current  
13 MRCI approach is adequate for calculation of vertical excitation energies.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Fig. 1 (around here)

### 3.2 Conical intersections

In Fig. 1 the MXS structures between ground state and the  $S_1$  state are presented. The comparison of selected geometrical parameters for the ring-puckered (Fig. 1a) and the NH-stretched (Fig. 1b) MXS structures reveals that they are in very good agreement with the benchmark MRCI values [27].

In Fig. 1a, the MXS between the valence  $\pi\pi^*$  state and the ground state shows an out-of-plane deformation with strong stretching of one of the CN bonds. We shall refer to this conical intersection as the ring-puckered MXS. The values of dihedral CCCN and CCCH dihedral angles are very close to the benchmark ones, being only by ca.  $2^\circ$  smaller. The length of the broken CN bond is 1.607 Å, thus being 0.007 Å shorter than the value obtained with the benchmark method. In Fig. 1b, the NH-stretched MXS is shown. It arises from the crossing

1  
2  
3 between the ground state and the lowest  $\pi\sigma_{\text{NH}}^*$  state. In comparison to the benchmark MRCI  
4 value, the NH distance using the current method is shorter by 0.007 Å.  
5  
6  
7

8  
9 As mentioned in the Introduction, based on previous findings for other five-membered  
10 heteroaromatic molecules [30, 31, 33], we have searched for a planar ring-opened MXS in  
11 pyrrole as well. The optimized structure, obtained at the MRCI level of theory, is presented in  
12 Fig. 1c. It should be pointed out that the MXS is planar and that the CN distance is 2.512 Å,  
13 which is by about 0.9 Å longer than the CN distance observed in the ring puckered MXS (Fig.  
14 1a). It is important to note that the ring-opened MXS is the lowest energy conical intersection  
15 identified in pyrrole so far and it arises from the crossing between the  $\pi\sigma_{\text{NC}}^*$  state and the  
16 ground state.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 Table 2. (around here)  
29  
30

31 Although the similarity of geometrical parameters suggests that the selected MR-  
32 CISD/SA-5-CASSCF(4,5)/BS level of theory is adequate, it is also of importance to compare  
33 the energies of the MXSs. MRCI and MRCI+Q energy values of pyrrole MXSs obtained by  
34 the MR-CISD(Q)/SA-5-CASSCF(4,5)/BS and benchmark MRCI values [27] are summarized  
35 in Table 2. The analysis of presented data shows that the energies of the MXSs are in very  
36 good agreement with the benchmark ones. The comparison among results reveals that the  
37 selected MRCI method is well suited for the description of both ring-puckering and NH-  
38 stretching mechanisms. In particular, the current MRCI and MRCI+Q energies of the ring-  
39 puckered MXS are by 0.06 eV higher and 0.07 eV lower than the benchmark MRCI and  
40 MRCI+Q values, respectively. For the NH-stretched MXS, the MRCI energy is by 0.04 eV  
41 higher, whereas the MRCI+Q value is by 0.18 eV lower than the benchmark values.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 3.3 Reaction paths

We have computed the reaction pathways between the ground state minimum and the three MXSs described in the previous section using the MR-CISD/SA-5-CASSCF(4,5)/BS level of theory. The resulting potential energy curves are shown in Fig. 2.

Fig. 2 (around here)

Comparison with the results obtained with the MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ method (Fig. 2 in [27]) reveals that NH-stretching potential energy curves (Fig. 2a) agree very well for the ground state  $S_0$  and the two Rydberg  $^1A_2$  and  $^1B_1$  states for the whole range of NH distances. The main difference is that the crossing between the lowest Rydberg state and the ground state occurs at around 1.9 Å in the present work instead of 2.1 Å found in our earlier study [27]. The other features of the potential energy curves exhibit the same behavior as observed earlier. Specifically, the lowest two Rydberg states show small energy barriers (0.24 eV for the  $^1A_2$  state and 0.12 for the  $^1B_1$  state) at the NH distance of 1.2 Å necessary to transform the  $\pi 3s$  orbital into the  $\pi\sigma_{\text{NH}}^*$  state as expected for a stretching of the NH bond. It should also be pointed out that the two valence  $^1A_1$  and  $^1B_2$  states show the same energy profile until the NH distance of 1.8 Å. After that, an intrusion of higher excited states occurs (not shown), which is presumably a direct consequence of the CAS(4,5) active space. However, in the NH stretching mechanism, the deactivation occurs via conical intersections among Rydberg states and the ground state and the differences in the valence states for large NH distances are of minor importance.

Fig. 2b shows that the LIIC path of the ring-puckering mechanism in the  $\pi\pi^*$  state occurs without barrier. Indeed, it is clearly seen that the lowest  $\pi\pi^*$  state is diabatically connected to the ground state, which may make it especially efficient for the internal conversion. The same result was observed in our previous study [27] thus providing additional

1  
2  
3 support for using the applied method. In the case of the ring-opening mechanism Fig. 2c  
4  
5 shows that the initially excited  $\pi\pi^*$  states can deactivate without barrier along this pathway.  
6  
7  
8 The character of the state should, however, change into  $\pi\sigma_{\text{NC}}^*$  in order to lead to the crossing  
9  
10 with the ground state.  
11

12  
13 Apart from the fact that the NH-stretching mechanism should dominate at low  
14  
15 excitation energies, it is difficult to draw general conclusions about the efficiency of each  
16  
17 mechanism based on the reaction paths alone in a clear cut way. When the excitation leads  
18  
19 into the spectral region of the  $\pi\pi^*$  state all mechanisms are energetically possible. In favor of  
20  
21 the NH-stretching mechanism is the fact that it requires the smallest deformations from the  
22  
23 Franck-Condon region in terms of mass-weighted distances (see Fig. 2). On the other hand, it  
24  
25 also requires the diabatic transformation from the  $\pi\pi^*$  state into the  $\pi\sigma_{\text{NH}}^*$  state, which  
26  
27 depends upon the activation of out-of-plane modes [18, 27]. The ring-opening mechanism  
28  
29 involves the lowest energy conical intersection, but it requires the largest deformations from  
30  
31 the Franck-Condon region and diabatic changes in the wave function at the same time.  
32  
33 Finally, the ring-puckering mechanism, as already mentioned, can directly proceed through a  
34  
35 diabatic connection. However, it involves the highest energy portions of the seam of conical  
36  
37 intersections.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

#### 4. Dynamics simulations of pyrrole

49  
50 The non-adiabatic excited state dynamics of pyrrole was started from the  $S_4$  state thus making  
51  
52 all pathways discussed in the previous section energetically available. The resulting average  
53  
54 adiabatic populations of the ground and excited states as a function of time are presented in  
55  
56 Fig. 3. Their analysis shows that the  $S_4$  state transfers its population to the  $S_3$  state in the first  
57  
58 10 fs. After ca. 50 fs, the  $S_4$  state is almost completely depopulated. The populations of  $S_3$  and  
59  
60  $S_2$  states reach a maximum at 10 fs and 20 fs, respectively. At about 75 fs, these states are

1  
2  
3 already depopulated. The  $S_2$  state shows a repopulation between 100 and 150 fs. The  
4  
5 population of the  $S_1$  state increases reaching a maximum at 75 fs. At 100 fs, the  $S_1$  and  $S_0$   
6  
7 states have approximately the same population. Between 100 fs and 200 fs, the simulation is  
8  
9 basically reduced to the  $S_1/S_0$  two-state dynamics, with the complete population transferred to  
10  
11 the ground state at about 200 fs.  
12  
13

14  
15  
16 Fig. 3 (around here)  
17

18 The  $S_1$  population shows a consecutive two-step first order decay type of behavior. By  
19  
20 fitting the  $S_1$  population curve with the function  
21

$$22 \quad f(t) = \frac{\tau_2}{\tau_2 - \tau_1} \left[ \exp\left(-\frac{t}{\tau_1}\right) - \exp\left(-\frac{t}{\tau_2}\right) \right], \quad (1)$$

23  
24  
25  
26  
27  
28  
29 two time constants  $\tau_1 = 44 \pm 2$  fs and  $\tau_2 = 80 \pm 2$  fs are obtained. Here,  $\tau_1$  measures the  
30  
31 population of  $S_1$  from the collection of states  $S_4$  to  $S_2$  and  $\tau_2$  describes the depopulation  
32  
33  $S_1 \rightarrow S_0$ . The approximate time constant for the overall population of the ground state can be  
34  
35 obtained by fitting the  $S_0$  population with the function  
36  
37

$$38 \quad f(t) = 1 - \exp\left(-\frac{t}{\tau_0}\right), \quad (2)$$

39  
40  
41  
42  
43  
44  
45 which gives  $\tau_0 = 139 \pm 2$  fs. Note that in these three time constants the error bars denote the  
46  
47 uncertainty of the fitting procedure and not of the process itself, which certainly is larger than  
48  
49 a few femtoseconds.  
50  
51

52  
53 In Fig. 4 a summary of the results of the dynamics simulation in terms of the fraction  
54  
55 of trajectories following each of the three mechanisms is given. The NH-stretching is the main  
56  
57 mechanism after excitation of pyrrole to the  $S_4$  state. This mechanism occurs in 80% of the  
58  
59 trajectories. Other 13% follow ring-deformation mechanisms (ring-opening and ring-  
60

1  
2  
3 puckering). 7% do not deactivate within the 200 fs of the dynamics simulation. Because of the  
4  
5  
6 uncertainties associated to the dynamics simulations and to the relatively small number of  
7  
8 trajectories, these fractions should be taken as qualitative trends of occurrence of each  
9  
10 mechanism, rather than a quantitative assessment of them. If trajectories starting in the  $\pi\pi^*$   
11  
12 and in the  $\pi\sigma_{\text{NH}}^*$  states are independently analyzed, these fractions remain essentially the  
13  
14 same, implying that the population of each mechanism depends on the excitation energy, but  
15  
16 not on the nature of the state. The fact that the fast H atom is formed along the NH stretching  
17  
18 pathway either excited in the  $\pi\pi^*$  or  $\pi\sigma^*$  states has also been observed in the photofragment  
19  
20 translational spectroscopy studies by Cronin et al. [26].  
21  
22  
23

24  
25 Experimental pump of pyrrole with 250 nm (4.96 eV) laser pulse followed by  
26  
27 ionization probe with 241 nm (5.15 eV) pulse reveals two time constants,  $\tau_f = 110 \pm 80$  fs and  
28  
29  $\tau_s = 1100 \pm 500$  fs [19]. These time constants correspond to the time for formation of fast and  
30  
31 slow H atoms, respectively. Since most of trajectories in our simulations finished in the  
32  
33 ground state of the dissociated pyrrolyl + H system, the deactivation time  $\tau_0$  should also  
34  
35 approximately give the time for the formation of the fast H atoms population. Indeed, the  
36  
37 comparison of  $\tau_0$  and  $\tau_f$  shows good agreement. Note, however, that the initial state in the  
38  
39 experiments (low energy  $\pi\sigma_{\text{NH}}^*$ ) and in the simulations (high energy  $\pi\pi^*$  and  $\pi\sigma_{\text{NH}}^*$ ) are not  
40  
41 the same. This is an indication that the fast H elimination occurs directly by the same process,  
42  
43 as soon as there is enough energy to overcome the  $\pi 3s/\pi\sigma_{\text{NH}}^*$  barrier in the  $S_1$  state.  
44  
45  
46  
47  
48  
49

50  
51 Fig. 4 (around here)

52  
53 Fig. 5 (around here)

54  
55  
56 The analysis of NH and CN bond distances was conducted for all trajectories and the  
57  
58 results are presented in Fig. 5. The top panel of this figure shows that in some cases the CN  
59  
60 distance is elongating during the dynamics. This behavior can be ascribed to the ring-opening

1  
2  
3 and ring-puckering deactivation mechanisms. Since the main deactivation channel is the NH-  
4 stretching, the majority of trajectories do not exhibit elongation of this specific bond. In the  
5  
6 stretching, the majority of trajectories do not exhibit elongation of this specific bond. In the  
7  
8 bottom panel of Fig. 5 the NH distance is monitored. In this figure three kinds of trajectories  
9  
10 can be distinguished. For part of the trajectories the NH distance remain constant at about 1  
11  
12 Å. They correspond to the trajectories following ring-distortion mechanisms. A minor fraction  
13  
14 of trajectories (3) has the NH distance oscillating at a medium distance of about 2 or 3 Å.  
15  
16 These are cases where the NH-stretching mechanism is activated, but instead finishing in  
17  
18 dissociation, the hot ground state of pyrrole is formed. In most of the trajectories the NH  
19  
20 distance is steadily increasing. In these cases, the NH-stretching mechanism is activated and  
21  
22 the H atom elimination is taking place. It should be mentioned that a cut-off value of 10 Å for  
23  
24 NH distance was used in Fig. 5 in order to simplify the data analysis. In some of the  
25  
26 trajectories, however, the NH distance was longer, up to 40 Å.  
27  
28  
29  
30  
31

32 Fig. 6 (around here)

33  
34  
35 Fig. 6a shows that the hydrogen dissociation starts on average 54 fs after the  
36  
37 photoexcitation. The kinetic energy of the dissociated hydrogen atom has a broad distribution  
38  
39 around the average value of 1.2 eV (Fig. 6b). This value is in very good agreement with the  
40  
41 experimental results, ~1 eV [11, 26], for the center of the fast H-elimination peak in the  
42  
43 kinetic energy release spectra. Note that, as expected, there is no formation of a slow H-  
44  
45 elimination peak, which should take place in the picosecond timescale [19], much longer than  
46  
47 the maximum simulation time (200 fs). The NH distance at the  $S_1 \rightarrow S_0$  hopping time is shown  
48  
49 in Fig. 6c for all trajectories that have returned to the ground state. The histogram shows two  
50  
51 distinct peaks. The first peak with average at 1.0 Å will be discussed below. The second peak  
52  
53 starts at 1.5 Å and presents a long tail for large distances up to 4 Å. This peak corresponds to  
54  
55 the trajectories deactivated by means of the NH-stretching mechanism. Its average value at  
56  
57  
58  
59  
60



1  
2  
3 2.1 Å is 0.2 Å larger than the NH distance for the crossing between the lowest  $\pi\sigma^*$  state and  
4  
5  
6 the ground state shown in Fig. 2 (left panel).  
7

8  
9 Fig. 7 (around here)  
10

11 Twelve out of ninety trajectories did not follow the NH-stretching mechanism. They  
12 appear in the short-distance peak in Fig. 6c. In order to understand which kind of mechanism  
13 they followed, it is useful to project them on the Cremer-Pople (CP) space  $Q-\phi$ . This is shown  
14 in Fig. 7 for all structures for which the  $S_1-S_0$  energy gap is smaller than 0.5 eV (open dots)  
15 and for structures at the hopping time (full dots). The ring-opened MXS is at  $Q = 0$  Å and the  
16 ring-puckered MXS is shown by a cross ( $E_1$  conformation). Since the ring-opened and the  
17 ring-puckered conical intersections correspond to distinct types of structures on the crossing  
18 seam with different electronic configurations, it could be expected that the structures at the  
19 hopping time would cluster in two disjoint regions around these MXSs. This, however, is not  
20 the case. Fig. 7 shows that the non-adiabatic events occur in a large continuous portion of the  
21 CP space, indicating that the crossing seam spans this entire region. The degree of puckering  
22 varies from almost planar ( $Q = 0.15$  Å) to the strongly puckered structures ( $Q = 0.75$  Å). Most  
23 of hopping events occur at  $E_1$ ,  ${}^2T_1$  and  ${}^2E$  conformations, indicating that not only the  $E_1$   
24 conformation of the ring-puckered MXS, but also other kinds of puckering can give rise to  
25 conical intersections in pyrrole.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 If we take  $Q = 0.3$  Å as an arbitrary threshold to distinguish between the ring-opening  
48 and ring-puckering mechanisms, nine trajectories deactivated at ring-puckered conformations  
49 and three trajectories deactivated at ring-opened conformations, thus corresponding to 10%  
50 and 3% of the total number of trajectories, respectively (see Fig. 4).  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 5. Conclusions

The photochemical processes in pyrrole were investigated using a high-level multireference configuration interaction method (MRCI) giving a balanced description of the four studied excited states, two of Rydberg character and two valence states. Cuts along the potential energy surfaces connecting the Franck-Condon region and three different minima on the crossing seam (MXS) (NH dissociation, ring puckering, and a planar ring-opened MXS) describe possible deactivation pathways. One of these intersection points, the ring-opened MXS, was characterized for the first time. Although it is the conical intersection of the lowest energy identified in pyrrole so far, its efficiency for the internal conversion process seems to be reduced by the required strong geometric deformations and by the diabatic change of the initially excited  $\pi\pi^*$  state into the  $\pi\sigma_{\text{NC}}^*$  state, which in turn crosses the ground state.

Non-adiabatic surface-hopping dynamics simulations of pyrrole were performed for 200 fs starting in the  $S_4$  state and using a high-level MR-CI approach for the electronic structure calculations. The dynamics simulations show that in fact all three types of conical intersections were accessed. The transfer of population from the initially excited  $S_4$  state to the ground state takes place in about 140 fs. This process occurs basically in two steps, with the  $S_1$  state being populated in about 44 fs and then being depleted in about 80 fs. Most of trajectories (80%) dissociated rapidly along the repulsive  $\pi\sigma_{\text{NH}}^*$  state giving rise to a population of fast H atoms. The computed deactivation time of 140 fs agrees very well with the experimentally measured time constant of 110 fs for the formation of fast hydrogen atoms. The computed average kinetic energy agrees very well with the experimentally observed average kinetic energy of the fast hydrogen atoms. A fraction of 13% of trajectories follows ring-deformation channels involving either ring puckering (10%) or planar ring opening (3%). These fractions did not depend on whether the initial state had  $\pi\pi^*$  or  $\pi\sigma_{\text{NH}}^*$  character.

1  
2  
3 Our calculations provide a detailed picture of the photodeactivation processes in  
4 pyrrole. Although the main objective of this work – the observation of the occurrence of the  
5  
6 different deactivation mechanisms – has been accomplished, it should be noted that the  
7  
8 participation of  $\pi\sigma_{\text{NH}}^*$  states in the initial conditions was much higher than what would be  
9  
10 expected from the oscillator strengths of these two transitions. This bias occurred because of  
11  
12 the relatively high vertical excitation energy of the  $^1\text{B}_1$  Rydberg state, which caused frequent  
13  
14 exchange of position with the  $\pi\pi^*$  in the Wigner sample. Interestingly, it turned out that the  
15  
16 observed percentages of the different mechanisms was insensitive to the initial character of  
17  
18  $\text{S}_4$ , consequently implying that that this bias is not so critical for the general interpretations.  
19  
20 Nevertheless, more investigations are needed to analyze the influence of excitation energies  
21  
22 on the product yields in order to explain the experimentally observed strong energy  
23  
24 dependence of the branching ratios for fast and slow hydrogen atoms.  
25  
26  
27  
28  
29  
30  
31  
32  
33

## 34 Acknowledgments

35  
36 This work was supported by the Austrian Science Fund within the framework of the Special  
37  
38 Research Program F16 (Advanced Light Sources) and Project P18411-N19. The calculations  
39  
40 were partially performed at the Linux PC cluster Schrödinger III of the computer center of the  
41  
42 University of Vienna. The work in Zagreb (M.E.M and M.V.) was supported by the Ministry  
43  
44 of Science, Education and Sport through the project 098-0982933-2920 and the COST D37  
45  
46 action.  
47  
48  
49  
50  
51  
52

## 53 References

- 54  
55  
56 [1] L. Serrano-Andrés, M. Merchán, I. Nebotgil, B. O. Roos, and M. Fulscher, J. Am. Chem.  
57  
58 Soc. **115**, 6184 (1993).  
59  
60 [2] M. H. Palmer, I. C. Walker, and M. F. Guest, Chem. Phys. **238**, 179 (1998).

- 1  
2  
3 [3] O. Christiansen, J. Gauss, J. F. Stanton, and P. Jorgensen, *J. Chem. Phys.* **111**, 525 (1999).  
4  
5 [4] D. J. Tozer, R. D. Amos, N. C. Handy, B. O. Roos, and L. Serrano-Andres, *Mol. Phys.*  
6 **97**, 859 (1999).  
7  
8 [5] J. Wan, J. Meller, M. Hada, M. Ehara, and H. Nakatsuji, *J. Chem. Phys.* **113**, 7853 (2000).  
9  
10 [6] B. O. Roos, P. A. Malmqvist, V. Molina, L. Serrano-Andres, and M. Merchan, *J. Chem.*  
11 *Phys.* **116**, 7526 (2002).  
12  
13 [7] C. G. Zhan, and D. A. Dixon, *J. Mol. Spectrosc.* **216**, 81 (2002).  
14  
15 [8] P. Celani, and H. J. Werner, *J. Chem. Phys.* **119**, 5044 (2003).  
16  
17 [9] M. H. Palmer, and P. J. Wilson, *Mol. Phys.* **101**, 2391 (2003).  
18  
19 [10] M. Pastore, C. Angeli, and R. Cimiraglia, *Chem. Phys. Lett.* **422**, 522 (2006).  
20  
21 [11] D. A. Blank, S. W. North, and Y. T. Lee, *Chem. Phys.* **187**, 35 (1994).  
22  
23 [12] A. B. Trofimov, H. Köppel, and J. Schirmer, *J. Chem. Phys.* **109**, 1025 (1998).  
24  
25 [13] A. L. Sobolewski, W. Domcke, C. Dedonder-Lardeux, and C. Jouvet, *PCCP* **4**, 1093  
26 (2002).  
27  
28 [14] J. Wei, A. Kuczmann, J. Riedel, F. Renth, and F. Temps, *PCCP* **5**, 315 (2003).  
29  
30 [15] V. Vallet, Z. G. Lan, S. Mahapatra, A. L. Sobolewski, and W. Domcke, *Faraday Discuss.*  
31 **127**, 283 (2004).  
32  
33 [16] V. Vallet, Z. G. Lan, S. Mahapatra, A. L. Sobolewski, and W. Domcke, *J. Chem. Phys.*  
34 **123** (2005).  
35  
36 [17] J. Wei, J. Riedel, A. Kuczmann, F. Renth, and F. Temps, *Faraday Discuss.* **127**, 267  
37 (2004).  
38  
39 [18] H. Köppel, E. V. Gromov, and A. B. Trofimov, *Chem. Phys.* **304**, 35 (2004).  
40  
41 [19] H. Lippert, H. H. Ritze, I. V. Hertel, and W. Radloff, *Chemphyschem* **5**, 1423 (2004).  
42  
43 [20] A. J. van den Brom, M. Kapelios, T. N. Kitsopoulos, N. H. Nahler, B. Cronin, and M. N.  
44 R. Ashfold, *PCCP* **7**, 892 (2005).  
45  
46 [21] I. Frank, and K. Damianos, *Journal of Chemical Physics* **126** (2007).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [22] Z. Lan, A. Dupays, V. Vallet, S. Mahapatra, and W. Domcke, *Journal of Photochemistry*  
4 and *Photobiology a-Chemistry* **190**, 177 (2007).  
5  
6  
7  
8 [23] Z. Lan, and W. Domcke, *Chem. Phys.* **350**, 125 (2008).  
9  
10 [24] A. Kumar, M. Kolaski, and K. S. Kim, *J. Chem. Phys.* **128** (2008).  
11  
12 [25] E. J. Shin, *Bull. Korean Chem. Soc.* **25**, 907 (2004).  
13  
14 [26] B. Cronin, M. G. D. Nix, R. H. Qadiri, and M. N. R. Ashfold, *PCCP* **6**, 5031 (2004).  
15  
16 [27] M. Barbatti, M. Vazdar, A. J. A. Aquino, M. Eckert-Maksic, and H. Lischka, *J. Chem.*  
17 *Phys.* **125**, 164323 (2006).  
18  
19 [28] V. Poterya, V. Profant, M. Farnik, P. Slavicek, and U. Buck, *J. Chem. Phys.* **127**, 064307  
20 (2007).  
21  
22 [29] M. Barbatti, B. Sellner, A. J. A. Aquino, and H. Lischka, in *Radiation Induced Molecular*  
23 *Phenomena in Nucleic Acid*, edited by M. K. Shukla, and J. Leszczynski (Springer, Netherlands,  
24 2008), pp. 209.  
25  
26 [30] S. Salzmann, M. Kleinschmidt, J. Tatchen, R. Weinkauff, and C. M. Marian, *PCCP* **10**, 380  
27 (2008).  
28  
29 [31] N. Gavrilov, S. Salzmann, and C. M. Marian, *Chem. Phys.* **349**, 269 (2008).  
30  
31 [32] M. Barbatti, H. Lischka, S. Salzmann, and C. M. Marian, *J. Chem. Phys.*, submitted  
32 (2008).  
33  
34 [33] S. Perun, A. L. Sobolewski, and W. Domcke, *Chem. Phys.* **313**, 107 (2005).  
35  
36 [34] M. Barbatti, and H. Lischka, *J. Am. Chem. Soc.* **130**, 6831 (2008).  
37  
38 [35] V. Profant, V. Poterya, M. Farnik, P. Slavicek, and U. Buck, *J. Phys. Chem. A* **111**, 12477  
39 (2007).  
40  
41 [36] A. L. Sobolewski, and W. Domcke, *Chem. Phys.* **259**, 181 (2000).  
42  
43 [37] I. Antol, M. Vazdar, M. Barbatti, and M. Eckert-Maksic, *Chem. Phys.* **349**, 308 (2008).  
44  
45 [38] M. Barbatti, M. Ruckebauer, J. J. Szymczak, A. J. A. Aquino, and H. Lischka, *PCCP* **10**,  
46 482 (2008).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [39] J. C. Tully, Faraday Discuss. **110**, 407 (1998).  
4  
5 [40] A. Ferretti, G. Granucci, A. Lami, M. Persico, and G. Villani, J. Chem. Phys. **104**, 5517  
6  
7 (1996).  
8  
9 [41] N. L. Doltsinis, and D. Marx, Phys. Rev. Lett. **88**, 166402 (2002).  
10  
11 [42] M. Barbatti, G. Granucci, M. Persico, M. Ruckebauer, M. Vazdar, M. Eckert-Maksic,  
12  
13 and H. Lischka, J. Photochem. Photobiol., A **190**, 228 (2007).  
14  
15 [43] E. Fabiano, T. W. Keal, and W. Thiel, Chem. Phys. **349**, 334 (2008).  
16  
17 [44] J. Pittner, H. Lischka, and M. Barbatti, Chem. Phys., doi:10.1016/j.chemphys.2008.10.013  
18  
19 (2008).  
20  
21 [45] J. C. Tully, J. Chem. Phys. **93**, 1061 (1990).  
22  
23 [46] L. Sun, and W. L. Hase, in *Reviews in Computational Chemistry*, edited by K. B. Lipkowitz *et*  
24  
25 *al.* (Wiley-VCH, New York, 2003), pp. 79.  
26  
27 [47] H. Lischka, R. Shepard, F. B. Brown, and I. Shavitt, Int. J. Quantum Chem. **S.15**, 91  
28  
29 (1981).  
30  
31 [48] H. Lischka, R. Shepard, R. M. Pitzer, I. Shavitt, M. Dallos, T. Müller, P. G. Szalay, M.  
32  
33 Seth, G. S. Kedziora, S. Yabushita, and Z. Y. Zhang, PCCP **3**, 664 (2001).  
34  
35 [49] H. Lischka, R. Shepard, I. Shavitt, R. M. Pitzer, M. Dallos, T. Mueller, P. G. Szalay, F. B.  
36  
37 Brown, R. Ahlrichs, H. J. Boehm, A. Chang, D. C. Comeau, R. Gdanitz, H. Dachsel, C. Ehrhardt,  
38  
39 M. Ernzerhof, P. Hoeschl, S. Irlle, G. Kedziora, T. Kovar, V. Parasuk, M. J. M. Pepper, P. Scharf,  
40  
41 H. Schiffer, M. Schindler, M. Schueler, M. Seth, E. A. Stahlberg, J.-G. Zhao, S. Yabushita, Z.  
42  
43 Zhang, M. Barbatti, S. Matsika, M. Schuurmann, D. R. Yarkony, S. R. Brozell, E. V. Beck, and J.-  
44  
45 P. Blaudeau, COLUMBUS, an ab initio electronic structure program, release 5.9.1,  
46  
47 [www.univie.ac.at/columbus](http://www.univie.ac.at/columbus) (2006).  
48  
49 [50] A. Bunge, J. Chem. Phys. **53**, 20 (1970).  
50  
51 [51] T. H. Dunning, J. Chem. Phys. **90**, 1007 (1989).  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [52] R. Shepard, H. Lischka, P. G. Szalay, T. Kovar, and M. Ernzerhof, *J. Chem. Phys.* **96**,  
4 2085 (1992).  
5  
6  
7  
8 [53] R. Shepard, in *Modern Electronic Structure Theory*, edited by D. R. Yarkony (World Scientific,  
9 Singapore, 1995), p. 345.  
10  
11  
12 [54] H. Lischka, M. Dallos, and R. Shepard, *Mol. Phys.* **100**, 1647 (2002).  
13  
14 [55] M. Dallos, H. Lischka, R. Shepard, D. R. Yarkony, and P. G. Szalay, *Journal of Chemical*  
15 *Physics* **120**, 7330 (2004).  
16  
17  
18 [56] H. Lischka, M. Dallos, P. G. Szalay, D. R. Yarkony, and R. Shepard, *Journal of Chemical*  
19 *Physics* **120**, 7322 (2004).  
20  
21  
22  
23 [57] S. R. Langhoff, and E. R. Davidson, *Int. J. Quantum Chem.* **8**, 61 (1974).  
24  
25 [58] P. J. Bruna, S. D. Peyerimhoff, and R. J. Buenker, *Chem. Phys. Lett.* **72**, 278 (1980).  
26  
27 [59] V. Bonacic-Koutecky, and R. Mitric, *Chem. Rev.* **105**, 11 (2005).  
28  
29 [60] R. Mitric, V. Bonacic-Koutecky, J. Pittner, and H. Lischka, *J. Chem. Phys.* **125** (2006).  
30  
31 [61] W. C. Swope, H. C. Andersen, P. H. Berens, and K. R. Wilson, *J. Chem. Phys.* **76**, 637  
32 (1982).  
33  
34 [62] G. Granucci, and M. Persico, *J. Chem. Phys.* **126**, 134114 (2007).  
35  
36 [63] D. Cremer, and J. A. Pople, *J. Am. Chem. Soc.* **97**, 1354 (1975).  
37  
38 [64] T. Helgaker, H. J. A. Jensen, P. Jørgensen, J. Olsen, K. Ruud, H. Ågren, T. Andersen, K.  
39 L. Bak, V. Bakken, O. Christiansen, P. Dahle, E. K. Dalskov, T. Enevoldsen, H. Heiberg, H.  
40 Hetta, D. Jonsson, S. Kirpekar, R. Kobayashi, H. Koch, K. V. Mikkelsen, P. Norman, M. J.  
41 Packer, T. Saue, P. R. Taylor, and O. Vahtras, DALTON, an ab initio electronic structure  
42 program, Release 1.0 (1997).  
43  
44  
45 [65] M. Barbatti, G. Granucci, M. Ruckebauer, M. Persico, and H. Lischka, NEWTON-X: a  
46 package for Newtonian dynamics close to the crossing seam, [www.univie.ac.at/newtonx](http://www.univie.ac.at/newtonx) (2007).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Tables

Table 1 – Vertical excitation energies of selected singlet states of pyrrole.

State	MRCI <sup>a</sup>	MRCI+Q <sup>b</sup>	TDDFT <sup>c</sup>	CASPT2	CC3 <sup>g</sup>	Exp
S <sub>0</sub> <sup>1</sup> A <sub>1</sub>	0.00	0.00	0.00	0.00	0.00	
π3s <sup>1</sup> A <sub>2</sub>	5.22	5.09	5.05	5.22 <sup>d</sup> /5.22 <sup>e</sup>	5.10	5.08 <sup>h</sup> /5.22 <sup>i</sup>
π3s <sup>1</sup> B <sub>1</sub>	6.21	5.86	5.88	5.85 <sup>f</sup> /5.92 <sup>e</sup>	5.99	6.22 <sup>h</sup>
ππ* <sup>1</sup> A <sub>1</sub>	6.55	6.39	6.29	5.82 <sup>d</sup> /5.98 <sup>e</sup>	6.37	
ππ* <sup>1</sup> B <sub>2</sub>	6.65	6.78	6.45	5.87 <sup>d</sup> /5.95 <sup>e</sup>	6.63	5.92 <sup>h</sup> /6.2-6.4 <sup>i</sup>

<sup>a</sup> Present results, MR-CISD/SA-5-CASSCF(4,5)/BS

<sup>b</sup> MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ, Reference [27].

<sup>c</sup> Reference [4].

<sup>d</sup> Reference [6].

<sup>e</sup> Reference [8].

<sup>f</sup> Reference [1].

<sup>g</sup> Reference [3].

<sup>h</sup> Assignments given in Reference [9].

<sup>i</sup> Assignments given in Reference [3].



Table 2 – Energy of pyrrole MXSs (in eV) relative to the minimum in the ground state.

MXS	MRCI <sup>a</sup>	MRCI+Q <sup>a</sup>	MRCI	MRCI+Q	MXS features
$\pi\pi^*/S_0$ ( $E_1$ )	4.95	4.86	4.89 <sup>b</sup>	4.93 <sup>b</sup>	ring puckering, Fig. 1a
$\pi\sigma_{\text{NH}}^*/S_0$	4.45	4.26	4.41 <sup>c</sup>	4.44 <sup>c</sup>	NH stretching, Fig. 1b
$\pi\sigma_{\text{NC}}^*/S_0$	4.11	3.86	-	-	ring opening, Fig. 1c

<sup>a</sup>Present results, MR-CISD/SA-5-CASSCF(4,5)/BS

<sup>b</sup>MR-CISD(Q)/SA-3-CAS(6,5)/6-31G(d), Reference [27].

<sup>c</sup>MR-CISD(Q)/SA-3-CAS(6,6)/6-31G(d), Reference [27].

For Peer Review Only

## Figure Captions

Fig. 1. Structures and selected geometric parameters for pyrrole MXSs obtained at the MRCI level. Distances are given in Å and dihedral angles in degrees. The number in brackets correspond to the benchmark MRCI value from Ref. [27].

Fig. 2. Potential energy curves calculated at the MRCI level along a) the rigid NH-stretching coordinate and along the LIIC path from the ground state minimum to b) the ring-puckered MXS and to c) the ring-opened MXS.

Fig. 3. Average adiabatic populations of trajectories for each state as a function of time after initial photoexcitation of pyrrole into the  $S_4$  state.


Fig. 4. Description and statistics of trajectory deactivation mechanisms.

Fig. 5. CN (top) and NH (bottom) distance variations as a function of time for all trajectories. The NH distance of 10 Å was used as a cut-off value (see text).


Fig. 6. Analysis of the trajectories showing NH dissociation. (a) Initial time of the dissociation, taking 2 Å for the NH bond as reference value. (b) Hydrogen kinetic energy. (c) NH distance for all trajectories at the time of the  $S_1 \rightarrow S_0$  hopping.

Fig. 7. Distribution of conformations in the Cremer-Pople  $Q-\phi$  space for trajectories following ring-deformation mechanisms. Full dots: conformations at the hopping time. Open dots: conformations with  $S_1-S_0$  energy gaps smaller than 0.5 eV. Cross: ring puckered MXS.


a) Ring-puckered MXS





c) Ring-opened MXS





b) NH-dissociation MXS













1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60







view Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Excited-state non-adiabatic dynamics simulations of pyrrole

Mario Vazdar,<sup>a</sup> Mirjana Eckert-Maksić,<sup>a\*</sup> Mario Barbatti,<sup>b\*</sup> Hans Lischka<sup>b\*</sup>

<sup>a</sup> *Laboratory for Physical-Organic Chemistry – Division of Organic Chemistry and Biochemistry, Rudjer Bošković Institute, 10002 Zagreb, Croatia;* <sup>b</sup> *Institute for Theoretical Chemistry – University of Vienna, Währingerstrasse 17, A 1090 Vienna, Austria.*

## Abstract

Non-adiabatic on-the-fly-dynamics simulations of the photodynamics of pyrrole were performed at multireference configuration interaction level involving five electronic states with a simulation time of 200 fs. The analysis of the time dependence of the average state occupations shows that the deactivation of pyrrole to the electronic ground state takes place in about 140 fs. This deactivation time agrees very well with the experimentally measured time constant of 110 fs for the formation of fast hydrogen atoms. After excitation into the S<sub>4</sub> state, 80% of the trajectories followed the NH-stretching mechanism giving rise to a population of fast H atoms. The computed average kinetic energy is in good accord with the experimentally observed average kinetic energy of the fast hydrogen atoms. It is found that 10% of trajectories followed the ring-puckering mechanism and 3% followed the ring-opening mechanism. This latter mechanism was characterized in pyrrole for the first time and involves the conical intersection of lowest energy of this molecule.

**Keywords:** non-adiabatic dynamics; conical intersection; photochemistry; pyrrole

\* Corresponding authors: H. Lischka (hans.lischka@univie.ac.at), M. Barbatti (mario.barbatti@univie.ac.at) and M. Eckert-Maksić (mmaksic@emma.irb.hr)

## 1. Introduction

Pyrrole is one of the simplest biologically relevant heteroaromatic compounds. For this reason, its electronic states have been intensively studied both experimentally and theoretically during the last decades with the emphasis on its UV spectrum [1-10] and its photodynamics [11-24]. In particular, it is known that the deactivation of UV-excited pyrrole to the ground state occurs at a very short (femtosecond) time scale [19] with low luminescence quantum yields [25], indicating the dominance of internal conversion processes. Sobolewski and coworkers [13] have proposed that the deactivation of pyrrole and related heteroatomic compounds occurs via the NH-stretching mechanism along a  ${}^1\pi\sigma_{\text{NH}}^*$  repulsive state. This mechanism, which has been examined in detail [15, 16, 22, 23] by means of wave packet dynamics simulations, can fully explain the presence of fast H atoms in the photofragmentation spectra [26]. Nevertheless, the mechanism responsible for the formation of slow H atoms and of other experimentally observed fragments such as HCN and  $\text{CNH}_2$  [11, 14, 17, 19, 20] is still subject of considerable debate [15, 19, 26-28].

Recently, we have suggested [27] that non-adiabatic deactivation of pyrrole may also proceed via a ring-puckering mechanism. This second kind of mechanism could not only be the source of heavy fragments, but also partially explain the slow H atoms [29]. Also recently, a third deactivation mechanism that can be relevant for pyrrole was identified in thiophene [30], furan [31], imidazole [32], and in the imidazole group of adenine [33]. In this mechanism the deactivation of five-membered rings proceeds by a planar ring-opening deformation. This process was observed to occur in a minor fraction of trajectories during dynamics simulations of adenine [34]. Based on these findings, we have currently attempted and succeeded to locate this type of mechanism in pyrrole, too.

1  
2  
3 Profant *et al.* [35] and Poterya *et al.* [28] have experimentally investigated the  
4 photolysis of pyrrole clusters. In addition, they have also performed theoretical calculations  
5 on the isolated pyrrole and on pyrrole complexes [28]. They have found that in presence of  
6 solvent molecules the NH stretching mechanism is inhibited, which results in a strong  
7 reduction of the fast H atom elimination process while keeping the slow H atom elimination.  
8 These are important results that on one hand once more confirm the role of the NH-stretching  
9 mechanism for the fast H atoms formation and on the other hand indicate that ring  
10 deformation mechanisms should be involved in the slow H atoms formation.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

22 The strong dependence of different fragment yields on the excitation energy [26]  
23 indicates that the individual mechanisms are in mutual competition and can play different  
24 roles depending on the initial conditions. Excited-state energy surfaces have been investigated  
25 in detail under special consideration of crossings between different energy surfaces as already  
26 mentioned above [1, 18, 23, 27, 28, 36] and reaction paths have been constructed  
27 subsequently. This information led to substantial progress in the understanding of the  
28 photochemical processes in pyrrole. However, it turned out to be very difficult to estimate the  
29 importance of individual intersections and related reaction pathways. In order to better  
30 understand how these mechanisms are activated, it is desirable to perform dynamics  
31 simulations. Such simulations exhibit a substantial complexity. For instance, as for selecting  
32 the proper quantum chemical methods, it needs to be taken into account that: first, the non-  
33 adiabatic dynamics of pyrrole involves multiple excited states showing often multireference  
34 character and, secondly, that it is essentially impossible to identify just a few important  
35 internal degrees of freedom by which the photochemical reaction mechanism can be  
36 described. Therefore, an essential condition is the usage of the full set of nuclear coordinates.  
37 These are usual requirements to be met e.g. in simulations of organic chromophores  
38 exhibiting high density of excited states [34, 37, 38]. One convenient way to satisfy especially  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 the second condition is to use mixed-quantum classical dynamics methods [39-44]. In this  
4  
5 work surface hopping dynamics is performed using the fewest-switches algorithm of Tully  
6  
7 [45]. The advantage of this approach is that it allows the application of an “on-the-fly”  
8  
9 strategy [42, 43, 46] where a pre-selection of certain internal degrees of freedom and any  
10  
11 fitting of pre-computed potential energy points is avoided by computing at each time step the  
12  
13 energies, the complete energy gradient and non-adiabatic coupling terms required for the  
14  
15 integration of Newton’s equations of motion and the time-dependent Schrödinger equation.  
16  
17 This on-the-fly strategy is computationally very expensive and requires analytical energy  
18  
19 gradients and non-adiabatic coupling vectors for computational efficiency. Due to the  
20  
21 stringent computational requirements most of the photodynamical simulations have been  
22  
23 performed so far at the relatively cost-effective complete active space self consistent field  
24  
25 level (CASSCF). Since in this case dynamical electron correlation effects are mostly  
26  
27 neglected, the relative balance of electronic states of different character can be strongly  
28  
29 violated. It should be stressed that the non-adiabatic dynamics simulations presented here  
30  
31 were carried out at a significantly higher level using the MR-CISD method including five  
32  
33 electronic states. This represents the state-of-the-art approach for this kind of simulations,  
34  
35 which has not been documented before for molecules of the size of pyrrole to the best of our  
36  
37 knowledge. The present calculations have been made possible by use of the analytic gradient  
38  
39 features of the program package COLUMBUS [47-49] as it will be described below.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 **2. Computational details**

51  
52 Multireference configuration interaction (MRCI) and complete active space self-consistent  
53  
54 field (CASSCF) calculations were performed for pyrrole. The CAS space was comprised of  
55  
56 four  $\pi$  electrons in five orbitals (two  $\pi$  orbitals, two  $\pi^*$  orbitals and one Rydberg 3s orbital).  
57  
58 This space will be conventionally designated as CAS(4,5) in the text. State averaging was  
59  
60

1  
2  
3 performed over five singlet states with equal weights (ground state, two valence  $\pi\pi^*$  states  
4 and two Rydberg  $\pi 3s$  states), which will be denoted as SA-5. MRCI calculations were  
5  
6 performed based on the orbitals computed by the SA-5-CASSCF(4,5) wave function. The  
7  
8 reference configurations for the MRCI were constructed within the CAS(4,5) by allowing  
9  
10 single and double excitations from the two  $\pi$  orbitals into the two  $\pi^*$  orbitals and the Rydberg  
11  
12  $3s$  orbital. The final configuration space was constructed by allowing all single and double  
13  
14 excitations from the reference configurations into the virtual orbital space (MR-CISD). All  
15  
16 core electrons and the lowest eight additional orbitals were frozen in the MRCI calculations  
17  
18 and the interacting space restriction [50] was applied. The basis set was composed of aug'-cc-  
19  
20 pVDZ type [51] on the nitrogen and carbon atoms (the prime indicates that d-aug functions  
21  
22 were removed). On the hydrogen atom connected to nitrogen, the cc-pVDZ basis set was  
23  
24 used, whereas for the remaining hydrogen atoms the cc'-pVDZ basis set was used (the prime  
25  
26 signifies that p-functions were deleted). This hybrid basis set will be denoted as BS.  
27  
28  
29  
30  
31  
32  
33

34  
35 The MRCI approach and the basis set were selected by balancing the accuracy  
36  
37 requirements of the calculations of four excited states of different character (see Table 1) and  
38  
39 the need for computational efficiency, since an on-the-fly approach requires several tens of  
40  
41 thousands of individual MRCI calculations to be carried out. Therefore, before starting the  
42  
43 dynamics simulations an extensive set of calculations had been performed, including the  
44  
45 Franck-Condon region, the seam of conical intersections, and reaction pathways. For the  
46  
47 determination of minima on the crossing seam (MXS), starting geometries were selected from  
48  
49 our previous MRCI calculations on pyrrole [27] and were reoptimized with the above-  
50  
51 described MRCI method. Reaction paths for the two ring-deformation processes were  
52  
53 constructed by the method of linear interpolation of internal coordinates (LIIC) between the  
54  
55 ground-state geometry and the corresponding ring-deformed conical intersections. The  
56  
57 reaction path for the NH-stretching process was constructed by rigidly stretching the NH  
58  
59  
60

1  
2  
3 distance in steps of 0.2 Å starting from the ground state equilibrium structure up to a NH  
4  
5 distance of 2.6 Å.  
6  
7

8 All energy calculations and MXS optimizations were performed by using analytical  
9  
10 gradient and non-adiabatic coupling procedures described in References [52-56]. For vertical  
11  
12 excitation energy calculations, the Davidson correction (+Q) [53, 57, 58] was used in order to  
13  
14 describe higher order excitation effects. For the  $C_{2v}$  labeling of the states, the  $x$  axis was  
15  
16 assumed to be oriented perpendicular to the ring plane.  
17  
18

19  
20 Mixed quantum-classical dynamics calculations were performed for pyrrole by using  
21  
22 an on-the-fly approach [42, 43, 46, 59, 60]. Energies, gradients, and non-adiabatic coupling  
23  
24 vectors were computed at each time step at the MR-CISD/SA-5-CASSCF(4,5)/BS level of  
25  
26 theory. The nuclear motion was represented by classical trajectories computed by numerical  
27  
28 integration of Newton's equations by the velocity-Verlet algorithm [61]. Non-adiabatic effects  
29  
30 were taken into account by means of the surface hopping approach [45]. Time-dependent  
31  
32 adiabatic populations were corrected for decoherence effects [62] ( $\alpha = 0.1$  hartree) and used to  
33  
34 calculate surface hopping probabilities in accordance to the Tully's fewest switches approach  
35  
36 [45]. In order to alleviate the computational costs, no coupling vectors were calculated  
37  
38 between non-consecutive states [44]. In total, 90 trajectories were computed. The initial  
39  
40 Cartesian coordinates and momenta were selected from a quantum harmonic oscillator  
41  
42 (Wigner) distribution in the ground state. The trajectories were started in the  $S_4$  state at these  
43  
44 geometries. This procedure gave rise to a composition of 60% of trajectories initially in the  
45  
46  $\pi\pi^*$  states and 40% in the  $\pi 3s/\pi\sigma_{NH}^*$  states. The minimum excitation energy was 6.36 eV  
47  
48 while the average was 6.76 eV with a standard deviation of 0.26 eV. The trajectories were  
49  
50 then propagated for a maximum time of 200 fs with a time step of 0.5 fs.  
51  
52  
53  
54  
55  
56  
57  
58

59 The structures of the puckered geometries were described in terms of the Cremer-  
60  
Pople parameters  $Q$  and  $\phi$  [63]. While the parameter  $Q$  measures the extent of puckering ( $Q =$

0 Å indicates a planar structure), the parameter  $\phi$  describes the kind of puckering. For 5-membered rings, there are only few kinds of puckered conformations available: envelope conformations with atom  $k$  above ( ${}^kE$ ) or below ( $E_k$ ) the ring plane and twisted conformations with atom  $k$  above the ring plane and atom  $k-1$  below the ring plane ( ${}^kT_{k-1}$ ). Because of the pyrrole symmetry,  $\phi$  can be reduced to the  $0^\circ - 90^\circ$  range by projecting all values on this quadrant.

All CASSCF and MR-CISD+Q calculations were performed with the COLUMBUS [47-49] program package. The atomic orbital (AO) integrals and AO gradient integrals have been calculated with program modules taken from DALTON [64]. The dynamic simulations were carried out using the NEWTON-X program [42, 65] with an interface to the COLUMBUS program package.

Table 1. (around here)

### 3. Analysis of the energy surfaces

In order to investigate the reliability of the MRCI method used in the dynamics study, we have performed a series of tests and comparisons with other previously published results. Specifically, we have compared vertical excitation energies, reaction paths, and MXS structures with results obtained with methods of higher level of theory.

#### 3.1 Vertical excitation energies

The theoretical computation of vertical excitation energies of pyrrole and the assignment of the experimental UV spectrum have been a matter of discussion for a long period of time [1, 3, 4, 8, 9]. The currently calculated values are compared to other available theoretical and experimental results in Table 1. The comparison reveals that vertical excitation energies computed by the MR-CISD/SA-5-CASSCF(4,5)/BS method are in good accordance with



1  
2  
3 results obtained previously by the MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ  
4 method [27] where the auxiliary (AUX) orbital represents the 3s Rydberg orbital into which  
5 single excitations from the valence CAS(6,5) are allowed. Most of the calculated vertical  
6 excitation energies differ by ca. 0.1-0.2 eV, except in the case of the  $^1B_1$  state where this  
7 difference is 0.35 eV. Furthermore, the present results for the  $^1A_2$  and  $^1B_1$  Rydberg states are  
8 in excellent agreement with experimental values assigned in Ref. [9]. The current energies of  
9 the  $\pi\pi^*$  valence states are higher than in most of the other methods with the deviation being  
10 particularly large in comparison to the CASPT2 results. Nevertheless, a series of different  
11 methods, like MRCI [2, 4], EOM-CCSD [8], CC3 [3], and TDDFT [4] indicates that CASPT2  
12 might be underestimating these transition energies. Therefore, we conclude that the current  
13 MRCI approach is adequate for calculation of vertical excitation energies.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

Fig. 1 (around here)

### 3.2 Conical intersections

33  
34  
35  
36  
37  
38 In Fig. 1 the MXS structures between ground state and the  $S_1$  state are presented. The  
39 comparison of selected geometrical parameters for the ring-puckered (Fig. 1a) and the NH-  
40 stretched (Fig. 1b) MXS structures reveals that they are in very good agreement with the  
41 benchmark MRCI values [27].  
42  
43  
44  
45  
46  
47  
48

49 In Fig. 1a, the MXS between the valence  $\pi\pi^*$  state and the ground state shows an out-of-  
50 plane deformation with strong stretching of one of the CN bonds. We shall refer to this  
51 conical intersection as the ring-puckered MXS. The values of dihedral CCCN and CCCH  
52 dihedral angles are very close to the benchmark ones, being only by ca.  $2^\circ$  smaller. The length  
53 of the broken CN bond is 1.607 Å, thus being 0.007 Å shorter than the value obtained with the  
54 benchmark method. In Fig. 1b, the NH-stretched MXS is shown. It arises from the crossing  
55  
56  
57  
58  
59  
60

1  
2  
3 between the ground state and the lowest  $\pi\sigma_{\text{NH}}^*$  state. In comparison to the benchmark MRCI  
4  
5 value, the NH distance using the current method is shorter by 0.007 Å.  
6  
7

8  
9 As mentioned in the Introduction, based on previous findings for other five-membered  
10  
11 heteroaromatic molecules [30, 31, 33], we have searched for a planar ring-opened MXS in  
12  
13 pyrrole as well. The optimized structure, obtained at the MRCI level of theory, is presented in  
14  
15 Fig. 1c. It should be pointed out that the MXS is planar and that the CN distance is 2.512 Å,  
16  
17 which is by about 0.9 Å longer than the CN distance observed in the ring puckered MXS (Fig.  
18  
19 1a). It is important to note that the ring-opened MXS is the lowest energy conical intersection  
20  
21 identified in pyrrole so far and it arises from the crossing between the  $\pi\sigma_{\text{NC}}^*$  state and the  
22  
23 ground state.  
24  
25  
26  
27

28 Table 2. (around here)  
29  
30

31 Although the similarity of geometrical parameters suggests that the selected MR-  
32  
33 CISD/SA-5-CASSCF(4,5)/BS level of theory is adequate, it is also of importance to compare  
34  
35 the energies of the MXSs. MRCI and MRCI+Q energy values of pyrrole MXSs obtained by  
36  
37 the MR-CISD(Q)/SA-5-CASSCF(4,5)/BS and benchmark MRCI values [27] are summarized  
38  
39 in Table 2. The analysis of presented data shows that the energies of the MXSs are in very  
40  
41 good agreement with the benchmark ones. The comparison among results reveals that the  
42  
43 selected MRCI method is well suited for the description of both ring-puckering and NH-  
44  
45 stretching mechanisms. In particular, the current MRCI and MRCI+Q energies of the ring-  
46  
47 puckered MXS are by 0.06 eV higher and 0.07 eV lower than the benchmark MRCI and  
48  
49 MRCI+Q values, respectively. For the NH-stretched MXS, the MRCI energy is by 0.04 eV  
50  
51 higher, whereas the MRCI+Q value is by 0.18 eV lower than the benchmark values.  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 3.3 Reaction paths

We have computed the reaction pathways between the ground state minimum and the three MXSs described in the previous section using the MR-CISD/SA-5-CASSCF(4,5)/BS level of theory. The resulting potential energy curves are shown in Fig. 2.

Fig. 2 (around here)

Comparison with the results obtained with the MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ method (Fig. 2 in [27]) reveals that NH-stretching potential energy curves (Fig. 2a) agree very well for the ground state  $S_0$  and the two Rydberg  $^1A_2$  and  $^1B_1$  states for the whole range of NH distances. The main difference is that the crossing between the lowest Rydberg state and the ground state occurs at around 1.9 Å in the present work instead of 2.1 Å found in our earlier study [27]. The other features of the potential energy curves exhibit the same behavior as observed earlier. Specifically, the lowest two Rydberg states show small energy barriers (0.24 eV for the  $^1A_2$  state and 0.12 for the  $^1B_1$  state) at the NH distance of 1.2 Å necessary to transform the  $\pi 3s$  orbital into the  $\pi\sigma_{\text{NH}}^*$  state as expected for a stretching of the NH bond. It should also be pointed out that the two valence  $^1A_1$  and  $^1B_2$  states show the same energy profile until the NH distance of 1.8 Å. After that, an intrusion of higher excited states occurs (not shown), which is presumably a direct consequence of the CAS(4,5) active space. However, in the NH stretching mechanism, the deactivation occurs via conical intersections among Rydberg states and the ground state and the differences in the valence states for large NH distances are of minor importance.

Fig. 2b shows that the LIIC path of the ring-puckering mechanism in the  $\pi\pi^*$  state occurs without barrier. Indeed, it is clearly seen that the lowest  $\pi\pi^*$  state is diabatically connected to the ground state, which may make it especially efficient for the internal conversion. The same result was observed in our previous study [27] thus providing additional

1  
2  
3 support for using the applied method. In the case of the ring-opening mechanism Fig. 2c  
4  
5 shows that the initially excited  $\pi\pi^*$  states can deactivate without barrier along this pathway.  
6  
7  
8 The character of the state should, however, change into  $\pi\sigma_{\text{NC}}^*$  in order to lead to the crossing  
9  
10 with the ground state.  
11

12  
13 Apart from the fact that the NH-stretching mechanism should dominate at low  
14  
15 excitation energies, it is difficult to draw general conclusions about the efficiency of each  
16  
17 mechanism based on the reaction paths alone in a clear cut way. When the excitation leads  
18  
19 into the spectral region of the  $\pi\pi^*$  state all mechanisms are energetically possible. In favor of  
20  
21 the NH-stretching mechanism is the fact that it requires the smallest deformations from the  
22  
23 Franck-Condon region in terms of mass-weighted distances (see Fig. 2). On the other hand, it  
24  
25 also requires the diabatic transformation from the  $\pi\pi^*$  state into the  $\pi\sigma_{\text{NH}}^*$  state, which  
26  
27 depends upon the activation of out-of-plane modes [18, 27]. The ring-opening mechanism  
28  
29 involves the lowest energy conical intersection, but it requires the largest deformations from  
30  
31 the Franck-Condon region and diabatic changes in the wave function at the same time.  
32  
33 Finally, the ring-puckering mechanism, as already mentioned, can directly proceed through a  
34  
35 diabatic connection. However, it involves the highest energy portions of the seam of conical  
36  
37 intersections.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

#### 4. Dynamics simulations of pyrrole

49 The non-adiabatic excited state dynamics of pyrrole was started from the  $S_4$  state thus making  
50  
51 all pathways discussed in the previous section energetically available. The resulting average  
52  
53 adiabatic populations of the ground and excited states as a function of time are presented in  
54  
55 Fig. 3. Their analysis shows that the  $S_4$  state transfers its population to the  $S_3$  state in the first  
56  
57 10 fs. After ca. 50 fs, the  $S_4$  state is almost completely depopulated. The populations of  $S_3$  and  
58  
59  $S_2$  states reach a maximum at 10 fs and 20 fs, respectively. At about 75 fs, these states are  
60

1  
2  
3 already depopulated. The  $S_2$  state shows a repopulation between 100 and 150 fs. The  
4  
5 population of the  $S_1$  state increases reaching a maximum at 75 fs. At 100 fs, the  $S_1$  and  $S_0$   
6  
7 states have approximately the same population. Between 100 fs and 200 fs, the simulation is  
8  
9 basically reduced to the  $S_1/S_0$  two-state dynamics, with the complete population transferred to  
10  
11 the ground state at about 200 fs.  
12  
13

14  
15  
16 Fig. 3 (around here)  
17

18  
19 The  $S_1$  population shows a consecutive two-step first order decay type of behavior. By  
20  
21 fitting the  $S_1$  population curve with the function  
22  
23

$$f(t) = \frac{\tau_2}{\tau_2 - \tau_1} \left[ \exp\left(-\frac{t}{\tau_1}\right) - \exp\left(-\frac{t}{\tau_2}\right) \right], \quad (1)$$

24  
25  
26  
27  
28  
29 two time constants  $\tau_1 = 44 \pm 2$  fs and  $\tau_2 = 80 \pm 2$  fs are obtained. Here,  $\tau_1$  measures the  
30  
31 population of  $S_1$  from the collection of states  $S_4$  to  $S_2$  and  $\tau_2$  describes the depopulation  
32  
33  $S_1 \rightarrow S_0$ . The approximate time constant for the overall population of the ground state can be  
34  
35 obtained by fitting the  $S_0$  population with the function  
36  
37  
38

$$f(t) = 1 - \exp\left(-\frac{t}{\tau_0}\right), \quad (2)$$

39  
40  
41  
42  
43  
44  
45 which gives  $\tau_0 = 139 \pm 2$  fs. Note that in these three time constants the error bars denote the  
46  
47 uncertainty of the fitting procedure and not of the process itself, which certainly is larger than  
48  
49 a few femtoseconds.  
50  
51

52  
53 In Fig. 4 a summary of the results of the dynamics simulation in terms of the fraction  
54  
55 of trajectories following each of the three mechanisms is given. The NH-stretching is the main  
56  
57 mechanism after excitation of pyrrole to the  $S_4$  state. This mechanism occurs in 80% of the  
58  
59 trajectories. Other 13% follow ring-deformation mechanisms (ring-opening and ring-  
60

1  
2  
3 puckering). 7% do not deactivate within the 200 fs of the dynamics simulation. Because of the  
4  
5 uncertainties associated to the dynamics simulations and to the relatively small number of  
6  
7 trajectories, these fractions should be taken as qualitative trends of occurrence of each  
8  
9 mechanism, rather than a quantitative assessment of them. If trajectories starting in the  $\pi\pi^*$   
10  
11 and in the  $\pi\sigma_{\text{NH}}^*$  states are independently analyzed, these fractions remain essentially the  
12  
13 same, implying that the population of each mechanism depends on the excitation energy, but  
14  
15 not on the nature of the state. The fact that the fast H atom is formed along the NH stretching  
16  
17 pathway either excited in the  $\pi\pi^*$  or  $\pi\sigma^*$  states has also been observed in the photofragment  
18  
19 translational spectroscopy studies by Cronin et al. [26].  
20  
21  
22  
23  
24

25  
26 Experimental pump of pyrrole with 250 nm (4.96 eV) laser pulse followed by  
27  
28 ionization probe with 241 nm (5.15 eV) pulse reveals two time constants,  $\tau_f = 110 \pm 80$  fs and  
29  
30  $\tau_s = 1100 \pm 500$  fs [19]. These time constants correspond to the time for formation of fast and  
31  
32 slow H atoms, respectively. Since most of trajectories in our simulations finished in the  
33  
34 ground state of the dissociated pyrrolyl + H system, the deactivation time  $\tau_0$  should also  
35  
36 approximately give the time for the formation of the fast H atoms population. Indeed, the  
37  
38 comparison of  $\tau_0$  and  $\tau_f$  shows good agreement. Note, however, that the initial state in the  
39  
40 experiments (low energy  $\pi\sigma_{\text{NH}}^*$ ) and in the simulations (high energy  $\pi\pi^*$  and  $\pi\sigma_{\text{NH}}^*$ ) are not  
41  
42 the same. This is an indication that the fast H elimination occurs directly by the same process,  
43  
44 as soon as there is enough energy to overcome the  $\pi 3s/\pi\sigma_{\text{NH}}^*$  barrier in the  $S_1$  state.  
45  
46  
47  
48  
49

50  
51 Fig. 4 (around here)  
52

53  
54 Fig. 5 (around here)  
55

56  
57 The analysis of NH and CN bond distances was conducted for all trajectories and the  
58  
59 results are presented in Fig. 5. The top panel of this figure shows that in some cases the CN  
60  
distance is elongating during the dynamics. This behavior can be ascribed to the ring-opening

1  
2  
3 and ring-puckering deactivation mechanisms. Since the main deactivation channel is the NH-  
4 stretching, the majority of trajectories do not exhibit elongation of this specific bond. In the  
5  
6 bottom panel of Fig. 5 the NH distance is monitored. In this figure three kinds of trajectories  
7  
8 can be distinguished. For part of the trajectories the NH distance remain constant at about 1  
9  
10 Å. They correspond to the trajectories following ring-distortion mechanisms. A minor fraction  
11  
12 of trajectories (3) has the NH distance oscillating at a medium distance of about 2 or 3 Å.  
13  
14 These are cases where the NH-stretching mechanism is activated, but instead finishing in  
15  
16 dissociation, the hot ground state of pyrrole is formed. In most of the trajectories the NH  
17  
18 distance is steadily increasing. In these cases, the NH-stretching mechanism is activated and  
19  
20 the H atom elimination is taking place. It should be mentioned that a cut-off value of 10 Å for  
21  
22 NH distance was used in Fig. 5 in order to simplify the data analysis. In some of the  
23  
24 trajectories, however, the NH distance was longer, up to 40 Å.  
25  
26  
27  
28  
29  
30  
31

32 Fig. 6 (around here)

33  
34  
35 Fig. 6a shows that the hydrogen dissociation starts on average 54 fs after the  
36  
37 photoexcitation. The kinetic energy of the dissociated hydrogen atom has a broad distribution  
38  
39 around the average value of 1.2 eV (Fig. 6b). This value is in very good agreement with the  
40  
41 experimental results, ~1 eV [11, 26], for the center of the fast H-elimination peak in the  
42  
43 kinetic energy release spectra. Note that, as expected, there is no formation of a slow H-  
44  
45 elimination peak, which should take place in the picosecond timescale [19], much longer than  
46  
47 the maximum simulation time (200 fs). The NH distance at the  $S_1 \rightarrow S_0$  hopping time is shown  
48  
49 in Fig. 6c for all trajectories that have returned to the ground state. The histogram shows two  
50  
51 distinct peaks. The first peak with average at 1.0 Å will be discussed below. The second peak  
52  
53 starts at 1.5 Å and presents a long tail for large distances up to 4 Å. This peak corresponds to  
54  
55 the trajectories deactivated by means of the NH-stretching mechanism. Its average value at  
56  
57  
58  
59  
60

1  
2  
3 2.1 Å is 0.2 Å larger than the NH distance for the crossing between the lowest  $\pi\sigma^*$  state and  
4  
5  
6 the ground state shown in Fig. 2 (left panel).  
7

8  
9 Fig. 7 (around here)

10  
11 Twelve out of ninety trajectories did not follow the NH-stretching mechanism. They  
12 appear in the short-distance peak in Fig. 6c. In order to understand which kind of mechanism  
13 they followed, it is useful to project them on the Cremer-Pople (CP) space  $Q-\phi$ . This is shown  
14 in Fig. 7 for all structures for which the  $S_1-S_0$  energy gap is smaller than 0.5 eV (open dots)  
15 and for structures at the hopping time (full dots). The ring-opened MXS is at  $Q = 0$  Å and the  
16 ring-puckered MXS is shown by a cross ( $E_1$  conformation). Since the ring-opened and the  
17 ring-puckered conical intersections correspond to distinct types of structures on the crossing  
18 seam with different electronic configurations, it could be expected that the structures at the  
19 hopping time would cluster in two disjoint regions around these MXSs. This, however, is not  
20 the case. Fig. 7 shows that the non-adiabatic events occur in a large continuous portion of the  
21 CP space, indicating that the crossing seam spans this entire region. The degree of puckering  
22 varies from almost planar ( $Q = 0.15$  Å) to the strongly puckered structures ( $Q = 0.75$  Å). Most  
23 of hopping events occur at  $E_1$ ,  ${}^2T_1$  and  ${}^2E$  conformations, indicating that not only the  $E_1$   
24 conformation of the ring-puckered MXS, but also other kinds of puckering can give rise to  
25 conical intersections in pyrrole.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 If we take  $Q = 0.3$  Å as an arbitrary threshold to distinguish between the ring-opening  
48 and ring-puckering mechanisms, nine trajectories deactivated at ring-puckered conformations  
49 and three trajectories deactivated at ring-opened conformations, thus corresponding to 10%  
50 and 3% of the total number of trajectories, respectively (see Fig. 4).  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## 5. Conclusions

The photochemical processes in pyrrole were investigated using a high-level multireference configuration interaction method (MRCI) giving a balanced description of the four studied excited states, two of Rydberg character and two valence states. Cuts along the potential energy surfaces connecting the Franck-Condon region and three different minima on the crossing seam (MXS) (NH dissociation, ring puckering, and a planar ring-opened MXS) describe possible deactivation pathways. One of these intersection points, the ring-opened MXS, was characterized for the first time. Although it is the conical intersection of the lowest energy identified in pyrrole so far, its efficiency for the internal conversion process seems to be reduced by the required strong geometric deformations and by the diabatic change of the initially excited  $\pi\pi^*$  state into the  $\pi\sigma_{\text{NC}}^*$  state, which in turn crosses the ground state.

Non-adiabatic surface-hopping dynamics simulations of pyrrole were performed for 200 fs starting in the  $S_4$  state and using a high-level MR-CI approach for the electronic structure calculations. The dynamics simulations show that in fact all three types of conical intersections were accessed. The transfer of population from the initially excited  $S_4$  state to the ground state takes place in about 140 fs. This process occurs basically in two steps, with the  $S_1$  state being populated in about 44 fs and then being depleted in about 80 fs. Most of trajectories (80%) dissociated rapidly along the repulsive  $\pi\sigma_{\text{NH}}^*$  state giving rise to a population of fast H atoms. The computed deactivation time of 140 fs agrees very well with the experimentally measured time constant of 110 fs for the formation of fast hydrogen atoms. The computed average kinetic energy agrees very well with the experimentally observed average kinetic energy of the fast hydrogen atoms. A fraction of 13% of trajectories follows ring-deformation channels involving either ring puckering (10%) or planar ring opening (3%). These fractions did not depend on whether the initial state had  $\pi\pi^*$  or  $\pi\sigma_{\text{NH}}^*$  character.

1  
2  
3 Our calculations provide a detailed picture of the photodeactivation processes in  
4 pyrrole. Although the main objective of this work – the observation of the occurrence of the  
5 different deactivation mechanisms – has been accomplished, it should be noted that the  
6 participation of  $\pi\sigma_{\text{NH}}^*$  states in the initial conditions was much higher than what would be  
7 expected from the oscillator strengths of these two transitions. This bias occurred because of  
8 the relatively high vertical excitation energy of the  $^1\text{B}_1$  Rydberg state, which caused frequent  
9 exchange of position with the  $\pi\pi^*$  in the Wigner sample. Interestingly, it turned out that the  
10 observed percentages of the different mechanisms was insensitive to the initial character of  
11  $\text{S}_4$ , consequently implying that that this bias is not so critical for the general interpretations.  
12 Nevertheless, more investigations are needed to analyze the influence of excitation energies  
13 on the product yields in order to explain the experimentally observed strong energy  
14 dependence of the branching ratios for fast and slow hydrogen atoms.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

## 34 Acknowledgments

35  
36 This work was supported by the Austrian Science Fund within the framework of the Special  
37 Research Program F16 (Advanced Light Sources) and Project P18411-N19. The calculations  
38 were partially performed at the Linux PC cluster Schrödinger III of the computer center of the  
39 University of Vienna. The work in Zagreb (M.E.M and M.V.) was supported by the Ministry  
40 of Science, Education and Sport through the project 098-0982933-2920 and the COST D37  
41 action.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

## 53 References

- 54  
55  
56 [1] L. Serrano-Andrés, M. Merchán, I. Nebotgil, B. O. Roos, and M. Fulscher, J. Am. Chem.  
57 Soc. **115**, 6184 (1993).  
58  
59 [2] M. H. Palmer, I. C. Walker, and M. F. Guest, Chem. Phys. **238**, 179 (1998).  
60

- 1  
2  
3 [3] O. Christiansen, J. Gauss, J. F. Stanton, and P. Jorgensen, *J. Chem. Phys.* **111**, 525 (1999).  
4  
5 [4] D. J. Tozer, R. D. Amos, N. C. Handy, B. O. Roos, and L. Serrano-Andres, *Mol. Phys.*  
6 **97**, 859 (1999).  
7  
8 [5] J. Wan, J. Meller, M. Hada, M. Ehara, and H. Nakatsuji, *J. Chem. Phys.* **113**, 7853 (2000).  
9  
10 [6] B. O. Roos, P. A. Malmqvist, V. Molina, L. Serrano-Andres, and M. Merchan, *J. Chem.*  
11 *Phys.* **116**, 7526 (2002).  
12  
13 [7] C. G. Zhan, and D. A. Dixon, *J. Mol. Spectrosc.* **216**, 81 (2002).  
14  
15 [8] P. Celani, and H. J. Werner, *J. Chem. Phys.* **119**, 5044 (2003).  
16  
17 [9] M. H. Palmer, and P. J. Wilson, *Mol. Phys.* **101**, 2391 (2003).  
18  
19 [10] M. Pastore, C. Angeli, and R. Cimiraglia, *Chem. Phys. Lett.* **422**, 522 (2006).  
20  
21 [11] D. A. Blank, S. W. North, and Y. T. Lee, *Chem. Phys.* **187**, 35 (1994).  
22  
23 [12] A. B. Trofimov, H. Köppel, and J. Schirmer, *J. Chem. Phys.* **109**, 1025 (1998).  
24  
25 [13] A. L. Sobolewski, W. Domcke, C. Dedonder-Lardeux, and C. Jouvet, *PCCP* **4**, 1093  
26 (2002).  
27  
28 [14] J. Wei, A. Kuczmann, J. Riedel, F. Renth, and F. Temps, *PCCP* **5**, 315 (2003).  
29  
30 [15] V. Vallet, Z. G. Lan, S. Mahapatra, A. L. Sobolewski, and W. Domcke, *Faraday Discuss.*  
31 **127**, 283 (2004).  
32  
33 [16] V. Vallet, Z. G. Lan, S. Mahapatra, A. L. Sobolewski, and W. Domcke, *J. Chem. Phys.*  
34 **123** (2005).  
35  
36 [17] J. Wei, J. Riedel, A. Kuczmann, F. Renth, and F. Temps, *Faraday Discuss.* **127**, 267  
37 (2004).  
38  
39 [18] H. Köppel, E. V. Gromov, and A. B. Trofimov, *Chem. Phys.* **304**, 35 (2004).  
40  
41 [19] H. Lippert, H. H. Ritze, I. V. Hertel, and W. Radloff, *Chemphyschem* **5**, 1423 (2004).  
42  
43 [20] A. J. van den Brom, M. Kapelios, T. N. Kitsopoulos, N. H. Nahler, B. Cronin, and M. N.  
44 R. Ashfold, *PCCP* **7**, 892 (2005).  
45  
46 [21] I. Frank, and K. Damianos, *Journal of Chemical Physics* **126** (2007).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [22] Z. Lan, A. Dupays, V. Vallet, S. Mahapatra, and W. Domcke, *Journal of Photochemistry*  
4 *and Photobiology a-Chemistry* **190**, 177 (2007).  
5  
6  
7  
8 [23] Z. Lan, and W. Domcke, *Chem. Phys.* **350**, 125 (2008).  
9  
10 [24] A. Kumar, M. Kolaski, and K. S. Kim, *J. Chem. Phys.* **128** (2008).  
11  
12 [25] E. J. Shin, *Bull. Korean Chem. Soc.* **25**, 907 (2004).  
13  
14 [26] B. Cronin, M. G. D. Nix, R. H. Qadiri, and M. N. R. Ashfold, *PCCP* **6**, 5031 (2004).  
15  
16 [27] M. Barbatti, M. Vazdar, A. J. A. Aquino, M. Eckert-Maksic, and H. Lischka, *J. Chem.*  
17 *Phys.* **125**, 164323 (2006).  
18  
19 [28] V. Poterya, V. Profant, M. Farnik, P. Slavicek, and U. Buck, *J. Chem. Phys.* **127**, 064307  
20 (2007).  
21  
22 [29] M. Barbatti, B. Sellner, A. J. A. Aquino, and H. Lischka, in *Radiation Induced Molecular*  
23 *Phenomena in Nucleic Acid*, edited by M. K. Shukla, and J. Leszczynski (Springer, Netherlands,  
24 2008), pp. 209.  
25  
26 [30] S. Salzmann, M. Kleinschmidt, J. Tatchen, R. Weinkauff, and C. M. Marian, *PCCP* **10**, 380  
27 (2008).  
28  
29 [31] N. Gavrilov, S. Salzmann, and C. M. Marian, *Chem. Phys.* **349**, 269 (2008).  
30  
31 [32] M. Barbatti, H. Lischka, S. Salzmann, and C. M. Marian, *J. Chem. Phys.*, submitted  
32 (2008).  
33  
34 [33] S. Perun, A. L. Sobolewski, and W. Domcke, *Chem. Phys.* **313**, 107 (2005).  
35  
36 [34] M. Barbatti, and H. Lischka, *J. Am. Chem. Soc.* **130**, 6831 (2008).  
37  
38 [35] V. Profant, V. Poterya, M. Farnik, P. Slavicek, and U. Buck, *J. Phys. Chem. A* **111**, 12477  
39 (2007).  
40  
41 [36] A. L. Sobolewski, and W. Domcke, *Chem. Phys.* **259**, 181 (2000).  
42  
43 [37] I. Antol, M. Vazdar, M. Barbatti, and M. Eckert-Maksic, *Chem. Phys.* **349**, 308 (2008).  
44  
45 [38] M. Barbatti, M. Ruckebauer, J. J. Szymczak, A. J. A. Aquino, and H. Lischka, *PCCP* **10**,  
46 482 (2008).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [39] J. C. Tully, *Faraday Discuss.* **110**, 407 (1998).  
4  
5 [40] A. Ferretti, G. Granucci, A. Lami, M. Persico, and G. Villani, *J. Chem. Phys.* **104**, 5517  
6  
7 (1996).  
8  
9 [41] N. L. Doltsinis, and D. Marx, *Phys. Rev. Lett.* **88**, 166402 (2002).  
10  
11 [42] M. Barbatti, G. Granucci, M. Persico, M. Ruckebauer, M. Vazdar, M. Eckert-Maksic,  
12  
13 and H. Lischka, *J. Photochem. Photobiol., A* **190**, 228 (2007).  
14  
15 [43] E. Fabiano, T. W. Keal, and W. Thiel, *Chem. Phys.* **349**, 334 (2008).  
16  
17 [44] J. Pittner, H. Lischka, and M. Barbatti, *Chem. Phys.*, doi:10.1016/j.chemphys.2008.10.013  
18  
19 (2008).  
20  
21 [45] J. C. Tully, *J. Chem. Phys.* **93**, 1061 (1990).  
22  
23 [46] L. Sun, and W. L. Hase, in *Reviews in Computational Chemistry*, edited by K. B. Lipkowitz *et*  
24  
25 *al.* (Wiley-VCH, New York, 2003), pp. 79.  
26  
27 [47] H. Lischka, R. Shepard, F. B. Brown, and I. Shavitt, *Int. J. Quantum Chem.* **S.15**, 91  
28  
29 (1981).  
30  
31 [48] H. Lischka, R. Shepard, R. M. Pitzer, I. Shavitt, M. Dallos, T. Müller, P. G. Szalay, M.  
32  
33 Seth, G. S. Kedziora, S. Yabushita, and Z. Y. Zhang, *PCCP* **3**, 664 (2001).  
34  
35 [49] H. Lischka, R. Shepard, I. Shavitt, R. M. Pitzer, M. Dallos, T. Mueller, P. G. Szalay, F. B.  
36  
37 Brown, R. Ahlrichs, H. J. Boehm, A. Chang, D. C. Comeau, R. Gdanitz, H. Dachsel, C. Ehrhardt,  
38  
39 M. Ernzerhof, P. Hocht, S. Irl, G. Kedziora, T. Kovar, V. Parasuk, M. J. M. Pepper, P. Scharf,  
40  
41 H. Schiffer, M. Schindler, M. Schueler, M. Seth, E. A. Stahlberg, J.-G. Zhao, S. Yabushita, Z.  
42  
43 Zhang, M. Barbatti, S. Matsika, M. Schuurmann, D. R. Yarkony, S. R. Brozell, E. V. Beck, and J.-  
44  
45 P. Blaudeau, COLUMBUS, an ab initio electronic structure program, release 5.9.1,  
46  
47 [www.univie.ac.at/columbus](http://www.univie.ac.at/columbus) (2006).  
48  
49 [50] A. Bunge, *J. Chem. Phys.* **53**, 20 (1970).  
50  
51 [51] T. H. Dunning, *J. Chem. Phys.* **90**, 1007 (1989).  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [52] R. Shepard, H. Lischka, P. G. Szalay, T. Kovar, and M. Ernzerhof, *J. Chem. Phys.* **96**,  
4 2085 (1992).  
5  
6  
7  
8 [53] R. Shepard, in *Modern Electronic Structure Theory*, edited by D. R. Yarkony (World Scientific,  
9 Singapore, 1995), p. 345.  
10  
11  
12 [54] H. Lischka, M. Dallos, and R. Shepard, *Mol. Phys.* **100**, 1647 (2002).  
13  
14  
15 [55] M. Dallos, H. Lischka, R. Shepard, D. R. Yarkony, and P. G. Szalay, *Journal of Chemical*  
16  
17 *Physics* **120**, 7330 (2004).  
18  
19  
20 [56] H. Lischka, M. Dallos, P. G. Szalay, D. R. Yarkony, and R. Shepard, *Journal of Chemical*  
21  
22 *Physics* **120**, 7322 (2004).  
23  
24  
25 [57] S. R. Langhoff, and E. R. Davidson, *Int. J. Quantum Chem.* **8**, 61 (1974).  
26  
27 [58] P. J. Bruna, S. D. Peyerimhoff, and R. J. Buenker, *Chem. Phys. Lett.* **72**, 278 (1980).  
28  
29 [59] V. Bonacic-Koutecky, and R. Mitric, *Chem. Rev.* **105**, 11 (2005).  
30  
31 [60] R. Mitric, V. Bonacic-Koutecky, J. Pittner, and H. Lischka, *J. Chem. Phys.* **125** (2006).  
32  
33 [61] W. C. Swope, H. C. Andersen, P. H. Berens, and K. R. Wilson, *J. Chem. Phys.* **76**, 637  
34 (1982).  
35  
36  
37 [62] G. Granucci, and M. Persico, *J. Chem. Phys.* **126**, 134114 (2007).  
38  
39  
40 [63] D. Cremer, and J. A. Pople, *J. Am. Chem. Soc.* **97**, 1354 (1975).  
41  
42 [64] T. Helgaker, H. J. A. Jensen, P. Jørgensen, J. Olsen, K. Ruud, H. Ågren, T. Andersen, K.  
43 L. Bak, V. Bakken, O. Christiansen, P. Dahle, E. K. Dalskov, T. Enevoldsen, H. Heiberg, H.  
44 Hetta, D. Jonsson, S. Kirpekar, R. Kobayashi, H. Koch, K. V. Mikkelsen, P. Norman, M. J.  
45 Packer, T. Saue, P. R. Taylor, and O. Vahtras, DALTON, an ab initio electronic structure  
46 program, Release 1.0 (1997).  
47  
48  
49 [65] M. Barbatti, G. Granucci, M. Ruckebauer, M. Persico, and H. Lischka, NEWTON-X: a  
50 package for Newtonian dynamics close to the crossing seam, [www.univie.ac.at/newtonx](http://www.univie.ac.at/newtonx) (2007).  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Tables

Table 1 – Vertical excitation energies of selected singlet states of pyrrole.

State	MRCI <sup>a</sup>	MRCI+Q <sup>b</sup>	TDDFT <sup>c</sup>	CASPT2	CC3 <sup>g</sup>	Exp
S <sub>0</sub> <sup>1</sup> A <sub>1</sub>	0.00	0.00	0.00	0.00	0.00	
π3s <sup>1</sup> A <sub>2</sub>	5.22	5.09	5.05	5.22 <sup>d</sup> /5.22 <sup>e</sup>	5.10	5.08 <sup>h</sup> /5.22 <sup>i</sup>
π3s <sup>1</sup> B <sub>1</sub>	6.21	5.86	5.88	5.85 <sup>f</sup> /5.92 <sup>e</sup>	5.99	6.22 <sup>h</sup>
ππ* <sup>1</sup> A <sub>1</sub>	6.55	6.39	6.29	5.82 <sup>d</sup> /5.98 <sup>e</sup>	6.37	
ππ* <sup>1</sup> B <sub>2</sub>	6.65	6.78	6.45	5.87 <sup>d</sup> /5.95 <sup>e</sup>	6.63	5.92 <sup>h</sup> /6.2-6.4 <sup>i</sup>

<sup>a</sup> Present results, MR-CISD/SA-5-CASSCF(4,5)/BS

<sup>b</sup> MR-CISD+Q/SA-5-CAS(6,5)+AUX(1)/d-aug-cc-pVDZ, Reference [27].

<sup>c</sup> Reference [4].

<sup>d</sup> Reference [6].

<sup>e</sup> Reference [8].

<sup>f</sup> Reference [1].

<sup>g</sup> Reference [3].

<sup>h</sup> Assignments given in Reference [9].

<sup>i</sup> Assignments given in Reference [3].

Table 2 – Energy of pyrrole MXSs (in eV) relative to the minimum in the ground state.

MXS	MRCI <sup>a</sup>	MRCI+Q <sup>a</sup>	MRCI	MRCI+Q	MXS features
$\pi\pi^*/S_0$ ( $E_1$ )	4.95	4.86	4.89 <sup>b</sup>	4.93 <sup>b</sup>	ring puckering, Fig. 1a
$\pi\sigma_{\text{NH}}^*/S_0$	4.45	4.26	4.41 <sup>c</sup>	4.44 <sup>c</sup>	NH stretching, Fig. 1b
$\pi\sigma_{\text{NC}}^*/S_0$	4.11	3.86	-	-	ring opening, Fig. 1c

<sup>a</sup>Present results, MR-CISD/SA-5-CASSCF(4,5)/BS

<sup>b</sup>MR-CISD(Q)/SA-3-CAS(6,5)/6-31G(d), Reference [27].

<sup>c</sup>MR-CISD(Q)/SA-3-CAS(6,6)/6-31G(d), Reference [27].



## Figure Captions

Fig. 1. Structures and selected geometric parameters for pyrrole MXSs obtained at the MRCI level. Distances are given in Å and dihedral angles in degrees. The number in brackets correspond to the benchmark MRCI value from Ref. [27].

Fig. 2. Potential energy curves calculated at the MRCI level along a) the rigid NH-stretching coordinate and along the LIIC path from the ground state minimum to b) the ring-puckered MXS and to c) the ring-opened MXS.

Fig. 3. Average adiabatic populations of trajectories for each state as a function of time after initial photoexcitation of pyrrole into the  $S_4$  state.

Fig. 4. Description and statistics of trajectory deactivation mechanisms.

Fig. 5. CN (top) and NH (bottom) distance variations as a function of time for all trajectories. The NH distance of 10 Å was used as a cut-off value (see text).

Fig. 6. Analysis of the trajectories showing NH dissociation. (a) Initial time of the dissociation, taking 2 Å for the NH bond as reference value. (b) Hydrogen kinetic energy. (c) NH distance for all trajectories at the time of the  $S_1 \rightarrow S_0$  hopping.

Fig. 7. Distribution of conformations in the Cremer-Pople  $Q-\phi$  space for trajectories following ring-deformation mechanisms. Full dots: conformations at the hopping time. Open dots: conformations with  $S_1-S_0$  energy gaps smaller than 0.5 eV. Cross: ring puckered MXS.