



HAL
open science

First synthesis of upper rim mono and dinitrone calix[4]arene derivatives

Sofiane Ben Sdira, Roselyne Baudry, Caroline Felix, Marie-Béatrice Giudicelli, Francis Vocanson, Monique Perrin, Roger Lamartine

► **To cite this version:**

Sofiane Ben Sdira, Roselyne Baudry, Caroline Felix, Marie-Béatrice Giudicelli, Francis Vocanson, et al.. First synthesis of upper rim mono and dinitrone calix[4]arene derivatives. *Supramolecular Chemistry*, 2007, 19 (6), pp.393-398. 10.1080/10610270601003932 . hal-00513494


HAL Id: hal-00513494

<https://hal.science/hal-00513494>

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



First synthesis of upper rim mono and dinitrone calix[4]arene derivatives

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2006-0047.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	07-Sep-2006
Complete List of Authors:	FELIX, Caroline; Université Lyon 1, chemistry VOCANSON, Francis; Université Lyon 1, UFR de chimie
Keywords:	calixarene, nitrone function, crystal structure
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p> <p>schema1.cdx</p>	

1
2
3
4 First Synthesis of Upper Rim Mono and Dinitrone Calix[4]arene Derivatives
5
6
78
9 SOFIANE BEN SDIRA, ROSELYNE BAUDRY, CAROLINE P. FELIX*, MARIE-BÉATRICE A. GIUDICELLI,10
11 FRANCIS VOCANSON, MONIQUE PERRIN AND ROGER J. LAMARTINE12
13 *Equipe Chimie Supramoléculaire Appliquée, CNRS UMR 5634, Université Claude Bernard Lyon 1,*14
15 *43 bd du 11 Novembre 1918, F-69622 Villeurbanne cedex, France.*
16
1718
19 **Abstract-** A new class of calix[4]arene derivatives bearing one or two nitrone groups on the upper rim
20 has been synthesized using the oxidation of chiral imines with hydrogen peroxide. The imine
21 intermediates were obtained *via* amination of the diformyl derivative. The ¹H NMR spectra and X-ray
22 data indicated a 1,3-disubstituted cone conformation for the imine derivatives and the existence of a
23 single (*E,E*) isomer. The structural identity of the nitrone derivatives was confirmed from NMR, IR and
24 ES-MS data. Both mono and dinitrone calix[4]arenes were in *E* configuration.
25
26
27
28
2930
31 *Keywords :* calixarene ; nitrone function ; crystal structure
32
3334
35 INTRODUCTION
36
3738
39 Calix[n]arenes are a family of macrocyclic molecules consisting of *n-para* substituted phenol units
40 connected through methylene bridges in their *ortho* position.¹ These molecules and their derivatives have
41 been extensively studied for the past ten years for their interesting and versatile properties *e.g.*
42 complexation, formation of supramolecular assemblies, design of biomimetics.
43
4445
46 Nitrones have been known for many years², but interest in their chemistry has grown mostly due to their
47 increasing use as synthons in cycloaddition reactions³, natural products synthesis⁴ and radical spin traps
48
49
50
51
52
53
54
55
56
57
58
59
6059
60 Nitrone derivatives are very efficient in trapping free radicals, especially for oxygen centered radicals in
biological media.⁹ For such applications it would be interesting to incorporate the nitrone function into
the calixarene core to achieve the elaboration of a three dimensional and conformationally rigid platform.

*Corresponding author. Tel.:+33-472 44 80 14;fax:+33-472 44 84 38;e-mail: caroline.felix@univ-lyon1.fr

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

It is known that the oxidation of imine derivatives with metachloroperbenzoic acid (MCPBA) or hydrogen peroxide led to the formation of the corresponding oxaziridines. The rearrangement of oxaziridine derivatives into nitrones, which involves the cleavage of the C-O bond, appears in the presence of strong Lewis acid as catalyst^{10,11} or high temperatures^{10,12}. Studies have shown that various factors can favour the transformation of an imine derivative into the corresponding nitron. The conformation of the imine derivative plays an important role. Nucleophilic attack of the peroxide at the imino carbon atom (orthogonal to the imino plane) followed by intramolecular nucleophilic displacement leads to the formation of the oxaziridine derivative.^{13,14} By contrast, nitron derivatives result from a concerted nucleophilic attack of the imino nitrogen atom on the peroxide (in the imino plane).^{14,15} Steric effects of substituted groups which hinder the approach of the peroxide to the imino carbon in a plane perpendicular to the C=N bond favour the nitron formation. The nitron formation is also observed with benzaldimine derivatives bearing electro donating groups at the para position.¹⁵ Finally, aprotic solvents and basic conditions favour the nitron formation.¹⁶

In this paper, we describe for the first time the synthesis of calix[4]arenes bearing nitron moieties at the upper rim.

RESULTS AND DISCUSSION

With the aim of synthesizing nitron substituted calix[4]arene derivatives, we have developed a strategy in three steps involving a di-formylation reaction of the di-propoxycalixarene **1** (scheme 1) and its conversion into chiral imines. The starting calixarene derivative **1** was chosen in order to block two distal hydroxyl groups. The compound **2** was synthesized in two steps.¹⁷ After recrystallisation (CHCl₃ / MeOH), the diformylcalix[4]arene **2** was obtained in quantitative yield. Single crystal X-ray of **2** has been obtained. The crystallographic data have been recently reported by our group¹⁸ and shown that this compound adopts a "pinched" cone conformation.

The chiral imine derivatives **3ab** were prepared from the corresponding chiral amines^{19,20} ((R)-1-cyclohexylethylamine and (S)-1-phenylethylamine) and are obtained with good yields (81 and 72 % respectively). Their constitution was established by ¹H NMR, ¹³C NMR, ES-MS and X-ray data. The ¹H

1
2
3
4 NMR spectra of **3ab** contain the signal of the imine function (8.56 ppm) instead of the aldehyde singlet
5 (9.81 ppm) identified for **2**. Such observations have been made on the ^{13}C NMR spectra with the imino
6 signal (160 ppm) instead of the aldehyde signal (190 ppm). The single signal of the imine function in
7 NMR suggests the presence of a single isomer which is the (*E,E*) isomer as confirmed by the X-ray data.
8
9 Beside the AB systems for the methylene bridges, two singlets at 7.47 and 7.53 ppm are identified for the
10 protons of the aromatic rings bearing the imine function which give rise to a 1,3-disubstituted cone
11 conformation.
12
13
14
15
16
17
18
19

20 The X-ray structure of **3a** (Fig. 1) confirms a cone conformation in the solid state in accordance with the
21 solution data. From X-ray analysis it is shown that the unit contains one macrocycle, one dichloromethane
22 and two methanol molecules. The calixarene compound and the dichloromethane of solvation have
23 crystallographically imposed twofold symmetry. The dichloromethane lies within the calixarene cup and
24 there are clear C-H... π interactions between the methylene C-H and an aromatic ring. The two
25 cyclohexyl groups orient the edges of the rings inside the cavity of the calixarene. We found two kinds of
26 hydrogen bonds : intramolecular hydrogen bond [OH (phenol)...O (ether)] and intermolecular hydrogen
27 bonds [OH (methanol)...N (imine)]. The X-ray structure unequivocally establishes that the absolute
28 configuration of the two asymmetric carbon atoms is *R* (the stereochemistry of the starting amine has been
29 conserved). It is also unambiguous that the stereochemistry of the imine bond is *E* confirming the
30 presence of a single (*E,E*) isomer as expected.
31
32
33
34
35
36
37
38
39
40
41

42 As observed for **3a**, the structure of **3b** (Fig. 2) shows a cone conformation. The calixarene compound
43 and the dichloromethane of solvation have crystallographically imposed twofold symmetry. The two phenol
44 moieties are orientated inside the cavity of the calixarene core whereas the methyl groups of the propyl
45 chain are directed outside the cavity. Intramolecular hydrogen bonds between the protons of phenolic
46 oxygen atoms and the adjacent ether bridges are observed, contributing to the cone conformation. As for
47 compound **3a**, the X-ray structure of **3b** unequivocally establishes the configuration of this derivative.
48
49 The absolute configuration of the two asymmetric carbon atoms is *S* (the stereochemistry of the starting
50 amine has been conserved). It also confirms the presence of a single (*E,E*) isomer.
51
52
53
54
55
56
57
58
59
60

1
2
3
4 The oxidation reaction described by Kraeīm et al.²¹ yielded the nitronne derivatives. The reaction of the
5 imines derivatives **3ab** with hydrogen peroxide, bicarbonate and benzonitrile in a CH₂Cl₂/MeOH mixture
6 has led to the formation of the expected dinitronne derivatives **4ab** and also to the mononitronne-
7 monoformyl derivatives **5ab** and to benzamide (scheme 1). These compounds have been separated by
8 chromatographic column. The structure of the four calixarene derivatives was determined by ¹H, ¹³C
9 NMR, IR and Mass spectrometry (Tables 1 and 2). For compound **4a**, the HMQC cross peaks indicate
10 that the proton at 7.21 ppm was correlated to the carbon at 133.92 ppm and the COSY cross peaks show
11 that the proton of the CH=N(O) group is correlated to the doublet (8.08 ppm) attributed to the aromatic
12 protons of the calixarene core. Thus 2D NMR analyses have confirmed the structure of the dinitronne
13 derivatives. A comparison of the ¹H NMR spectra of compounds **4** and **5** (Table 1) reveals that the
14 majority of the signals are doubled for **5** indicating a disymmetry in the molecule. Moreover, the
15 characteristic singlets of the nitronne and of the aldehyde have been observed for compounds **5**. This
16 phenomenon of starting aldehyde returned from the imine oxidation as already been described.²²
17
18 The ¹H NMR spectra of all four calixarene-nitronne substituted **4** and **5** exhibit a single singlet for the
19 CH=N(O) proton which demonstrates that the hydrogen peroxide oxidation didn't racemize the starting
20 configuration. Both mono and di-nitronne are still in *E* configuration. The ¹³C NMR chemical shifts of the
21 methylene bridges indicated a cone conformation for the calixarene core in solution.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 CONCLUSION

41
42
43 This work confirms that steric and mesomeric effects have a marked influence upon the oxydation
44 reaction of imine function as described in the literature.¹⁵ The conformation of the calixarene imine
45 derivatives may favoured the nitronne formation. ¹H NMR spectra and X-ray data have indicated a 1,3-
46 disubstituted cone conformation of the imine derivatives **3** and the existence of a single (*E,E*) isomer. To
47 our knowledge, a unique exemple of calixarene bearing nitronne radicals on the lower rim has been
48 reported.²³ No calixarene functionalized with nitronne at the upper rim has been described. Thus, we
49 presented the first synthesis of mono and di-nitronne substituted calix[4]arene derivatives. These
50 compounds represent new rigid supports for further applications such as radicals trapping.
51
52
53
54
55
56
57
58
59
60

EXPERIMENTAL SECTION

General methods. Solvents were purified and dried by standard methods prior to use. All reactions were carried out under nitrogen. Column chromatography was performed with silica gel 60 (0.040-0.063 nm). Melting points are uncorrected. ^1H NMR and ^{13}C NMR spectra were obtained at 300.13 and 75 MHz (CDCl_3 , TMS as internal standard, chemical shifts in ppm, J in Hertz). Mass spectra were obtained by electrospray technique (positive mode). As verified by other authors²⁴ elemental analyses of calixarenes are very often incorrect because of the inclusion of solvent molecules and thus cannot be considered as an appropriate criterion of purity. Nevertheless, the identities of the reported compounds have been confirmed by their structural data.

11,23-Diformyl-26,28-di-propoxy-25,27-di-hydroxycalix[4]arene (2). A solution of di-propyloxycalix[4]arene **1** (2.1 g, 4.17 mmol) in 70 mL of CHCl_3 was cooled to 15°C . Then, SnCl_4 (5.01 mL, 42.72 mmol) and 1,1-dichlorodimethylether (0.963 mL, 10.62 mmol) were quickly added. The mixture was stirred at room temperature for 30 min. The organic layer was washed with water, dried with Na_2SO_4 and the solvent was removed *in vacuo*. The crude product was then recrystallized from $\text{CHCl}_3/\text{MeOH}$ to give 1.95 g of **2** as a white powder. Yield: 92 %. NMR data are in accordance with those described in the literature.¹⁷

5,17-Bis-[N-(S)-cyclohexylethylimino]-25,27-dipropyloxycalix[4]arene (3a). In a 100 mL three-necked flask equipped with stirrer and condenser, 0.156 g (0.27 mmol) of aldehyde **2** was dissolved in 40 mL of CH_2Cl_2 under nitrogen. (S)-cyclohexylamine (0.1 mL, 0.69 mmol) was added. The mixture containing molecular sieves (4\AA) was stirred at 40°C for 48h. The suspension was filtered off on celite and the solvent was removed *in vacuo*. The crude solid was recrystallized from $\text{CHCl}_3/\text{MeOH}$ to give 0.189 g of a white powder. Yield: 81 %. Mp = $220\text{--}222^\circ\text{C}$. $[\alpha]_D^{20} = +63.3$ (c = 0.3, CHCl_3). ^1H NMR: 0.88-1.85 (m + d, 34H, H-cy + CH_3CH , CH_3CH_2), 2.04-2.10 (m, 4H, CH_2CH_3), 2.90-2.96 (m, 2H, CHCH_3), 3.47 (AB, 4H, $J_{\text{AB}} = 12.81$, ArCH_2Ar), 4.00 (bt, 4H, CH_2O), 4.30 (AB, 4H, $J_{\text{AB}} = 13.02$, ArCH_2Ar), 6.77 (t, 2H, J = 7.53, H-Ar), 6.96-7.00 (bt, 4H, H-Ar), 7.48 (sb, 4H, H-Ar), 8.08 (s, 2H, OH), 8.57 (s, 2H, $\text{CH}=\text{N}$). RMN ^{13}C : 11.28 (CH_3CH_2), 20.46 (CH_3CH), 23.89 (CH_2CH_3), 26.63, 26.84, 27.03, 30.30, 30.72 (CH_2 -

cy), 31.64 (ArCH₂Ar), 44.14 (CH-cy), 72.67 (CHN), 78.87 (CH₂O), 125.74, 128.04, 128.92, 129.54, 133.38, 152.26, 156.19, 159.05 (CH=N). IR: 1637 (CH=N), 3221 (OH). ES-MS(+) for C₅₂H₆₆N₂O (783.1) $m/z = 805.4$ [M+Na]⁺, 783.6 [M+H]⁺, 392.3 [M+2H]²⁺.

5,17-Bis-[N-(L)-phenylethylimino]-25,27-dipropoxyxycalix[4]arene (3b). Same procedure with 0.156 g of aldehyde **2** (0.27 mmol) and 0.1 mL of chiral amine (0.81 mmol). Yield: 72 %. [α]_D²⁰ = -5.7 (c = 0.3, CHCl₃). ¹H NMR: 1.32 (t, 6H, J = 7.35, CH₃CH₂), 1.62 (d, 6H, J = 6.57, CHCH₃(NH)), 2.05-2.12 (m, 4H, CH₂CH₃), 3.47 (AB, 4H, J_{AB} = 12.99, ArCH₂Ar), 4.01 (t, 4H, J = 6.42, CH₂O), 4.31 (AB, 4H, J_{AB} = 12.99, ArCH₂Ar), 4.52 (q, 2H, J = 6.57, CHCH₃), 6.76 (t, 2H, J = 7.53, H-Ar), 6.95 (d, 4H, J = 7.35, H-Ar), 7.36 (t, 6H, J = 7.14, H-Ph), 7.45 (d, 4H, J = 7.17, H-Ph), 7.53 (s, 4H, H-Ar), 8.25 (s, 2H, OH), 8.58 (s, 2H, CH=N); ¹³C NMR: 11.28 (CH₃CH₂), 23.90 (CH₂CH₃), 24.86 (CH₃CH(N)), 31.64 (ArCH₂Ar), 69.77 (CHN), 78.87 (CH₂O), 125.72, 126.11, 127.12, 127.93, 128.50, 128.91, 129.17, 129.92, 132.67, 133.29, 145.76, 152.15, 156.48, 160.02 (CH=N). IR: 1637 (CH=N). ES-MS(+) for C₅₂H₅₄N₂O₄ (771.01) $m/z = 771.5$ [M+H]⁺, 386.2 [M+2H]²⁺.

General procedure for the preparation of the nitrene derivatives

In a 100 mL flask, a mixture of dichloromethane (10 mL), methanol (30 mL), sodium bicarbonate (100 mg, 1.2 mmol), benzonitrile (1 mmol, 0.1 mL), imine (0.13 mmol), hydrogen peroxide (11 mmol) was stirred at room temperature for 72h. The mixture was then treated with 25 mL of water, extracted with chloroform (3 × 20 mL). The organic layer was separated, dried with Na₂SO₄ and the solvent was removed *in vacuo*. The crude product was then eluted from a column of silica gel to give the mono or disubstituted nitrene derivative.

5,17-Bis-[N-(S)-cyclohexylethyl nitronyl]-25,27-dipropoxyxycalix[4]arene (4a). The crude product was then eluted from a column of silica gel with AcOEt/hexane (1/1) to give 0.034 mmol of a white solid (Rf = 0.11). Yield: 26 %. Mp = 277-280 °C. [α]_D²⁰ = +22.8 (c = 0.39, CHCl₃). ¹H NMR: 1.28-2.10 (m, 38H, CH₃CH + H-cy + CH₃CH₂ + CH₂CH₃), 3.44-3.48 (dd, J_{AB} = 13.17, 4H, ArCH₂Ar), 3.55 (m, 2H, CHCH₃), 4.00 (t, 4H, J = 6.21, CH₂O), 4.30 (AB, 4H, J_{AB} = 13.17, ArCH₂Ar), 6.77 (t, 2H, J = 7.71, H-Ar), 7.01 (d, 4H, J = 7.71, H-Ar), 7.21 (s, 2H, N=CH), 8.08 (bs, 4H, H-Ar), 8.83 (s, 2H, OH); ¹³C NMR: 11.29

(CH_3CH_2), 17.31 (CH_3CH), 23.88 (CH_2CH_3), 26.05, 26.44, 26.64, 29.82, 30.52 (CH_2 -cy); 31.77 ($ArCH_2Ar$), 40.77 (CH -Cy), 77.39 ($CHN(O)$), 78.79 (CH_2O), 122.29, 125.83, 128.47, 130.04, 133.30, 133.90 ($CH=N(O)$), 152.13, 156.03 (C, Ar). ES-MS(+) for $C_{52}H_{66}N_2O_6$ (815.1) $m/z = 837.4 [M+Na]^+$, 815.5 $[M+H]^+$, 853.5 $[M+K]^+$, 408.4 $[M+2H]^{2+}$.

5,17-Bis-[N-(L)-phenylethylnitronyl]-25,27-dipropylcalix[4]arene (4b). The crude product was then eluted from a column of silica gel with AcOEt/hexane (1/1) to give 0.046 mmol of a white solid ($R_f = 0.11$). Yield: 35 %. Mp = 224-227°C. $[\alpha]_D^{20} = +16.9$ (c = 0.51, $CHCl_3$). 1H NMR: 0.90 (t, 6H, J = 7.17, CH_3CH_2), 1.90 (d, 6H, J = 6.78, CH_3CH), 2.01-2.08 (m, 4H, CH_2CH_3), 3.42 (AB, 4H, $J_{AB} = 13.17$, $ArCH_2Ar$), 3.98 (t, 4H, J = 6.24, CH_2O), 4.24 (AB, 4H, $J_{AB} = 13.17$, $ArCH_2Ar$), 5.13 (q, 2H, J = 6.78, $CHCH_3$), 6.72 (t, 2H, J = 7.53, $H-Ar$), 6.95 (t, 4H, J = 7.53, $H-Ar$), 7.35(s, 2H, $CHN(O)$), 7.36-7.52 (m, 6H, $H-Ar$), 7.53 (d, 4H, J = 7.53, $H-Ar$), 8.01 (s, 2H, $H-Ar$), 8.11 (s, 2H, $H-Ar$), 8.76 (s, 2H, OH); ^{13}C NMR: 11.25 (CH_3CH_2), 19.46 (CH_3CH), 23.87 (CH_2CH_3), 31.69 ($ArCH_2Ar$), 74.65 (CHN), 78.75 (OCH_2), 122.33, 125.76, 127.71, 128.46, 129.13, 129.67, 130.08, 133.14 (C, Ar), 133.58 ($CH=N$), 139.25, 152.09, 156.18 (C, Ar). ES-MS(+) for $C_{52}H_{54}N_2O_6$ (802.01) $m/z = 803.3 [M+H]^+$, 825.2 $[M + Na]^+$.

5-Formyl-17-[N-(S)-cyclohexylethylnitronyl]-25,27-dipropoxycalix[4]arene (5a). The crude product was then eluted from a column of silica gel with AcOEt/hexane (1/1) to give 0.051 mmol of a white solid ($R_f = 0.36$). Yield: 39 %. Mp = 247-249°C. 1H NMR: 1.27-2.10 (m, 24H, H-cy + CH_3CH_2 + CH_2CH_3 + CH_3CH), 3.46-3.56 (m, 5H, $CHCH_3$ + $ArCH_2Ar$), 3.98-4.04 (m, 4H, CH_2O), 4.28-4.35 (dd, 4H, $J_{AB} = 13.20$, 4H, $ArCH_2Ar$), 6.80 (t, 2H, J = 7.53, $H-Ar$), 6.94 (d, 2H, J = 7.71, $H-Ar$), 7.04 (d, 2H, J = 7.53, $H-Ar$), 7.22 (s, 1H, $CH=N(O)$), 7.64 (s, 2H, $H-Ar$), 8.05 (s, 1H, $H-Ar$), 8.14 (s, 1H, $H-Ar$), 8.83 (s, 1H, OH), 9.30 (s, 1H, OH), 9.80 (s, 1H, CHO); ^{13}C NMR: 11.31 (CH_3CH_2), 17.32 (CH_3CH), 23.89 (CH_2CH_3), 26.03, 26.62, 29.82, 30.10, 30.51 (CH_2 -cy), 40.78 (CH -cy), 77.42 ($CHN(O)$), 78.90 (CH_2O), 122.42, 125.98, 128.89, 129.16, 129.50, 130.02, 131.30, 132.69 (C, Ar), 133.42 ($CH=N$) 152.10, 155.94 (C, Ar), 191.37 (CHO). IR: 1672 (CHO). ES-MS(+) for $C_{44}H_{51}NO_6$ (689.89) $m/z = 690.3 [M+H]^+$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

5-Formyl-17-[N-(L)-phenylethylnitronyl]-25,27-dipropoxycalix[4]arene (5b). The crude product was then eluted from a column of silica gel with AcOEt/hexane (1/1) to give 0.041 mmol of a white solid ($R_f = 0.60$). Yield: 31 %. Mp = 212-215°C. $^1\text{H NMR}$: 128-136 (t, 6H, $J = 7,35$, CH_3CH_2), 1.90 (d, 3H, $J = 6,78$, CH_3CH), 2.01-2.1 (m, 4H, CH_2CH_3), 3.47 (AB, 4H, $J_{\text{AB}} = 13.35$, ArCH_2Ar), 3.99-4.03 (m, 4H, CH_2O), 4.26-4.34 (dd, 4H, $J = 12.81$, ArCH_2Ar); 5.13 (q, 1H, $J = 6.78$, CHCH_3), 6.72-6.77 (m, 2H, $H\text{-Ar}$), 6.93-7.12 (m, 4H, $H\text{-Ar}$), 7.36 (s, 1H, $\text{CH}=\text{N}(\text{O})$), 7.38-7.51 (m + s, 3H, $H\text{-Ar}$), 7.53 (d, 2H, $J = 6,96$, $H\text{-Ar}$), 7.64 (s, 2H, $H\text{-Ar}$), 8.04 (s, 1H, $H\text{-Ar}$), 8.11 (s, 1H, $H\text{-Ar}$), 8.81 (s, 1H, OH), 9.26 (s, 1H, OH), 9.81 (s, 1H, CHO); $^{13}\text{C NMR}$: 11.29 (CH_3CH_2), 19.43 (CH_3CH), 31.69 (ArCH_2Ar), 74.67 ($\text{CHN}(\text{O})$), 78.88 (OCH_2), 122.46, 125.93, 127.68, 128.33, 128.87, 129.13, 129.94, 130.17, 131.33, 132.63 (C, Ar), 133.38 ($\text{CH}=\text{N}(\text{O})$), 139.22, 150.26, 152.09, 156.08, 160.15 (C, Ar), 191.38 (CHO). IR: 1673 (CHO). ES-MS(+) for $\text{C}_{44}\text{H}_{45}\text{NO}_6$ (683.84) $m/z = 684.2$ $[\text{M}+\text{H}]^+$, 706.2 $[\text{M} + \text{Na}]^+$.

X-ray analysis of 3ab

Single crystals of **3a** and **3b** were obtained from a saturated dichloromethane / methanol solution. Data were collected at 293 K for **3a** and 173 K for **3b** on a Nonius Kappa CCD with Mo- $\text{K}\alpha$ radiation. Final unit cell parameters were obtained by means of a least-squares refinement. The structure has been solved by direct methods using SHELXS97. ²⁵ Usual non-H atoms were refined isotropically except for some disordered solvents. Hydrogen were calculated at theoretical positions and refined riding. The crystallographic data were reported on Table 1.

CCDC reference numbers 239937 for **3a** and 239938 for **3b**.

Crystal data for derivative 3a : $\text{C}_{55}\text{H}_{76}\text{Cl}_2\text{N}_2\text{O}_6$, $M = 930.51$, $T = 293(2)$ K, $a = 16.224(2)$, $b = 16.224(3)$, $c = 20.088(4)$ Å, $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$, $V = 5287.3(15)$ Å³, space group P4(1)2(1)2, $Z = 4$, $\mu = 0.188$ mm⁻¹, $D_c = 1.171$ g.cm⁻³, $wR_2 = 0.1639$, RI [$I > 2\sigma(I)$] = 0.0667.

Crystal data for derivative 3b : $\text{C}_{55}\text{H}_{56}\text{Cl}_2\text{N}_2\text{O}_4$, $M = 878.36$, $T = 173(2)$ K, $a = 15.297(2)$, $b = 15.297(2)$, $c = 19.770(4)$ Å, $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$, $V = 4626.1(13)$ Å³, space group P4(1)2(1)2, $Z = 4$, $\mu = 0.188$ mm⁻¹, $D_c = 1.229$ g.cm⁻³, $wR_2 = 0.1982$, RI [$I > 2\sigma(I)$] = 0.0739.

References

- [1] See, for example: (a) Gutsche, C. D. (1998) *Calixarenes Revisited*, Cambridge, Royal Society of Chemistry; (b) Gutsche, C. D. (2001) *Calixarenes 2001*, Dordrecht, Kluwer Academic Publishers.

- 1
2
3
4
5 [2] (a) Hamer, J.; Macaluso, A. *Chem Rev.* **1964**, *64*, 473. (b) Delpierre, G. R.; Lamchen, M. *Quart. Rev.*,
6 **1965**, 329. (c) Breuer, E. (1982) *The Chemistry of Amino Nitroso and Nitro Compounds and Their*
7 *Derivatives*, Chichester, West Sussex, Wiley. (d) Stamm, H. *Methodicum Chemicum* **1975**, *6*, 329.
- 8
9
10 [3] (a) Black, D. St. C.; Crozier, R. F.; Davis, V. C. *Synthesis* **1975**, 205. (b) Tufariello, J. J. (1984) *1,3-*
11 *Dipolar Cycloaddition Chemistry*, New York, Wiley. (c) Padwa, A. *Angew. Chem., Int. Ed. Engl.*
12 **1976**, *15*, 123. (d) Oppolzer, W. *Angew. Chem., Int. Ed. Engl.* **1977**, *16*, 10. (e) Balasubramanian, N.
13 *Org. Prep. Proc. Int.* **1985**, *17*, 23.
- 14
15
16 [4] Tufariello, J. J. *Acc. Chem. Res.* **1979**, *12*, 396.
- 17
18 [5] Janzen, E. G. *Acc. Chem. Res.* **1971**, *4*, 31.
- 19
20 [6] Marchand-Brynaert, J.; Bounkhala-Khrouz, Z.; H. Vanlierde, Z.; Ghosez, L. *Heterocycles* **1990**, *30*,
21 971.
- 22
23 [7] Warshaw, J. A.; Gallis, D. E.; Acken, B. J.; Gonzales, O. J.; Crist, D. R. *J. Org. Chem.* **1989**, *54*,
24 1736.
- 25
26 [8] Bourguet, E.; Baneres, J. L.; Girard, J. P.; Parello, J.; Vidal, J. P.; Lusinchì, X.; Declercq, J. P. *Org.*
27 *Lett.* **2001**, *3*, 3067.
- 28
29 [9] (a) Davies, M.; Timmins, G. *Paramagn. Reson.* **2000**, *17*, 1. (b) Bardelang, D.; Rockenbauer, A.;
30 Karoui, H.; Finet, J. P.; Tordo, P. *J. Phys. Chem. B.* **2005**, *109*, 105.
- 31
32 [10] Emmons, W. D. *Chem. Heterocycl. Compds.* **1964**, *19*, 624.
- 33
34 [11] Milliet, P.; Lusinchì, X. *Tetrahedron Lett.* **1971**, *40*, 3763.
- 35
36 [12] Splitter, J. S.; Su, T. M.; Ono, H.; Calvin, M. *J. Am. Chem. Soc.* **1971**, *93*, 4075.
- 37
38 [13] (a) Ogata, Y.; Sawaki, Y. *J. Am. Chem. Soc.*, 1973, **95**, 4687. (b) Ogata, Y.; Sawaki, Y. *J. Am. Chem.*
39 *Soc.* **1973**, *95*, 4692.
- 40
41 [14] Boyd, D. R.; Coulter, P. B.; Sharma, N. D.; Jennings, W. B.; Wilson, V. E. *Tetrahedron Lett.* **1985**,
42 26, 1673.
- 43
44 [15] Børgo, J.; Boyd, D. R. *J. Chem. Soc., Perkin Trans.2* **1973**, 1575.
- 45
46 [16] Boyd, A. J.; Boyd, D. R.; Malone, J. F.; Sharma, N. D.; Dev, K.; Jennigs, W. B. *Tetrahedron Lett.*
47 **1984**, *25*, 2497.
- 48
49 [17] Arduini, A.; Fabbi, M.; Mantovani, M.; Mirone, L.; Pochini, A.; Secchi, A.; Ungaro, R. *J. Org.*
50 *Chem.* **1995**, *60*, 1454.
- 51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 [18] Seigle-Ferrand, P.; Ben Sdira, S.; Felix, C.; Lamartine, R.; Bavoux, C.; Fenet, B.; Bayard, F.;
5
6 Vocanson, F. *Mater. Sci. Eng. C* **2006**, *26*, 181.
7
- 8 [19] Struck, O.; Chrisstoffels, L. A.; Lugtenberg, R. J.; Verboom, W.; Van Hummel, G. J.; Harkema, S.;
9
10 Reinhoudt, D. N. *J. Org. Chem.* **1997**, *62*, 2487.
11
- 12 [20] Solovyov, A.; Cherenok, S.; Tsymbal, I.; Failla, S.; Consiglio, G. A.; Finocchiaro, P.; Kalchenko, V.
13
14 *Heteroatom. Chem.* **2001**, *12*, 58.
15
- 16 [21] Kraïem, J.; Kacem, Y.; Khiari, J.; Ben Hassine, B. *Synth. Commun.* **2001**, *2*, 263.
17
- 18 [22] Kloc, K.; Kubicz, E.; Młochowski, J.; Syper, L. *Synthesis* **1987**, *12*, 1084.,
19
- 20 [23] Ulrich, G.; Turek, P.; Ziesel, R. *Tetrahedron Lett.* **1996**, *37*, 8755.
21
- 22 [24](a) Böhmer, V.; Jung, K.; Schön, M.; Wolff, A. *J. Org. Chem.* **1992**, *57*, 790. (b) Gutsche, C. D.; See,
23
24 K. A. *J. Org. Chem.* **1992**, *57*, 4527. (c) Lazzarotto, M.; Sansone, F.; Baldini, V.; Casnati, A.; Cozzini,
25
26 P.; Ungaro, R. *Eur. J. Org. Chem.* **2001**, 595.
27
- 28 [25] SHELX97 (includes SHELXS97, SHELXL97) – Programs for Crystal Structure Analysis (release 97-
29
30 2) : G. M. Sheldrick, Institut für Anorganische Chemie der Universität, Göttingen, Germany, 1998
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 1. Selected ^1H NMR chemical shifts (ppm) of **4ab** and **5ab** in CDCl_3 .

	4a	4b	5a	5b
<i>CH=N(O)</i>	7.21	7.35	7.22	7.36
<i>OH</i>	8.83	8.76	8.83 and 9.30	8.81 and 9.26
<i>CH</i> ₂ (bridges)	3.48 (d) and 4.30 (d)	3.42 (d) and 4.24 (d)	3.46-3.56 (dd) 4.28-4.35 (dd)	3.47 (dd) 4.26-4.34 (dd)
	AB system	AB system	2 AB systems	2 AB systems
<i>CHO</i>	/	/	9.80	9.81

Table 2. Selected ^{13}C NMR chemical shifts (ppm) of **4ab** and **5ab** in CDCl_3 .

	4a	4b	5a	5b
CH=N(O)	133.92	133.58	133.42	133.38
CH ₂	31.69	31.77	30.51	31.69
(bridges)				
CHO	/	/	191.37	191.38




Fig. 1. X-ray crystal structure of **3a** included dichloromethane and methanol. The hydrogen bonds are indicated by dash lines.




Fig. 2. X-ray crystal structure of **3b**. The hydrogen bonds are indicated by dash lines.