



**HAL**  
open science

## Synthesis of highly functionalized fluorinated porphyrins

Tomas Briza, Robert Kaplánek, Vladimír Král

► **To cite this version:**

Tomas Briza, Robert Kaplánek, Vladimír Král. Synthesis of highly functionalized fluorinated porphyrins. *Supramolecular Chemistry*, 2008, 20 (03), pp.237-242. 10.1080/10610270601120371 . hal-00513500


**HAL Id: hal-00513500**

**<https://hal.science/hal-00513500>**

Submitted on 1 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Synthesis of highly functionalized fluorinated porphyrins

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2006-0074.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	06-Nov-2006
Complete List of Authors:	Briza, Tomas; ICT - Prague Kaplánek, Robert; ICT-Prague Král, Vladimír; ICT Prague, Analytical Chem
Keywords:	fluorinated dendritic structure, polyfluoroalkylated porphyrin, fluorous, porphyrin core



# Synthesis of highly functionalized fluorinated porphyrins

TOMÁŠ BŘÍZA<sup>a,b\*</sup>, ROBERT KAPLÁNEK<sup>a,b\*</sup>, MARTIN HAVLÍK<sup>a</sup>, BOHUMIL DOLENSKÝ<sup>a</sup>, ZDENĚK KEJÍK<sup>a</sup>, PAVEL MARTÁSEK<sup>b</sup> and VLADIMÍR KRÁL<sup>a</sup>

<sup>a</sup>*Department of Analytical Chemistry, Institute of Chemical Technology in Prague, Technická 5, Prague 6, 16628, Czech Republic*

<sup>b</sup>*First Faculty of Medicine, Charles University in Prague, Kateřinská 32, Prague 2, 12108, Czech Republic*

\* corresponding authors: ftor@seznam.cz (T. Bříza), robert.kaplanek@vscht.cz (R. Kaplánek);  
fax: (+420) 224 310 352

**Highly functionalized fluorinated porphyrins were synthesized by a convergent strategy. Nucleophilic substitution using fluorinated branched unit and 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin afforded highly functionalized fluorinated porphyrin 8 containing twenty-four fluorinated chains in the peripheral region.**

*Keywords:* fluorinated dendritic structure, polyfluoroalkylated porphyrin, fluorous, porphyrin core

## INTRODUCTION

Fluorinated highly branched structures are a specific class of highly functionalized compounds possessing remarkable properties. Several fluorinated dendrimers have been synthesized to date [1]. These fluorinated macromolecules can be utilized, for example, as surfactants in biphasic systems [2], phase transfer catalysts [3], and nanoreactors [4]. In the field of material science they have been used as electron transport materials for organic light-emitting diodes [5], materials for non-linear optic [6] and, in the field of biomedical chemistry, as drug delivery systems [7]. Known fluorinated highly branched structures reported previously contain core such as aryls [8], alkylsilanes [9], aminoalkylmethanes [2], azacrowns [10], thiophosphates [11], fluorophenyls [12], cations of transition metals [13], etc. To the best of our knowledge, there are only a few polyfluoralkylated porphyrins.

In 1997 Pozzi et al. prepared highly fluorinated porphyrin [14]. It contained eight polyfluorinated chains and was used as a ligand for fluorous biphasic catalysis [14]. In 2002 we prepared porphyrin substituted with four polyfluorinated chains, which showed complexation ability for fluorinated aromatic compounds [15]. Thus, our aim is to develop synthetic routes for the preparation of highly functionalized fluorinated porphyrins which could serve in novel investigations.

## RESULTS AND DISCUSSION

Our strategy for preparation of highly functionalized fluorinated porphyrins is based on, analogous to syntheses of dendrimers, convergent approach. Suitable dendrons are synthesized and then connected to the porphyrin skeleton. The design utilizes highly functionalized benzene derivatives.

Initially, we looked for a suitable building block for the construction of the branched fluorinated unit. We chose methyl gallate, because it enables the connection of three fluorinated units *via* hydroxyl groups. Moreover, it bears an ester moiety, which can be employed after another transformation (reduction) as a focal site for monomer connectivity.

As a model dendritic unit we chose a gallate substituted with three 3-(perfluorohexyl)propyl chains. Similar gallates with three fluorinated chains was already prepared [16], but the fluorinated chains were separated from the gallate group by a four carbon spacer. Our method enables one to prepare dendrons whereby perfluoralkylated chains are insulated by three carbon spacers only. As a fluoroalkylating agent we used 3-(perfluorohexyl)propyl triflate prepared under similar conditions to those of 2-(perfluorohexyl)ethyl triflate [17], [18]. The triflate was reacted with the trisodium salt of gallate and the reaction afforded corresponding polyfluorinated gallate **2**. The gallate ester **2** was reduced to alcohol **3** by  $\text{LiAlH}_4$ . The alcohol **3** was oxidized by mangan (IV) oxide to aldehyde **4**. Figure 1




FIGURE 1 Synthesis of model dendritic units

For final synthesis to attain dendritic fluorinated porphyrins we chose two synthetic approaches. In the first approach we tried to utilize the common method for the construction of porphyrins from an aldehyde and pyrrole [19]. Thus, we reacted aldehyde **4** with pyrrole under conditions for preparation of the porphyrins [19]. Unfortunately, the reaction afforded only a complex and intractable mixture of unidentified compounds.

Therefore, we applied the second approach, based on the nucleophilic substitution of 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin by fluorinated alcohol **3**. Alcohol **3** was converted to corresponding sodium salt with NaH and reacted with the porphyrin in DMF. The reaction afforded polyfluorinated porphyrin **5** with twelve perfluorohexyl chains in the peripheral region. Figure 2.




FIGURE 2 Synthesis of porphyrin **5**

Because the model synthesis of fluorinated porphyrin **5** was successful, we decided to utilize this approach for the construction of larger fluorinated dendritic porphyrins with double the number of fluorinated chains. Initially, it was necessary to prepare a suitable dendron with the required number of fluorinated chains. At this point, we were able to use to our great advantage our recently published branched fluorinated triflate [20]. According to our published results we prepared corresponding fluorinated gallate **6** with six fluorinated chains from gallate **1** and branched triflate. The ester group was reduced with  $\text{LiAlH}_4$  to alcohol **7**.

Alcohol **7** was used according to the model reaction with compound **3** described above, for the construction of a fluorinated porphyrin **5**. Compound **7** was transformed to the sodium salt with NaH and reacted with 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin. The reaction afforded highly functionalized fluorinated porphyrin **8** with twenty-four fluorinated chains in the peripheral region. Figure 3.




FIGURE 3 Synthesis of porphyrin **8**

The solubility of both porphyrins **5** and **8** is dramatically different. The porphyrin **5** is soluble in common organic solvents as dichloromethane, chloroform, diethyl ether, acetone etc. On the other hand, porphyrin **8** with twofold number of fluorinated chains dramatically decreased its solubility in common organic solvents with the exception of diethyl ether in which is surprisingly well soluble. The solubility of porphyrin **8** can be enhanced by adding freon  $\text{CCl}_3\text{F}$  (F 113) to common solvents. Thus, NMR spectra were successfully obtained in a mixture of  $\text{CDCl}_3$  and F 113.

The detailed structure of porphyrin **8** is shown below. Figure 4.




FIGURE 4 Porphyrin **8** ( $C_{321}H_{206}F_{328}N_4O_{16}$ )

## CONCLUSION

We have synthesized fluorinated highly functionalized porphyrins **5** and **8** by a convergent synthesis. The synthesis was based on a nucleophilic substitution between dendritic units **3**, **7** and 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin. Porphyrins **5** and **8** will be studied as a potential ligands, drug delivery systems, and as a possible shuttle for the transport of gases.

## EXPERIMENTAL

### Measurements and materials

All chemicals were purchased from Aldrich and used without further purification. All reactions required anhydrous conditions were performed using dried solvents and under argon atmosphere. NMR spectra were recorded on a Varian Gemini 300 HC (FT,  $^1\text{H}$  at 300MHz,  $^{13}\text{C}$  at 75 MHz,  $^{19}\text{F}$  at 281 MHz) instrument using TMS and  $\text{CFCl}_3$  as the internal standards. Chemical shifts are quoted in ppm ( $\delta$ -scale; s-singlet, bs-broad singlet, d-doublet, t-triplet, q-quadruplet, m-multiplet), coupling constants J in Hz, solvents  $\text{CDCl}_3$ . IR spectrometry analyses were performed on a Bruker IFS 66v/S instrument. Mass spectrometry analyses were performed on a Hewlett-Packard GC HP5890 instrument.

### Compound 2

A flask was charged with sodium hydride (60% suspension in mineral oil, 160 mg, 3.96 mmol) and THF (3 mL). The mixture was cooled to 0 °C and solution of methyl gallate **1** (220 mg, 1.19 mmol) in THF (5 mL) was slowly added. When addition was complete, next portion of THF (30 mL) was added to the mixture. Then cooling bath was removed and the mixture was stirred at laboratory temperature for 30 min. After this time, solution of [3-(perfluorohexyl)propyl] triflate (2.00 g, 3.92 mmol) in THF (12 mL) was dropped to the reaction mixture. When addition was complete, the mixture was refluxed for 1 h. After this time, the mixture was diluted with water (10 mL) and brine (50 mL). The mixture was extracted with diethyl ether (3×50 mL). The organic phase was dried over  $\text{MgSO}_4$ . Drying agent was removed by filtration and solvent was evaporated by means of vacuum rotary evaporator. The product was separated by column chromatography on silica (2.5×25 cm,

1  
2  
3 eluent – dichloromethane). The product **2** was obtained as yellowish viscous oil (1.210 g, 80  
4  
5  
6 %).

7  
8  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 2.01 (m, 2H), 2.14 (m, 4H), 2.38 (m, 6H), 3.89 (s, 3H), 4.59 (t, 2H,  $^3J_{\text{HH}}$   
9  
10 = 5.6), 4.12 (t, 4H,  $^3J_{\text{HH}}$  = 5.9), 7.29 (s, 2H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 20.6 (s, 2C), 21.5 (s, 1C),  
11  
12 27.8 (t, 3C,  $^2J_{\text{CF}}$  = 22), 52.0 (s, 1C), 67.2 (s, 2C), 71.9 (s, 1C), 108.2 (s, 2C), 107.0-120.0 (m,  
13  
14 18C), 125.7 (s, 1C), 141.6 (s, 1C), 152.3 (s, 2C), 166.4 (s, 1C);  $^{19}\text{F}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = -81.3  
15  
16 (m, 9 F), -115.2 (m, 6F), -122.5 (m, 6F), -123.5 (m, 6F), -124.4 (m, 6F), -126.8 (m, 6F). Anal.  
17  
18 Calcd for  $\text{C}_{35}\text{H}_{23}\text{F}_{39}\text{O}_5$ : C, 33.25; H, 1.83; Found: C, 33.74; H, 2.34.  
19  
20  
21  
22  
23  
24

### 25 **Compound 3**

26  
27 A flask was charged with gallate **2** (100 mg, 79  $\mu\text{mol}$ ) and diethyl ether (5 mL). The mixture  
28  
29 was cooled to  $-78\text{ }^\circ\text{C}$  and  $\text{LiAlH}_4$  (1 M solution in diethyl ether, 9 mg, 237  $\mu\text{mol}$ ) was slowly  
30  
31 added. Then, the cooling bath was removed and the mixture was stirred at laboratory  
32  
33 temperature for 1 h. The reaction was quenched with MeOH (2 mL), diluted with water (50  
34  
35 mL) and mixture was extracted with diethyl ether (3 $\times$ 50 mL). The organic phase was dried  
36  
37 over  $\text{MgSO}_4$ . Drying agent was removed by filtration and solvent was evaporated by means of  
38  
39 vacuum rotary evaporator. The product was separated by column chromatography on silica  
40  
41 (2.5 $\times$ 20 cm, eluent – diethyl ether). The product **3** was obtained as yellowish viscous oil (91  
42  
43 mg, 93 %).  
44  
45  
46  
47  
48

49  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 2.00 (m, 2H), 2.12 (m, 4H), 2.35 (m, 6H), 3.95 (t, 2H), 4.08 (t, 4H),  
50  
51 4.62 (s, 2H), 6.60 (s, 2H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 20.6 (s, 2C), 21.3 (s, 1C), 27.8 (t, 2C,  $^2J_{\text{CF}}$  =  
52  
53 22), 27.9 (t, 1C,  $^2J_{\text{CF}}$  = 22), 65.1 (s, 1C), 67.5 (s, 2C), 71.8 (s, 1C), 105.3 (s, 2C), 107.0-122.0  
54  
55 (m, 18C), 136.7 (s, 1C), 137.1 (s, 2C), 152.6 (s, 1C);  $^{19}\text{F}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = -81.4 (m, 9 F),  
56  
57 -114.9 (m, 6F), -122.5 (m, 6F), -123.4 (m, 6F), -124.1 (m, 6F), -126.8 (m, 6F). Anal. Calcd  
58  
59 for  $\text{C}_{34}\text{H}_{23}\text{F}_{39}\text{O}_4$ : C, 33.03; H, 1.87; Found: C, 33.54; H, 2.49.  
60

**Compound 4**

A flask was charged with gallate **3** (1.01 g, 0.82 mmol), dichloromethane (20 mL) and mangan (IV) oxide (248 mg, 2.85 mmol). The mixture was stirred under reflux for 4 h and then next portion of mangan (IV) oxide (250 mg, 2.88 mmol) was added, heating bath was removed and the mixture was stirred at laboratory temperature overnight. Then the mixture was evaporated to dryness, diluted with diethyl ether and filtered. The product was separated by column chromatography (2.5×25 cm, eluent – dichloromethane). The isolation afforded product **4** as viscous colorless oil (260 mg, 26 %) and starting alcohol (430 mg).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 2.03 (m, 2H), 2.15 (m, 4H), 2.35 (m, 6H), 4.11 (t, 2H,  $^3J_{\text{HH}} = 5.8$ ), 4.14 (t, 4H,  $^3J_{\text{HH}} = 5.8$ ), 7.16 (s, 2H), 9.84 (s, 1H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 20.6 (s, 2C), 21.5 (s, 1C), 27.8 (t, 2C,  $^2J_{\text{CF}} = 22$ ), 67.7 (s, 2C), 72.0 (s, 1C), 108.1 (s, 2C), 107.5-121.7 (m, 18C), 132.1 (s, 2C), 143.0 (s, 1C), 153.0 (s, 1C), 190.7 (s, 1C);  $^{19}\text{F}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = -81.58 (t, 6F,  $^3J_{\text{FF}} = 10$ ), -81.64 (t, 3F,  $^3J_{\text{FF}} = 10$ ) -115.0 (m, 6F), -122.5 (m, 6F), -123.5 (m, 6F), -124.1 (m, 6F), -126.8 (m, 6F). Anal. Calcd for  $\text{C}_{34}\text{H}_{21}\text{F}_{39}\text{O}_4$ : C, 33.08; H, 1.71; Found: C, 33.31; H, 1.92.

**Porphyrin 5**

A flask was charged with compound **3** (50 mg, 40  $\mu\text{mol}$ ), anhydrous DMF (2 mL) and sodium hydride (60 % suspension in mineral oil, 2 mg, 84  $\mu\text{mol}$ ). The mixture was stirred at laboratory temperature for 20 min. The 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin was added and the mixture was heated to 90 °C overnight. Then the mixture was evaporated to dryness and product was separated by column chromatography on silica (2×20 cm, eluent – dichloromethane/petroleum ether 1:1). The product **5** was obtained as dark red wax (7 mg, 11 %)

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ = -2.91 (2H, s), 2.08 (8H, m), 2.19 (16H, m), 2.3-2.5 (24H, m), 4.10 (8H, m), 4.19 (12H, t, <sup>3</sup>J<sub>HH</sub> = 5.5 Hz), 5.51 (8H, s), 6.88 (8H, s), 8.85 (8H, s); <sup>19</sup>F NMR (CDCl<sub>3</sub>) δ = -81.4 (9F, t, J<sub>FF</sub> = 10 Hz), -115.0 (6F, m), -122.5 (6F, m), -123.4 (6F, m), -124.1 (6F, m), -126.7 (6F, m), -139.0 (2F, dd, J<sub>FF</sub> = 7 Hz, J<sub>FF</sub> = 22 Hz), -156.3 (2F, dd, J<sub>FF</sub> = 7 Hz, J<sub>FF</sub> = 22 Hz); MS (MALDI): for C<sub>180</sub>H<sub>98</sub>F<sub>172</sub>N<sub>4</sub>O<sub>16</sub> Calculated: 5840.5, founded: 5839.5.

### Compound 7

A flask was charged with compound **6** (880 mg, 0.36 mmol) and THF (40 mL). The mixture was cooled to 0 °C and LiAlH<sub>4</sub> (1 M solution in diethyl ether, 20 mg, 0.53 mmol) was slowly dropped under vigorous stirring. Then the mixture was stirred for 30 minutes at this temperature and then was slowly warmed to laboratory temperature within 15 min. The reaction was quenched with MeOH (3 mL), diluted with water (50 mL) and mixture was extracted with diethyl ether (3×50 mL). The organic phase was dried over MgSO<sub>4</sub>. Drying agent was removed by filtration and solvent was evaporated by means of vacuum rotary evaporator. The product was separated by column chromatography on silica (2.0×20 cm, eluent – hexane/dichloromethane – 1:1). The product **7** was obtained as yellowish viscous oil (760 mg, 87 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.50 – 1.74 (m, 24H), 1.82-1.90 (m, 3H), 1.98-2.14 (m, 12H), 3.83 (d, 1H, <sup>3</sup>J<sub>HH</sub> = 5.49), 3.92 (d, 2H, <sup>3</sup>J<sub>HH</sub> = 4.94), 4.63 (s, 2H), 6.59 (s, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 17.7 (s, 12C), 30.9 (t, 6C, <sup>2</sup>J<sub>CF</sub> = 26), 38.2 (s, 2C), 38.9 (s, 1C), 65.4 (s, 1C), 70.6 (s, 2C), 75.2 (s, 1C), 104.9 (s, 2C), 107.4-121.7 (m, 36C), 136.7 (s, 2C), 136.9 (s, 1C), 152.9 (s, 1C). <sup>19</sup>F NMR (CDCl<sub>3</sub>) δ -81.6 (m, 18F), -115.1 (m, 12F), -122.6 (m, 12F), -123.6 (m, 12F), -124.3 (m, 12F), -126.9 (m, 12F). MS (LRMS): for C<sub>67</sub>H<sub>50</sub>F<sub>78</sub>O<sub>4</sub> Calculated: 2401, Found: 2400.

## Porphyrin 8

A flask was charged with compound **7** (270 mg, 0.11 mmol), anhydrous DMF (10 mL) and sodium hydride (60 % suspension in mineral oil, 10 mg, 0.25 mmol). The mixture was stirred at laboratory temperature for 20 min. The 5,10,15,20-tetrakis(pentafluorophenyl)porphyrin (10 mg, 0.01 mmol) was added and the mixture was heated to 90 °C overnight. The mixture was diluted with water (10 mL) and was extracted with trifluoromethyl toluene (2 ×10 ml). The organic phase was dried over MgSO<sub>4</sub>. Drying agent was removed by filtration and solvent was evaporated by means of vacuum rotary evaporator. The product was separated by column chromatography on silica (2.5×20 cm, eluent – petroleum ether/dichloromethane 1:1). The product was obtained in a form of dark red semisolid wax (45 mg). The second purification was accomplished by preparative TLC on silica (20 × 20 cm, 2000 microns, eluent – petroleum ether/dichloromethane 3:1). The pure product **8** was obtained as red dark wax (30 mg, 28%).

<sup>1</sup>H NMR (CDCl<sub>3</sub> / CCl<sub>3</sub>F) δ -2.82 (s, 2H), 1.50 – 2.20 (m, 156H), 3.98 (d, 8H, <sup>3</sup>J<sub>HH</sub> = 5.23), 4.07 (d, 16H, <sup>3</sup>J<sub>HH</sub> = 4.95), 5.52 (s, 8H), 6.90 (s, 8H), 8.87 (s, 8H); <sup>19</sup>F NMR (CDCl<sub>3</sub> / CCl<sub>3</sub>F) δ -81.6 (t, 72F, <sup>3</sup>J<sub>FF</sub> = 10), -115.1 (m, 48F), -122.6 (m, 48F), -123.6 (m, 48F), -124.3 (m, 48F), -126.9 (m, 48F), -138.9 (8F, dd, <sup>3</sup>J<sub>FF</sub> = 7.3, <sup>4</sup>J<sub>FF</sub> = 22.0), -156.9 (8F, dd, <sup>3</sup>J<sub>FF</sub> = 7.3, <sup>4</sup>J<sub>FF</sub> = 22.0); IR (KBr) cm<sup>-1</sup> 2924, 1499, 1368, 1241, 1206, 1145, 1121. MS (MALDI): for C<sub>321</sub>H<sub>206</sub>F<sub>328</sub>N<sub>4</sub>O<sub>16</sub> Calculated: 10498.61, Found: 10499.52.

## Acknowledgement

The authors thank the Grant agency of Czech Academy of Science for financial support of this project, Grant No. KJB 401 280 501, and MSMT (CAG, 1M6837805002), MSMT (0021620806).

**Reference**

- [1] Caminade, A.-M.; Turrin, C.-O.; Sutra, P.; Majoral, J.-P. *Current Opinion in Colloid and Interface Science* **2003**, *8*, 282.
- [2] Cooper, A. I.; Londono, J. D.; Wignall, G.; McClain, J. B.; Samulski, E. T.; Lin, J. S.; Obrynin, A.; Rubinstein, M.; Burke, A. L. C.; Fréchet, J. M. J.; DeSimone, J. M. *Nature* **1997**, *389*, 368.
- [3] Goetheer, E. L. V.; Baars, M. W. P. L.; van den Broeke, L. J. P.; Meijer, E. W.; Keurentjes, J. T. F. *Ind. Eng. Chem. Res* **2000**, *39*, 4634.
- [4] Yeung, L. K.; Crooks, R. M. *Nano Lett.* **2001**, *1*, 14.
- [5] Sakamoto, Y.; Sužuji, T.; Miura, A.; Fujikawa, H.; Tokito, S.; Taga, Y. *J. Am. Chem. Soc.* **2000**, *122*, 1832.
- [6] Luo, J.; Ma, H.; Haller, M.; Jen, A. K. Y.; Barto, R. R. *Chem. Commun.* **2002**, 888.
- [7] Zhuo, R. X.; Du, B.; Lu, Z. R. *J. Control. Release* **1999**, *57*, 249.
- [8] Miller, T. M.; Neenan, T. X.; Zayas, R.; Bair, H. E. *J. Am. Chem. Soc.* **1992**, *114*, 1018.
- [9] Lorenz, K.; Frey, H.; Stuehn, B.; Muelhaupt, R. *Macromolecules* **1997**, *30*, 6860.
- [10] Zhuo, R. X.; Du, B.; Lu, Z. R. *J. Control. Release* **1999**, *57*, 249.
- [11] Turrin, C. O.; Maraval, V.; Leclaire, J.; Dantras, E.; Lacabanne, C.; Caminade, A. M.; Majora, J.-P. *Tetrahedron* **2003**, *59*, 3965.
- [12] Sakamoto, Y.; Suzuki, T.; Miura, A.; Fujikawa, H.; Tokito, S.; Taga, Y. *J. Am. Chem. Soc.* **2000**, *122*, 1832.
- [13] Pitois, C.; Vestberg, R.; Rodlert, M.; Malmstrom, E.; Hult, A.; Lindgren, M. *Opt. Mater.* **2003**, *21*, 499.
- [14] Pozzi, G.; Montanari, F.; Quici, S. *Chem. Comm.* **1997**, 69.
- [15] Paleta, O., Beneš, M., Koutníková, J., Král, V. *Tetrahedron Lett.* **2002**, *43*, 6827.
- [16] a) Johansson, G.; Percec, V.; Ungar, G.; Zhou, J. P. *Macromolecules* **1996**, *29*, 646. b) Markowicz, M. W.; Dembinski, R. *Org. Lett.* **2002**, *22*, 3785. c) Percec, V.; Glodde, M.; Johansson, G.; Balagurusamy, V. S. K.; Heiney, P. A. *Angew. Chem. Int. Ed.* **2003**, *42*, 4338.
- [17] a) Nohira, H.; Sakaigawa, A.; Yokoi, S. *Japan Patent Application*, JP 06264058 A2 19940920, **1994**. b) Negoro, M.; Kawata, K. *American Patent Application*, US 2006068998 A1 20060330, **2006**.
- [18] Bříza, T.; Kvíčala, J.; Mysík, P.; Paleta, O.; Čermák, J. *Synthetic Lett.* **2001**, *5*, 685.
- [19] Feirong, L.; Kexin, Y.; John, S.; Tyhonas, K.; MacCrum, A.; Lindsey, J. S. *Tetrahedron* **1997**, *53*, 12339.

1  
2  
3 [20] Kaplánek, R.; Bříza, T.; Havlík, M.; Dolenský, B.; Kejík, Z.; Martásek, P.; Král, V. *J. Fluor. Chem.* **2006**,  
4  
5 127, 386.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only

## Figures




FIGURE 1 Synthesis of model dendritic units




FIGURE 2 Synthesis of porphyrin 5




FIGURE 3 Synthesis of porphyrin 8




FIGURE 4 Porphyrin **8** ( $C_{321}H_{206}F_{328}N_4O_{16}$ )