



**HAL**  
open science

## Cyclophanes Or Cyclodextrins: What Is The Best Host For Aromatic Volatile Organic Compounds ?

Paul Ionut Dron, Sophie Fourmentin, Francine Cazier, David Landy,  
Gheorghe Surpateanu

► **To cite this version:**

Paul Ionut Dron, Sophie Fourmentin, Francine Cazier, David Landy, Gheorghe Surpateanu. Cyclophanes Or Cyclodextrins: What Is The Best Host For Aromatic Volatile Organic Compounds?. *Supramolecular Chemistry*, 2008, 20 (05), pp.473-477. 10.1080/10610270701358525 . hal-00513512

**HAL Id: hal-00513512**

**<https://hal.science/hal-00513512>**

Submitted on 1 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Cyclophanes Or Cyclodextrins: What Is The Best Host For Aromatic Volatile Organic Compounds ?

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2007-0044.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	13-Mar-2007
Complete List of Authors:	Ionut Dron, Paul; LSOE Fourmentin, Sophie; LSOE, MREI Cazier, Francine; LSOE Landy, David; LSOE Surpateanu, Gheorghe; LSOE
Keywords:	Cyclophanes, Cyclodextrins, Inclusion compounds, Static headspace, Molecular modelling
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p> <p>Figure1.cdx</p>	



1  
2  
3 Cyclophanes Or Cyclodextrins: What Is The Best Host For Aromatic Volatile Organic  
4  
5  
6 Compounds ?  
7  
8  
9

10 Paul Ionut Dron, Sophie Fourmentin<sup>1</sup>, Francine Cazier, David Landy and Gheorghe  
11  
12 Surpateanu  
13

14  
15  
16  
17 Laboratoire de Synthèse Organique et Environnement (EA2599), Université du Littoral  
18  
19 Côte d'Opale, 145 Avenue Maurice Schumann, 59140 Dunkerque, France.  
20  
21

22  
23  
24 **Abstract**  
25  
26  
27

28  
29 The first results of the complexing ability of cyclobis(paraquat-*p*-phenylene) as  
30  
31 supramolecular host with different aromatic volatile organic compounds are presented.  
32  
33 The formation constants of cyclobis(paraquat-*p*-phenylene) with toluene and  
34  
35 halogenobenzenes were determined in aqueous solution by static headspace associated  
36  
37 with gas chromatography and compared with the ones obtained by cyclodextrins. The  
38  
39 data indicated the formation of 1:1 inclusion compounds in both cases. The results  
40  
41 underlined a greater complexation ability for cyclobis(paraquat-*p*-phenylene) which was  
42  
43 confirmed by a theoretical study.  
44  
45  
46  
47  
48  
49

50  
51 Keywords: Cyclophanes, Cyclodextrins, Inclusion compounds, Static headspace,  
52  
53 Molecular modelling  
54  
55  
56  
57

58 **Introduction**  
59  
60

---

<sup>1</sup>Corresponding author email: lamotte@univ-littoral.fr

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Cyclobis(paraquat-*p*-phenylene) (CBPQT<sup>4+</sup>) (Figure 1) and its inclusion complexes have drawn much attention because of their many interesting properties and applications especially in the design of nanoscale molecular devices [1,2]. This cyclophane can be employed as a binding probe for studies in molecular recognition. Our interest lies in volatile organic compounds (VOCs) recognition by supramolecular hosts like modified cyclodextrins (CDs) (Figure 1) or cyclophanes [3,4]. Indeed, the internal cavity of CBPQT<sup>4+</sup> has a shape and hydrophobicity that make it a selective inclusion host for aromatic residues [5,6].

### Insert Figure 1

Association of CBPQT<sup>4+</sup> with an electron rich aromatic guest produces a colored, charge-transfer complex. Binding constants are then usually determined by monitoring the appearance of this complex using standard spectrometric methods. These methods are not appropriate in the case of VOCs as guest because of the low solubility and the high volatility of these compounds. We have recently developed a method based on static headspace gas chromatography (HSGC) to study the inclusion complex formed between CDs and VOCs [7]. This method has been applied to the determination of the formation constant of the inclusion complexes formed between CBPQT<sup>4+</sup> tetrachloride salt,  $\alpha$ -cyclodextrin ( $\alpha$ -CD),  $\beta$ -cyclodextrin ( $\beta$ -CD) and randomly-methylated- $\beta$ -cyclodextrin (RAMEB) with toluene and halogenobenzenes in water.

### Results and discussions

1  
2  
3 In order to evaluate the inclusion ability of the supramolecular hosts, we used a static  
4 headspace method. Such a method is based on the variation of the peak area of the VOC  
5 observed upon addition of each host in a solution containing the VOC.  
6  
7

8  
9  
10 The stability constants for the inclusion of VOCs with CBPQT<sup>4+</sup>,  $\alpha$ -CD,  $\beta$ -CD and  
11 RAMEB in aqueous solution were determined at 30°C with four host concentrations  
12 (35mM, 20mM, 10mM and 5mM for RAMEB; 7mM, 4mM, 2mM and 1mM for  $\alpha$ -CD  
13 and  $\beta$ -CD; 0.3mM, 0.2mM, 0.1mM and 0.03mM for CBPQT<sup>4+</sup>). The effect of an  
14 increased concentration of CBPQT<sup>4+</sup> on the chromatographic peak of toluene is shown  
15 in Figure 2.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 **Insert Figure 2**  
29  
30  
31

32 The obtained variations are in good agreement with the 1:1 host/guest ratio, as could be  
33 expected from analogous results for other substituted benzene derivatives for  
34 cyclophanes [8,9] or cyclodextrins [10-13]. The experimental data obtained for  
35 iodobenzene and CBPQT<sup>4+</sup> are given in Figure 3 with the theoretical curve for a 1:1  
36 complex with a formation constant of 9459 M<sup>-1</sup>. As one can see, the experimental points  
37 fit well with the theoretical curve.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 **Insert Figure 3**  
51  
52

53 The formation constants are calculated by an algorithmic procedure [14] and are  
54 reported in Table 1 in comparison with literature data (in bracket) [11-13].  
55  
56  
57  
58  
59  
60

**Insert Table 1**

1  
2  
3  
4  
5  
6 There are few references in literature in which HSGC is used for determining stability  
7 constants of different compounds with CDs [15-20] and especially for VOCs  
8 [13,21,22].  
9

10  
11  
12 The obtained formation constants are in good agreement with the corresponding ones  
13 found in literature for CDs [11-13], while, in the case of CBPQT<sup>4+</sup>, the values are in the  
14 same order of magnitude than for other aromatic guests in aqueous media [8,9]. The  
15 formation constants obtained with CBPQT<sup>4+</sup> are 10 to 100-fold stronger than for CDs,  
16 which demonstrates a greater complexation ability of the cyclophane.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 In addition, if one plots the logarithm of the formation constant ( $\log K_f$ ) as a function of  
28 the molar refractivity (MR) of the substituent (Figure 4), a good linear relationship is  
29 obtained for both CBPQT<sup>4+</sup> and  $\beta$ -CD. This linear variation with the size of the  
30 substituent shows that Van der Waals interactions are important for the two  
31 supramolecular hosts. In addition, it has to be underlined that nearly identical slopes are  
32 obtained for the two hosts, whereas a greater intercept is observed for CBPQT<sup>4+</sup>. This  
33 leads to think that the greater complexation ability of the cyclophane is a consequence  
34 of a better encapsulation of the phenyl ring.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 **Insert Figure 4**  
54  
55  
56  
57  
58  
59  
60

53 In order to get better insights into the mechanisms of complexation, we have also  
54 realised a comparative docking study of iodobenzene into CBPQT<sup>4+</sup> and  $\beta$ -CD. The  
55 initial CBPQT<sup>4+</sup> structure was in agreement with literature data (geometrical parameters  
56  
57  
58  
59  
60

1  
2  
3 strictly equal to already published PM3 results) [23-24]. The optimal docked structures  
4  
5 are illustrated in Figure 5.  
6  
7  
8  
9

### 10 **Insert Figure 5**

11  
12  
13  
14  
15 It can be seen that a very close proximity exists between iodobenzene and CBPQT<sup>4+</sup>,  
16  
17 leading to a greater stabilisation by Van der Waals forces than for  $\beta$ -CD inclusion  
18  
19 compound. Indeed, a value of -15.7 kcal/mol is obtained in the case of CBPQT<sup>4+</sup>,  
20  
21 against -10.6 kcal/mol for  $\beta$ -CD. Moreover, the existence of charges on the cyclophane  
22  
23 structure implies greater electrostatic interactions with the guest than those taking place  
24  
25 with  $\beta$ -CD (-8.7 kcal/mol against -0.8 kcal/mol). Thus, experimental and theoretical  
26  
27 data suggest that both steric complementarity and charge-transfer mechanisms are  
28  
29 operating inside the cyclophane complex, whereas CD complex is mainly influenced by  
30  
31 Van der Waals interactions. As a consequence, the total stabilisation energy is largely in  
32  
33 favour of CBPQT<sup>4+</sup> (-24.4 kcal/mol against -11.6 kcal/mol), which is in good qualitative  
34  
35 agreement with the determined formation constants.  
36  
37  
38  
39  
40  
41  
42  
43

### 44 **Materials and methods**

#### 45 a) Chemicals

46  
47  
48  
49  
50  
51  
52  
53 Toluene and halogenobenzenes (Aldrich) were all of analytical reagent grade and were  
54  
55 used as received.  $\alpha$ -CD,  $\beta$ -CD and RAMEB were purchased from Wacker Chimie S.A.  
56  
57 (Lyon). Cyclobis(paraquat-*p*-phenylene) was synthesised in our laboratory as described  
58  
59 [6]. Tetrachloride salt of CBPQT<sup>4+</sup> was used because of its greater solubility (~2mM)  
60

1  
2  
3 compared with tetrakis(hexafluorophosphate) salt (~0.01mM). Deionised water was  
4  
5 used throughout this work.  
6  
7  
8  
9

10 b) Static headspace  
11

12  
13  
14  
15 Headspace sampling is employed with gas chromatography (GC) in numerous fields  
16 and with a variety of applications [25, 26]. A headspace sample is in principle a gas  
17 sample which has been previously in contact with a liquid or solid sample from which  
18 volatile compounds were released into the gas with subsequent analysis by gas  
19 chromatography. Headspace gas chromatography is therefore a technique of gas  
20 extraction and can be carried out comparable to a solvent extraction as a one-step  
21 extraction (static or equilibrium headspace) or as a continuous extraction (dynamic  
22 headspace) [27]. Measurements were conducted with a Agilent headspace autosampler.  
23  
24 Sample solutions of 5 mL containing various concentrations of pollutant (10-100 ppm)  
25 were introduced into 10 mL headspace vials and sealed using silicone septa and  
26 aluminium foil. The vials were then thermostated at  $30 \pm 0.1^\circ\text{C}$ . After the equilibrium  
27 was established (30 min), 1 mL of vapor from the above solution was drawn out from  
28 the vial using a gas-tight syringe and injected directly in the chromatographic column  
29 via a transfer line. This sample was then analyzed by gas chromatography (Perkin Elmer  
30 Autosystem XL) equipped with a flame-ionization detector using a DB624 column. The  
31 GC settings were programmed as follows: detector temperature,  $280^\circ\text{C}$ ; column  
32 temperature,  $120^\circ\text{C}$  during 8 min.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54  
55 Full equilibrium of the chemical solute between liquid and gas phases is required. For  
56 the chemicals tested, the equilibrium time was 30 min. Linear response of the GC  
57 detector to the chemical concentration range tested is another requirement for the  
58  
59  
60



successful application of the method. For all compounds, we verify that the GC detector responds linearly. The correlation coefficients are all better than 0.998.

#### c) Formation constants

The host/guest system (H/VOC) is characterised by a titration method. Different concentrations of host are used while the initial concentration of the VOC is constant. Assuming that 1:1 complexes are formed, the calculation of formation constant  $K_f$  is developed as follows:



$$K_f = [H/VOC]/[H][VOC] = [H/VOC]/([H]_T - [H/VOC]) * ([VOC]_T - [H/VOC]) \quad (2)$$

$$[H/VOC] = -1/2 \sqrt{[(1/K_f + [H]_T + [VOC]_T)^2 - 4[H]_T [VOC]_T]} + 1/2(1/K_f + [H]_T + [VOC]_T) \quad (3)$$

where  $K_f$  and T stand for formation constant and total, respectively. For a given value of  $K_f$ ,  $[H/VOC]$  is known. An algorithmic treatment developed in our laboratory is used to calculate the formation constant  $K_f$  from the experimental data [14]. Only one concentration of VOC in water is needed [7].

#### d) Molecular modeling

All the calculations were performed with CaChe [28]. Each symmetric host structure (CBPQT<sup>4+</sup> and  $\beta$ -CD) was first subjected to an optimization with PM3 hamiltonian.

1  
2  
3 Partial charges for both hosts and guest were then obtained from electrostatic potentials  
4 with a single point calculation (B88-LYP energy functional with the DZVP basis sets).  
5  
6 The docking of iodobenzene into each host was then realised by means of dummy  
7 atoms, with the MM3 force field. The distance between host and guest and the rotation  
8 of the guest inside the host were the two intermolecular variables investigated with a  
9 regular increment (0.1Å for distance, 5° for rotation), the others being implicitly  
10 submitted to Newton-Raphson optimisation. Complexation energy was defined as the  
11 difference between energy of the optimal docked structure and energy of the individual  
12 components.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

### 28 **Conclusion**

29  
30  
31 This study explores for the first time the complexation ability of CBPQT<sup>4+</sup> towards  
32 aromatic VOCs. The theoretical and experimental results underlined a greater capacity  
33 of CBPQT<sup>4+</sup> to complex VOCs if compared with CDs due to a better encapsulation of  
34 the phenyl ring by CBPQT<sup>4+</sup>. Nevertheless CDs, and especially modified CDs, present a  
35 greater solubility than CBPQT<sup>4+</sup>. Further work in our laboratory will investigate the  
36 capacity of CBPQT<sup>4+</sup> as a molecular sensor for VOCs by cyclic voltamperometry.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- [1] Saha, S.; Johansson, E.; Flood, A.H.; Tseng, H.-R.; Zink, J.I.; Stoddart, J.F. *Chemistry - A European Journal* **2005**, *11*(23), 6846.
- [2] Cooke, G.; Garety, J.F.; Hewage, S.G.; Rabani, G.; Rotello, V.M.; Woisel, P. *Chem. Comm.* **2006**, *39*, 4119.
- [3] Fourmentin, S.; Surpateanu, G.G.; Blach, P.; Landy, D.; Decock, P.; Surpateanu, G. *J. Incl. Phenom. Macro. Chem.* **2006**, *55*(3-4), 263.
- [4] Surpateanu G. G., Becuwe M., Lungu N. C., Dron P. I., Fourmentin S., Landy D.; Surpateanu G. *J. Photochem. Photobiol. A: Chemistry* **2007**, *185*(2-3), 312.
- [5] Castro, R.; Nixon, K. R.; Evanseck, J.D.; Kaifer, A.E., *J. Org. Chem.* **1996**, *61*(21) 7298
- [6] Odell, B.; Reddington, M.V.; Slawin, A.M.Z.; Spencer R., Stoddart, J.F.; Williams, D.J., *Angew. Int. Ed. Engl.* **1988**, *27*, 1547.
- [7] Fourmentin, S.; Outirite, M.; Blach, P.; Landy, D.; Ponchel, A.; Monflier, E.; Surpateanu, G. *J. Haz. Mat.* **2007**, *141*, 92.
- [8] Staley, S.A.; Smith, B.D., *Tet. Lett.* **1996**, *37*(3), 283.
- [9] Bria, M.; Cooke, G.; Cooper A.; Garety, J.F.; Hewage, S.G.; Nutley, M.; Rabani, G.; Woisel, P., *Tet. Lett.* **2007**, *48*, 301.
- [10] Guo, Q.-X.; Luo, S.-H.; Liu, Y.-C., *J. Incl. Phenom. Macro. Chem.* **1998**, *30*, 173.
- [11] Sanemasa, I.; Akamine, Y. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 2059.
- [12] Takuma, T.; Deguchi, T.; Sanemasa, I. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 1246.
- [13] Szaniszlo N., Fenyvesi E., Balla J. *J. Incl. Phenom. Macro. Chem.* **2005**, *53*, 241.
- [14] Landy, D.; Fourmentin, S.; Salome, M.; Surpateanu, G. *J. Incl. Phenom. Macro. Chem.* **2000**, *38*, 187.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- [15] Saito, Y.; Yoshihara, K.; Tanamura, I.; Ueda, H.; Sato, T. *Chem. Pharm. Bull.* **1997**, *45*, 1711.
- [16] Saito, Y.; Tanamura, I.; Ueda, H.; Sato, T. *Chem. Pharm. Bull.* **1998**, *46*, 1777.
- [17] Saito, Y.; Tanamura, I.; Ueda, H.; Sato, T. *Int. J. Cosmet. Sci.* **1999**, *21*, 189.
- [18] Saito, Y.; Misawa, K.; Hashizaki, K.; Taguchi, H.; Ogawa, N.; Ueda, H. *Chem. Pharm. Bull.* **2004**, *52*, 259.
- [19] Tanamura, I.; Saito, Y.; Ueda, H.; Sato, T. *Chem. Pharm. Bull.* **1998**, *46*, 540.
- [20] Wu, J.-S.; Zheng, J.-Z.; Toda, K.; Sanamasa, I. *Anal. Sci.* **1999**, *15*, 701.
- [21] Misawa, K.; Saito, Y.; Hashizaki, K.; Taguchi, H.; Ogawa, N.; Ueda, H. *J. Incl. Phenom. Macro. Chem.* **2005**, *53*, 237.
- [22] Lantz, A.W.; Wetterer, S.M.; Armstrong, D.W. *Anal. Bioanal. Chem.* **2005**, *383*, 160.
- [23] Zhang, K.-C.; Liu, L.; Mu, T.-W.; Guo, Q.-X. *Chem. Phys. Lett.* **2001**, *333*, 195.
- [24] Zhang, K.-C.; Liu, L.; Mu, T.-W.; Guo, Q.-X. *J. Incl. Phenom. Macro. Chem.* **2001**, *40*, 189.
- [25] Kolb, B. *J. Chromatogr. A* **1999**, *842*, 163.
- [26] Snow, N.H.; Slack, G.C. *Trends Anal. Chem.* **2002**, *21*, 608.
- [27] Kolb, B.; Ettre, L.S. *Static Headspace-Gas Chromatography Theory and Practice*, Wiley-VCH, New York, USA, 1997.
- [28] CAChe, Work System, 6.01, **2003**.

1  
2  
3 Figure Captions  
4  
5  
6  
7

8 Figure 1: Chemical structure of (1) cyclobis(paraquat-*p*-phenylene) tetrachloride salt  
9 and (2)  $\alpha$ -CD: if  $n=1$  and  $R=H$ ;  $\beta$ -CD: if  $n=2$  and  $R = H$ : RAMEB; if  $n=2$  and  $R = H$  or  
10  $CH_3$  (1.8 OH groups modified per glucopyranose unit).  
11  
12  
13  
14

15  
16  
17 Figure 2: Representation of the variation of the chromatogram of toluene 10 ppm (a)  
18 with various concentrations of  $CBPQT^{4+}$ : 0.03mM (b), 0.1mM (c), 0.2mM (d) and  
19 0.3mM (e).  
20  
21  
22  
23  
24

25  
26  
27 Figure 3: Representation of the experimental point ( $\blacklozenge$ ) obtained for iodobenzene and  
28  $CBPQT^{4+}$  compare with theoretical titration curve (- -) for a 1:1 complex ( $K_f=9459 M^{-1}$ )  
29 with the algorithmic treatment.  
30  
31  
32  
33


34  
35  
36 Figure 4: Variation of  $\log K_f$  in function of molar refractivity (MR) of the substituent for  
37  $\beta$ -CD ( $\blacktriangle$ ) and  $CBPQT^{4+}$  ( $\blacklozenge$ ).  
38  
39  
40  
41

42  
43 Figure 5: Optimized MM3 structures for inclusion of iodobenzene in  $CBPQT^{4+}$  (1), and  
44 in  $\beta$ -CD (2).  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


1  
2  
3 Table Captions  
4  
5  
6  
7

8 Table 1: Formation constants ( $M^{-1}$ ) obtained at 30°C for the different VOCs with  
9  
10 CBPQT<sup>4+</sup>,  $\alpha$ -CD,  $\beta$ -CD and RAMEB  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only




**Figure 1:** Chemical structure of (1) cyclobis(paraquat-*p*-phenylene) tetrachloride salt and (2)  $\alpha$ -CD: if  $n=1$  and  $R=H$ ;  $\beta$ -CD: if  $n=2$  and  $R = H$ ; RAMEB; if  $n=2$  and  $R = H$  or  $CH_3$  (1.8 OH groups modified per glucopyranose unit).




**Figure 2:** Representation of the variation of the chromatogram of toluene 10 ppm (a) with various concentrations of CBPQT<sup>4+</sup>: 0.03mM (b), 0.1mM (c), 0.2mM (d) and 0.3mM (e).






**Figure 3:** Representation of the experimental point (◆) obtained for iodobenzene and CBPQT<sup>4+</sup> compare with theoretical titration curve (- -) for a 1:1 complex ( $K_f=9459 \text{ M}^{-1}$ ) with the algorithmic treatment.



**Figure 4:** Variation of logK<sub>f</sub> in function of molar refractivity (MR) of the substituent for β-CD (▲) and CBPQT<sup>4+</sup> (◆).




**Figure 5:** Optimized MM3 structures for inclusion of iodobenzene in CBPQT<sup>4+</sup> (1), and in  $\beta$ -CD (2).

	CBPQT <sup>4+</sup>	$\alpha$ -CD	$\beta$ -CD	RAMEB
Toluene	3872	38 (29 <sup>a</sup> )	142 (140 <sup>b</sup> - 172 <sup>a</sup> )	171 (144 <sup>a</sup> )
Fluorobenzene	1083	31 (34 <sup>c</sup> )	64 (70 <sup>c</sup> )	90
Chlorobenzene	2300	128 (100 <sup>c</sup> )	174 (160 <sup>c</sup> )	157
Bromobenzene	4312	333 (51 <sup>c</sup> )	237 (310 <sup>c</sup> )	244
Iodobenzene	9459	917 (1100 <sup>c</sup> )	622 (800 <sup>c</sup> )	785

In bracket literature data; See references: <sup>a</sup> [13], <sup>b</sup> [11], <sup>c</sup> [12]

**Table 1:** Formation constants ( $M^{-1}$ ) obtained at 30°C for the different VOCs with CBPQT<sup>4+</sup>,  $\alpha$ -CD,  $\beta$ -CD and RAMEB



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

