



HAL
open science

Protonation of Tetrapropoxy-4-tert-butylcalix[4]arene: NMR Study of Interaction and Probable Structures of the Product

Jaroslav Kriz, Jiri Dybal, Emanuel Makrlik, Jan Budka, Petr Vanura

► **To cite this version:**

Jaroslav Kriz, Jiri Dybal, Emanuel Makrlik, Jan Budka, Petr Vanura. Protonation of Tetrapropoxy-4-tert-butylcalix[4]arene: NMR Study of Interaction and Probable Structures of the Product. *Supramolecular Chemistry*, 2008, 20 (05), pp.487-494. 10.1080/10610270701422065 . hal-00513513


HAL Id: hal-00513513

<https://hal.science/hal-00513513>

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Protonation of Tetrapropoxy-4-tert-butylcalix[4]arene: NMR Study of Interaction and Probable Structures of the Product

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2007-0055.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	19-Apr-2007
Complete List of Authors:	Kriz, Jaroslav; Institute of Macromolecular Chemistry, Structure Analysis Dybal, Jiri; Institute of Macromolecular Chemistry, Structure Analysis Makrlik, Emanuel; University of West Bohemia Budka, Jan; Prague Institute of Chemical Technology Vanura, Petr; Prague Institute of Chemical Technology
Keywords:	Calixarene complex, calixarene protonation, NMR, DFT
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p>	
Scheme-1.CDX	



Protonation of Tetrapropoxy-4-*tert*-butylcalix[4]arene: NMR Study of Interaction and Probable Structures of the Product

JAROSLAV KRÍŽ^a, JIŘÍ DYBAL^a, EMANUEL MAKRLÍK^b, JAN BUDKA^c and PETR VAŇURA^c

^a *Institute of Macromolecular Chemistry, Academy of Sciences of the Czech Republic, Heyrovského sq. 2, 162 06 Prague, Czech Republic. E-mail: kriz@imc.cas.cz*

^b *Faculty of Applied Sciences, University of West Bohemia, Husova 11, 306 14 Pilsen, Czech Republic.*

^c *Prague Institute of Chemical Technology, Technická 5, 166 28 Prague, Czech Republic.*

Using ¹H and ¹³C NMR spectroscopy, the interaction of tetrapropoxy-*p-tert*-butylcalix[4]arene (1) with H₃O⁺ ions produced by hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC) and traces of water was studied in nitrobenzene-*d*₅. It was shown that 1 readily forms an equimolecular complex with H₃O⁺. The equilibrium constant *K* of its formation is 2.6 at 296 K. Exchange between bound and free 1 is fast even under mild excess of HDCC, the correlation time τ_{ex} being about 0.13 ms. NMR shows that H₃O⁺ is bound to the aryl-oxygen atoms and this binding forces the calixarene cup to adopt a more open and symmetrical conformation. This conclusion is in a full accord with high precision quantum DFT calculations which find one structure of the complex corresponding to a global energy minimum, in which the H₃O⁺ ion is bound to three of the oxygen atoms by strong hydrogen bonds and to the remaining oxygen by two weaker hydrogen bonds. The calixarene part is forced into a C₄ symmetrical opened form.

When stored for weeks, the complex gradually transforms into other forms, most probably its hydrates, according to spectral evidence and DFT calculations.

Keywords: calixarene complex; calixarene protonation; NMR; DFT

INTRODUCTION

Calixarene-based molecules have received intense attention in the recent years [1,2]. Many studies have focused on the binding ability of calixarene derivatives with carbonyl or analogous groups at their lower rims toward metal ions [3-12]. In our recent studies [13-15], we have shown that calixarenes carrying coordinating groups, such as dimethylthioamide, ester or keto groups on their lower rim, bind a hydroxonium ion H_3O^+ in equimolecular and very stable complexes. NMR spectra and high-level quantum calculations revealed that H_3O^+ binds partly to the attached coordinating groups but also to the calixarene aryl-oxygen atoms. Quantum calculations even indicated that binding almost exclusively to the Ar-O- atoms could be even preferred in some cases. It is thus interesting to examine H_3O^+ binding to a calixarene containing only Ar-O-R groups where R is an alkyl such as *n*-propyl in our case.

Like in our previous studies [13-15, 17, 18], hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC)[16] was used as a reliable source of protons, which were converted to hydroxonium ions H_3O^+ by a 3.1 mol/mol excess of water. Combining NMR spectral evidence with DFT quantum mechanical calculations, we suggest the most probable structure of the protonated calixarene.

RESULTS AND DISCUSSION

NMR measurements. Our experiments with **1** (see Scheme 1) were carried out in nitrobenzene- d_5 as one of the sufficiently polar solvents (needed for full ionization of HDCC), which do not form inclusion complexes with calixarenes. After mixing **1** with increasing amounts of HDCC, its ^1H NMR signals gradually shift to a various degree as illustrated in Figure 1. The signal assignment corresponds to Scheme 1, where the protons are numbered in the same way as the carbons they are attached to. The signal shifts are analogous to those

observed with other calixarene derivatives [13-15] as a sign of conformation change due to binding of H_3O^+ ion at the lower rim.

As the signals shift immediately and without splitting, the protonation of **1** must be very fast. However, the persistent shift at larger excess of HDCC (cf. Figure 1) indicates that the product formation is not quantitative in the 1:1 mixture. We are thus dealing with a dynamic equilibrium, which is only moderately favorable to the complex.

Figure 1: 300.13 MHz NMR spectra of calixarene **1** (A) and its 1:2 (B) and 1:4 mol/mol (C) mixtures with HDCC (nitrobenzene- d_5 , 296 K). Signal assignment in A (cf. Scheme 1), extrapolated maximal shifts $\Delta\delta_{\text{max}}$ (in ppm) in C, correlation times of exchange τ_{ex} (in ms) in B.

Scheme 1

The equilibrium should follow the usual relation

$$K = \frac{[\mathbf{1} \cdot \text{H}^+][\text{A}^-]}{[\mathbf{1}][\text{A}^-\text{H}^+]}, \quad (1)$$

where A^-H^+ and A^- stand for HDCC or any other suitable proton donor and its anion. Under fast exchange between free and coordinated **1**, the actual chemical shift of any signal is a weighted average of the shifts corresponding to the exchanging sites. Thus the actual relative shift is $\Delta\delta = \alpha\Delta\delta_{\text{max}}$, where $\alpha = [\mathbf{1} \cdot \text{H}^+]/[\mathbf{1}]_0$ and $\Delta\delta_{\text{max}}$, obtained by extrapolation, is the maximum relative shift corresponding to the pure complex. Using Eq.1, $\Delta\delta$ can be expressed as

$$\Delta\delta = \Delta\delta_{\text{max}} \frac{K(1 + \beta) - \sqrt{K^2(1 + \beta)^2 - 4(K - 1)K\beta}}{2(K - 1)}, \quad (2)$$

where $\beta = [A^-H^+]_0/[1]_0$. Thus K can be obtained by fitting Eq.2 to the experimental values of $\Delta\delta$. Figure 2 shows $\Delta\delta$ of individual signals in the proton spectra of **1**/HDCC mixtures and the fitting curves.

Figure 2: Relative shifts $\Delta\delta$ of signals in ^1H NMR spectrum of **1** and its mixtures with HDCC in nitrobenzene- d_5 at 296 K and their fitting with Equation 2.

The value of K to which all the actual relative shifts fit is 2.61 ± 0.03 . This is a much lower value than we observed in protonation of calixarenes bearing additional coordinating groups [13-15].

The apparent swiftness of exchange between free and bound **1** gives us the opportunity to examine its dynamics quantitatively. One of the possible methods is the measurement of transverse relaxation rate using the Carr-Purcell-Meiboom-Gill sequence (CPMG) [19] with a varied delay t_p between the π pulses. As shown by Luz and Meiboom [20], the actually measured relaxation rate $R_2(t_p)$ can be expressed as

$$R_2(t_p) = R_2^0 + p_1 p_2 \delta\omega^2 \tau_{ex} [1 - (\tau_{ex}/t_p) \tanh(t_p/\tau_{ex})] \quad (3)$$

where R_2^0 is the part of relaxation rate independent of exchange, p_1 and p_2 are the probabilities of finding the nucleus at the sites 1 and 2, respectively, $\delta\omega$ is the relative chemical shift between the two sites in rads and τ_{ex} is the correlation time of exchange. For short τ_{ex} , Equation 3 can be approximated (slightly modifying the relations in Ref. 21) by

$$R_2(t_p) = \zeta - \xi \tau_{ex}^2 / t_p \quad (4)$$

with

$$\zeta = R_2^0 + \xi \tau_{ex} \quad (4a)$$

and

$$\xi = 4\pi^2 \alpha (1 - \alpha) (\Delta\delta_{\max})^2 \quad (4b)$$

1
2
3 where $\Delta\delta_{\max}$ is the relative shift (in Hz) of the given signal between free **1** and its complex.
4
5
6 Thus the dependence of $R_2(t_p)$ on $1/t_p$ is linear with the slope $\xi\tau_{ex}^{-2}$. As $\Delta\delta_{\max}$ is obtained by
7
8
9 extrapolation from the spectra and α can be calculated from actual $\Delta\delta$ (see above), obtaining
10
11 τ_{ex} from the slope is straightforward. Figure 3 shows dependences for three most mobile
12
13 signals in the proton spectrum of the **1**/HDCC mixture 1:2 mol/mol, namely 4+8, 10 and 9e.
14
15 The values of τ_{ex} obtained from the individual slopes are 1.33×10^{-4} , 1.36×10^{-4} , and 1.24×10^{-4} s,
16
17
18 respectively. These values are sufficiently close to each other to have a physical meaning of
19
20 the correlation time of exchange between free and bound forms of **1**. Evidently, the exchange
21
22 is quite fast, its rate constant being about 7.6×10^3 s⁻¹.
23
24
25
26
27
28
29

30
31 **Figure 3:** Dependence of the measured net increase in the transverse relaxation rate of the
32 indicated most mobile signals in the 300.13 MHz ¹H NMR spectrum of the
33 mixture of **1** with HDCC 1:2 (mol/mol) in nitrobenzene-*d*₅ at 296 K
34
35

36
37 Returning now to the shape of ¹H NMR spectra, the 0.323 ppm mutual approach of the
38
39 signals 9a and 9e (axial and equatorial CH₂ protons) relative to free **1** is a clear sign that the
40
41 aromatic part of **1** adopts a more open and symmetric conformation in the complex [12]. This
42
43 conclusion is corroborated by the significant deshielding of aromatic protons 4, 8. Such a
44
45 change can be forced only by a H₃O⁺ binding at the lower rim of **1**, i.e. to the aryl-oxygen
46
47 atoms. In agreement with this, protons 10 (O-CH₂-) are significantly deshielded, too.
48
49 Assuming that H₃O⁺ is bound by three strong hydrogen bonds [13-15], one would expect that
50
51 one of the four O-CH₂- groups should be perceptibly different from the other three. The shape
52
53 of the proton (and also carbon, see below) spectrum indicates a C₄ symmetry, however. There
54
55 are two possible explanations of this behavior: either the H₃O⁺ ion adopts multiple weaker
56
57 bonds to the fourth oxygen atom and is thus placed near to the center of the lower rim or it
58
59
60

1
2
3 rotates very fast within the molecule averaging thus its structure in the NMR time window.
4
5 Deciding between these two alternatives is beyond the abilities of NMR.
6
7

8 Most of the just described features can also be read out from the ^{13}C NMR spectra.
9
10 Figure 4 reproduces DEPT45 spectra (showing only carbons with attached protons) of **1** and
11 its 1:4 mol/mol mixture with HDCC, the latter having been measured 6 to 18 hours after
12 mixing. Large downfield shifts of signals 4,8 and 10 leave no doubt that a complex was
13 formed between **1** and H_3O^+ , with the ion probably bound to the aryl-oxygen atoms. In
14 analogy to proton spectra, the signals are single (except for small accompanying signals)
15 suggesting thus again either C_4 symmetry of the complex or fast averaging of the structure by
16 internal motion.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Figure 4:** 75.5 MHz ^{13}C NMR DEPT45 spectra of 5 mmol/L calixarene **1** (A) and its mixture
33 with 4.0 mol/mol excess of HDCC (B) 6 – 18 h after mixing (nitrobenzene- d_5 , 296
34 K). Assignment of signals in A (cf. Scheme 1), relative shifts (in ppm) in B.
35
36

37 However, measurement of long-range DEPT spectra designed to show signals of quaternary
38 carbons shown in Figure 5 (which was more time-consuming and reflects thus a later stage of
39 the development of the system) revealed splitting of carbon signals 5, 6 and 7 suggesting
40 either lower symmetry of the complex or coexistence of two different structures. In addition
41 to it, the smaller companions of each of the main signals, which already appeared in the
42 normal DEPT45 spectra, somewhat increased in relative intensity. Evidently, a new variant of
43 the complex is evolving very slowly from its original form and, as one can expect, the
44 exchange between both forms is quite slow, too. Such multiple complex forms were already
45 observed in the case of some calixarenes with attached ester and keto groups [14, 15] where
46 the formation of the H_3O^+ complex was very slow. In contrast to it, the formation of the
47 present original complex is very fast although its equilibrium is less favorable.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10 **Figure 5:** Parts of combined 75.5 MHz ^{13}C NMR DEPT45 and LR-DEPT45 spectra of 5
11 mmol/L calixarene **1** (A) and its mixture with 4.0 mol/mol excess of HDCC (B)
12 10 – 12 days after mixing (nitrobenzene- d_5 , 296 K)
13
14


15
16
17
18 **Figure 6:** Parts of 300.13 MHz ^1H NMR (above) and 75.5 MHz ^{13}C NMR DEPT45 (below)
19 spectra of the mixture of 5 mmol/L calixarene **1** with 4.0 mol/mol excess of
20 HDCC measured 3 weeks after mixing (nitrobenzene- d_5 , 296 K).
21
22

23
24 Inspection of the spectra of **1** – HDCC 1:4 mixture (see Figure 6, where the assignment of
25 signals to individual forms was made using a combination of ^1H NOESY and ^1H - ^{13}C HSQC
26 spectra) after 3 weeks of storage reveals two different new forms of the complex, B and C,
27 produced from the original form A. The same new forms can be observed in a lower relative
28 population in systems with a lower excess of HDCC, even in the 1:1 mixture. Therefore, it is
29 not probable that the new forms are complexes with a higher number of H_3O^+ ions. However,
30 a 0.23 ppm downfield relative shift of the broad H_2O signal (not shown) in the long-stored 1:4
31 mixture indicates that water takes part in the new forms. As it can be seen in Figure 6, the
32 more prominent form B among the new products no longer exhibits C_4 symmetry. Quite to the
33 contrary, there is magnetic nonequivalence between at least two doubles of opposite aromatic
34 rings (in the proton spectrum of B, even all four *tert*-butyl groups are nonequivalent).
35 Experience with NMR spectra of calixarenes teaches us that such nonequivalence should not
36 be overvalued: due to their strong long-range shielding effects of the aromatic rings, even
37 subtle changes in their relative orientation can lead to remarkable shifts of the signals of
38 adjacent nuclei. Nevertheless, the observed nonequivalence cannot be ignored for two
39 reasons: (i) it reflects lowering of the molecular symmetry from apparent C_4 to C_2 or even
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 lower and (ii) it shows that the residence time in the exchange between nonequivalent
4 analogous nuclei must be larger than 0.05 s, i.e. the internal motion in the molecule B is at
5
6
7
8 least two orders of magnitude slower than that in the form A. Therefore, the most plausible
9
10 conclusion appears to be that the form B (and probably also C) is a hydrate of the complex A
11
12 with a water molecule placed inside the calixarene cavity. The molecule has to sit near
13
14 enough to the lower rim in order to interact with the bound H_3O^+ ion, which necessarily
15
16 means that it interacts with at least two opposite aromatic rings (probably by a weak hydrogen
17
18 bond) lowering thus the symmetry of the complex and slowing down its internal motion.
19
20
21

22 **Quantum chemical calculations.** Geometry optimizations of calixarene and its complex with
23
24 H_3O^+ were performed by the quantum chemical calculations at the density functional level of
25
26 theory (DFT, B3LYP functional) using the Gaussian 03 suite of programs [22]. The 6-31G(d)
27
28 basis set was used and the optimization was unconstrained. The optimized structure of free
29
30 calixarene is shown in Figure 7A. The most stable conformation forms a pinched cone
31
32 structure with a C_2 symmetry.
33
34
35

36 The structure obtained by optimization of the calixarene - H_3O^+ complex is shown in
37
38 Figure 7B, together with the lengths of hydrogen bonds (in Å). Compared to free molecule the
39
40 calixarene part of the complex is distorted so that its structure is very close to the C_4
41
42 symmetry. The hydroxonium ion is placed in the cage formed by calixarene and is bound by
43
44 strong hydrogen bonds to oxygen atoms of three C-O-C groups and by two somewhat weaker
45
46 hydrogen bonds to the last C-O-C group. The stabilization energy of the complex calculated
47
48 as the difference between the energies of the optimized complex and optimized free
49
50 calixarene molecule and hydroxonium ion is -394.72 kJ/mol. Stabilization energy of the
51
52 complex can be expressed also with respect to the aqueous hydroxonium ion $[\text{H}_3\text{O}^+(\text{H}_2\text{O})_3]$
53
54 and free calixarene (**1**) according to the formula
55
56
57
58
59



1
2
3 Then it is -8.20 kJ/mol. This value, corrected further for entropy change, probably
4
5 corresponds to the value of K established above by NMR.
6
7

8 The gradual development of another form of the complex detected by NMR led us to
9
10 explore its further possible changes or interactions. As the structure described above
11
12 corresponds to the global energy minimum found by unconstrained geometry optimization, a
13
14 new form could scarcely be produced by an internal rearrangement of the structure. We thus
15
16 explored possible interactions of the complex with other components of the system.
17
18 Nitrobenzene, which is largely prevalent in the system, does not appear do form any stable
19
20 formation with the complex. However, water does. The calculated geometry of the complex
21
22 hydrate is shown in Figure 8. The stabilization energy corresponding to such hydration is
23
24 predicted to be - 19.75 kJ/mol, i.e. sufficiently high for the hydrate to be stable.
25
26
27
28

29 The interesting feature of the hydrate is that the water molecule is bound by the
30
31 already bonded H_3O^+ ion by a system of three medium-strong hydrogen bonds indicated in
32
33 the Figure. In order to accomplish that, the water molecule has to penetrate the rather rigid
34
35 and hydrophobic calixarene cavity (no other way apparently leads to a stable product), which
36
37 can be expected to lead to rather long reaction times as observed. In addition, the water
38
39 molecule also interacts with the aromatic systems by two weak hydrogen bonds (some of the
40
41
42
43
44
45

46 **Figure 7:** Optimized geometry of free **1** (A) and the **1**· H_3O^+ complex (B) calculated at the
47
48 B3LYP/6-31G(d) level (positions of hydrogen atoms in **1** are not shown)

49
50
51 O-H- C_{ar} distances being as short as 2.55 Å). In agreement with NMR spectra, these bonds
52
53 without doubt slow down internal motions in the calixarene unit.
54

55 In contrast to NMR, however, we were not able to find any alternative stable form of
56
57 the hydrate. Taking into account the fact that our calculations were done *in vacuo* (as no
58
59 sufficiently reliable correction for the influence of polar medium has been found for this level
60

1
2
3 of precision), we cannot be quite sure that other structural forms of the complex hydrate are
4
5 excluded by theory.
6
7
8
9

10
11
12 **Figure 8:** Optimized geometry of the hydrated $\mathbf{1} \cdot \text{H}_3\text{O}^+$ complex calculated at the
13 B3LYP/6-31G(d) level (positions of hydrogen atoms in $\mathbf{1}$ are not shown).
14
15

16
17 **Conclusions.** According to the evidence of ^1H and ^{13}C NMR spectra consistent with the
18 predictions of high-precision DFT calculations, calixarene $\mathbf{1}$ binds the H_3O^+ ion into an
19 equimolecular complex. The formation of the main form of the complex, similarly as the
20 exchange between free and bound $\mathbf{1}$, is very fast but the equilibrium is only moderately
21 favorable to the complex. In the main form, H_3O^+ is probably bound by three strong hydrogen
22 bonds to three aryl-oxygen atoms of $\mathbf{1}$ and by two weaker hydrogen bonds to the remaining C-
23 O-C group. In contrast to free $\mathbf{1}$, the complex has almost C_4 symmetry. Under prolonged
24 storage of the complex-containing mixtures, small amounts of another form of the complex
25 appear. It is evidently formed from the original complex and probably is the product of its
26 hydration.
27
28
29
30
31
32
33
34
35
36
37
38
39

40
41 When compared with the calixarenes with strongly coordinating groups in addition to
42 aryl-oxygen atoms at their lower rim [13-15], the present calixarene type clearly is a markedly
43 less efficient complexing agent. From this one can conclude that the C-O-C groups, although
44 they clearly participate in the complex H_3O^+ binding in the former types [13-15], are as such
45 less effective than in cooperation with e.g. dimethylamide, dimethylthioamide, ester and keto-
46 groups.
47
48
49
50
51
52
53

54 55 **EXPERIMENTAL**

56
57 **Materials and samples.** Nitrobenzene- d_5 was purchased by Aldrich. Tetrapropoxy-*p*-*tert*-
58 butylcalix[4]arene ($\mathbf{1}$) was prepared by using the procedure published elsewhere [23].
59
60

1
2
3 Preparation of hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC) was described earlier [16].
4
5 For NMR samples, 0.5×10^{-5} mol of **1** was dissolved in a mixture of appropriate amounts of
6
7 nitrobenzene-*d*₅ and 0.002 mol/L solution of HDCC in the same solvent. The mixing and all
8
9 subsequent measurements were done at 296 K.
10
11

12 **NMR spectra.** ¹H and ¹³C NMR spectra were measured in a quadrature detection mode at
13
14 300.13 and 75.45 MHz, respectively, with an upgraded Bruker Avance DPX300 spectrometer.
15
16 32 and 64 kpoints were measured for ¹H and ¹³C NMR, respectively. ¹H NMR spectra were
17
18 measured using a self-devised diffusion-filtering stimulated echo pulse sequence $-d_1-\pi/2_\phi-$
19
20 $GR-\pi/2_\sigma-d_2-\pi/2_y-GR-REC_x$ in order to remove obscuring broad signals of water and
21
22 nitrobenzene; *d*₁ and *d*₂ were 10 s and 60 ms, respectively, the *z*-gradient GR was 50 G/cm
23
24 and phase cycling was $\phi = x, y, -x, -y, \sigma = \phi + \pi/2$. 128 scans were collected for each
25
26 spectrum. ¹³C NMR measurements were performed using either the ¹H-¹³C DEPT45 sequence
27
28 (collecting 8000 scans) or its combination with its long-range variant optimized for the
29
30 coupling constant 6 Hz (collecting at least 20000 scans); exponential weighting (*lb* = 1 Hz)
31
32 was used before Fourier transform. In ¹H – ¹³C 2D HSQC and HMBC spectra used for signal
33
34 assignment, 1028 points in F2 and 256 increments in F1 dimensions were measured using a *z*-
35
36 gradient inverse-detection probe.
37
38
39
40
41
42

43 **Quantum mechanical calculations.** *Ab initio* molecular orbital calculations were performed
44
45 using the GAUSSIAN 03 suite of programs [16]. Molecular geometry was fully optimized at
46
47 the B3LYP level of density functional theory (DFT) with the 6-31G(d) basis set. The
48
49 optimization was unrestrained. Several local configurations near the achieved energy
50
51 minimum were examined. As the renewed optimizations converged to the same molecular
52
53 geometry, we believe the achieved energy minimum to be the global one.
54
55
56
57
58
59
60

Acknowledgements. This work was supported by the Academy of Sciences of the Czech Republic, Project T400500402, and the Czech Ministry of Education, Youth and Sports, Projects MSM 4977751303 and MSM 6046137307.

References

1. Gutsche, C. D. *Calixarenes Revisited*. The Royal Society of Chemistry, Cambridge, 1998.
2. Böhmer, V. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 713.
3. Arduini, A.; Pochini, A.; Reverberi, S.; Ungaro, R. *Tetrahedron* **1986**, *42*, 2089.
4. Arduini, A.; Ghidini, E.; Pochini, A.; Ungaro, R.; Andreetti, G.D.; Calestani, G.; Ugozzoli, F. *J. Inclusion Phenom.* **1988**, *6*, 119.
5. Arnaud-Neu, F.; Collins, E.M.; Deasy, M.; Ferguson, G.; Harris, S.J.; Kaitner, B.; Lough, A.J.; McKervey, M.A.; Marques, E.; Ruhl, B.L.; Schwing-Weill, M.J.; Seward, E.M. *J. Am. Chem. Soc.* **1989**, *111*, 8681.
6. Arnaud-Neu, F.; Barrett, G.; Harris, S.J.; Owens, M.; McKervey, M.A.; Schwing-Weill, M.J.; Schwinté, P. *Inorg. Chem.* **1993**, *32*, 2644.
7. Ohto, K.; Murakami, E.; Shinohara, T.; Shiratsuchi, K.; Inoue, K.; Iwasaki, M. *Anal. Chim. Acta* **1997**, *341*, 275.
8. Ye, Z.; He, W.; Shi, X.; Zhu, L. *J. Coord. Chem.* **2001**, *54*, 105.
9. Danil de Namor, A.F.; Chahine, S.; Kowalska, D.; Castellano, E.E.; Piro, O.E. *J. Am. Chem. Soc.* **2002**, *124*, 12824.
10. Marcos, P. M.; Ascenso, J.R.; Segurado, M.A.P.; Pereira, J.L.C. *J. Inclusion Phenom.* **2002**, *42*, 281.
11. Marcos, P.M.; Félix, S.; Ascenso, J.R.; Segurado, M.A.P.; Pereira, J.L.C.; Khazaeli-Parsa, P.; Hubscher-Bruder, V.; Arnaud-Neu, F. *New J.Chem.* **2004**, *28*, 748.
12. Gutsche, C. D. *Aldrichimica Acta* **1995**, *28*, 3.
13. Kříž, J.; Dybal, J.; Makrlík, E.; Vaňura, P. *Supramol. Chem.* in print.
14. Kříž, J.; Dybal, J.; Makrlík, E.; Vaňura, P. *Supramol. Chem.* in print.
15. Kříž, J.; Dybal, J.; Makrlík, E.; Budka, J.; Vaňura, P. *Monatsh. Chem.* in print.
16. Makrlík, E.; Vaňura, P. *Talanta* **1985**, *32*, 423.
17. Kříž, J.; Makrlík, E.; Vaňura, P. *Biopolymers* **2006**, *81*, 104.
18. Kříž, J.; Dybal, J.; Makrlík, E. *Biopolymers* **2006**, *82*, 536.
19. Meiboom, S.; Gill, D. *Rev. Sci. Instrum.* **1958**, *29*, 688.
20. Luz, L.; Meiboom, S. *J. Chem. Phys.* **1963**, *39*, 366.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. Canet, D.; Robert, J.B. in *Dynamics of Solutions and Fluid Mixtures by NMR* (J. J. Delpuech Ed.), Wiley, Chichester 1995, p. 127.
22. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, Jr., J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; and Pople, J. A. *Gaussian 03, Revision C.02*; Gaussian, Inc.: Wallingford CT, 2004.
23. Kelderman, E.; Derhaeg, L.; Heesink, G.J.T.; Verboom, W.; Engbersen, J.F.J.; van Hulst, N.F.; Persoons, A.; Reinhoudt, D.N. *Angew. Chem.* **1992**, *104*, 1107.

Protonation of Tetrapropoxy-4-*tert*-butylcalix[4]arene: NMR Study of Interaction and Probable Structures of the Product

JAROSLAV KRÍŽ^a, JIŘÍ DYBAL^a, EMANUEL MAKRLÍK^b, JAN BUDKA^c and PETR VAŇURA^c

^a *Institute of Macromolecular Chemistry, Academy of Sciences of the Czech Republic, Heyrovského sq. 2, 162 06 Prague, Czech Republic. E-mail: kriz@imc.cas.cz*

^b *Faculty of Applied Sciences, University of West Bohemia, Husova 11, 306 14 Pilsen, Czech Republic.*

^c *Prague Institute of Chemical Technology, Technická 5, 166 28 Prague, Czech Republic.*

Using ¹H and ¹³C NMR spectroscopy, the interaction of tetrapropoxy-*p-tert*-butylcalix[4]arene (1) with H₃O⁺ ions produced by hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC) and traces of water was studied in nitrobenzene-*d*₅. It was shown that 1 readily forms an equimolecular complex with H₃O⁺. The equilibrium constant *K* of its formation is 2.6 at 296 K. Exchange between bound and free 1 is fast even under mild excess of HDCC, the correlation time τ_{ex} being about 0.13 ms. NMR shows that H₃O⁺ is bound to the aryl-oxygen atoms and this binding forces the calixarene cup to adopt a more open and symmetrical conformation. This conclusion is in a full accord with high precision quantum DFT calculations which find one structure of the complex corresponding to a global energy minimum, in which the H₃O⁺ ion is bound to three of the oxygen atoms by strong hydrogen bonds and to the remaining oxygen by two weaker hydrogen bonds. The calixarene part is forced into a C₄ symmetrical opened form.

When stored for weeks, the complex gradually transforms into other forms, most probably its hydrates, according to spectral evidence and DFT calculations.

Keywords: calixarene complex; calixarene protonation; NMR; DFT

INTRODUCTION

Calixarene-based molecules have received intense attention in the recent years [1,2]. Many studies have focused on the binding ability of calixarene derivatives with carbonyl or analogous groups at their lower rims toward metal ions [3-12]. In our recent studies [13-15], we have shown that calixarenes carrying coordinating groups, such as dimethylthioamide, ester or keto groups on their lower rim, bind a hydroxonium ion H_3O^+ in equimolecular and very stable complexes. NMR spectra and high-level quantum calculations revealed that H_3O^+ binds partly to the attached coordinating groups but also to the calixarene aryl-oxygen atoms. Quantum calculations even indicated that binding almost exclusively to the Ar-O- atoms could be even preferred in some cases. It is thus interesting to examine H_3O^+ binding to a calixarene containing only Ar-O-R groups where R is an alkyl such as *n*-propyl in our case.

Like in our previous studies [13-15, 17, 18], hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC)[16] was used as a reliable source of protons, which were converted to hydroxonium ions H_3O^+ by a 3.1 mol/mol excess of water. Combining NMR spectral evidence with DFT quantum mechanical calculations, we suggest the most probable structure of the protonated calixarene.

RESULTS AND DISCUSSION

NMR measurements. Our experiments with **1** (see Scheme 1) were carried out in nitrobenzene-*d*₅ as one of the sufficiently polar solvents (needed for full ionization of HDCC), which do not form inclusion complexes with calixarenes. After mixing **1** with increasing amounts of HDCC, its ¹H NMR signals gradually shift to a various degree as illustrated in Figure 1. The signal assignment corresponds to Scheme 1, where the protons are numbered in the same way as the carbons they are attached to. The signal shifts are analogous to those

observed with other calixarene derivatives [13-15] as a sign of conformation change due to binding of H_3O^+ ion at the lower rim.

As the signals shift immediately and without splitting, the protonation of **1** must be very fast. However, the persistent shift at larger excess of HDCC (cf. Figure 1) indicates that the product formation is not quantitative in the 1:1 mixture. We are thus dealing with a dynamic equilibrium, which is only moderately favorable to the complex.

Figure 1: 300.13 MHz NMR spectra of calixarene **1** (A) and its 1:2 (B) and 1:4 mol/mol (C) mixtures with HDCC (nitrobenzene- d_5 , 296 K). Signal assignment in A (cf. Scheme 1), extrapolated maximal shifts $\Delta\delta_{\text{max}}$ (in ppm) in C, correlation times of exchange τ_{ex} (in ms) in B.

Scheme 1

The equilibrium should follow the usual relation

$$K = \frac{[\mathbf{1} \cdot \text{H}^+][\text{A}^-]}{[\mathbf{1}][\text{A}^-\text{H}^+]}, \quad (1)$$

where A^-H^+ and A^- stand for HDCC or any other suitable proton donor and its anion. Under fast exchange between free and coordinated **1**, the actual chemical shift of any signal is a weighted average of the shifts corresponding to the exchanging sites. Thus the actual relative shift is $\Delta\delta = \alpha\Delta\delta_{\text{max}}$, where $\alpha = [\mathbf{1} \cdot \text{H}^+]/[\mathbf{1}]_0$ and $\Delta\delta_{\text{max}}$, obtained by extrapolation, is the maximum relative shift corresponding to the pure complex. Using Eq.1, $\Delta\delta$ can be expressed as

$$\Delta\delta = \Delta\delta_{\text{max}} \frac{K(1 + \beta) - \sqrt{K^2(1 + \beta)^2 - 4(K - 1)K\beta}}{2(K - 1)}, \quad (2)$$

where $\beta = [A^-H^+]_0/[1]_0$. Thus K can be obtained by fitting Eq.2 to the experimental values of $\Delta\delta$. Figure 2 shows $\Delta\delta$ of individual signals in the proton spectra of **1**/HDCC mixtures and the fitting curves.

Figure 2: Relative shifts $\Delta\delta$ of signals in ^1H NMR spectrum of **1** and its mixtures with HDCC in nitrobenzene- d_5 at 296 K and their fitting with Equation 2.

The value of K to which all the actual relative shifts fit is 2.61 ± 0.03 . This is a much lower value than we observed in protonation of calixarenes bearing additional coordinating groups [13-15].

The apparent swiftness of exchange between free and bound **1** gives us the opportunity to examine its dynamics quantitatively. One of the possible methods is the measurement of transverse relaxation rate using the Carr-Purcell-Meiboom-Gill sequence (CPMG) [19] with a varied delay t_p between the π pulses. As shown by Luz and Meiboom [20], the actually measured relaxation rate $R_2(t_p)$ can be expressed as

$$R_2(t_p) = R_2^0 + p_1 p_2 \delta\omega^2 \tau_{ex} [1 - (\tau_{ex}/t_p) \tanh(t_p/\tau_{ex})] \quad (3)$$

where R_2^0 is the part of relaxation rate independent of exchange, p_1 and p_2 are the probabilities of finding the nucleus at the sites 1 and 2, respectively, $\delta\omega$ is the relative chemical shift between the two sites in rads and τ_{ex} is the correlation time of exchange. For short τ_{ex} , Equation 3 can be approximated (slightly modifying the relations in Ref. 21) by

$$R_2(t_p) = \zeta - \xi \tau_{ex}^2 / t_p \quad (4)$$

with

$$\zeta = R_2^0 + \xi \tau_{ex} \quad (4a)$$

and

$$\xi = 4\pi^2 \alpha (1 - \alpha) (\Delta\delta_{\max})^2 \quad (4b)$$

1
2
3 where $\Delta\delta_{\max}$ is the relative shift (in Hz) of the given signal between free **1** and its complex.
4
5 Thus the dependence of $R_2(t_p)$ on $1/t_p$ is linear with the slope $\xi\tau_{ex}^{-2}$. As $\Delta\delta_{\max}$ is obtained by
6
7 extrapolation from the spectra and α can be calculated from actual $\Delta\delta$ (see above), obtaining
8
9 τ_{ex} from the slope is straightforward. Figure 3 shows dependences for three most mobile
10
11 signals in the proton spectrum of the **1**/HDCC mixture 1:2 mol/mol, namely 4+8, 10 and 9e.
12
13 The values of τ_{ex} obtained from the individual slopes are 1.33×10^{-4} , 1.36×10^{-4} , and 1.24×10^{-4} s,
14
15 respectively. These values are sufficiently close to each other to have a physical meaning of
16
17 the correlation time of exchange between free and bound forms of **1**. Evidently, the exchange
18
19 is quite fast, its rate constant being about 7.6×10^3 s⁻¹.
20
21
22
23
24
25
26
27
28
29
30

31 **Figure 3:** Dependence of the measured net increase in the transverse relaxation rate of the
32 indicated most mobile signals in the 300.13 MHz ¹H NMR spectrum of the
33 mixture of **1** with HDCC 1:2 (mol/mol) in nitrobenzene-*d*₅ at 296 K
34
35

36
37 Returning now to the shape of ¹H NMR spectra, the 0.323 ppm mutual approach of the
38 signals 9a and 9e (axial and equatorial CH₂ protons) relative to free **1** is a clear sign that the
39 aromatic part of **1** adopts a more open and symmetric conformation in the complex [12]. This
40 conclusion is corroborated by the significant deshielding of aromatic protons 4, 8. Such a
41 change can be forced only by a H₃O⁺ binding at the lower rim of **1**, i.e. to the aryl-oxygen
42 atoms. In agreement with this, protons 10 (O-CH₂-) are significantly deshielded, too.
43
44 Assuming that H₃O⁺ is bound by three strong hydrogen bonds [13-15], one would expect that
45 one of the four O-CH₂- groups should be perceptibly different from the other three. The shape
46 of the proton (and also carbon, see below) spectrum indicates a C₄ symmetry, however. There
47 are two possible explanations of this behavior: either the H₃O⁺ ion adopts multiple weaker
48 bonds to the fourth oxygen atom and is thus placed near to the center of the lower rim or it
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 rotates very fast within the molecule averaging thus its structure in the NMR time window.
4
5
6 Deciding between these two alternatives is beyond the abilities of NMR.
7

8 Most of the just described features can also be read out from the ^{13}C NMR spectra.
9
10 Figure 4 reproduces DEPT45 spectra (showing only carbons with attached protons) of **1** and
11 its 1:4 mol/mol mixture with HDCC, the latter having been measured 6 to 18 hours after
12 mixing. Large downfield shifts of signals 4,8 and 10 leave no doubt that a complex was
13 formed between **1** and H_3O^+ , with the ion probably bound to the aryl-oxygen atoms. In
14 analogy to proton spectra, the signals are single (except for small accompanying signals)
15 suggesting thus again either C_4 symmetry of the complex or fast averaging of the structure by
16 internal motion.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Figure 4:** 75.5 MHz ^{13}C NMR DEPT45 spectra of 5 mmol/L calixarene **1** (A) and its mixture
33 with 4.0 mol/mol excess of HDCC (B) 6 – 18 h after mixing (nitrobenzene- d_5 , 296
34 K). Assignment of signals in A (cf. Scheme 1), relative shifts (in ppm) in B.
35
36

37 However, measurement of long-range DEPT spectra designed to show signals of quaternary
38 carbons shown in Figure 5 (which was more time-consuming and reflects thus a later stage of
39 the development of the system) revealed splitting of carbon signals 5, 6 and 7 suggesting
40 either lower symmetry of the complex or coexistence of two different structures. In addition
41 to it, the smaller companions of each of the main signals, which already appeared in the
42 normal DEPT45 spectra, somewhat increased in relative intensity. Evidently, a new variant of
43 the complex is evolving very slowly from its original form and, as one can expect, the
44 exchange between both forms is quite slow, too. Such multiple complex forms were already
45 observed in the case of some calixarenes with attached ester and keto groups [14, 15] where
46 the formation of the H_3O^+ complex was very slow. In contrast to it, the formation of the
47 present original complex is very fast although its equilibrium is less favorable.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10 **Figure 5:** Parts of combined 75.5 MHz ^{13}C NMR DEPT45 and LR-DEPT45 spectra of 5
11 mmol/L calixarene **1** (A) and its mixture with 4.0 mol/mol excess of HDCC (B)
12 10 – 12 days after mixing (nitrobenzene- d_5 , 296 K)
13
14


15
16
17
18 **Figure 6:** Parts of 300.13 MHz ^1H NMR (above) and 75.5 MHz ^{13}C NMR DEPT45 (below)
19 spectra of the mixture of 5 mmol/L calixarene **1** with 4.0 mol/mol excess of
20 HDCC measured 3 weeks after mixing (nitrobenzene- d_5 , 296 K).
21
22

23
24 Inspection of the spectra of **1** – HDCC 1:4 mixture (see Figure 6, where the assignment of
25 signals to individual forms was made using a combination of ^1H NOESY and ^1H - ^{13}C HSQC
26 spectra) after 3 weeks of storage reveals two different new forms of the complex, B and C,
27 produced from the original form A. The same new forms can be observed in a lower relative
28 population in systems with a lower excess of HDCC, even in the 1:1 mixture. Therefore, it is
29 not probable that the new forms are complexes with a higher number of H_3O^+ ions. However,
30 a 0.23 ppm downfield relative shift of the broad H_2O signal (not shown) in the long-stored 1:4
31 mixture indicates that water takes part in the new forms. As it can be seen in Figure 6, the
32 more prominent form B among the new products no longer exhibits C_4 symmetry. Quite to the
33 contrary, there is magnetic nonequivalence between at least two doubles of opposite aromatic
34 rings (in the proton spectrum of B, even all four *tert*-butyl groups are nonequivalent).
35 Experience with NMR spectra of calixarenes teaches us that such nonequivalence should not
36 be overvalued: due to their strong long-range shielding effects of the aromatic rings, even
37 subtle changes in their relative orientation can lead to remarkable shifts of the signals of
38 adjacent nuclei. Nevertheless, the observed nonequivalence cannot be ignored for two
39 reasons: (i) it reflects lowering of the molecular symmetry from apparent C_4 to C_2 or even
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 lower and (ii) it shows that the residence time in the exchange between nonequivalent
4 analogous nuclei must be larger than 0.05 s, i.e. the internal motion in the molecule B is at
5
6
7
8 least two orders of magnitude slower than that in the form A. Therefore, the most plausible
9
10 conclusion appears to be that the form B (and probably also C) is a hydrate of the complex A
11
12 with a water molecule placed inside the calixarene cavity. The molecule has to sit near
13
14 enough to the lower rim in order to interact with the bound H_3O^+ ion, which necessarily
15
16 means that it interacts with at least two opposite aromatic rings (probably by a weak hydrogen
17
18 bond) lowering thus the symmetry of the complex and slowing down its internal motion.
19
20

21
22 **Quantum chemical calculations.** Geometry optimizations of calixarene and its complex with
23
24 H_3O^+ were performed by the quantum chemical calculations at the density functional level of
25
26 theory (DFT, B3LYP functional) using the Gaussian 03 suite of programs [22]. The 6-31G(d)
27
28 basis set was used and the optimization was unconstrained. The optimized structure of free
29
30 calixarene is shown in Figure 7A. The most stable conformation forms a pinched cone
31
32 structure with a C_2 symmetry.
33
34

35
36 The structure obtained by optimization of the calixarene - H_3O^+ complex is shown in
37
38 Figure 7B, together with the lengths of hydrogen bonds (in Å). Compared to free molecule the
39
40 calixarene part of the complex is distorted so that its structure is very close to the C_4
41
42 symmetry. The hydroxonium ion is placed in the cage formed by calixarene and is bound by
43
44 strong hydrogen bonds to oxygen atoms of three C-O-C groups and by two somewhat weaker
45
46 hydrogen bonds to the last C-O-C group. The stabilization energy of the complex calculated
47
48 as the difference between the energies of the optimized complex and optimized free
49
50 calixarene molecule and hydroxonium ion is -394.72 kJ/mol. Stabilization energy of the
51
52 complex can be expressed also with respect to the aqueous hydroxonium ion $[\text{H}_3\text{O}^+(\text{H}_2\text{O})_3]$
53
54 and free calixarene (**1**) according to the formula
55
56
57
58



1
2
3 Then it is -8.20 kJ/mol. This value, corrected further for entropy change, probably
4
5 corresponds to the value of K established above by NMR.
6
7

8 The gradual development of another form of the complex detected by NMR led us to
9
10 explore its further possible changes or interactions. As the structure described above
11
12 corresponds to the global energy minimum found by unconstrained geometry optimization, a
13
14 new form could scarcely be produced by an internal rearrangement of the structure. We thus
15
16 explored possible interactions of the complex with other components of the system.
17
18 Nitrobenzene, which is largely prevalent in the system, does not appear do form any stable
19
20 formation with the complex. However, water does. The calculated geometry of the complex
21
22 hydrate is shown in Figure 8. The stabilization energy corresponding to such hydration is
23
24 predicted to be - 19.75 kJ/mol, i.e. sufficiently high for the hydrate to be stable.
25
26
27
28

29 The interesting feature of the hydrate is that the water molecule is bound by the
30
31 already bonded H_3O^+ ion by a system of three medium-strong hydrogen bonds indicated in
32
33 the Figure. In order to accomplish that, the water molecule has to penetrate the rather rigid
34
35 and hydrophobic calixarene cavity (no other way apparently leads to a stable product), which
36
37 can be expected to lead to rather long reaction times as observed. In addition, the water
38
39 molecule also interacts with the aromatic systems by two weak hydrogen bonds (some of the
40
41
42
43
44
45

46 **Figure 7:** Optimized geometry of free **1** (A) and the **1**· H_3O^+ complex (B) calculated at the
47
48 B3LYP/6-31G(d) level (positions of hydrogen atoms in **1** are not shown)

49
50
51 O-H- C_{ar} distances being as short as 2.55 Å). In agreement with NMR spectra, these bonds
52
53 without doubt slow down internal motions in the calixarene unit.
54

55 In contrast to NMR, however, we were not able to find any alternative stable form of
56
57 the hydrate. Taking into account the fact that our calculations were done *in vacuo* (as no
58
59 sufficiently reliable correction for the influence of polar medium has been found for this level
60

of precision), we cannot be quite sure that other structural forms of the complex hydrate are excluded by theory.

Figure 8: Optimized geometry of the hydrated $\mathbf{1} \cdot \text{H}_3\text{O}^+$ complex calculated at the B3LYP/6-31G(d) level (positions of hydrogen atoms in $\mathbf{1}$ are not shown).

Conclusions. According to the evidence of ^1H and ^{13}C NMR spectra consistent with the predictions of high-precision DFT calculations, calixarene $\mathbf{1}$ binds the H_3O^+ ion into an equimolecular complex. The formation of the main form of the complex, similarly as the exchange between free and bound $\mathbf{1}$, is very fast but the equilibrium is only moderately favorable to the complex. In the main form, H_3O^+ is probably bound by three strong hydrogen bonds to three aryl-oxygen atoms of $\mathbf{1}$ and by two weaker hydrogen bonds to the remaining C-O-C group. In contrast to free $\mathbf{1}$, the complex has almost C_4 symmetry. Under prolonged storage of the complex-containing mixtures, small amounts of another form of the complex appear. It is evidently formed from the original complex and probably is the product of its hydration.

When compared with the calixarenes with strongly coordinating groups in addition to aryl-oxygen atoms at their lower rim [13-15], the present calixarene type clearly is a markedly less efficient complexing agent. From this one can conclude that the C-O-C groups, although they clearly participate in the complex H_3O^+ binding in the former types [13-15], are as such less effective than in cooperation with e.g. dimethylamide, dimethylthioamide, ester and keto-groups.

EXPERIMENTAL

Materials and samples. Nitrobenzene- d_5 was purchased by Aldrich. Tetrapropoxy-*p*-*tert*-butylcalix[4]arene ($\mathbf{1}$) was prepared by using the procedure published elsewhere [23].

1
2
3 Preparation of hydrogen bis(1,2-dicarbollyl) cobaltate (HDCC) was described earlier [16].
4
5 For NMR samples, 0.5×10^{-5} mol of **1** was dissolved in a mixture of appropriate amounts of
6
7 nitrobenzene-*d*₅ and 0.002 mol/L solution of HDCC in the same solvent. The mixing and all
8
9 subsequent measurements were done at 296 K.
10
11

12 **NMR spectra.** ¹H and ¹³C NMR spectra were measured in a quadrature detection mode at
13
14 300.13 and 75.45 MHz, respectively, with an upgraded Bruker Avance DPX300 spectrometer.
15
16 32 and 64 kpoints were measured for ¹H and ¹³C NMR, respectively. ¹H NMR spectra were
17
18 measured using a self-devised diffusion-filtering stimulated echo pulse sequence $-d_1-\pi/2_\phi-$
19
20 $GR-\pi/2_\sigma-d_2-\pi/2_y-GR-REC_x$ in order to remove obscuring broad signals of water and
21
22 nitrobenzene; *d*₁ and *d*₂ were 10 s and 60 ms, respectively, the z-gradient GR was 50 G/cm
23
24 and phase cycling was $\phi = x, y, -x, -y$, $\sigma = \phi + \pi/2$. 128 scans were collected for each
25
26 spectrum. ¹³C NMR measurements were performed using either the ¹H-¹³C DEPT45 sequence
27
28 (collecting 8000 scans) or its combination with its long-range variant optimized for the
29
30 coupling constant 6 Hz (collecting at least 20000 scans); exponential weighting (*lb* = 1 Hz)
31
32 was used before Fourier transform. In ¹H – ¹³C 2D HSQC and HMBC spectra used for signal
33
34 assignment, 1028 points in F2 and 256 increments in F1 dimensions were measured using a z-
35
36 gradient inverse-detection probe.
37
38
39
40
41
42


43 **Quantum mechanical calculations.** *Ab initio* molecular orbital calculations were performed
44
45 using the GAUSSIAN 03 suite of programs [16]. Molecular geometry was fully optimized at
46
47 the B3LYP level of density functional theory (DFT) with the 6-31G(d) basis set. The
48
49 optimization was unrestrained. Several local configurations near the achieved energy
50
51 minimum were examined. As the renewed optimizations converged to the same molecular
52
53 geometry, we believe the achieved energy minimum to be the global one.
54
55
56
57
58
59
60


Acknowledgements. This work was supported by the Academy of Sciences of the Czech Republic, Project T400500402, and the Czech Ministry of Education, Youth and Sports, Projects MSM 4977751303 and MSM 6046137307.


References


1. Gutsche, C. D. *Calixarenes Revisited*. The Royal Society of Chemistry, Cambridge, 1998.
2. Böhmer, V. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 713.
3. Arduini, A.; Pochini, A.; Reverberi, S.; Ungaro, R. *Tetrahedron* **1986**, *42*, 2089.
4. Arduini, A.; Ghidini, E.; Pochini, A.; Ungaro, R.; Andreotti, G.D.; Calestani, G.; Ugozzoli, F. *J. Inclusion Phenom.* **1988**, *6*, 119.
5. Arnaud-Neu, F.; Collins, E.M.; Deasy, M.; Ferguson, G.; Harris, S.J.; Kaitner, B.; Lough, A.J.; McKervey, M.A.; Marques, E.; Ruhl, B.L.; Schwing-Weill, M.J.; Seward, E.M. *J. Am. Chem. Soc.* **1989**, *111*, 8681.
6. Arnaud-Neu, F.; Barrett, G.; Harris, S.J.; Owens, M.; McKervey, M.A.; Schwing-Weill, M.J.; Schwinté, P. *Inorg. Chem.* **1993**, *32*, 2644.
7. Ohto, K.; Murakami, E.; Shinohara, T.; Shiratsuchi, K.; Inoue, K.; Iwasaki, M. *Anal. Chim. Acta* **1997**, *341*, 275.
8. Ye, Z.; He, W.; Shi, X.; Zhu, L. *J. Coord. Chem.* **2001**, *54*, 105.
9. Danil de Namor, A.F.; Chahine, S.; Kowalska, D.; Castellano, E.E.; Piro, O.E. *J. Am. Chem. Soc.* **2002**, *124*, 12824.
10. Marcos, P. M.; Ascenso, J.R.; Segurado, M.A.P.; Pereira, J.L.C. *J. Inclusion Phenom.* **2002**, *42*, 281.
11. Marcos, P.M.; Félix, S.; Ascenso, J.R.; Segurado, M.A.P.; Pereira, J.L.C.; Khazaeli-Parsa, P.; Hubscher-Bruder, V.; Arnaud-Neu, F. *New J.Chem.* **2004**, *28*, 748.
12. Gutsche, C. D. *Aldrichimica Acta* **1995**, *28*, 3.
13. Kříž, J.; Dybal, J.; Makrlík, E.; Vaňura, P. *Supramol. Chem.* in print.
14. Kříž, J.; Dybal, J.; Makrlík, E.; Vaňura, P. *Supramol. Chem.* in print.
15. Kříž, J.; Dybal, J.; Makrlík, E.; Budka, J.; Vaňura, P. *Monatsh. Chem.* in print.
16. Makrlík, E.; Vaňura, P. *Talanta* **1985**, *32*, 423.
17. Kříž, J.; Makrlík, E.; Vaňura, P. *Biopolymers* **2006**, *81*, 104.
18. Kříž, J.; Dybal, J.; Makrlík, E. *Biopolymers* **2006**, *82*, 536.
19. Meiboom, S.; Gill, D. *Rev. Sci. Instrum.* **1958**, *29*, 688.
20. Luz, L.; Meiboom, S. *J. Chem. Phys.* **1963**, *39*, 366.


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. Canet, D.; Robert, J.B. in *Dynamics of Solutions and Fluid Mixtures by NMR* (J. J. Delpuech Ed.), Wiley, Chichester 1995, p. 127.
22. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, Jr., J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; and Pople, J. A. *Gaussian 03, Revision C.02*; Gaussian, Inc.: Wallingford CT, 2004.
23. Kelderman, E.; Derhaeg, L.; Heesink, G.J.T.; Verboom, W.; Engbersen, J.F.J.; van Hulst, N.F.; Persoons, A.; Reinhoudt, D.N. *Angew. Chem.* **1992**, *104*, 1107.





















For Peer Review Only