

Structural characterisation of trimethylsilyl-protected DNA bases

Kieran Molloy, Gareth Price, Doug Smith, Gabriele Kociok-Khön

▶ To cite this version:

Kieran Molloy, Gareth Price, Doug Smith, Gabriele Kociok-Khön. Structural characterisation of trimethylsilyl-protected DNA bases. Supramolecular Chemistry, 2008, 20 (08), pp.697-707. 10.1080/10610270701732885. hal-00513518

HAL Id: hal-00513518 https://hal.science/hal-00513518

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Supramolecular Chemistry



Structural characterisation of trimethylsilyl-protected DNA bases

Journal:	Supramolecular Chemistry
Manuscript ID:	GSCH-2007-0133.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	05-Oct-2007
Complete List of Authors:	Molloy, Kieran; University of Bath, Chemistry Price, Gareth; University of Bath, Chemistry Smith, Doug; University of Bath, Chemistry Kociok-Khön, Gabriele; University of Bath, Chemistry
Keywords:	silylated DNA bases, adenine, guanine, cytosine, thymine





C22

C21

Si2





Figure 2. The asymmetric unit of 2; ellipsoids are at the 50% probability level. Hydrogen atoms of the Me3Si groups omitted for clarity. 262x107mm (300 x 300 DPI)



Figure 3 The helical chain structure of 2 viewed (a) orthogonal to, and (b) along the axis of propagation (c); Symmetry operator 3/2-x, 1-y, 1/2+z. 444x329mm (96 x 96 DPI)





Figure 4 The asymmetric unit of 3 showing the two independent molecules in the asymmetric unit; ellipsoids are at the 50% probability level. Hydrogen atoms of the Me3Si groups omitted for clarity. Symmetry operator 1-x, y,1/2-z. 266x164mm (300 x 300 DPI)





Figure 5 Two views of the dimerisation of one of the independent molecules on the asymmetric unit of 3. 444x329mm (96 x 96 DPI)



444x329mm (96 x 96 DPI)





Figure 7 The lattice structure of 4 viewed (a) orthogonal to, and (b) along the axis of propagation 444x329mm (96 x 96 DPI)





Figure 1. The asymmetric unit of 1; ellipsoids are at the 50% probability level. Hydrogen atoms of the Me3Si groups omitted for clarity. 210x129mm (300 x 300 DPI)

URL: http:/mc.manuscriptcentral.com/tandf/gsch Email: suprachem@mail.cm.utexas.edu



Figure 2. The asymmetric unit of 2; ellipsoids are at the 50% probability level. Hydrogen atoms of the Me3Si groups omitted for clarity. 262x107mm (300 x 300 DPI)



Figure 3 The helical chain structure of 2 viewed (a) orthogonal to, and (b) along the axis of propagation (c); Symmetry operator 3/2-x, 1-y, 1/2+z. 444x329mm (96 x 96 DPI)





Figure 4 The asymmetric unit of 3 showing the two independent molecules in the asymmetric unit; ellipsoids are at the 50% probability level. Hydrogen atoms of the Me3Si groups omitted for clarity. Symmetry operator 1-x, y,1/2-z. 266x164mm (300 x 300 DPI)

URL: http:/mc.manuscriptcentral.com/tandf/gsch Email: suprachem@mail.cm.utexas.edu





Figure 5 Two views of the dimerisation of one of the independent molecules on the asymmetric unit of 3. 444x329mm (96 x 96 DPI)



444x329mm (96 x 96 DPI)

URL: http:/mc.manuscriptcentral.com/tandf/gsch Email: suprachem@mail.cm.utexas.edu





Figure 7 The lattice structure of 4 viewed (a) orthogonal to, and (b) along the axis of propagation 444x329mm (96 x 96 DPI)

