



HAL
open science

Symmetry properties of Penrose type tilings

Nicolae Cotfas

► **To cite this version:**

Nicolae Cotfas. Symmetry properties of Penrose type tilings. Philosophical Magazine, 2008, 88 (13-15), pp.2017-2023. 10.1080/14786430802035691 . hal-00513883


HAL Id: hal-00513883

<https://hal.science/hal-00513883>

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Symmetry properties of Penrose type tilings

Journal:	<i>Philosophical Magazine & Philosophical Magazine Letters</i>
Manuscript ID:	TPHM-07-Sep-0272.R1
Journal Selection:	Philosophical Magazine
Date Submitted by the Author:	19-Feb-2008
Complete List of Authors:	Cotfas, Nicolae; University of Bucharest, Faculty of Physics
Keywords:	aperiodic materials, quasicrystals
Keywords (user supplied):	Penrose tiling, self-similarities
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p> <p>cotfas-revised1.tex</p>	



Philosophical Magazine

Vol. 00, No. 00, 21 December 2008, 1–6

Symmetry properties of Penrose type tilings

N. COTFAS*

Faculty of Physics, University of Bucharest, Romania

(.....)

The Penrose tiling is directly related to the atomic structure of certain decagonal quasicrystals and, despite its aperiodicity, is highly symmetric. It is known that the numbers $1, -\tau, (-\tau)^2, (-\tau)^3, \dots$, where $\tau = (1 + \sqrt{5})/2$, are scaling factors of the Penrose tiling. We show that the set of scaling factors is much larger, and for most of them the number of the corresponding inflation centers is infinite.

1. Introduction

The best known and by far most studied aperiodic tiling of plane is the Penrose tiling [1, 2]. This tiling plays an important role in the quasicrystal structure understanding and has some remarkable mathematical properties. Direct structure observation, using high-resolution transmission electron microscopy, shows that the atomic arrangement in the tenfold symmetry plane of certain decagonal quasicrystals can simply be interpreted in terms of the Penrose tiling [3, 4]. Many aspects concerning the mathematical properties have been studied and are found in the literature; see for example [5–7].

In this paper, we present some results concerning the inflation symmetries of the Penrose type tilings. Our approach, based on the use of the three projectors corresponding to the usual decomposition of the five-dimensional superspace used in the cut and project construction, is an extension of the method used by Katz and Duneau in the description of self-similarity properties of icosahedral quasicrystals [8, 9]. There are also two important differences. In the case analysed by Katz and Duneau, the representation of the icosahedral group in the superspace \mathbb{R}^6 is a sum of only two irreducible representations, and the orthogonal projection of the lattice \mathbb{Z}^6 on the internal subspace is a dense subset. In our case, the representation of the symmetry group C_5 in the superspace \mathbb{R}^5 is a sum of three real irreducible representations, and the orthogonal projection of the lattice \mathbb{Z}^5 on the internal three-dimensional subspace is not a dense subset. The vertex set of the Penrose tiling is a multi-component model set [7, 10, 11], and we have adapted the Katz and Duneau method to such a case.

The set Λ of inflation factors obtained by using our method is a union of four cut and project sets. To our knowledge, the known inflation factors of the Penrose tiling are only those forming the multiplicative semigroup $\mathcal{I} = \{(-\tau)^k \mid k \in \mathbb{N}\}$. Since the elements of this semigroup are not uniformly distributed in \mathbb{R} we can not have $\Lambda \subset \mathcal{I}$. We have found new scaling factors and inflation centers, but very probably our list is not exhaustive. It is known that the inflation factor τ is directly related to a construction of the Penrose tiling by tile substitutions [5, 6]. We do

*Email: ncotfas@yahoo.com

not know if the new found scaling factors are related to some substitution rules or not.

The importance of inflation symmetries in quasicrystal physics is comparable with that of translation symmetries in the description of crystals. A fractal shape of the energy spectrum may be a direct consequence of the self-similarity properties of the quasicrystal. A method to study the self-similarity properties of the cut and project sets has been presented in [12], but as the authors show at the end of the paper, the method does not apply to the Penrose type tilings.

2. Penrose type tilings

In this section we review certain results concerning the Penrose type tilings defined by projection and introduce certain notations. The relation

$$a : \mathbb{R}^5 \longrightarrow \mathbb{R}^5, \quad a(x_1, x_2, x_3, x_4, x_5) = (x_2, x_3, x_4, x_5, x_1) \quad (1)$$

defines an orthogonal representation of the cyclic group $C_5 = \langle a \mid a^5 = e \rangle = \{e, a, a^2, a^3, a^4\}$ in the usual five-dimensional Euclidean space \mathbb{R}^5 . The space \mathbb{R}^5 can be decomposed into a sum of orthogonal C_5 -invariant subspaces $\mathbb{R}^5 = \mathbf{E} \oplus \mathbf{E}' \oplus \mathbf{E}''$, where

$$\mathbf{E} = \left\{ \alpha \left(1, \cos \frac{2\pi}{5}, \cos \frac{4\pi}{5}, \cos \frac{6\pi}{5}, \cos \frac{8\pi}{5} \right) + \beta \left(0, \sin \frac{2\pi}{5}, \sin \frac{4\pi}{5}, \sin \frac{6\pi}{5}, \sin \frac{8\pi}{5} \right) \mid \alpha, \beta \in \mathbb{R} \right\}$$

$$\mathbf{E}' = \left\{ \alpha \left(1, \cos \frac{4\pi}{5}, \cos \frac{8\pi}{5}, \cos \frac{2\pi}{5}, \cos \frac{6\pi}{5} \right) + \beta \left(0, \sin \frac{4\pi}{5}, \sin \frac{8\pi}{5}, \sin \frac{2\pi}{5}, \sin \frac{6\pi}{5} \right) \mid \alpha, \beta \in \mathbb{R} \right\}$$

$$\mathbf{E}'' = \left\{ \alpha(1, 1, 1, 1, 1) \mid \alpha \in \mathbb{R} \right\}$$

It is known that $\cos \frac{\pi}{5} = \frac{\tau}{2}$ and $\cos \frac{2\pi}{5} = -\frac{\tau'}{2}$, where τ, τ' are the irrational numbers $\tau = (1 + \sqrt{5})/2$ and $\tau' = (1 - \sqrt{5})/2$.

The vectors $e_1 = (1, 0, 0, 0, 0)$, $e_2 = (0, 1, 0, 0, 0)$, ..., $e_5 = (0, 0, 0, 0, 1)$ form the canonical basis of \mathbb{R}^5 , and the matrices of the orthogonal projectors corresponding to \mathbf{E} , \mathbf{E}' and \mathbf{E}'' in this basis are

$$\pi = \frac{1}{5} \mathcal{M}(2, -\tau', -\tau), \quad \pi' = \frac{1}{5} \mathcal{M}(2, -\tau, -\tau'), \quad \pi'' = \frac{1}{5} \mathcal{M}(1, 1, 1) \quad (2)$$

where

$$\mathcal{M}(\alpha, \beta, \gamma) = \begin{pmatrix} \alpha & \beta & \gamma & \gamma & \beta \\ \beta & \alpha & \beta & \gamma & \gamma \\ \gamma & \beta & \alpha & \beta & \gamma \\ \gamma & \gamma & \beta & \alpha & \beta \\ \beta & \gamma & \gamma & \beta & \alpha \end{pmatrix}. \quad (3)$$

The projections on the space \mathbf{E} of the endpoints of the vectors e_1, e_2, \dots, e_5 are the vertices of a regular pentagon. Any point $x \in \mathbb{Z}^5$ has ten arithmetical neighbours, namely, $x \pm e_1, x \pm e_2, x \pm e_3, x \pm e_4, x \pm e_5$, and the projections on \mathbf{E} of these points are the vertices of a regular decagon. If we project on \mathbf{E} the 2-faces of the unit hypercube $(0, 1)^5$ we get, up to certain rotations, only two rhombs.

We embed the physical two-dimensional space into \mathbb{R}^5 by identifying it with the subspace \mathbf{E} , and regard the complementary space $\mathbf{E}^\perp = \mathbf{E}' \oplus \mathbf{E}''$ which corresponds to the projector $\pi^\perp = \pi' + \pi''$ as an internal space. By using the strip

$$\mathbf{S} = \{ x \in \mathbb{R}^5 \mid \pi^\perp x \in \mathbf{W} \} = \mathbf{E} + (0, 1)^5 \quad (4)$$

1 corresponding to the window $\mathbf{W} = \pi^\perp((0, 1)^5)$ we define the pattern [8]

$$2 \mathcal{P} = \pi(\mathbb{Z}^5 \cap \mathbf{S}) = \{ \pi x \mid x \in \mathbb{Z}^5, \pi^\perp x \in \mathbf{W} \}. \quad (5)$$

3
 4
 5
 6 The window \mathbf{W} defined as the projection on the three-dimensional space \mathbf{E}^\perp of the
 7 open unit hypercube $(0, 1)^5$ is a polyhedron and one can remark that its frontier
 8 contains some points belonging to $\pi^\perp(\mathbb{Z}^5)$. The pattern \mathcal{P} is interesting, but it is
 9 a particular one. For any vector $v \in \mathbf{E}'$ we can consider the translated strip $v + \mathbf{S}$
 10 which corresponds to the translated window $v + \mathbf{W}$, and define the pattern

$$11 \mathcal{P}_v = \pi(\mathbb{Z}^5 \cap (v + \mathbf{S})) = \{ \pi x \mid x \in \mathbb{Z}^5, \pi^\perp x \in v + \mathbf{W} \}. \quad (6)$$

12
 13 We say that \mathcal{P}_v is a *non-singular pattern* if v is such that the frontier of $v + \mathbf{W}$
 14 does not contain any element of $\pi^\perp(\mathbb{Z}^5)$. In the opposite case we say that \mathcal{P}_v is a
 15 *singular pattern*. One can prove that the points of any non-singular pattern \mathcal{P}_v are
 16 the vertices of a tiling of the plane \mathbf{E} formed by only two tiles, each of them in a
 17 finite number of orientations [8].

18 The hyperplane

$$19 \mathcal{E} = \mathbf{E} \oplus \mathbf{E}' = \{ (x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5 \mid x_1 + x_2 + x_3 + x_4 + x_5 = 0 \} \quad (7)$$

20 is orthogonal to the one-dimensional space $\mathbf{E}'' = \{ (\alpha, \alpha, \alpha, \alpha, \alpha) \mid \alpha \in \mathbb{R} \}$, and the
 21 lattice \mathbb{Z}^5 is contained in the family of parallel equidistant spaces $\{ \mathcal{E}_n \mid n \in \mathbb{Z} \}$,
 22 where

$$23 \mathcal{E}_n = nw + \mathcal{E} = \{ (x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5 \mid x_1 + x_2 + x_3 + x_4 + x_5 = n \} \quad (8)$$

24 and $w = (0.2, 0.2, 0.2, 0.2, 0.2)$. If $x = (x_1, x_2, x_3, x_4, x_5)$ belongs to the open unit
 25 hypercube $(0, 1)^5$ then $0 < x_1 + x_2 + x_3 + x_4 + x_5 < 5$, and therefore only the
 26 hyperplanes $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ and \mathcal{E}_4 meet $(0, 1)^5$. This means that the set $\mathbb{Z}^5 \cap (v + \mathbf{S})$
 27 occurring in the definition of \mathcal{P}_v is contained in $\mathcal{E}_1 \cup \mathcal{E}_2 \cup \mathcal{E}_3 \cup \mathcal{E}_4$, whence

$$28 \mathcal{P}_v = \bigcup_{n=1}^4 \{ \pi x \mid x \in \mathbb{Z}^5 \cap \mathcal{E}_n, \pi^\perp x \in v + \mathbf{W}_n \} \quad (9)$$

29 where $\mathbf{W}_n = \pi'(\mathcal{E}_n \cap (0, 1)^5) = -nw + (\mathcal{E}_n \cap \mathbf{W})$. By direct computation, one can
 30 prove that the set \mathbf{W}_1 is the interior of the regular pentagon with the vertices $\pi'e_1,$
 31 $\pi'e_3, \pi'e_5, \pi'e_2, \pi'e_4$ (considered in this order), $\mathbf{W}_2 = -\tau\mathbf{W}_1, \mathbf{W}_3 = \tau\mathbf{W}_1$ and
 32 $\mathbf{W}_4 = -\mathbf{W}_1$.

31 3. Self-similarities of the singular pattern \mathcal{P}

32 The translation corresponding to the vector $5w = (1, 1, 1, 1, 1) \in \mathbb{Z}^5 \cap \mathbf{E}''$

$$33 \mathbb{Z}^5 \cap \mathcal{E}_n \longrightarrow \mathbb{Z}^5 \cap \mathcal{E}_{n+5} : x \mapsto x + 5w \quad (10)$$

34 is a one-to-one transformation, $\pi(x + 5w) = \pi x$ and $\pi'(x + 5w) = \pi'x$. If we put
 35 $\mathbf{W}_{5k} = \emptyset, \mathbf{W}_{5k+1} = \mathbf{W}_1, \mathbf{W}_{5k+2} = -\tau\mathbf{W}_1, \mathbf{W}_{5k+3} = \tau\mathbf{W}_1$ and $\mathbf{W}_{5k+4} = -\mathbf{W}_1$

for any $k \in \mathbb{Z}$ then we can re-write the definition of \mathcal{P}_v as

$$\mathcal{P}_v = \bigcup_{n \in \mathbb{Z}} \{ \pi x \mid x \in \mathbb{Z}^5 \cap \mathcal{E}_n, \pi' x \in v + \mathbf{W}_n \}. \tag{11}$$

We say [8, 12] that $\lambda \in \mathbb{R}$ is an *inflation factor* of \mathcal{P}_v if there exists a point $y \in \mathbf{E}$ (called an *inflation center*) such that \mathcal{P}_v is invariant under the mapping $\mathbf{E} \rightarrow \mathbf{E} : x \mapsto y + \lambda(x - y)$, that is, $x \in \mathcal{P}_v \implies y + \lambda(x - y) \in \mathcal{P}_v$.

Theorem 3.1: *The matrix $A = \lambda\pi + \lambda'\pi' + \lambda''\pi''$ has integer entries if and only if $(\lambda, \lambda', \lambda'')$ belongs to the set*

$$\mathcal{L} = \left\{ \left(\frac{\alpha - \beta}{2} + \beta\tau, \frac{\alpha - \beta}{2} + \beta\tau', \frac{\alpha + \gamma}{2} + 2\gamma \right) ; \alpha, \beta, \gamma \in \mathbb{Z}, (-1)^\alpha = (-1)^\beta = (-1)^\gamma \right\}.$$

Proof: From (2) we get

$$A = \mathcal{M} \left(\frac{2}{5}\lambda + \frac{2}{5}\lambda' + \frac{1}{5}\lambda'', \frac{-1 + \sqrt{5}}{10}\lambda + \frac{-1 - \sqrt{5}}{10}\lambda' + \frac{1}{5}\lambda'', \frac{-1 - \sqrt{5}}{10}\lambda + \frac{-1 + \sqrt{5}}{10}\lambda' + \frac{1}{5}\lambda'' \right).$$

If this matrix has integer entries then $2\lambda + 2\lambda' + \lambda'' \in 5\mathbb{Z}$, $-\lambda - \lambda' + 2\lambda'' \in 5\mathbb{Z}$ and $\lambda - \lambda' \in \mathbb{Z}\sqrt{5}$, whence $\lambda + \lambda' \in \mathbb{Z}$. Denoting $\lambda + \lambda' = \alpha$ and $\lambda - \lambda' = \beta\sqrt{5}$ we obtain $\lambda = \frac{\alpha - \beta}{2} + \beta\tau$, $\lambda' = \frac{\alpha - \beta}{2} + \beta\tau'$ and $-\alpha + 2\lambda'' \in 5\mathbb{Z}$. If we put $-\alpha + 2\lambda'' = 5\gamma$ then $\lambda'' = \frac{\alpha + 5\gamma}{2}$ and $A = \mathcal{M} \left(\frac{\alpha + \gamma}{2}, \frac{\gamma + \beta}{2}, \frac{\gamma - \beta}{2} \right)$. In order to have integer entries the numbers α , β and γ must have the same parity. \square

Theorem 3.2: *If $(\lambda, \lambda', \lambda'')$ belongs to the subset*

$$\tilde{\mathcal{L}} = \{ (\lambda, \lambda', \lambda'') \in \mathcal{L} \mid \lambda' \mathbf{W}_n \subseteq \mathbf{W}_{\lambda'' n}, \text{ for any } n \in \{1, 2, 3, 4\} \} \tag{12}$$

then λ is an inflation factor of the singular pattern \mathcal{P} .

Proof: If $(\lambda, \lambda', \lambda'') \in \tilde{\mathcal{L}}$ then the corresponding mapping $A : \mathbb{R}^5 \rightarrow \mathbb{R}^5$, $A = \lambda\pi + \lambda'\pi' + \lambda''\pi''$ has the property

$$\left. \begin{array}{l} x \in \mathbb{Z}^5 \cap \mathcal{E}_n \\ \pi' x \in \mathbf{W}_n \end{array} \right\} \implies \left\{ \begin{array}{l} Ax \in \mathbb{Z}^5 \cap \mathcal{E}_{\lambda'' n} \\ \pi'(Ax) \in \mathbf{W}_{\lambda'' n} \end{array} \right. \tag{13}$$

In view of the definition (6) considered for $v = (0, 0, 0, 0, 0)$, this means that $\pi x \in \mathcal{P} \implies \pi(Ax) \in \mathcal{P}$. But $\pi(Ax) = \lambda\pi x$, and hence, we have $y \in \mathcal{P} \implies \lambda y \in \mathcal{P}$. \square

Theorem 3.3: *Each element of the set*

$$\begin{aligned} \Lambda = & \left\{ 1 - \frac{\beta + 5\gamma}{2} + \beta\tau \mid \beta, \gamma \in \mathbb{Z}, (-1)^\beta = (-1)^\gamma, -\frac{\tau}{2} \leq 1 - \frac{\beta + 5\gamma}{2} + \beta\tau' \leq 1 \right\} \\ & \cup \left\{ 2 - \frac{\beta + 5\gamma}{2} + \beta\tau \mid \beta, \gamma \in \mathbb{Z}, (-1)^\beta = (-1)^\gamma, -\frac{1}{2} \leq 2 - \frac{\beta + 5\gamma}{2} + \beta\tau' \leq \frac{1}{\tau} \right\} \\ & \cup \left\{ 3 - \frac{\beta + 5\gamma}{2} + \beta\tau \mid \beta, \gamma \in \mathbb{Z}, (-1)^\beta = (-1)^\gamma, -\frac{1}{\tau} \leq 3 - \frac{\beta + 5\gamma}{2} + \beta\tau' \leq \frac{1}{2} \right\} \\ & \cup \left\{ 4 - \frac{\beta + 5\gamma}{2} + \beta\tau \mid \beta, \gamma \in \mathbb{Z}, (-1)^\beta = (-1)^\gamma, -1 \leq 4 - \frac{\beta + 5\gamma}{2} + \beta\tau' \leq \frac{\tau}{2} \right\} \end{aligned}$$

is an inflation factor of \mathcal{P} .

Proof: It is sufficient to analyse the cases $\lambda'' \in \{0, 1, 2, 3, 4, 5\}$. In the case $\lambda'' = 0$ we do not obtain any inflation factor. The ratio between the distance from the

1 centre of a regular pentagon to one of its sides and the distance from the centre to
 2 one vertex is $\cos\frac{\pi}{5} = \frac{\tau}{2}$. By imposing the conditions

$$\begin{aligned} &\lambda'W_1 \subseteq W_1, \quad \lambda'W_2 \subseteq W_2, \quad \lambda'W_3 \subseteq W_3, \quad \lambda'W_4 \subseteq W_4 \text{ in the case } \lambda'' = 1, \\ &\lambda'W_1 \subseteq W_2, \quad \lambda'W_2 \subseteq W_4, \quad \lambda'W_3 \subseteq W_1, \quad \lambda'W_4 \subseteq W_3 \text{ in the case } \lambda'' = 2, \\ &\lambda'W_1 \subseteq W_3, \quad \lambda'W_2 \subseteq W_1, \quad \lambda'W_3 \subseteq W_4, \quad \lambda'W_4 \subseteq W_2 \text{ in the case } \lambda'' = 3, \\ &\lambda'W_1 \subseteq W_4, \quad \lambda'W_2 \subseteq W_3, \quad \lambda'W_3 \subseteq W_2, \quad \lambda'W_4 \subseteq W_1 \text{ in the case } \lambda'' = 4 \end{aligned}$$

10 we get the set Λ . □

11 The well-known inflation factor $\lambda = -\tau$ belongs to Λ , and is obtained for $\gamma = 1$,
 12 $\beta = -1$. One can remark that Λ is a union of four cut and project sets. Since the
 13 elements of the multiplicative semigroup $\mathcal{I} = \{(-\tau)^k \mid k \in \mathbb{N}\}$ are not uniformly
 14 distributed in \mathbb{R} , we can not have $\Lambda \subset \mathcal{I}$.

15 **Theorem 3.4:** *If $(\lambda, \lambda', \lambda'')$ belongs to the subset*

$$16 \tilde{\mathcal{L}}_0 = \{ (\lambda, \lambda', \lambda'') \in \mathcal{L} \mid \lambda' \overline{W}_n \subset W_{\lambda''n} \text{ for any } n \in \{1, 2, 3, 4\} \} \quad (14)$$

17 then there is an infinite number of inflation centers $y \in \mathbf{E}$ such that

$$18 \mathbf{E} \longrightarrow \mathbf{E} : x \mapsto y + \lambda(x - y) \quad (15)$$

19 is an affine self-similarity of \mathcal{P} , that is, such that $x \in \mathcal{P} \implies y + \lambda(x - y) \in \mathcal{P}$.

20 **Proof:** If $(\lambda, \lambda', \lambda'') \in \tilde{\mathcal{L}}_0$ then there is a neighbourhood $\mathcal{V}_{(\lambda, \lambda', \lambda'')}$ of the origin in
 21 \mathbf{E}' such that the relation $(1 - \lambda')u + \lambda'W_n \subset W_{\lambda''n}$ is verified for any $n \in \{1, 2, 3, 4\}$
 22 and any $u \in \mathcal{V}_{(\lambda, \lambda', \lambda'')}$. Since the projection $\pi'(\mathbb{Z}^5 \cap \mathcal{E})$ is dense in \mathbf{E}' , the set
 23 $\mathcal{C}_{(\lambda, \lambda', \lambda'')} = \{t \in \mathbb{Z}^5 \cap \mathcal{E} \mid \pi't \in \mathcal{V}_{(\lambda, \lambda', \lambda'')}\}$ is an infinite set. For each $t \in \mathcal{C}_{(\lambda, \lambda', \lambda'')}$
 24 the mapping $A_t : \mathbb{R}^5 \longrightarrow \mathbb{R}^5$, $A_t x = t + (\lambda\pi + \lambda'\pi' + \lambda''\pi'')(x - t)$ satisfies the
 25 relations $\pi'(A_t x) = \pi't + \lambda'\pi'(x - t) = (1 - \lambda')\pi't + \lambda'\pi'x$, and

$$26 \left. \begin{aligned} x &\in \mathbb{Z}^5 \cap \mathcal{E}_n \\ \pi'x &\in W_n \end{aligned} \right\} \implies \left\{ \begin{aligned} A_t x &\in \mathbb{Z}^5 \cap \mathcal{E}_{\lambda''n} \\ \pi'(A_t x) &\in W_{\lambda''n} \end{aligned} \right. \quad (16)$$

27 This means that $\pi x \in \mathcal{P} \implies \pi(A_t x) \in \mathcal{P}$, that is, $\pi x \in \mathcal{P} \implies \pi t + \lambda(\pi x - \pi t) \in \mathcal{P}$.
 28 □

29 4. Self-similarities of the patterns \mathcal{P}_v

30 In order to avoid certain ‘defects’ in tiling we have to translate the strip \mathbf{S} by
 31 a vector $v \in \mathbf{E}'$ such that the frontier of $v + \mathbf{S}$ contains no point of \mathbb{Z}^5 , and
 32 to define $\mathcal{P}_v = \pi(\mathbb{Z}^5 \cap (v + \mathbf{S}))$. The translated strip $v + \mathbf{S}$ corresponds to the
 33 translated window $v + W = \pi^\perp(v + (0, 1)^5)$. In this case $\pi'(\mathcal{E}_n \cap (v + (0, 1)^5)) =$
 34 $-nw + \mathcal{E}_n \cap (v + W) = v + W_n$, for any $n \in \{1, 2, 3, 4\}$.

35 **Theorem 4.1:** *If $(\lambda, \lambda', \lambda'') \in \tilde{\mathcal{L}}_0$ then there is an infinite number of inflation*
 36 *centers $y \in \mathbf{E}$ such that*

$$37 \mathbf{E} \longrightarrow \mathbf{E} : x \mapsto y + \lambda(x - y) \quad (17)$$

38 is an affine self-similarity of \mathcal{P}_v , that is, such that $x \in \mathcal{P}_v \implies y + \lambda(x - y) \in \mathcal{P}_v$.

Proof: If $(\lambda, \lambda', \lambda'') \in \tilde{\mathcal{L}}_0$ then there is a neighbourhood $\mathcal{U}_{(\lambda, \lambda', \lambda'')}$ of v in \mathbf{E}' such that the relation $(1 - \lambda')(u - v) + \lambda' \mathbf{W}_n \subset \mathbf{W}_{\lambda''_n}$ is verified for any $n \in \{1, 2, 3, 4\}$ and any $u \in \mathcal{U}_{(\lambda, \lambda', \lambda'')}$. Since the projection $\pi'(\mathbb{Z}^5 \cap \mathcal{E})$ is dense in \mathbf{E}' , the set $\mathcal{D}_{(\lambda, \lambda', \lambda'')} = \{t \in \mathbb{Z}^5 \cap \mathcal{E} \mid \pi't \in \mathcal{U}_{(\lambda, \lambda', \lambda'')} \}$ is an infinite set. For each $t \in \mathcal{D}_{(\lambda, \lambda', \lambda'')}$ the mapping $A_t : \mathbb{R}^5 \rightarrow \mathbb{R}^5$, $A_t x = t + (\lambda\pi + \lambda'\pi' + \lambda''\pi'')(x - t)$ satisfies the relations $\pi'(A_t x) = \pi't + \lambda'\pi'(x - t) = v + (1 - \lambda')(\pi't - v) + \lambda'(\pi'x - v)$ and

$$\left. \begin{array}{l} x \in \mathbb{Z}^5 \cap \mathcal{E}_n \\ \pi'x \in v + \mathbf{W}_n \end{array} \right\} \implies \left\{ \begin{array}{l} A_t x \in \mathbb{Z}^5 \cap \mathcal{E}_{\lambda''_n} \\ \pi'(A_t x) \in v + \mathbf{W}_{\lambda''_n} \end{array} \right. \quad (18)$$

This means that $\pi x \in \mathcal{P}_v \implies \pi(A_t x) \in \mathcal{P}_v$, that is, $\pi x \in \mathcal{P}_v \implies \pi t + \lambda(\pi x - \pi t) \in \mathcal{P}_v$.
□

5. Concluding remark

In the existing models, generally, the pentagonal/decagonal quasicrystals are considered to be two-dimensional, the third dimension being treated separately. Recently, Ben-Abraham *et. al.* [13] have shown that strip projection method offers a natural way to describe these quasicrystals as three-dimensional quasicrystals. It is sufficient to choose in the above decomposition \mathbf{E}^\perp as a physical space instead of \mathbf{E} . By a simple change of interpretation, our mathematical results concerning the Penrose type tilings become mathematical results concerning the model proposed by Ben-Abraham *et. al.* These new structures are invariant under certain transformations which act as inflation but with different constants in quasiperiodic layers and orthogonal direction.

Acknowledgment

The author is grateful to one of the referees for some very useful suggestions. This research was supported by the grant CEx05-D11-03.

References

- [1] R. Penrose, *Mathematical Intelligencer* **2** 32 (1979).
- [2] N. G. de Bruijn, *Kon. Nederl. Akad. Wetensch. Proc. A* **84** 38 (1981).
- [3] P. J. Steinhardt, H.-C. Jeong, K. Saitoh, M. Tanaka, E. Abe and A. P. Tsai, *Nature* **396** 55 (1998).
- [4] E. Abe, T. J. Sato and A. P. Tsai, *Phys. Rev. Lett.* **82** 5269 (1999).
- [5] C. Janot, *Quasicrystals: A Primer* (Oxford University Press, Oxford, 1992).
- [6] M. Senechal, *Quasicrystals and Geometry* (Cambridge University Press, Cambridge, 1995).
- [7] R. V. Moody and J. Patera, *Can. J. Phys.* **72** 442 (1994).
- [8] A. Katz and M. Duneau, *J. Phys. (France)* **47** 181 (1986).
- [9] N. Cotfas, *J. Phys. A: Math. Gen.* **31** 7273 (1998).
- [10] M. Baake and R. V. Moody, in *Aperiodic 97*, edited by M. de Boissieu, J.-L. Verger-Gaugry and R. Currat (World Scientific, Singapore, 1998) p. 9.
- [11] N. Cotfas, *J. Phys. A: Math. Gen.* **37** 3125 (2004).
- [12] Z. Masáková, J. Patera and E. Pelantová, *J. Phys. A: Math. Gen.* **31** 1443 (1998).
- [13] S. I. Ben-Abraham and A. Quandt, *Acta Cryst. A* **63** 177 (2007).