



**HAL**  
open science

## Novel swine influenza virus subtype H3N1 in italy

Ana Moreno, Ilaria Barbieri, Enrica Sozzi, Andrea Luppi, Davide Lelli,  
Guerino Lombardi, Maria Grazia Zanoni, Paolo Cordioli

► **To cite this version:**

Ana Moreno, Ilaria Barbieri, Enrica Sozzi, Andrea Luppi, Davide Lelli, et al.. Novel swine influenza virus subtype H3N1 in italy. *Veterinary Microbiology*, 2009, 138 (3-4), pp.361. 10.1016/j.vetmic.2009.04.007 . hal-00514608

**HAL Id: hal-00514608**

**<https://hal.science/hal-00514608>**

Submitted on 3 Sep 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript

Title: Novel swine influenza virus subtype H3N1 in italy

Authors: Ana Moreno, Ilaria Barbieri, Enrica Sozzi, Andrea Luppi, Davide Lelli, Guerino Lombardi, Maria Grazia Zanoni, Paolo Cordioli



PII: S0378-1135(09)00181-3  
DOI: doi:10.1016/j.vetmic.2009.04.007  
Reference: VETMIC 4408

To appear in: *VETMIC*

Received date: 9-12-2008  
Revised date: 18-3-2009  
Accepted date: 3-4-2009

Please cite this article as: Moreno, A., Barbieri, I., Sozzi, E., Luppi, A., Lelli, D., Lombardi, G., Zanoni, M.G., Cordioli, P., Novel swine influenza virus subtype H3N1 in italy, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2009.04.007

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 NOVEL SWINE INFLUENZA VIRUS SUBTYPE H3N1 IN ITALY:

2 Ana Moreno, Ilaria Barbieri, Enrica Sozzi, Andrea Luppi, Davide Lelli, Guerino Lombardi, Maria Grazia Zanoni, Paolo Cordioli

3 *Department of Animal Health, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna (IZSLER), Via Bianchi, 9,*

4 *25124-Brescia, Italy*

5

6 \* Corresponding author. Mailing address: Laboratory of Virology, Department of Animal Health, IZSLER, Via Bianchi, 9, 25124 – Brescia

7 (Italy). Phone: +39 030 2290361. Fax: +39 030 2290535. E-mail: [anamaria.morenomartin@bs.izs.it](mailto:anamaria.morenomartin@bs.izs.it)

8

Accepted Manuscript

9 Abstract

10 To date, three subtypes of swine influenza viruses, H1N1, H1N2, and H3N2 have been isolated in Italy. In 2006, a novel swine influenza  
11 virus subtype (H3N1) was isolated from coughing pigs. RT-PCR performed on lung tissues, experimental infection in pigs with the novel  
12 isolate, and cloning the virus by plaque assay confirmed this unique H and N combination. The novel isolate was also antigenically and  
13 genetically characterized. Genetic and phylogenetic analysis showed that the complete HA gene of the H3N1 strain has the highest  
14 nucleotide identity to three Italian H3N2 strains, one isolated in 2001 and two in 2004, whereas the full length NA sequence is closely  
15 related to three H1N1 subtype viruses isolated in Italy in 2004. The remaining genes are also closely related to respective genes found  
16 in H1N1 and H3N2 SIVs currently circulating in Italy. This suggests that the novel SIV could be a reassortant between the H3N2 and  
17 H1N1 SIVs circulating in Italy.

18

19 Keywords: Swine, novel subtype H3N1, genetic characterization, Italy

20

Accepted Manuscript

## 21 1- Introduction

22 Swine influenza is an economically important respiratory disease of swine caused by influenza A virus. Currently, three predominant  
23 subtypes (H1N1, H1N2, and H3N2) are prevalent in pigs throughout the world. However, subtype H1N7, H4N6, H9N2, and H3N3  
24 viruses have also been isolated sporadically from pigs without becoming established (Karasin et al, 2004; Lipatov et al, 2004; Yu et al,  
25 2008). In Europe over the last years, the epidemiology of swine influenza viruses has considerably changed. The H1N1 viruses now  
26 prevalent in European countries are antigenically distinct from the classical H1N1 strains and apparently came from the introduction of  
27 an avian virus in toto (Pensaert et al, 1981). These "avian-like" H1N1 viruses that emerged in European mainland pigs in 1979 replaced  
28 the previously circulating classical H1N1 strains. The H3N2 viruses present in Europe since 1984 are human-avian reassortants  
29 possessing six internal genes from the avian-like H1N1 viruses and HA and NA genes from the earlier isolated human-like H3N2 viruses  
30 (Castrucci et al, 1993). The third subtype (H1N2) was first isolated in Great Britain in 1994 (Brown et al, 1995) and subsequently spread  
31 to the swine population of continental Europe (Marozin et al, 2002). The H1N2 viruses originally resulted from multiple reassortment  
32 events initially involving human H1N1 and H3N2 viruses, followed by reassortment with avian-like swine viruses (Brown et al, 1998). In  
33 Italy, three subtypes (H1N1, H1N2, and H3N2) of swine influenza viruses have been reported in the pig population (Castrucci et al,  
34 1993; De Jong et al, 2007; Marozin et al, 2002). Phylogenetic analysis of Italian strains isolated during the period 1998-2007 were  
35 recently conducted (Moreno Martin et al. 2008 a, 2008b) and revealed that recent Italian H1N1 SIVs isolated in 2001-2007 were closely  
36 related to the A/Sw/IV/1455/99-like H1N1 swine viruses whereas Italian H1N1 strains isolated in the nineties clustered with the  
37 A/Sw/Fin/2899/82 SIV. Italian H1N2 strains were genetically similar to H1N2 viruses from the UK but were divided into two distinct  
38 clusters, one regarding strains isolated in 1998-2003 closely related to contemporary strains isolated in North Europe and the other  
39 related to recent Italian strains. Very interesting was the identification of three reassortant strains. One strain A/Sw/It/2064/99 H1N2 was  
40 closely related to H1N1 Italian SIVs (Marozin et al, 2002), the other strain A/Sw/It/5433/01 H1N1 showed 95-97% homology to recent  
41 Italian H1N2 SIVs, and the last reassortant A/Sw/It/11271/03 H1N2 showed a high similarity to the Italian H3N2 strains (Moreno Martin et  
42 al, 2008a). Regarding the subtype H3N2, De Jong et al (2007) reported that Italian strains isolated in the eighties and nineties displayed  
43 antigenic and genetic changes similar to those observed in Northern European viruses in the same period. Phylogenetic analysis  
44 showed that the Italian strains were located in the Eurasian virus lineage called A/Port Chalmers/1/73 (PCh lineage) where a gradual  
45 evolution of swine viruses starting from A/PCh/1/73-like human influenza virus is observed. This virus is considered the most probable  
46 ancestor of the swine PC73 lineage.

47 Recently the isolation of a novel swine influenza virus H3N1 was reported in Taiwan, USA and Korea (Lekcharoensuk et al, 2006; Ma et  
48 al, 2006; Shin et al, 2006) but it has never been isolated in Europe. This paper describes the isolation and characterization of a novel  
49 H3N1 swine influenza virus from pigs in Italy. Antigenic typing and genetic characterization suggested that the new isolate was a  
50 recombination between the H1N1 and H3N2 viruses circulating in Italy.

## 51 2- Materials and methods

### 52 2.1- Clinical samples

53 In March 2006, a severe respiratory disease, characterized by coughing and increased mortality, was observed in a feeder to finisher  
54 swine farm with 10000 head, located in the province of Cremona (North Italy). SIV vaccination had not been applied. Two dead pigs  
55 were submitted to the laboratory for respiratory disease diagnostic tests. Gross lesions consisted in purple areas of consolidation in the  
56 apical and cardiac lobes of the lungs, interlobular edema, mediastinal lymph nodes enlarged and pleuritis.

### 57 2.2- Virus isolation and subtype determination

58 The novel SIV isolate was obtained from a 10% lung homogenate applied onto Madin-Darby canine kidney (MDCK) cells and further  
59 inoculated through the allantoic sac route of 9-11 day old SPF chicken embryonated eggs (CEE). Culture supernatant (CS) after  
60 observed cytopathic effect and allantoic fluid (AF) were tested with an haemagglutination assay (HA) using chicken erythrocytes

61 performed as described (OIE Manual, 2005). The presence of influenza A was detected by using a double antibody sandwich ELISA  
62 (DAS-ELISA) with an anti-NPA Mab (ATCC n. HB65 H16-L10-4R5) carried out as previously described (Siebinga and de Boer, 1988).  
63 The subtype of the isolate 66945/06 was determined from culture supernatant, allantoic fluid and lung tissues by two multiplex RT-PCR  
64 assays (Chiapponi et al, 2003).

### 65 2.3- Anti-sera

66 For the antigenic characterization, hyper immune anti-sera collected 21 days after experimental infection of SPF chickens were used.  
67 The strains used to produce the anti-sera were the following: Reference strains A/Sw/Fin/2899/82 (2899/82) H1N1, A/Sw/CA/3633/84  
68 (3633/84) H3N2 and swine field strains A/Sw/It/1521/98 (1521/98), A/Sw/It/2064/99 (2064/99) H1N2, A/Sw/It/125746/05 (125746/05)  
69 H1N1 and A/Sw/It/79604/06 (79604/06) H3N2 isolated in Italy. The selection of the Italian field strains to produce chicken anti-sera was  
70 based on the genomic characteristics and phylogenetic analysis previously conducted, strains 1521/98, 2064/99 (Marozin et al, 2002)  
71 and 125746/05 and 79604/06 (Moreno Martin et al, 2008a, 2008b); The partial nucleotide sequence of HA gene of the last two strains is  
72 available at GenBank database (<http://www.ncbi.nlm.nih.gov/>).

### 73 2.4- Antigenic characterization

74 The SIV strain 66945/06 was antigenically characterized by HI, NI and virus neutralization (VN) as previously described (OIE manual,  
75 2005; Van Deusen et al, 1983; Van Reeth et al, 2003) using the anti-sera above described.

### 76 2.6- Genomic sequencing and phylogenetic analysis

77 Viral RNA was extracted from lung homogenate, AF and CS by using the Trizol reagent (Invitrogen, Carlsbad, CA, USA) as described  
78 by the manufacturer. Viral RNA isolation by QIAamp® ViralRNA Mini Kit (Qiagen, Hilden, Germany) and RT-PCR by OneStep RT-PCR  
79 Kit (Qiagen, Hilden, Germany) were performed either directly on clinical specimens (tissue homogenate and nasal swabs collected after  
80 the experimental infection) or AF and CS as described previously (Bragstad et al, 2005). Clinical specimens were screened for the  
81 presence of influenza A viruses by partial amplification of the M gene according to standard methods (Fouchier et al, 2000). Partial  
82 amplification of HA, NA, PB1, PB2, PA, NP, M and NS1 genes of the described strain and other circulating Italian SIVs was carried out  
83 under standard conditions using gene-specific primers (sequence available on request). On the basis of this partial characterization, the  
84 full length HA and NA genes of the novel strain and some Italian strains were amplified using universal primers as described (Hoffman  
85 et al, 2001). For sequencing, amplified products were separated onto agarose gel and then purified with Qiaquick® gel extraction kit  
86 (Qiagen, Inc, Valencia, CA, USA). Sequencing reactions were performed by means of BigDye® Terminator Cycle Sequencing kit v1.1  
87 (Applied Biosystems, Foster City, CA, USA). Both strands of the amplicons were sequenced using the same forward and reverse  
88 primers used for the amplification. Full length HA and NA amplicons were also sequenced with internal primers (Chiapponi et al, 2003).  
89 Sequences were resolved by using an ABI 3130 DNA automatic sequencer (Applied Biosystems, Foster City, CA, USA). DNA  
90 sequences were combined and edited using the Lasergene sequencing analysis software package (DNASTAR, Madison, WI). Multiple  
91 sequence alignments were made using ClustalW and maximum parsimony phylogenetic trees were created using MEGA4 (Tamura et  
92 al, 2007). Each tree is a consensus of 1,000 bootstrap replicates.

### 93 2.7- Plaque assay

94 In order to confirm the presence of the novel subtype H3N1, the novel SIV isolate was plaque cloned as previously described (Youil et  
95 al, 2004) and retested by HI and NI tests and again subtyped by RT-PCR.

### 96 2.8- Experimental swine infection

97 Two 8-week-old SPF pigs were infected intra-tracheally with  $10^{7.5}$  EID<sub>50</sub>/swine of the novel strain prepared in CEE. Nasal swabs were  
98 collected 2, 5, 7, 9 and 12 days after infection and analyzed by RT-PCR using gene M specific primers to evaluate virus shedding. RT-  
99 PCR positive samples were then inoculated into a monolayer of MDCK cells and SPF CEE. Serum samples were collected 15, 27, and

100 35 days after infection and were tested for antibodies anti-NPA, and against H1 and H3 subtypes. A Mab-based (Mab HB65)  
 101 competitive ELISA was performed to detect anti-NPA antibodies as described (De Boer et al, 1990). Antibodies against H1 and H3 were  
 102 detected by HI test performed as described (OIE Manual, 2005), using the SIVs 66945 H3N1, 79604/06 H3N2, and 125746/05 H1N1,  
 103 1513/98 H1N1, 1521/98 H1N2 (Marozin et al, 2002) as antigen.

#### 104 2.9- Detecting swine respiratory pathogens

105 The lung homogenate was cultured using different media for the most common swine respiratory bacterial pathogens. The presence of  
 106 Porcine reproductive and respiratory syndrome virus (PRRSV), Porcine circoviruses type 2 (PCV2) and *Mycoplasma hyopneumoniae*  
 107 was determined by using either RT-PCR, multiplex PCR and PCR assays respectively, as previously described (Calsamiglia et al,  
 108 1999; Ouardani et al, 1999; Suárez et al, 1994).

#### 109 2.10- Nucleotide sequence accession numbers

110 GenBank accession nos. of the complete HA genes are: [EU037014](#), [EU045362](#), [EU045365](#), [EU982298](#), [EU045371](#), [EU982299](#),  
 111 [EU982297](#), [EU982296](#), [EU982295](#). GenBank accession nos. of the complete NA genes are: [EU037015](#), [EU045388](#), [EU045389](#),  
 112 [EU045393](#). GenBank accession nos. of partial nucleotide sequences of internal genes are: NP gene: [EU091685](#), [EU091690](#),  
 113 [EU091695](#). PB1 gene: [EU091687](#), [EU091692](#), [EU091697](#). PB2 gene: [EU091688](#), [EU091693](#), [EU091698](#). PA gene: [EU091686](#),  
 114 [EU091691](#), [EU091696](#). M gene: [EU091684](#), [EU091689](#), [EU091694](#). NS gene: [EU091704](#), [EU091705](#), [EU091707](#).

### 115 3- Results

#### 116 3.1- Virus isolation and subtype determination

117 Allantoic fluid collected 5 days post incubation of CEE and MDCK CS agglutinated chicken erythrocytes in HA (HA titer of 7 and 6  
 118  $\log_2/25\mu\text{l}$  respectively). The presence of influenza A antigen was detected by sandwich ELISA using the Mab anti-NPA HB65 in both AF  
 119 and MDCK CS. A multiplex RT-PCR specific for HA and NA was further used to subtype both AF and CS that unexpectedly resulted to  
 120 be H3N1. Neither H1 nor N2 specific bands were observed. The presence of H3N1 virus in the lung tissues was also confirmed by RT-  
 121 PCR and was the only H-N combination detected in the sample. The isolate was named A/Sw/It/66945/06 (66945/06). Respiratory  
 122 pathogens such as bacteria, *M. hyopneumoniae* or PRRSV that could produce lesions similar to SIV were not observed whereas PCV2  
 123 was detected by PCR. The virus resulting from the plaque assay and cloning by limited dilution was identified as subtype H3N1 by RT-  
 124 PCR and HI and NI tests.

#### 125 3.2- Antigenic characterization

126 The HI and VN tests were conducted using a panel of six sera from SPF chickens experimentally infected with two early reference SIV  
 127 strains (2899/82 H1N1 and 3633/84 H3N2), two well known SIV strains H1N2 (Marozin et al, 2002) isolated in Italy at the end of the  
 128 nineties (1521/98 closely related to the HA of the European H1N2 viruses and 2064/99 closely related to the HA of the European H1N1  
 129 viruses) and, finally, with two recent Italian strains, 125746/05 H1N1 and 79604/06 H3N2, that exhibited different HA genetic  
 130 characteristics compared to early European isolates (Moreno Martin et al, 2008a, 2008b). The same panel of sera was used to  
 131 determine the antigenic reactivity of the neuraminidase in NI tests. The HI tests showed that the strain 66945 reacted with post-infection  
 132 chicken antisera to H3N2 SIVs at high titers (8  $\log_2$  using antiserum CA/3633/84 and 9  $\log_2$  using antiserum It/79604/06) and did not  
 133 react with chicken antisera towards H1N2. Cross reactions at low titers were observed against H1N1 SIVs 2899/82 (4  $\log_2$ ) and  
 134 125746/05 (5  $\log_2$ ), probably due to the interference in HI with N1 antibodies. The novel strain H3N1 reacted only with chicken anti-sera  
 135 to CA/3633/84 H3N2 (SIVs using VN, whereas no reactions were observed with H1N2 and H1N1 anti-sera. In NI assay, N activity of the  
 136 novel strain was inhibited only by antisera produced towards H1N1 SIVs. Results of the HI and NI tests confirmed the presence of a  
 137 novel subtype H3N1. Results are shown in table n.1.

#### 138 3.4- Experimental swine infection

139 Two SPF pigs were infected intra-tracheally with the isolate H3N1 and were reared in isolation for 35 days. No respiratory distress was  
 140 reported during the observation period except for an increase in body temperature up to 40,5°C for 2-3 days. Virus shedding was  
 141 demonstrated testing nasal swabs by RT-PCR only 3 and 5 days PI in the two experimentally infected pigs, and these samples were  
 142 then inoculated in SPF CEE and into MDCK cells. Infected allantoic fluids and cell culture supernatants were characterized by HI and NI  
 143 and also by RT-PCR confirming that the subtype H3N1 was the only one to be detected. Serum samples were tested for antibodies  
 144 against NPA and HA of subtypes 1 and 3. Both antibodies against NPA and H3 were detected in all the serum samples collected 15, 27  
 145 and 35 days after infection whereas no specific antibodies towards H1 were evidenced. HI titers, expressed as  $\log_2$ , were: for pig n.1:7,  
 146 6, and 5  $\log_2$  against 66945/06 H3N1 and 7, 6 and 6  $\log_2$  against 79604/06 H3N2 for sera collected 15, 27 and 35 days PI respectively;  
 147 for pig n.2: 5, 6, and 6  $\log_2$  against 66945/06 H3N1 and 5, 6 and 6  $\log_2$  against 79604/06 H3N2 for sera collected 15, 27 and 35 days  
 148 PI. HI titer against SIV of subtypes H1N1 and H1N2 resulted to be always  $<3 \log_2$ .

### 149 3.5- Genomic sequence and phylogenetic analysis

150 Initially the partial nucleotide sequences of the H3N1 isolate genes were compared to those of the SIVs isolated in Italy in the last 10  
 151 years. Phylogenetic analysis indicated that the HA gene of strain H3N1 was closely related to three Italian H3N2 strains, one isolated in  
 152 2001, A/Sw/It/7680/01 (7680/01) and two in 2004, A/Sw/It/297576/04 (297576), A/Sw/It/302219-3/04 (302219-3/04). The NA, NP, PB1,  
 153 PB2, PA, and M genes showed the highest % of homology to three Italian H1N1 strains isolated in 2004, A/Sw/It/53949/04 (53949/04),  
 154 A/Sw/It/65296/04 (65296/04), A/Sw/It/247578/04 (247578/04). The remaining NS gene was closely related to other three SIVs currently  
 155 circulating in Italy (A/Sw/It/1484/02 H3N2, A/Sw/It/4230/02 H1N1, A/Sw/It/207828/02 H1N1).

156 For a further analysis, BLAST search in the GenBank database demonstrated that the full length HA and NA genes exhibited 96% and  
 157 98% nucleotide similarity to those of the A/Sw/Sp/42386/2002 H3N2 and A/Sw/Sp/53207/04 H1N1 respectively. The partial internal  
 158 genes M, NP, PB1, PB2, and PA showed 98% nucleotide identity with the same Spanish SIVs (53207/04) while the NS gene showed  
 159 the highest identity (97%) with isolate A/Sw/It/1081/00 H1N2.

160 The comparison of the full HA and NA genomic sequences of the isolate 66945/06 with some Italian SIVs confirmed the close  
 161 relationship observed when comparing the partial HA and NA nucleotide sequence between the novel strain and the six Italian SIVs  
 162 described above (99,0-98,1% at HA and 98,1-98,3 % at NA nucleotide level respectively).

163 The HA and NA sequences were then aligned to those of other SIVs retrieved from GenBank. Since the highest number of sequences  
 164 available in GenBank resulted to be mostly partial, the phylogeny reconstruct was limited to only the HA1 region of the H3 SIVs. The  
 165 resultant maximum parsimony phylogenetic tree indicated that the HA1 region of the HA gene of the novel strain and other recent Italian  
 166 H3N2 isolates were placed into the antigenic cluster 2 belonging to the PC73-like lineage (De Jong et al, 2007) together with  
 167 contemporary H3N2 SIVs isolated in South Europe. The H3N1 66945 resulted to belong to a different cluster that the other H3N1 strains  
 168 isolated in USA and Korea that were closely related (figure n.1). The NA gene of the strain 66945 appeared in the phylogenetic tree  
 169 (figure n. 2) closely related to Italian and Spanish H1N1 SIVs isolated in the last years but placed in a cluster different from other  
 170 contemporary H1N1 Italian SIVs related to the Sw/IV/1455/99-like viruses (Marozin et al, 2002).

171 The comparison of the deduced amino acid sequence of the HA gene of the isolate 66945 with that of other H3N2 and H3N1 SIVs  
 172 indicates that most of the residues are highly conserved, especially those associated with the sialoside receptor-binding region, Y98,  
 173 W153, H183, E190, and L194 (Kaverin et al, 2004), and those responsible for host range specificity at positions 226 and 228 (Vines et  
 174 al, 1998). The strain 66945 H3N1 and the recent Italian isolates H3N2 exhibited the same aa residues, Y98, W153, H183, E190, L194,  
 175 L226, and S228, which L226 and S228 are mainly responsible for sialyl  $\alpha$ 2,6-galactose (SA $\alpha$ 2,6Gal) specificity. Also the amino acids  
 176 within the antigenic sites of the HA gene of the Italian H3 SIVs, H3N1 isolate included, resulted to be highly conserved. In addition the  
 177 novel strain and the related Italian and Spanish H3N2 strains showed the aa residues S137, N145 and N278 characteristic of the  
 178 antigenic cluster 2.

## 179 4- Discussion

180 The first isolation and characterization of the novel subtype H3N1 in Europe was described. One strain was isolated in 2006 from a  
181 feeder to finisher pig farm located in North Italy where specific respiratory symptoms were observed. The antigenic and genetic  
182 characterization confirmed the isolation of a novel H3N1 SIV, never described in Europe previously. The genetic characterization of all  
183 genes of the new isolate revealed that it was closely related to the circulating Italian SIV strains. The complete HA gene presented the  
184 highest identity to three Italian H3N2 SIVs isolated in 2001 and 2004, whereas the NA gene was instead closely related to three Italian  
185 H1N1 SIVs isolated in 2004. The deduced aa sequence of the HA gene showed the aa residues (L226 and S228) responsible for the  
186 receptor specificity typical of swine and human influenza viruses. In contrast, the residues involved in the SA $\alpha$ 2,3Gal linkage, typical of  
187 avian viruses, are Q226 and G228. The partial sequencing of internal genes NP, PB1, PB2, PA and M genes showed the high identity to  
188 the same H1N1 Italian strains isolated in 2004. Finally the NS gene exhibited the highest identity to one H3N2 and two H1N1 Italian  
189 SIVs isolated in 2002. This suggested that the novel SIV was a reassortant between the H3N2 and H1N1 SIVs circulating in Italy. The  
190 experimental infection of two SPF pigs was performed in order to further confirm the presence of the novel subtype. The H3N1 SIV  
191 could replicate in the respiratory tract and was shed in nasal secretions although only for a few days. Both virus isolation and typing  
192 through serological tests demonstrated the presence of the unique H3N1 subtype.

193 Monitoring programs performed in the last two years on pig farms with respiratory symptoms brought to the isolation of several strains of  
194 the three circulating subtypes but not of subtype H3N1. It is worth notice that the subtype H3N1 has been rarely observed throughout  
195 the world whereas the isolation of H1N2 reassortant viruses results frequently. Mitnaul et al (2000) revealed that HA and NA proteins  
196 recognized the same molecule (sialic acid) with conflicting activities and a balance of protein activities were essential to ensure an  
197 efficient replication of the virus. Drastic changes in this equilibrium could affect viral replication. Even if the reason of the low frequency  
198 of H3N1 SIV isolation can not be explained, it could be hypothesized that the activities of H3 and N1 proteins are not optimally balanced  
199 compared to the H1 and N2 proteins. This novel subtype does not appear to be so wide spread nor successfully maintained in the  
200 Italian swine population. On the contrary, reassortment strains among swine viruses of H3N2, H1N1 and H1N2 were frequently  
201 observed in Europe (Marozin et al, 2002; Moreno Martin et al., 2008a, Zell et al, 2008). Swine are also considered as a vessel for  
202 reassortment of human and avian viruses because they possess both SA $\alpha$ 2,6Gal and SA $\alpha$ 2,3Gal receptors characteristic of human and  
203 avian viruses respectively (Ito et al, 1998; Lekcharoensuk et al, 2006; Ma et al, 2006). Indeed there is good evidence that pigs are more  
204 frequently involved in interspecies transmission of influenza A viruses than are other animals (Pensaert et al 1981; Karasin et al 2004,  
205 Yu et al 2008, Zhou et al, 1999). All these findings highlight the need for a stringent surveillance in pig populations in order to better  
206 understand the circulation of atypical swine influenza viruses.

207

## 208 Acknowledgements

209 We would like to thank Dr. Leonardo James Vinco and Manola Adella for excellent technical assistance.

210

- 211 References
- 212 Avian Influenza. May 2005, posting date. Chapter 2.7.14, OIE Manual of diagnostic tests and vaccines for terrestrial animals (mammals,
- 213 birds and bees), Fifth Ed. 2004. [http://www.oie.int/eng/normes/mmanual/A\\_summry.htm](http://www.oie.int/eng/normes/mmanual/A_summry.htm).
- 214 Bragstad, K., Jørgensen, P. H., Handberg, K. J., Møllergaard, S., Corbet, S., Fomsgaard, A., 2005. New avian influenza A virus subtype
- 215 combination H5N7 identified in Danish mallard ducks. *Virus Res.* 109, 181-190.
- 216 Brown, I. H., Harris, P.A., McCauley, J. W., Alexander, D. J., 1998. Multiple genetic reassortment of avian and human influenza A
- 217 viruses in European pigs, resulting in the emergence of an H1N2 virus of novel genotype. *J Gen Virol.* 79, 2947- 2955.
- 218 Brown, I. H., Chakraverty, P., Harris, P. A., Alexander, D. J., 1995. Disease outbreaks in pigs in Great Britain due to an influenza A virus
- 219 of H1N2 subtype. *Vet Rec.* 136, 328-329.
- 220 Calsamiglia, M., Pijoan, C., Trigo, A., 1999. Application of a nested polymerase chain reaction assay to detect *Mycoplasma*
- 221 *hyopneumoniae* from nasal swabs. *J Vet Diagn Invest.* 11, 246-251.
- 222 Castrucci, M. R., Donatelli, I., Sidoli, L., Barigazzi, G., Kawaoka, Y., Webster, R. G., 1993. Genetic reassortant between avian and
- 223 human influenza A viruses in Italian pigs. *Virology.* 232, 310-318.
- 224 Chiapponi, C., Fallacara, F., Foni, E., 2003. Subtyping of H1N1, H1N2 and H3N2 swine influenza viruses by two multiplex RT-PCR, p.
- 225 257-258. In Paolo Martelli, Sandro Cavirani, Antonio Lavazza (ed.), PRRS PMWS swine influenza. Proceedings of the 4th International
- 226 Symposium on emerging and re-emerging pig disease: Rome, Italy. <http://www.unipr.it/arpa/facvet/dip/dipsa/ric/prrs2003/258-259.pdf>.
- 227 De Boer, G.E., Back, W., Osterhaus, A.D.M.E., 1990. An ELISA for detection of antibodies against influenza A nucleoprotein in humans
- 228 and various animal species. *Arch Virol.* 115, 47-61.
- 229 De Jong, J.C., Smith, D. J., Lapedes, A. S., Donatelli, I., Campitelli, L., Barigazzi, G., Van Reeth, K., Jones, T.C., Rimmelzwaan, G. F.,
- 230 Osterhaus, A. D. M. E., Fouchier, R.A.M., 2007. Antigenic and genetic evolution of swine influenza A (H3N2) viruses in Europe. *J Virol.*
- 231 81(8), 4315-4322.
- 232 Fouchier, R. A. M., Bestebroer, T.M., Herfst, S., van der Kemp, L., Rimmelzwaan, G. F., Osterhaus, A. D. M. E., 2000. Detection of
- 233 influenza A viruses from different species by PCR amplification of conserved sequences in the Matrix gene. *J Clin Microbiol.* 38,
- 234 4096:4101.
- 235 Hoffman, E., Stech, J., Guan, Y., Webster, R. G., Perez, D. R., 2001. Universal primer set for the full-length amplification of all influenza
- 236 A viruses. *Arch Virol.* 146, 2275-2289.
- 237 Ito T., Couceiro J.N.S.S., Kelm S., Baum L.G., Krauss S., Castrucci M.R., Donatelli I., Kida H., Paulson J.C., Webster R.G., Kawaoka
- 238 Y., 1998. Molecular basis for the generation in pigs of Influenza A viruses with pandemic potential. *J. Virol.* 72, 7367-7373.
- 239 Karasin A. I., West, K., Carman, S., Olsen, C. W., 2004. Characterization of avian H3N3 and H1N1 influenza A viruses isolated from pigs
- 240 in Canada. *J Clin Microbiol.* 42, 4349-4354.
- 241 Kaverin N. V., Rudneva, I. A., Ilyushina, N. A., Lipatov, A. S., Krauss, S., Webster, R. G., 2004. Structural differences among
- 242 haemagglutinins of influenza A virus subtypes are reflected in their antigenic architecture: analysis of H9 escape mutants. *J Virol.*
- 243 78,240-249.
- 244 Lekcharoensuk, P., Lagger, K. M., Vemulapalli, R., Woodruff, M., Vincent, A. L., Richt, J. A., 2006. Novel swine influenza virus subtype
- 245 H3N1, United States. *Emerg Inf Dis.* 12(5), 787-794.
- 246 Lipatov, A. S., Govorkova, E. A., Webby, R. J., Ozaki, H., Peiris, M., Guan, Y., Poon, L., Webster, R. G., 2004. Influenza: emergence
- 247 and control. *J Virol.* 78, 8951-8959.
- 248 Ma, W., Gramer, M., Rossow, K., Yoon, K. J., 2006. Isolation and genetic characterization of new reassortant H3N1 swine influenza
- 249 virus from pigs in the Midwestern United States. *J Virol.* 80 (10), 5092:5096.

- 250 Marozin, S., Gregory, V., Cameron, K., Bennett, M., Valette, M., Aymard, M., Foni, E., Barigazzi, G., Lin, Y., Hay, A., 2002. Antigenic  
 251 and genetic diversity among swine influenza A H1N1 and H1N2 viruses in Europe. *J Gen Virol.* 83, 735-745.
- 252 Mitnaul, L.J., Matrosovich, M. N., Castrucci, M. R., Tuzikov, A. B., Bovin, N. V., Kobasa, D., Kawaoka, Y., 2000. Balanced hemagglutinin  
 253 and neuraminidase activities are critical for efficient replication of influenza A virus. *J Virol.* 74, 6015-6020.
- 254 Moreno Martin, A., Barbieri, I., Chiapponi, C., Foni, E., Sozzi, E., Canelli, E., Luppi, A., Cordioli, P., 2008a. Genetic characterization of  
 255 H1N1 and H1N2 swine influenza viruses isolated in Italy in 1998-2007, abstr. p 99. Second annual meeting EPIZONE (Network of  
 256 Excellence for Epizootic Disease Diagnosis and Control) "Need for Speed", 4-6 June 2008, Brescia(Italy), [http: www.epizone.net](http://www.epizone.net).
- 257 Moreno Martin, A., Barbieri, I., Sozzi, E., Luppi, A., Canelli, E., Foni, E., Alborali, L., Cordioli, P., 2008b. Genetic characterization of  
 258 swine influenza H3N2 subtype in Italy, abstr. P 1.025, p 25. Proceedings of 20<sup>th</sup> International Pig Veterinary Society Congress, 22-26  
 259 June, Durban, South Africa.
- 260 Ouardani, M., Wilson, L., Jetté, R., Montpetit, C., Dea, S., 1999. Multiplex PCR for detection and typing of porcine circoviruses. *J Clin*  
 261 *Microbiol.* 37(12), 3917-3924.
- 262 Pensaert, M., Ottis, K., Vandeputte, J., Kaplan, M. M., Bachmann, P.A., 1981. Evidence for the natural transmission of influenza A virus  
 263 from wild ducks to swine and its potential for man. *Bull World Health Organ.* 59, 75-78.
- 264 Shin, J. Y., M. S. Song, E. H. Lee, Y. M. Lee, S. Y. Kim, H. K. Kim, J. K. Choi, C.J. Kim, R. J. Webby, and Y. K. Choi. 2006. Isolation  
 265 and characterization of novel H3N1 swine influenza viruses from pigs with respiratory diseases in Korea. *J Clin Microbiol.* 44, 3923-  
 266 3927.
- 267 Siebinga, J.T., and G. F. de Boer. 1988. Influenza A viral nucleoprotein detection in isolates from human and various animal species.  
 268 *Arch Virol* 100, 75-87.
- 269 Suárez, P., R. Zardoya, C. Prieto, A. Solana, E. Tabarés, J.M. Bautista, and J.M. Castro. 1994. Direct detection of the porcine  
 270 reproductive and respiratory syndrome (PRRS) virus by reverse polymerase chain reaction (RT-PCR). *Arch Virol* 135, 89-99.
- 271 Tamura, K, Dudley, J., Nei, M., Kumar, S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol*  
 272 *Biol Evol* 24, 1596-1599.
- 273 Van Deusen, R. A., V. S. Hinshaw, D. A. Senne, and D. Pellacani. 1983. Micro neuraminidase-inhibition assays for classification of  
 274 influenza A virus neuraminidases. *Av Dis.* 27, 745-750.
- 275 Van Reeth, K., Gregory, V., Hay, A., Pensaert, M. 2003. Protection against a European H1N2 swine influenza virus in pigs previously  
 276 infected with H1N1 and/or H3N2 subtypes. *Vaccine.* 21, 1375-1381.
- 277 Vines, A., K. Wells, M. Matrosovich, M. R. Castrucci, T. Ito, Y. Kawaoka. 1998. The role of influenza A virus hemagglutinin residues 226  
 278 and 228 in receptor specificity and host range restriction. *J Virol.* 72, 7626:7631.
- 279 Youil, R., Q. Su, T. J. Toner, C. Szymkowiak, W-S. Kwan, B. Rubin, L. Petrukhin, I. Kiseleva, A.R. Shaw, and D. Di Stefano. 2004.  
 280 Comparative study of influenza virus replication in Vero and MDCK cell lines. *J Virol Meth.* 120, 23-31.
- 281 Yu H., Hua R-H., Wei T-C., Zhou Y-J., Tian Z-J., Li G-X., Liu T-Q., Tong G-Z., 2008. Isolation and genetic characterization of avian  
 282 origin H9N2 influenza viruses from pigs in China. *Vet Microbiol.*, 131, 82-92.
- 283 Zell R., Motzke S., Krumbholz A., Wutzler P., Herwig V., Dürwald R., 2008. Novel reassortant of swine influenza H1N2 virus in  
 284 Germany. *J Gen Virol* 89, 271-276.
- 285 Zhou N.N., Senne D.A., Landgraf J.S., Swenson S.L., Erickson G., Rossow K., Liu L., Yoon K-J., Krauss S., Webster R.G., 1999.  
 286 Genetic reassortment of avian, swine and human influenza A viruses in American pigs. *J. Virol.* 73, 8851-8856.
- 287

288

	<b>Antigen</b>	<b>It/66945/06</b>		
	<b>subtype</b>	<b>H3N1</b>		
	test	VN	HI	NI
	Fin/2899/82 H1N1	<10	16	+
	It/125746/05 H1N1	<10	32	+
<b>Antiserum</b>	It/1521/98 H1N2	<10	<2	-
	It/2064/99 H1N2	<10	<2	-
	CA/3633/84 H3N2	640	256	-
	It/79604/06 H3N2	1280	512	-

289  
290  
291

Table n.1- Results of the antigenic characterization of the novel strain performed by serological tests. Virus neutralization titres are expressed as ND<sub>50</sub> reciprocal. Using HI test results as HI titre reciprocal.

292 Figure n. 1. Phylogenetic relationship of the HA1 region of HA gene from the newly identified H3N1 swine influenza virus  
293 (A/swine/Italy/66945/2006(H3N1)) with other Italian and database collected H3N2 and H3N1 SIV strains. The tree was created by  
294 Maximum Parsimony method and bootstrapped with 1,000 replicate. Only bootstrap values higher than 70% are shown.  
295

296 Figure n. 2. Phylogenetic relationship of the complete NA gene from the newly identified H3N1 swine influenza virus  
297 (A/swine/Italy/66945/2006(H3N1)) to NA gene of other Italian and database collected H1N1 SIV strains. Tree was generated as  
298 described in the legend to Fig.n.1.

Accepted Manuscript

Figure n.1



