



HAL
open science

Multi-particle sampling in Monte Carlo simulations on fluids: Efficiency and extended implementations

Ivo Nezbeda

► **To cite this version:**

Ivo Nezbeda. Multi-particle sampling in Monte Carlo simulations on fluids: Efficiency and extended implementations. *Molecular Simulation*, 2009, 35 (08), pp.660-672. 10.1080/08927020902725572 . hal-00515075

HAL Id: hal-00515075

<https://hal.science/hal-00515075>

Submitted on 4 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Multi-particle sampling in Monte Carlo simulations on fluids:
Efficiency and extended implementations**

Journal:	<i>Molecular Simulation</i> / <i>Journal of Experimental Nanoscience</i>
Manuscript ID:	GMOS-2008-0189.R1
Journal:	Molecular Simulation
Date Submitted by the Author:	15-Dec-2008
Complete List of Authors:	Nezbeda, Ivo; ICPF, Acad. Sci., E. Hala Lab.; J. E. Purkinje Univ., Faculty of Science
Keywords:	Multi-particle move MC, Polarizable models of water, Reaction field, Parallelization



1
2
3
4
5
6
7
8
9
10
11
12
13 **Multi-particle sampling in Monte Carlo simulations on fluids:**
14 **Efficiency and extended implementations**
15
16

17
18 Filip Moučka^a and Ivo Nezbeda^{a,b,1}
19

20 ^aFaculty of Science, J. E. Purkinje University, 400 96 Ústí nad Labem, Czech Republic
21

22 ^bE. Hala Laboratory of Thermodynamics, Institute of Chemical Process Fundamentals,
23 Academy of Sciences, 165 02 Prague 6, Czech Republic
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

59 ¹Author for correspondence. E-mail: IvoNez@icpf.cas.cz
60

Abstract

Various technical aspects affecting efficiency of a recently proposed novel Monte Carlo simulation scheme based on biased simultaneous displacements/rotations of *all* particles of the system are investigated using two polarizable models of water, the Chialvo-Cummings and Brodholt-Sampoli-Vallauri models, as a test case. Necessary expressions for polarizable site-site interaction models are derived along with a novel smoothing of the potential at the cutoff distance. In addition to the common thermodynamic and structural properties, the mean squared displacements, rotation relaxation, speed of equilibration (translational order parameter), and autocorrelation coefficients have been computed as well in order to assess efficiency of the method. Gain in speed by parallelization has also been examined.

Performance of the method is compared with both the standard one-particle move method and available approximate methods. It is shown that the multi-particle move method performs about by a factor of 10 faster for the systems considered when compared with the common Monte Carlo scheme, and several times faster when compared with approximate methods. Parallelized codes of the multi-particle move method may then perform about seventy times faster than the conventional Monte Carlo. These conclusions hold true for the system size simulated ($N=256$) because the efficiency of the multi-particle method depends on the size of the system: its efficiency even increases with increasing number of particles.

Keywords: Multi-particle move MC; Parallelization; Reaction field; Polarizable models of water

1. Introduction

In general, when studying molecular systems with a pairwise interaction potential by computer simulations, the choice of the method, Monte Carlo (MC) or molecular dynamics (MD), is in most cases only a matter of convenience and/or personal preference [1, 2]. The preferred use of either method is dictated either by the studied system, e.g., models with a discontinuous potential favor the MC method, or the properties to be evaluated, e.g., kinetic properties are naturally obtainable from MD simulations. This sort of equivalence between the two simulation methods is however immediately lost if the pairwise additivity does not apply. An example are polarizable models. In this case the interaction between *all* N particles comprising the system must be evaluated in every single timestep (configuration). Since in MD simulations all particles change their positions in every timestep anyhow, this need not bring about any dramatic increase of the CPU time, but at least by a factor 2-3 in dependence on the method used to reach self-consistency of induced dipoles [3, 4]. On the other hand, in common MC simulations only the change of energy due to the change of the position or orientation of the particle attempting a move is required. Thus, for polarizable models, additional computations to determine the total interaction energy make the MC method by the factor $\mathcal{O}(N)$ slower in comparison with MC simulations of nonpolarizable models. Somewhere in between these methods there are hybrid MD/MC methods; for a discussion of their advantages and disadvantages see, e.g., [1].

To overcome the inefficiency of MC simulations of polarizable models (and, in general, for models with non-additive interactions) we have recently proposed a novel MC method [5]. The method, called multi-particle move (MPM) MC scheme, considers simultaneous translations/rotations of all particles at once in the spirit of molecular dynamics. It means that the moves are biased according to the forces acting between the molecules. To demonstrate feasibility and efficiency of the method, in [5] we considered the polar-

1
2
3
4
5
6
7 izable Stockmayer fluid. Since, however, its implementation for more complex many-site
8
9 interaction models need not be straightforward, we consider in this paper the most com-
10
11 mon class of polarizable models of water and derive and present all necessary expressions.
12
13 Furthermore, it turns out that the efficiency of the method is also rather sensitive to the
14
15 treatment of the long-range Coulombic interactions; more accurately, to their approxima-
16
17 tion in vicinity of the cutoff distance, and special attention has therefore been paid to this
18
19 problem. A novel smoothing of the potential at the cutoff distance is proposed and used.
20
21 Finally, technical aspects affecting the efficiency of simulations, as e.g., acceptance ratio,
22
23 sensitivity of the correlation length to the type of move, etc., are addressed as well along
24
25 with a potential gain in speed by (simple) parallelization. For comparison with currently
26
27 available other MC methods we consider both the standard one-particle MC method and
28
29 more efficient, but only approximate, PAPI [3, 4] and ANES [6] methods. Although both
30
31 latter methods have been shown to yield, if implemented with careful choice of simulation
32
33 parameters, results identical to those calculated rigorously through matrix minimization
34
35 [3, 4, 7], both are suitable for polarizable models only and cannot be used for simulations
36
37 considering other kinds of non-additive interactions.

38
39 After presenting the method in the next Section, and deriving all necessary expressions
40
41 and providing all technical details for its implementation in Section 3, we then consider
42
43 the Chialvo-Cummings [8] and Brodholt-Sampoli-Vallauri [9] polarizable models of water
44
45 and present and discuss results in Section 4.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. The biased multi-particle move method

To move all particles at once in MC method without any bias would be very inefficient. The MPM method starts from the known force-bias method [10], in which trial moves are biased in the direction of forces or torques acting on the molecules, and extends it to a simultaneous displacements (or rotations) of all particles. Thus, if \mathbf{F}_k is the force acting on particle k to be displaced and t_m is the maximum length of any component of the translational vector, \mathbf{t}_k , then its components t_k^α , $\alpha = 1, 2, 3$, are generated according to the probability distribution

$$\pi(t_k^\alpha) = \frac{\exp[\lambda\beta F_k^\alpha t_k^\alpha]}{\int_{-t_m}^{t_m} \exp[\lambda\beta F_k^\alpha t_k^\alpha] dt_k^\alpha} = \frac{\exp[\lambda\beta F_k^\alpha t_k^\alpha]}{2 \sinh[\lambda\beta F_k^\alpha t_m] / \lambda\beta F_k^\alpha}, \quad (1)$$

where $k = 1 \dots N$, and λ is a parameter whose value lies between $\frac{1}{2}$ and 1 [10]; we use $\lambda = \frac{1}{2}$. The acceptance probability of this move is

$$\text{Prob} = \min \left\{ 1, \exp[-\beta\Delta U] \frac{\omega^{new \rightarrow old}}{\omega^{old \rightarrow new}} \right\}, \quad (2)$$

where

$$\omega^{old \rightarrow new} = \prod_{\alpha=1}^3 \frac{\lambda\beta F_k^{\alpha,old} \exp[\lambda\beta F_k^{\alpha,old} t_k^\alpha]}{2 \sinh[\lambda\beta F_k^{\alpha,old} t_m]} \quad (3)$$

and

$$\omega^{new \rightarrow old} = \prod_{\alpha=1}^3 \frac{\lambda\beta F_k^{\alpha,new} \exp[\lambda\beta F_k^{\alpha,new} (-t_k^\alpha)]}{2 \sinh[\lambda\beta F_k^{\alpha,new} t_m]}, \quad (4)$$

where ΔU is the associated change in the internal energy, and $\beta = 1/k_B T$ where k_B is the Boltzmann constant and T is the absolute temperature.

The above expressions, written for the conventional one-particle translational move, are easily extended to multi-particle moves (for details see the original paper [5]):

$$\omega^{old \rightarrow new} = \prod_{\alpha=1}^3 \prod_{k=1}^N \omega_k^{\alpha,old \rightarrow new}, \quad (5)$$

where again

$$\omega_k^{\alpha,old \rightarrow new} = \frac{\lambda\beta F_k^{\alpha,old} \exp[\lambda\beta F_k^{\alpha,old} t_k^\alpha]}{2 \sinh[\lambda\beta F_k^{\alpha,old} t_m]}, \quad (6)$$

and

$$\omega^{new \rightarrow old} = \prod_{\alpha=1}^3 \prod_{k=1}^N \omega_k^{\alpha, new \rightarrow old}, \quad (7)$$

A formula for generating the translation vector is obtained by inverting Eqn. (1),

$$t_k^\alpha = \frac{\ln[\exp(-\lambda\beta F_k^{\alpha, old} t_m) + 2u_k^\alpha \sinh(\lambda\beta F_k^{\alpha, old} t_m)]}{\lambda\beta F_k^{\alpha, old}}, \quad (8)$$

where u_k^α is a random number within the interval $(0, 1)$.

A similar modification applies to the rotational move which can be implemented in a number of different ways. In this work particle k attempts the rotation around a randomly chosen vector \mathbf{r}_k whose length determines the value of the rotation angle. Thus, if \mathbf{M}_k is the torque acting on particle k to be rotated and r_m is the maximum length of any component of the rotational vector \mathbf{r}_k , then its components are generated according to the formula

$$r_k^\alpha = \frac{\ln[\exp(-\lambda\beta M_k^{\alpha, old} r_m) + 2u_k^\alpha \sinh(\lambda\beta M_k^{\alpha, old} r_m)]}{\lambda\beta M_k^{\alpha, old}}. \quad (9)$$

When deriving this formula we have followed the same reasoning as for Eqn. (8) considering the relation $\Delta U \approx (\mathbf{M}_k \cdot \mathbf{r}_k)$ instead of $\Delta U \approx (\mathbf{F}_k \cdot \mathbf{t}_k)$. The acceptance probability is given by the same general formula as for translations, Eqn. (2), with F_k^α being replaced by M_k^α , t_k^α by r_k^α , and t_m by r_m in Eqns (3) and (4).

When generating a trial configuration, one can follow two possible scenarios: (i) let each particle to decide, with a certain probability p_r , whether to attempt rotation or translation (each-particle-decides scenario), or (ii) all particles are subject to rotation at once with probability p_r , otherwise all of them are displaced (all-particles-same scenario). In both cases, each particle is translated or rotated independently of translations and rotations of other particles by vectors \mathbf{t}_k or \mathbf{r}_k , respectively, according to Eqns (8) or (9).

3. Implementation and simulation details

Implementation of the MPM method for polarizable site-site models is considerably different from that considered in the original paper - the polarizable Stockmayer potential. To keep the considerations as general as possible but keeping simultaneously in mind the most common models of water, TIPiP models [11, 12, 13], we will consider here a general site-site potential model with one Lennard-Jones (LJ) site referred to as the oxygen (O) site, several Coulombic sites, and an induced dipole moment located in the center of mass (P). The total configurational energy is thus given by

$$U = U_{\text{LJ}} + U_{\text{qq}} + U_{\text{pol}} , \quad (10)$$

where U_{LJ} is the energy due to the LJ interactions between the O-sites, U_{qq} stands for the charge-charge interactions, and U_{pol} is the contribution due to the polarization:

$$U_{\text{LJ}} = \sum_{i<j} u_{\text{LJ},ij} = 4\epsilon \sum_{i<j} \left(\left(\frac{\sigma}{R_{i\text{O},j\text{O}}} \right)^{12} - \left(\frac{\sigma}{R_{i\text{O},j\text{O}}} \right)^6 \right) \quad (11)$$

$$U_{\text{qq}} = \sum_{i<j} u_{\text{qq},ij} = \frac{1}{4\pi\epsilon_0} \sum_{i<j} \sum_{\alpha,\beta} \frac{q_\alpha q_\beta}{R_{i\alpha,j\beta}} \quad (12)$$

$$U_{\text{pol}} = -\frac{1}{2} \sum_i (\mathbf{p}_i \cdot \mathbf{E}_{i\text{P}}^q) . \quad (13)$$

In the above equations $R_{i\alpha,j\beta} = |\mathbf{R}_{i\alpha,j\beta}|$ is the site-site separation between site α of molecule i and site β of molecule j , \mathbf{p}_i is the induced dipole moment of molecule i , and $\mathbf{E}_{i\text{P}}^q$ is the electric field at the center of the induced dipole of molecule i originating in the Coulombic sites of the other molecules of the system,

$$\mathbf{E}_{i\text{P}}^q = \frac{1}{4\pi\epsilon_0} \sum_{j \neq i} \sum_{\alpha} \frac{q_\alpha \mathbf{R}_{i\text{P},j\alpha}}{R_{i\text{P},j\alpha}^3} . \quad (14)$$

The induced dipoles are assumed to be proportional to the total electric field acting at their centers,

$$\mathbf{p}_i = \alpha (\mathbf{E}_{i\text{P}}^q + \mathbf{E}_{i\text{P}}^p) , \quad (15)$$

where α is the scalar polarizability and \mathbf{E}_{iP}^p is the electric field at the center of the induced dipole of molecule i originating in the induced dipoles of the other molecules of the system,

$$\mathbf{E}_{iP}^p = \frac{1}{4\pi\epsilon_0} \sum_{j \neq i} \left(\frac{(3\mathbf{p}_j \cdot \mathbf{R}_{iP,jP})\mathbf{R}_{iP,jP}}{R_{iP,jP}^5} - \frac{\mathbf{p}_j}{R_{iP,jP}^3} \right). \quad (16)$$

To determine the contribution of the polarizable part to the total energy, one has to evaluate first the value of the induced dipoles from equation (15).

To handle the long-range interactions we use the reaction field method [14]. It means that the interaction energy of particle i is given by two contributions: (i) its interaction with all particles within the sphere at the center of mass (i.e., site P) of particle i and the cutoff radius R_c , and (ii) the interaction with the dielectric continuum representing the particles outside this sphere. Because of the large value of permittivity of water and the fact that the prefactor $(\epsilon_{\text{RF}} - 1)/(2\epsilon_{\text{RF}} + 1)$ tends to $1/2$ for large ϵ_{RF} , we set $\epsilon_{\text{RF}} = \infty$. This formalism gives rise to additional terms and the equations must therefore be modified accordingly:

$$\begin{aligned} U_{\text{LJ,cut}} &= \sum_{i < j} H(R_c - R_{iO,jO}) u_{\text{LJ},ij} + U_{\text{LJcorr}} \\ &= 4\epsilon \sum_{i < j} H(R_c - R_{iO,jO}) \left(\left(\frac{\sigma}{R_{iO,jO}} \right)^{12} - \left(\frac{\sigma}{R_{iO,jO}} \right)^6 \right) + U_{\text{LJcorr}} \end{aligned} \quad (17)$$

$$\begin{aligned} U_{\text{qq,RF}} &= \frac{1}{4\pi\epsilon_0} \sum_{i < j} H(R_c - R_{iP,jP}) \sum_{\alpha,\beta} \frac{q_{i\alpha} q_{j\beta}}{R_{i\alpha,j\beta}} \left(1 + \frac{1}{2} \left(\frac{R_{i\alpha,j\beta}}{R_c} \right)^3 \right) \\ &\quad - \frac{N}{8\pi\epsilon_0 R_c^3} \left(\sum_{\alpha} q_{\alpha} \mathbf{R}_{i\alpha} \right)^2 \end{aligned} \quad (18)$$

$$U_{\text{pol,RF}} = -\frac{1}{2} \sum_i \mathbf{p}_{i,\text{RF}} \cdot \mathbf{E}_{iP,\text{RF}}^q \quad (19)$$

$$\begin{aligned} \mathbf{E}_{iP,\text{RF}}^q &= \frac{1}{4\pi\epsilon_0} \sum_{j \neq i} H(R_c - R_{iP,jP}) \sum_{\alpha} \frac{q_{\alpha} \mathbf{R}_{iP,j\alpha}}{R_{iP,j\alpha}^3} \left(1 - \left(\frac{R_{iP,j\alpha}}{R_c} \right)^3 \right) \\ &\quad + \frac{1}{4\pi\epsilon_0 R_c^3} \sum_{\alpha} q_{\alpha} \mathbf{R}_{i\alpha} \end{aligned} \quad (20)$$

$$\mathbf{p}_{i,\text{RF}} = \alpha \left(\mathbf{E}_{iP,\text{RF}}^q + \mathbf{E}_{iP,\text{RF}}^p \right) \quad (21)$$

$$\mathbf{E}_{iP,\text{RF}}^p = \frac{1}{4\pi\epsilon_0} \sum_{j \neq i} H(R_c - R_{iP,jP}) \left(\frac{3\mathbf{p}_{j,\text{RF}} \cdot \mathbf{R}_{iP,jP} \mathbf{R}_{iP,jP}}{R_{iP,jP}^5} - \frac{\mathbf{p}_{j,\text{RF}}}{R_{iP,jP}^3} \left(1 - \left(\frac{R_{iP,jP}}{R_c} \right)^3 \right) \right)$$

$$+ \frac{\mathbf{P}_{i,\text{RF}}}{4\pi\epsilon_0 R_c^3}, \quad (22)$$

where H is the Heaviside step function.

The above equations constitute the standard method used in the conventional MC simulations and they can also be used in the MPM method. Their implementation in the MPM method is straightforward but rather inefficient. The problem lies in that when a particle is displaced, the number of molecules within its reaction sphere may change considerably giving rise to a large change in energy which cannot be predicted via the linearization of the configuration energy and which is the principal of the force bias method. In other words, the configuration energy can change more or less independently of the forces acting on the particles only because of the varying number of the particles inside of the reaction spheres. This fact plays a significant role, especially at higher densities and lower temperatures (ambient conditions). Consequently, one is forced to use much smaller values of the translational vector to maintain a reasonable acceptance ratio which evidently lowers efficiency of the entire method. To demonstrate the problem with the discontinuity in the internal energy, we measured the internal energy as a function of the displacement along any randomly selected direction of a randomly selected particle. A typical result demonstrating the step behavior of the energy is shown in figure 1 (conventional RF curve). It is discernible that even for very small displacements significant differences in the energies are observed. To overcome this problem, the potential must be smoothed at the cutoff distance. For simple potentials this is straightforward whereas for the site-site models it is not so simple and various methods can be found in literature (for a discussion on the cutoff and smoothing see, e.g., [15]). For the purpose of this study we have developed and used another method based on an averaging of $U_{ij,\text{qq,RF}}$ over a certain range about the cutoff distance. The idea behind the averaging of the potential is rather simple. The cutoff radius R_c of the dielectric cavity of the conventional RF approach can be chosen arbitrarily, more or less without a significant influence on the structural properties, from the interval from about 8\AA to $L/2$ [16], where L is the simulation box size.

It means, all the forms of the cutoff of the intermolecular reaction field potential with different R_c , which is taken from the appropriately chosen interval, can be considered as proper. An averaged potential, obtained by blending together these proper forms of the conventional reaction field potential should be thus proper, too. Thus, to get a smooth potential, instead of $U_{ij,qq,RF}$ we have used an intermolecular potential $U_{ij,qq,BRF}$ obtained by averaging $U_{ij,qq,RF}$ potentials over an interval (R_{c1}, R_{c2}) :

$$U_{ij,qq,BRF} = \frac{1}{R_{c2} - R_{c1}} \int_{R_{c1}}^{R_{c2}} U_{ij,qq,RF}(R_c; \dots) dR_c . \quad (23)$$

In figure 1 we show (smooth U_{qq} ; conventional U_{pol} curve) that the jumps in the energy are smaller, in contrast to those of the conventional reaction field curve, but they still persist because there is another discontinuity embodied in the polarizable term of energy $U_{pol,RF}$ caused by the discontinuities of $\mathbf{E}_{iP,RF}^q$ and $\mathbf{E}_{iP,RF}^p$. To remove them, the same procedure can be applied. Hence,

$$\mathbf{E}_{iP,BRF}^q = \frac{1}{R_{c2} - R_{c1}} \int_{R_{c1}}^{R_{c2}} \mathbf{E}_{iP,RF}^q(R_c; \dots) dR_c \quad (24)$$

$$\mathbf{E}_{iP,BRF}^p = \frac{1}{R_{c2} - R_{c1}} \int_{R_{c1}}^{R_{c2}} \mathbf{E}_{iP,RF}^p(R_c; \dots) dR_c \quad (25)$$

$$\mathbf{p}_{i,BRF} = \alpha (\mathbf{E}_{iP,BRF}^q + \mathbf{E}_{iP,BRF}^p) \quad (26)$$

and the polarizable energy is then

$$U_{pol,BRF} = -\frac{1}{2} \sum_i (\mathbf{p}_{i,BRF} \cdot \mathbf{E}_{iP,BRF}^q) . \quad (27)$$

Few words on smoothing the energy (forces) at the cutoff seem now appropriate. In general, one should prove that the polarization energy is minimized if the smoothed functions $\mathbf{E}_{iP,BRF}^q$ and $\mathbf{E}_{iP,BRF}^p$ are used for the evaluation of equation (27) instead of the conventional RF electrostatics. However, every method for handling the long range electrostatic interactions is only an approximation. Moreover, the systematic error due to the smoothing is, in most cases, negligible in comparison with the statistical one. The behavior of the internal energy in dependence on the displacement after application of the

above smoothing procedures for both U_{qq} and U_{pol} is also shown in Figure 1 (smooth U_{qq} ; smooth U_{pol} curve) for comparison. As one can see, there are no energy discontinuities during the translations of the particles and the energy changes can be estimated by the products of the forces and the translational vectors, $(\mathbf{F}_i \cdot \Delta \mathbf{r}_i)$, much better.

Finally, to implement the MPM method, the forces and torques acting on the molecules have to be determined. In fact, it is not necessary to know their exact values because they affect only the choice of subsequent configurations (walk in the configuration space) but not the thermodynamic values at equilibrium. Performing simple algebra one can derive the equations for the forces and torques. For the force acting on the particle i we use

$$\mathbf{F}_i = \mathbf{F}_{i,\text{LJ,cut}} + \mathbf{F}_{i,\text{qq,BRF}} + \mathbf{F}_{i,\text{pq,BRF}} + \mathbf{F}_{i,\text{qp,BRF}} + \mathbf{F}_{i,\text{pp,BRF}} \quad (28)$$

where $\mathbf{F}_{i,\text{LJ,cut}}$ is the force on the oxygen site from other oxygens, $\mathbf{F}_{i,\text{qq,BRF}}$ the force on the partial charge sites from the partial charges of the other molecules, $\mathbf{F}_{i,\text{pq,BRF}}$ the force on the induced dipole site from the partial charges of the other molecules, $\mathbf{F}_{i,\text{qp,BRF}}$ the force on the partial charge sites from the induced dipoles of the other molecules, and $\mathbf{F}_{i,\text{pp,BRF}}$ the force on the induced dipole site from the induced dipoles of the other molecules. Deriving the expressions for the forces one has to keep in mind that the site-site interactions are not central due to the potential smoothing; the contributions to $\mathbf{F}_{i,\text{pq,BRF}}$ can be calculated during the evaluation of $\mathbf{F}_{i,\text{qp,BRF}}$ using Newton's third law.

Since the local fields at all particle positions are dependent on each other, an iterative technique is the common method to solve the set of equations (15) and (16) to obtain the self-consistent contribution to the total energy. This is usually fast converging process with the criterion [17]

$$\max_{k=1,\dots,N} \frac{|\mathbf{p}_k(n) - \mathbf{p}_k(n-1)|}{|\mathbf{p}_k(n)|} < 10^{-m} \quad (29)$$

used to terminate the iteration; we always used $m = 5$. The number of iteration loops was about 12 for the MPM MC and 8 for the conventional MC. It is worth recalling that Jedlovsky and Richardi [18] used much softer criterion to terminate the iteration, $m = 3$,

in order to speed up the simulation, and Přeboda et al. [4] used some times even softer one.

In the NVT simulations we evaluated both the common thermodynamic properties, the internal energy and the dielectric constant, and the relevant structural properties, the site-site correlation functions g_{ij} and the dipole-dipole correlation functions $G_l, l = 1, 2$ [16]. The latter functions, called also ‘local g factors’, provide information on the mutual alignment of dipoles; moreover, they may also be used as an alternative route to the dielectric constant, see e.g. [19]. In this paper we evaluate the static dielectric constant, ε_r , from the expression [4]:

$$\varepsilon_r = 1 + \frac{\rho\tau^2}{3\varepsilon_0 k_B T}, \quad (30)$$

where τ^2 is the fluctuation of the total dipole moment $\mathbf{M} = \sum_{i=1}^N (\boldsymbol{\mu}_i + \mathbf{p}_i)$ of the simulation box per molecule,

$$\tau^2 = \frac{\langle \mathbf{M}^2 \rangle - \langle \mathbf{M} \rangle^2}{N}. \quad (31)$$

To be able to assess efficiency of the method, in addition to the above quantities we computed also three measures, the translational order parameter, TOP [20], the rotational relaxation parameter of the first-order (re-orientation of a molecule-fixed unit vector), RRP [21], and the mean-squared displacement, MSD [2]. The measures are defined as follows:

$$\text{TOP}(t) = \frac{1}{N} \sum_{i=1}^N \sqrt{\langle \cos(\mathbf{k} \cdot \mathbf{r}_i(t)) \rangle^2 + \langle \sin(\mathbf{k} \cdot \mathbf{r}_i(t)) \rangle^2}, \quad (32)$$

$$\text{RRP}(t) = \frac{1}{N} \sum_{i=1}^N \langle \boldsymbol{\nu}_i(t) \boldsymbol{\nu}_i(0) \rangle \quad (33)$$

$$\text{MSD}(t) = \frac{1}{N} \sum_{i=1}^N \frac{\langle |\mathbf{r}_i(t) - \mathbf{r}_i(0)|^2 \rangle}{t}, \quad (34)$$

where $\mathbf{k} = (-1, 1, -1)$, $\mathbf{r}(t)$ is the position of the particle at time (configuration) t , and averaging is performed over all particles of the sample. We remark that the used TOP given by Eqn. (32) differs from that given in [2]; the latter is not origin-independent and may yield even unphysical results [20]. Finally, for more accurate efficiency measurements we

calculated also the autocorrelation coefficients of equilibrium thermodynamic properties according to the following formula [22]

$$c(k) = \frac{\frac{1}{n-k} \sum_{i=1}^{n-k} \Delta X_i \Delta X_{i+k}}{\frac{1}{n} \sum_{i=1}^n \Delta X_i^2}, \quad (35)$$

where n is the data count and X_i is the i -th measured value of the considered thermodynamic property. The faster the autocorrelation coefficient converges to zero with increasing k , the more efficient the simulation is because the time necessary to generate two subsequent uncorrelated configurations is shorter.

The results reported in the next Section were obtained using 256 and 500 particles. Most simulations were however carried out with the former number of particles whereas the latter one served mainly for checking purposes only. In all the simulations we used the Lennard-Jones interaction cutoff $R_{c,LJ} = 9\text{\AA}$. In the conventional MC simulations we used the reaction field without any smoothing and cutoff $R_c = 9\text{\AA}$. This cutoff value was used by Pěrdota et al. [3, 4] and is therefore used also here for a fair comparison. For MPM MC we used the same cutoffs, however, in the case of smoothing, the electrostatic contributions were smoothed over the interval $R_c \in (8.8\text{\AA}, 9\text{\AA})$. We found that the efficiencies of the all-particles-same and each-particle-decides concepts are similar. Thus, the all-particles-same scenario has been used in most simulations because it provides the possibility to check separately the acceptance ratios of the moves. In the case of the MPM MC in an NPT ensemble, the volume change was realized by the standard way leading to the acceptance ratio of 0.3. The MC steps consisted of a volume change with probability 0.02, or a rotation or translation of all particles with equal probability (using $p_r = 0.5$, $t_m = 0.05\text{\AA}$, $r_m = 0.12$). The thermodynamic properties were measured after every 100 steps. In the case of the MPM MC in an NVT ensemble, there were 60 attempts between two subsequent measurements (using $p_r = 0.5$, $t_m = 0.07\text{\AA}$, $r_m = 0.12$ for state point A, and $p_r = 0.5$, $t_m = 0.12\text{\AA}$, $r_m = 0.23$ for state point C; states A and C are defined in the following section). In the conventional MC, all the properties were measured after every $3N$ molecule displacement/rotation attempts with acceptance ratios about 0.3. The

1
2
3
4
5
6
7 maximum possible length of any component of the conventional MC translation vector
8 was 0.5Å for state A, and 0.9Å for state C. The particles were rotated around uniformly
9 randomly selected axes by uniformly generated random angles not greater than 1.2rad for
10 state A and 1.8rad for state C. To measure the computation speed, and hence to compare
11 efficiency, all the computations ran on the same hardware and all simulations started from
12 the same configuration: FCC lattice with random orientations of the molecules, $\mu_i = \mu\nu_i$.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3. Results and discussion

We have implemented the above outlined MPM MC method for the Chialvo-Cummings (CC) polarizable model of water [8] which is based on the SPC/E model [23], and the Brodholt-Sampoli-Vallauri (BSV) polarizable model [9] based on the TIP4P [24], both of them with the scalar polarizability $\alpha = 1.444\text{\AA}^3$. For further details and the parameters of the models we refer the reader to the original papers [8, 9]. We performed both NVT and NPT simulations at two thermodynamic states: at ambient conditions, $T = 298\text{K}$ and number density $\rho = 0.0334\text{\AA}^{-3}$ (NVT) or pressure $p = 1\text{bar}$ (NPT), and a slightly subcritical temperature (of real water), $T = 573\text{K}$ and $\rho = 0.0241\text{\AA}^{-3}$ (NVT) or $p = 100\text{bar}$ (NPT). Following Jedlovszky and Richardi [18], we will refer to these states as A and C, respectively. These states were also considered by Předota et al. [3, 4] and they were therefore chosen for the sake of comparison. With respect to the complexity of the problem, both physical and computational, and the novelty of cutoff smoothing, in the following we will first demonstrate correctness of the proposed implementation of the MPM method by presenting the results for the thermodynamic and structural properties and their comparison with those reported in [3, 4, 18]. Then the problem of efficiency of the method and its assessment will be addressed.

3.1. Thermodynamic and structural properties

In tables 1 and 2 we list the results for the internal energy, the average values of the total dipole moment and the induced dipole moment of the molecules, and the dielectric constant. It seems necessary first to recall that in both references, [4] and [18], the authors report error estimates only for the internal energy but not for other properties. It is also necessary to remark that the latter results seem to be subject of rather large errors. In state A, the MPM MC results for the average values of both the permanent and total dipoles, and for the dielectric constant agree with literature data. As for the energy, the agreement between the results reported by Předota et al. and those by Jedlovszky and Richardi is at the edge of their combined errors. Our average value of U lies between

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

theirs and agrees with both of them also within the combined errors. Practically the same applies also to the energy in state C with much less differences between the average values. Předota et al. do not report any results for other quantities at this state. We agree with Jedlovsky and Richardi for the averaged dipoles and the only difference we find is for the dielectric constant. However, with respect to the usual large uncertainty of this quantity this is not surprising (see also the discussion below).

The equilibrium number densities obtained by the NPT simulations are shown in table 3. In state C, the results agree, more or less, with those reported in [18]. However, in state A, they differ significantly. When examining more carefully the table in [18] we find that the listed results for non-polarizable models disagree with those reported in literature by other authors. For instance, the number density for SPC/E water in state A reported in [18] is 0.0352\AA^{-3} which corresponds to 1053kg m^{-3} and is quite off the experimental value to which the potential was fitted. On the other hand, the value we get using the MPM MC method is 0.033431\AA^{-3} (i.e., 1000.08kg m^{-3}) which is much closer to the experimental measurements and agrees also with other simulation data, see e.g. [25, 26]. We therefore tend to question reliability of the data given in [18].

The site-site correlation functions are shown in figures 2 and 3 where they are compared with those obtained by Předota et al. This comparison confirms again that the structure generated by the conventional and MPM MC methods are identical. In addition to the site-site correlation functions $g_{\alpha\beta}$, we also computed the first two dipole-dipole correlation functions G_l ; they are shown in figure 4. These functions are not available for the studied model in literature but were reported in [19] for its parent model TIP4P, and in [27] for other polar fluids. For nonpolarizable models, function G_1 is characterized by a fast decay after the first coordination shell with a small but persistent negative value over a large range of separations. It means that after the first coordination shell a certain uniform alignment can be observed [19]. For the polarizable model, G_1 also decays fast to unity after the first peak but then it exhibits an oscillatory behavior characteristic for cutoff

1
2
3
4
5
6
7 models.

8 9 *3.2. Efficiency of the method*

10
11 As mentioned above, all simulations started from the same crystalline phase. To assess
12 efficiency of different methods, we focussed on the rate with which the systems went over
13 from the crystalline phase to an equilibrated disordered phase. The results are shown in
14 figure 5 for the TOP and RRP. As we can see, in the MPM MC method the system
15 melts very fast (note the different time scales) whereas using the conventional MC it
16 takes about by one order more time. It may be also of interest to remark that, in contrast
17 to our previous work on the Stockmayer fluid [5], the RRP converges more slowly than
18 the TOP because of the highly directional intermolecular potential. In Figure 6 we show
19 the development of the total internal energy. Its rate of convergence coincides with that
20 of TOP, suggesting the computational convenience of the biased MPM MC method and
21 verifies that all the simulated systems reached equilibria.
22
23
24
25
26
27
28
29
30
31
32

33 A closely related question is how fast the system loses its memory. Thus, in order to
34 eliminate the possibility of spurious fast convergence we have followed the previous paper
35 [5] and applied the blocking method of Flyvbjerg and Petersen ([28]) to the total energy.
36 In figure 7 we present an example of the resulting graphs which were analyzed in order
37 to determine the correlation lengths and associated computational times. In addition,
38 we present a comparison of the autocorrelation coefficients of the configurational energy
39 in figure 8. We find that the computational correlation times, i.e., the machine time
40 necessary to generate an independent block of configurations, is about ten times shorter
41 for the MPM method in comparison with the conventional single-particle MC. However,
42 it is also fair to say that these estimates are rather approximate due to the requirement
43 of having extremely large amount of data for the analysis and the presence of long range
44 correlations embodied in the polarizable models itself.
45
46
47
48
49
50
51
52
53
54
55
56

57 The last quantity of interest is the MSD shown in figure 9. This quantity is related to
58 diffusivity and reflects mobility of the molecules. As it is seen, in the biased MPM MC
59
60

1
2
3
4
5
6
7 method particles move much faster and thus, again, sample the configuration space much
8 more efficiently.
9

10 We did not carry out any efficiency measurements in the NPT simulations because the
11 only difference between the NPT and NVT simulations is an additional volume change
12 step, which is naturally all-particle based and is thus not affected by the MPM MC.
13
14

15
16 In addition to the efficiency discussed above in terms of various measures, a direct
17 comparison of the CPU time required by different methods would be also quite useful
18 information. However, such a comparison can hardly be made in a fair way and only
19 approximate or indirect estimates can be given. The approximate PAPI method, designed
20 to perform efficiently, is governed by a number of parameters. It was reported in [4] that
21 the combination which is accurate and simultaneously also efficient performs about five
22 times faster in comparison with the common (exact) method. In the case of the MPM
23 MC method the gain in speed is one order which means that it may require only one half
24 of the CPU time required by (only approximate!) PAPI method. As regards the ANES
25 method, we are not aware of any representative comparison with the conventional MC;
26 however, computing-time requirements of this method increase only as N^2 under certain
27 conditions (e.g. small nuclear displacements and preferential sampling) [6]
28
29
30
31
32
33
34
35
36
37
38
39
40

41 *3.3. Parameter dependence*

42
43 There are several simulation parameters that must be set to run the simulation and
44 that also affect its efficiency. Dependence of efficiency of the MPM MC on these param-
45 eters was investigated in an NVT ensemble using the polarizable BSV model at ambient
46 conditions with 256 particles. To assess the parameter dependence we focused on the
47 autocorrelation coefficients of the configurational energy, U , its individual contributions
48 (U_{LJ} , U_{qq} , U_{pol}), and the mean value of the induced dipole moment μ . Both U_{qq} and
49 U_{pol} were smoothed as described above. In order to make the measurement of accep-
50 tance ratio of translations a_t and rotations a_r possible, trial configurations were gener-
51 ated using the all-particles-same scenario. It was practically impossible to examine the
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 multiplicity of the combinations of parameters t_m , r_m and p_r . Thus, a simple iterative
8 procedure was applied. At first, we made simulations with $r_m = 0.12$ (corresponding
9 to the acceptance ratio of rotations of all the particles $a_r = 0.35$), $p_r = 0.5$ and $t_m =$
10 $\{0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1\}$ Å. The resulting autocorrelation coefficients re-
11 vealed that the most convenient choice for t_m lies between 0.05Å and 0.07Å, whereas the
12 corresponding acceptance ratios a_t are between 0.3 and 0.55. The efficiency of the MPM
13 MC simulations is almost constant in this interval. Then we carried out simulations with
14 parameters $t_m = 0.05$ Å, $p_r = 0.5$ and $r_m = \{0.04, 0.06, 0.08, 0.09, 0.10, 0.11, 0.13, 0.14, 0.15, 0.17\}$.
15 The fastest drop of the autocorrelation coefficients was observed for r_m between 0.09 and
16 0.13 corresponding to a_r between 0.28 and 0.58. The efficiency dependence on p_r was ex-
17 plored by simulations with $t_m = 0.05$ Å, $r_m = 0.12$ and $p_r = \{0.1, 0.2, 0.25, 0.3, 0.35, 0.45, 0.55, 0.6, 0.65, 0.7\}$.
18 The most convenient values of p_r seem to be between 0.4 and 0.6. This analysis shows that
19 all the parameters chosen (intuitively) at the beginning of the whole procedure fall to the
20 intervals of the most convenient values. In figure 10 we show selected convergence profiles
21 of the autocorrelation coefficients of the configurational energy resulting from these MPM
22 MC simulations using different combinations of parameters.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 In table 4 we give the parameters used for the translational and rotational moves for
39 different smoothing techniques leading to the acceptance ratios about 0.3. As it is seen,
40 the smoothing does play the significant role for the translations in the most interesting
41 low temperature state A. Increase of temperature diminishes differences in the energy
42 change as it is demonstrated by the parameters for state C. Rotation does not lead to
43 any change in the number of particles within the cutoff sphere and does not affect thus
44 the energy change to such an extent regardless of the thermodynamic state.
45
46
47
48
49
50
51
52

53 3.4. Parallelization

54 The MPM MC scheme is a natural candidate of parallelization, i.e., another possibility
55 to speed up the computations. To this end, the MPM MC program code was parallelized
56 and simulations of BSV polarizable model of water at ambient conditions were performed.
57
58
59
60

1
2
3
4
5
6
7 The OpenMP directives and Intel Fortran Compiler 10.1. were used for this purpose. The
8 computer hardware available was 14 nodes SGI AltixXE 310 (7 pieces of 1U server AltixXE
9 310, each containing 2 nodes) with 28 processors Intel Xeon Quad-core 5365 (frequency
10 3,00 GHz) and 16 GB shared memory DDR2 (667MHz) at each node. Our goal was
11 to parallelize the code for one computer node, i. e., 8 computational cores with shared
12 memory. With respect to the program structures already contained in the MPM MC
13 there was no problem to add a few OpenMP directives and distribute the calculation job
14 to the individual cores efficiently. The calculations of distances and vectors between the
15 interaction sites of molecules, calculations of nonpolarizable energies and forces, iteration
16 of induced dipoles, calculation of forces resulting from the polarizability of molecules and
17 the initializations of long memory fields were parallelized in this way. On the other hand,
18 neither the computations of the MSD, TOP, RRP, and the subroutine generating trial
19 configurations, nor the pseudo random number generator were parallelized.
20
21
22
23
24
25
26
27
28
29
30
31

32 To compare the speed of computations, the program speed running on 8 cpu cores
33 was about 7 times higher than that on a single cpu (for both $N = 256$ and $N = 500$).
34 Thus, combining the MPM MC scheme with parallelization using a common 8-core com-
35 putational node, the factor of the gain in speed has been about 70 in comparison with
36 a common one-particle MC. Finally, to avoid any confusion we must mention in passing
37 that all the results concerning the efficiency and discussed in the preceding subsections
38 were obtained without any parallelization.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4. Conclusions

We have recently proposed a novel MC method, the MPM MC scheme, in which trial moves of all particles are attempted simultaneously. The main gain in efficiency is due to biasing the moves with respect to the forces acting between the molecules. There is no question concerning validation of the method but its implementation need not be straightforward for complex systems. In this paper we have considered a site-site interaction model with the Coulombic interactions and an isotropic polarizability exemplified by the Chialvo-Cummings and Brodholt-Sampoli-Vallauri models of water. When implementing the MPM MC method for such systems one may also encounter problems with cutoff corrections which affect the efficiency. Whereas in the case of the common one-particle MC method no smoothing need even be applied, in the case of the MPM MC this becomes necessity to make the simulation really efficient (i.e., to change configurations one from another significantly). We have therefore introduced and tested in this paper a novel method, a sort of an average correction; but in fact, this is only a matter of a personal taste and other routines may be applied as well. If the more time-consuming Ewald summation is employed, the convenience of the MPM MC should be similar or better because using the MPM MC means greater configurational change for the same number of calculations of the configurational energy.

In contrast to the well known Hybrid MC/MD method, the MPM MC does not require any time consuming fine-tuning of a number of parameters and the implementation of the MPM MC is rather simple. Originally, the three parameters governing the efficiency, t_m , r_m , and p_r , were set by intuition following the recommendations for the common MC simulations. It has been however shown that the range of optimal values of these parameters is rather broad and that the use of intuition is sufficient. Moreover, the MPM MC may be naturally parallelized which is not the case of the common one-particle MC scheme. In spite of using just the simplest OpenMP directives and parallelizing the most

1
2
3
4
5
6
7 important program sections only, we obtained a very efficient tool for studying systems
8 not only with polarizable models but, in general, with non-additive forces which is a path
9 that will be likely followed in future development of the liquid state theory.
10

11
12 As for an application of the MPM MC to molecular systems with pairwise additive
13 interactions, we have found out from our Lennard-Jones fluid simulations that the MPM
14 MC is about seven times less efficient than the conventional MC. It is not surprising
15 because the MPM MC has not been devised for simple systems and does not utilize
16 additivity of the configurational energy at all. That is the reason why we decided not
17 to present such results regarding these simple models. On the other hand, we think that
18 the MPM MC could be an ideal candidate for nowadays massively parallel Graphics
19 processing unit computing, which is the case where the ability of efficient parallelization
20 can potentially overcome other aspects of the algorithm.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgement

This work was supported by the Grand Agency of the Academy of Sciences of the Czech Republic (Grant No. IAA400720802). Valuable discussions with Dr. M. Předota, University of South Bohemia, are also greatly acknowledged.

References

- [1] D. Frenkel, B. Smit, *Understanding Molecular Simulation* (Academic Press, San Diego, 2002).
- [2] Allen, M. P., Tildesley, D. J., 1987, *The Computer Simulation of Liquids* (Clarendon Press, Oxford).
- [3] M. Předota, P.T. Cummings, and A.A. Chialvo, *Mol. Phys.* 99 (2001), p. 349.
- [4] M. Předota, P.T. Cummings, and A.A. Chialvo, *Mol. Phys.* 100 (2002), p. 2703.
- [5] F. Moučka, M. Rouha, and I. Nezbeda, *J. Chem. Phys.* 126 (2007), p. 224106.
- [6] B. Chen, J.I. Siepmann, *Theoret. Chem. Accounts* 103 (1999), p. 87.
- [7] B. Chen, J. I. Siepmann *J. Phys. Chem. B* 104 (2000), p. 2378.
- [8] A.A. Chialvo, P.T. Cummings, *J. Chem. Phys.* 105 (1996), p. 8274.
- [9] J. Brodholt, M. Sampoli, R. Vallauri, *Mol. Phys.* 86 (1995), p. 149.
- [10] C. Pangali, M. Rao, and B.J. Berne, *Chem. Phys. Lett.* 55 (1978), p. 413.
- [11] B. Guillot, *J. Mol. Liq.* 101 (2002), p. 219.
- [12] S.W. Rick, *J. Chem. Phys.* 120 (2004), p. 6085.
- [13] J.L.F. Abascal, C. Vega, *J. Chem. Phys.* 123 (2005), p. 234505.
- [14] M. Neumann, *Mol. Phys.* 39 (1980), p. 437.
- [15] I.P. Omelyan, *Phys. Lett. A* 223 (1996), p. 295.
- [16] O. Steinhauser, *Molec. Phys.* 45 (1982), p. 335.
- [17] F.J. Vesely, *J. Comput. Phys.* 24 (1977), p. 361.

- 1
2
3
4
5
6
7 [18] P. Jedlovszky, J. Richardi, J. Chem. Phys. 110 (1999), p. 8019.
8
9
10 [19] J. Kolafa, I. Nezbeda, Molec. Phys. 98 (2000), p. 1505.
11
12 [20] [http://www.ccl.net/cca/software/SOURCES/FORTRAN/allen-tildesley-](http://www.ccl.net/cca/software/SOURCES/FORTRAN/allen-tildesley-book/f.25.shtml)
13 [book/f.25.shtml](http://www.ccl.net/cca/software/SOURCES/FORTRAN/allen-tildesley-book/f.25.shtml)
14
15
16
17 [21] R. Witt, L. Sturz, A. Doelle, and F. Mueller-Plathe, J. Phys. Chem. A 104 (2000),
18 p. 5716.
19
20
21 [22] J. Kolafa, J. W. Perram, Mol. Simul. 9 (1992), p. 351.
22
23
24 [23] H.J.C. Berendsen, J.R. Griega, and T.P. Straatsma, J. phys. Chem. 91 (1987), p.
25 6269.
26
27
28
29 [24] W.L. Jorgensen, J. Chandrasekhar, J.D. Madura, R.W. Impey, and M.L. Klein, J.
30 Chem. Phys. 79 (1983), p. 926.
31
32
33 [25] W.L. Jorgensen, C. Jenson, J. Comput. Chem. 19 (1998), p. 1179.
34
35
36 [26] E. Sanz, C. Vega, J.L.F. Abascal, and L.G. MacDowell Phys. Rev. Lett. 92 (2004),
37 p. 255701.
38
39
40
41 [27] J. Kolafa, I. Nezbeda, M. Lisal, M., Molec. Phys. 99 (2001), p. 1751.
42
43
44 [28] H. Flyvbjerg, H.G. Petersen, J. Chem. Phys. 91 (1989), p. 461.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Thermodynamic properties of the CC model obtained using different simulation methods.

	$U/ \text{kJ mol}^{-1}$	$\langle \mathbf{p} \rangle$	$\langle \mathbf{p} + \boldsymbol{\mu} \rangle$	ε_r
state point A				
MPM				
$U_{\text{qq,BRF}}; U_{\text{pol,BRF}}$	-40.62 ± 0.03	0.998 ± 0.001	2.81 ± 0.01	99 ± 10
$U_{\text{qq,BRF}}; U_{\text{pol,RF}}$	-40.61 ± 0.05	0.999 ± 0.001	2.82 ± 0.01	87 ± 15
$U_{\text{qq,RF}}; U_{\text{pol,RF}}$	-40.62 ± 0.05	0.997 ± 0.002	2.81 ± 0.01	86 ± 20
Ref. [4]	-40.71 ± 0.06	1.00	2.82	90
Ref. [17]	-40.31 ± 0.35	0.978	2.792	104.8
state point C				
MPM				
$U_{\text{qq,BRF}}; U_{\text{pol,BRF}}$	-23.48 ± 0.02	0.582 ± 0.001	2.374 ± 0.001	21.5 ± 0.3
$U_{\text{qq,BRF}}; U_{\text{pol,RF}}$	-23.48 ± 0.02	0.582 ± 0.001	2.374 ± 0.001	21.7 ± 0.5
$U_{\text{qq,RF}}; U_{\text{pol,RF}}$	-23.51 ± 0.02	0.583 ± 0.001	2.376 ± 0.001	21.8 ± 0.5
Ref. [4]	-23.53 ± 0.02	-	-	-
Ref. [17]	-23.48 ± 0.58	0.585	2.378	17.4

Table 2: Thermodynamic properties of the BSV model (MPM MC with $U_{\text{qq,BRF}}$; $U_{\text{pol,BRF}}$).

	$U/ \text{kJ mol}^{-1}$	$\langle \mathbf{p} \rangle$	$\langle \mathbf{p} + \boldsymbol{\mu} \rangle$	ϵ_r
state point A				
MPM	-41.93 ± 0.02	0.988 ± 0.001	2.808 ± 0.001	150 ± 15
Ref. [18]	-41.70 ± 0.41	0.966	2.778	132.2
state point C				
MPM	-24.305 ± 0.008	0.580 ± 0.001	2.376 ± 0.001	23.5 ± 0.3
Ref. [18]	-24.20 ± 0.55	0.576	2.365	18.4

Table 3: Number densities obtained by NPT simulations for the CC and BSV models (MPM MC with $U_{\text{qq,BRF}}$; $U_{\text{pol,BRF}}$).

	$\rho / 10^{-2} \text{ \AA}^{-3}$			
	state point A		state point C	
	CC	BSV	CC	BSV
MPM	2.863 ± 0.001	3.377 ± 0.003	0.1485 ± 0.0002	0.1569 ± 0.0004
Ref. [18]	2.956 ± 0.046	3.622 ± 0.045	0.140 ± 0.005	0.152 ± 0.007

Table 4: Parameters of the MPM MC moves adjusted so as to maintain the acceptance ratio about 0.3 using different smoothing methods and the CC model of water in an NVT ensemble.

	$t_m / \text{\AA}$	r_m / rad
state point A		
MPM		
$U_{\text{qq,BRF}}; U_{\text{pol,BRF}}$	0.07	0.13
$U_{\text{qq,BRF}}; U_{\text{pol,RF}}$	0.05	0.13
$U_{\text{qq,RF}}; U_{\text{pol,RF}}$	0.03	0.13
state point C		
MPM		
$U_{\text{qq,BRF}}; U_{\text{pol,BRF}}$	0.12	0.23
$U_{\text{qq,BRF}}; U_{\text{pol,RF}}$	0.12	0.23
$U_{\text{qq,RF}}; U_{\text{pol,RF}}$	0.12	0.23

Figure Captions

Figure 1. Typical behavior of the internal energy during the translation of a randomly chosen molecule along any randomly selected constant direction using different reaction field smoothings and the linearized energy: large scale (top), detail (bottom).

Figure 2. Comparison of the site-site correlation functions of the CC model in state A: MPM MC method with the smoothed U_{qq} and U_{pol} (this work; solid line) and the common MC method (Ref. [4]; circles).

Figure 3. The same as Fig. 2 for state C.

Figure 4. The dipole-dipole correlation functions of the CC model in state A (left column), and state B (right column).

Figure 5. Comparison of the development of the translational order parameters (TOP) and the rotational relaxation parameters (RRP) in the CC model in an NVT simulation and state A (left column) and state B (right column): the conventional MC (upper row) and the MPM-MC with the smoothed U_{qq} and U_{pol} (lower row).

Figure 6. The same as in Fig. 5 for the internal energy.

Figure 7. Block analysis of the total internal energy for state C using the CC model in the NVT ensemble: conventional MC method (left), MPM MC method with the smoothed U_{qq} and U_{pol} (right). $\sigma(m)$ is the standard deviation.

Figure 8. Autocorrelation coefficients of the total internal energy using the CC model and different methods in the NVT simulation: state A (top), state C (bottom). MPM MC is with the smoothed U_{qq} and U_{pol} .

Figure 9. The same as in figure 5 for the mean square displacement (MSD).

1
2
3
4
5
6
7 **Figure 10.** Comparison of the autocorrelation coefficients of U for the BSV model in
8 state A and NVT simulations using the MPM MC with the smoothed U_{qq} and U_{pol}
9 and different parameters.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Fig. 1 / Moučka





Fig. 2 / Moučka




Fig. 3 / Moučka





Fig. 4 / Moučka



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 5 / Moučka




Fig. 6 / Moučka






Fig. 7 / Moučka



Review Only

Fig. 8 / Moučka





Fig. 9 / Moučka



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 10 / Moučka

