



HAL
open science

Does cytogenetic evolution have any prognostic relevance in myelodysplastic syndromes? A study on 153 patients from a single institution

Paolo Bernasconi, Catherine Klersy, Marina Boni, Paola Maria Cavigliano, Ilaria Giardini, Barbara Rocca, Rita Zappatore, Irene Dambruoso, Celeste Calvello, Marilena Caresana, et al.

► To cite this version:

Paolo Bernasconi, Catherine Klersy, Marina Boni, Paola Maria Cavigliano, Ilaria Giardini, et al.. Does cytogenetic evolution have any prognostic relevance in myelodysplastic syndromes? A study on 153 patients from a single institution. *Annals of Hematology*, 2010, 89 (6), pp.545-551. 10.1007/s00277-010-0927-z . hal-00516157

HAL Id: hal-00516157

<https://hal.science/hal-00516157>

Submitted on 9 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Editorial Manager(tm) for Annals of Hematology
Manuscript Draft

Manuscript Number: AOHE-D-09-00174R2

Title: DOES CYTOGENETIC EVOLUTION HAVE ANY PROGNOSTIC RELEVANCE IN MYELODYSPLASTIC SYNDROMES? A STUDY ON 153 PATIENTS FROM A SINGLE INSTITUTION.

Article Type: Original Article

Keywords: Myelodysplastic syndromes 1, cytogenetics 2, prognostic factors 3

Corresponding Author: Prof. P. Bernasconi,

Corresponding Author's Institution:

First Author: P. Bernasconi

Order of Authors: P. Bernasconi; Catherine Klersy; Marina Boni; Paola Maria Cavigliano; Ilaria Giardini; Barbara Rocca; Rita Zappatore; Irene Dambruoso; Celeste Calvello; Marilena Caresana; Mario Lazzarino, Prof

Abstract: The present study was designed to establish the incidence of cytogenetic evolution (CE), defined as the acquisition of chromosomal defects during the course of MDS, in order to correlate it with the WHO classification and IPSS score; and to assess its impact on overall survival (OS) and risk of MDS/AML evolution (progression-free interval, PFI) by means of Cox models for time-dependent covariates. Adjustments for known risk factors were achieved by performing a bi-variable analysis. The study was carried out in 153 MDS patients who were followed for a median period of 45.2 months. Disease progression occurred in 42.4% of patients after a 65.2 month median PFI, while CE occurred in 30.7% of patients.

Our study shows that i) CE was more common in advanced than in early MDS and advanced MDS presented secondary chromosomal defects distinct from those of early MDS; ii) CE significantly effected OS and PFI independently of other prognostic variables; iii) del(7)(q31q34) was the only secondary chromosomal defect which significantly effected PFI; trisomy 8 had only a moderate influence.

Response to Reviewers: Replies to Reviewers' comments

As requested:

1. FAB classification has been completely omitted from the manuscript (text and tables). However, in table 1 it has been underlined that 11 of the 153 patients, classified as RAEB-t according to FAB classification, were classified as AML according to WHO. Furthermore, in table 2 and 3 it is underlined that 5 of the 40 patients who experienced cytogenetic evolution were classified as RAEB-t according to FAB but as AML according to WHO.
2. Peripheral blood counts have been completely omitted from the tables.
3. The WHO classification at diagnosis and cytogenetic evolution with percentage of marrow blasts has been provided (table 2).
4. IPSS has been deleted from the tables and the manuscript has concentrated on cytogenetic evaluation

5. The number of tables has been reduced to four. In particular:
The previous tab.2 has been fused to tab.1; the previous tab.3 has been omitted and substituted with the new tab.2 which reports the clinical and cytogenetic features of the 40 patients who experienced CE; the previous tables 4 and 5 has been fused together in the new table 3 which maintains their relevant information; the previous table 6, now table 4, has been maintained; the previous tables 7 and 8 has been omitted.

1 **DOES CYTOGENETIC EVOLUTION HAVE ANY PROGNOSTIC RELEVANCE IN**
2 **MYELOYDYSPLASTIC SYNDROMES? A STUDY ON 153 PATIENTS FROM A SINGLE**
3 **INSTITUTION.**

4 Paolo Bernasconi¹, Catherine Klersy², Marina Boni¹, Paola Maria Cavigliano¹, Ilaria Giardini¹,
5 Barbara Rocca¹, Rita Zappatore¹, Irene Dambruoso¹, Celeste Calvello¹, Marilena Caresana¹, Mario
6 Lazzarino¹.

7 ¹Division of Hematology, Fondazione IRCCS Policlinico San Matteo, University of Pavia, Italy;

8 ²Scientific Direction, Clinical Epidemiology and Biometry Service, Fondazione IRCCS Policlinico
9 San Matteo, Italy.

10
11
12
13
14
15
16
17
18
19
20
21
22
23 **Running title:** Cytogenetic evolution in MDS
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 Prof. Paolo Bernasconi
48 Hematology Division,
49 Fondazione IRCCS Policlinico San Matteo
50 University of Pavia
51 27100-Pavia, Italy
52 Phone +39 0382 503065
53 Fax +39 0382 502250
54 E-mail:p.bernasconi@smatteo.pv.it
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Summary The present study was designed to establish the incidence of cytogenetic evolution (CE), defined as the acquisition of chromosomal defects during the course of MDS, in order to correlate it with the WHO classification and IPSS score; and to assess its impact on overall survival (OS) and risk of MDS/AML evolution (progression-free interval, PFI) by means of Cox models for time-dependent covariates. Adjustments for known risk factors were achieved by performing a bi-variable analysis. The study was carried out in 153 MDS patients who were followed for a median period of 45.2 months. Disease progression occurred in 42.4% of patients after a 65.2 month median PFI, while CE occurred in 30.7% of patients.

Our study shows that i) CE was more common in advanced than in early MDS and advanced MDS presented secondary chromosomal defects distinct from those of early MDS; ii) CE significantly effected OS and PFI independently of other prognostic variables; iii) del(7)(q31q34) was the only secondary chromosomal defect which significantly effected PFI; trisomy 8 had only a moderate influence.

Keywords: Myelodysplastic syndromes, cytogenetics, prognostic factors

Introduction

1
2 Myelodysplastic syndromes (MDS) are heterogeneous clonal stem cell disorders characterized by a
3
4 hypercellular marrow showing ineffective hematopoiesis which is responsible for one or more
5
6 peripheral blood cytopenia(s) [1,2]. Currently, MDS is diagnosed according to the World Health
7
8 Organization (WHO) classification which identifies eight subtypes [3]. This last classification has
9
10 been validated in a large number of patients [4-8] and is continually subject to improvement [9-14].
11
12 The WHO classification system provides relevant prognostic information for a disease with an
13
14 extremely variable natural course [1-15]. MDS patients experience morbidity and mortality rates
15
16 higher than those of age-matched normal subjects because of complications related to peripheral
17
18 cytopenia [16] and AML progression which occurs in about 30% of cases [17]. In 1997, the
19
20 International Prognostic Scoring System (IPSS) revealed that the number of peripheral blood
21
22 cytopenias, cytogenetic pattern and blast cell percentage are the most significant prognostic factors
23
24 in MDS [18]. Additional studies have demonstrated that IPSS power may be improved by better
25
26 defining the prognostic relevance of the chromosomal defects included within the intermediate IPSS
27
28 cytogenetic category [19-22] and this assumption has been further validated [23]. Even more
29
30 recently, the prognostic power of the cytogenetic pattern has been proven by the newly developed
31
32 WHO classification-based prognostic system (WPSS) [24]. However, all these studies have
33
34 analysed the clinical relevance of chromosomal abnormalities revealed only at clinical diagnosis
35
36 (primary defects) [25,26]. In contrast, very few studies have evaluated the incidence and the
37
38 prognostic significance of cytogenetic evolution (CE), defined as the acquisition of either an
39
40 abnormal clone in a karyotypically normal patient or additional defects in patients with an already
41
42 abnormal chromosomal pattern (secondary abnormalities) [27-36]. The majority of these studies,
43
44 which include small numbers of patients with a short follow-up period, have simply reported that
45
46 the acquisition of secondary defects occurred at the time of AML progression in approximately 60%
47
48 of MDS patients, but none have assessed how many patients experience CE before clinical
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 progression and whether this CE is predictive of overall survival (OS) and progression-free interval
2 (PFI) independently of WHO classification, primary defects and IPSS cytogenetic categories.
3

4 These are the goals of the present study, which included 153 patients, observed for a sufficiently
5 long follow-up period, in whom the significant effect of primary cytogenetic defects on disease
6 outcome has already been revealed [21]. An additional goal was to establish which secondary defect
7 significantly predicts leukemic transformation.
8
9
10
11
12
13

14 **Materials and methods**

15 *Patients*

16 All 153 *de novo* MDS patients analysed in the present study were included in a previous series and
17 were diagnosed at the Division of Haematology, Foundation IRCCS Policlinico San Matteo, Pavia
18 between January 1990 and December 2004. All patients were classified according to WHO
19 classification. Our study does not include any patient with either secondary MDS or with a white
20 blood cell (WBC) count above $12 \times 10^9/L$. The diagnostic procedures, performed at clinical diagnosis
21 and during follow-up, and clinical monitoring were carried out as already reported [22]. Moreover,
22 since the IPSS was developed for patients undergoing treatments not affecting the natural course of
23 MDS and since the MDS natural course may be significantly changed by intensive chemotherapy
24 and allogeneic bone marrow transplantation, patients submitted to these therapeutic procedures
25 were evaluated only until the time of such treatments.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 *Cytogenetic studies*

49 Chromosome studies were performed as previously reported [21]. Chromosome identification and
50 karyotype description were performed according to the International System for Chromosome
51 Nomenclature [37]. Karyotypes were defined as complex when they included ≥ 3 chromosomal
52 abnormalities.
53
54
55
56
57
58
59
60
61
62
63
64
65

Statistical analysis

1
2 Data were described as the mean and standard deviation (SD) or median and 25th-75th percentiles
3
4 (IQR) if continuous and described as a count and a percentage if categorical. The role of CE on
5
6 survival and MDS/AML evolution-free survival was assessed by Cox models for time-dependent
7
8 covariates. A bi-variable analysis was also performed, in order to adjust for known risk factors. In a
9
10 predefined subgroup analysis the following risk factors were investigated: cytopenias, blast cell
11
12 percentage, WHO classification, karyotype pattern at clinical diagnosis, and IPSS cytogenetic
13
14 categories. Time to event or to censoring was computed from diagnosis. The role of specific
15
16 secondary defects on survival and on MDL/AML evolution-free survival after CE was assessed by
17
18 Cox regression, with the time to event or censoring computed from CE. Hazard ratios (HR) and
19
20 their 95% confidence intervals were estimated. The Kaplan Meier method was used to compute the
21
22 cumulative event-free survival (PFI) and its 95% confidence interval (CI) at two and five years
23
24 from diagnosis (for all patients) or CE onset (for patients with CE).
25
26
27
28
29
30
31 Stata 10.1 (StataCorp, College Station, TX, USA) was used for computation. A 2-sided p-
32
33 value<0.05 was considered statistically significant.
34
35
36
37
38

Results

Patients

39
40
41 Patients' clinical and haematological features at clinical diagnosis are listed in table 1. Considering
42
43 all 153 patients, the median follow-up was 45.2 months (IQR=23.8-75.7), whereas the median
44
45 survival was not reached (IQR=34.3-not reached). At the time of analysis, 107 (69.9%) patients had
46
47 survived and 46 (30%) had died. Disease progression occurred in a total of 65 (42.4%) patients, 11
48
49 evolved to a more advanced MDS subtype and 54 to AML. The median progression-free interval
50
51 (PFI) was 65.2 months (IQR 16.60-not reached).
52
53
54
55
56
57

58 Treatment was as follows: 128 patients received supportive care (transfusions and hematopoietic
59
60 growth factors); 3 differentiation-inducing agents; 2 immunosuppressive agents; 5 low-dose
61
62
63
64
65

chemotherapy aimed at reducing WBC counts and 15 various regimens of intensive chemotherapy.

Twelve patients were submitted to allogeneic bone marrow transplantation (allo-BMT): 11 had previously received supportive treatment only, whereas one had been submitted to intensive chemotherapy which induced a partial remission. The decision to administer intensive chemotherapy or allo-BMT was exclusively based on either the patients' age or the fact that progression to a more advanced MDS or to AML had just occurred.

Since our patients were not uniformly treated, in order to avoid any effect of intensive chemotherapy or allo-BMT on our results, the 15 and 12 patients submitted to the respective procedures were censored at the start of both treatments. So, we are quite confident that, by excluding these patients, we did not affect the impact of CE on OS or PFI.

Cytogenetic results

At diagnosis, all 153 patients had successful cytogenetic analyses (tab. 1). Cytogenetic evolution occurred in a total of 47 (30.7%) patients. However, since six patients (one 5q- syndrome, one RA, one RCMD and 3 RAEB-2) developed secondary abnormalities after disease evolution, they were considered chromosomally stable when assessing whether CE was predictive of OS or PFI. Thus, a total of 40 (26.1%) patients acquired secondary abnormalities during the course of the disease, before disease progression. Their WHO subtype and blast cell percentage at clinical diagnosis and at cytogenetic re-evaluation are listed in table 2. In these patients, the median time from clinical diagnosis to the acquisition of secondary defects was 12.9 months (IQR=6.6-27.3). The secondary abnormalities included 5q- in four patients, monosomy 7 in three, del(7)(q31q35) in eight, trisomy 8 in five, del(17)(p13) in four and various defects (single and double abnormalities) in sixteen. The interstitial long arm deletion of chromosome 5 more commonly occurred in patients who did not experience disease progression, instead -7, del(7)(q31q35) and del(17p) more frequently developed in patients with disease progression. Patients surviving with or without CE according to various

1 parameters and those with either a stable disease or evolving to more advanced MDS/AML with or
2 without CE according to the same parameters are reported in table 3.
3
4
5

6 **Prognostic relevance of cytogenetic evolution**

7
8
9 The significant impact of cytogenetic evolution on OS and MDS/AML progression was analysed by
10 time-dependent Cox regression. Patients who experienced CE during the follow-up period increased
11 their risk of dying by seven-fold and their risk of disease progression (PFI) by thirty-six-fold with
12 respect to patients who remained karyotypically stable ($p < 0.001$ in both cases) (tab.4). Two- and 5-
13 year survivals were respectively 40% (95% CI: 19-60) and 10% (95% CI: 0.7-35) in patients with
14 CE, versus 93% (95% CI: 86-96) and 70% (95% CI: 58-70) in patients without CE. After adjusting
15 the effect of CE for other known risk factors (as listed in the methods section) in a bi-variable
16 analysis, the prognostic power of CE was proven in all cases for both death and PFI (tab.4).
17 Furthermore, CE had a statistically relevant influence on the OS of patients who at clinical
18 diagnosis were either chromosomally normal or abnormal ($p = 0.004$ and $p < 0.001$ respectively). In
19 addition, when the impact of CE on the risk of MDS/AML evolution was adjusted for the presence
20 of primary defects, which significantly effected PFI when analysed as a single variable (Hazard
21 ratio= 2.3 [85% CI= 1.07-4.94], $p = 0.02$), CE maintained its statistical power ($p < 0.001$), whereas
22 primary defects lost their statistical power ($p = 0.501$) (data not shown).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

61 Finally, the OS and the risk of MDS/AML evolution for each of the most common single
62 chromosomal defects acquired during the follow-up period before disease evolution were compared
63 to those of the remaining set of cases with secondary defects. This analysis showed that no single
64 secondary defect effected OS. In contrast, $\text{del}(7)(\text{q}31\text{q}34)$ was the only chromosomal abnormality
65 that significantly effected the risk of MDS/AML progression ($p = 0.002$), while trisomy 8 had only a
66 moderate effect ($p = 0.052$).

67 **Discussion**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

To the best of our knowledge, this is one of the few studies which evaluates the incidence of secondary chromosomal defects (CE) in an adequate number of MDS patients with a sufficient follow-up period of 45.2 months (IQR=23.8-75.7). These patients have been cytogenetically analysed at least twice before MDS/AML evolution. Former studies, which often included small numbers of patients with short follow-up periods, have established that CE occurs during the course of the disease in about 14-46% of MDS patients, more commonly in those with a sudden evolution to AML [27-35]. In a more recent study, 24% of MDS patients developed disease progression which, in the eight patients cytogenetically examined during transformation, was associated with CE [36]. None of these studies has evaluated the incidence of secondary defects in relation to other biological and clinical parameters. In the present series, 47 patients (30.7%) acquired additional chromosomal defects during the follow-up period, 40 of them (25.4%) before disease progression (tab. 2). The frequency of this event increased with the worsening of the MDS clinical stage (tab. 3). All these data confirm current evidence which suggest that MDS evolution is accompanied by a steady increase in genetic instability and by an expansion of genomic alterations [17, 38-39].

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

This is the first study which reveals that the acquisition of secondary chromosomal defects during the follow-up period before disease evolution predicts OS and PFI in MDS patients, an issue never addressed by previously published follow-up studies [27-36]. However, the clinical relevance of CE could be affected by the fact that the decision of performing a new bone marrow aspiration along with a cytogenetic re-evaluation might have been dictated by other clinical indicators of disease progression. These last include a worsening of anemia [16], neutropenia [40], thrombocytopenia [41], a sudden need for red blood cell (RBC) transfusions [24] and an abrupt increase in lactic dehydrogenase levels [42]. Recently, a study has evaluated the impact of cytopenias on MDS outcomes [43]. This study has established that in IPSS intermediate-1 and -2 categories, the haemoglobin value only, has an additive prognostic value regarding IPSS for evaluation of OS, but not for time to AML evolution. Another retrospective analysis has pointed out that in low-risk MDS the total number of RBC transfusions is an independent negative prognostic factor for OS and PFI

1 [24]. In the present study, the 40 patients who experienced CE before disease progression
2 underwent a cytogenetic re-assessment because of a reduction in peripheral blood values, especially
3 haemoglobin values. Thus, most of our patients presented a worsening of anemia that could be due
4 either to an increase in bone marrow apoptosis, the feature dissociating MDS from AML [17, 39],
5 or to an increase in marrow blast cell percentage. However, this last event was excluded. In fact, at
6 the time of CE no patient changed the subtype of MDS diagnosed at the onset of the disease (tab.
7 2).

8
9
10
11
12
13
14
15
16
17 In our series, patients who experienced CE presented a risk of dying seven-fold superior and a risk
18 of transformation to advanced MDS and AML thirty-six-fold superior than those of chromosomally
19 stable patients (tab. 4). However, since many studies have demonstrated that primary chromosomal
20 defects and other clinical-biological factors significantly effect MDS natural history [18-23], it was
21 questioned whether after adjusting for these last variables the statistical power of cytogenetic
22 evolution was maintained. In a first step, the bi-variable analysis applied revealed that the impact of
23 CE on OS was independent of the number and type of peripheral blood cytopenias, the blast cell
24 percentage, the WHO classification, the initial chromosomal pattern and IPSS cytogenetic
25 categories.

26
27
28
29
30
31
32
33
34
35
36
37
38
39 In the subsequent bi-variable analysis it was revealed that after adjusting for all other prognostic
40 variables, the relevance of CE on PFI was still significant (tab. 4). In particular, the acquisition of
41 secondary defects remained significant even after adjusting for the prognostic influence of the
42 initial chromosomal pattern ($p < 0.001$ in patients with either a normal or an abnormal karyotype at
43 clinical diagnosis), which impact on PFI was lost after performing this analysis. We speculate that
44 this effect could be due to the fact that 28 of the 94 patients (30%) with an abnormal karyotype at
45 clinical diagnosis acquired additional defects during the follow-up period (tab. 2).

46
47
48
49
50
51
52
53
54
55
56 As far as leukemic evolution is concerned, until now no study has established which numerical or
57 structural abnormality is required for such an event. It has been proposed that some chromosomal
58 alterations may be responsible for each step in the evolution of the disease, but since the pathway of
59
60
61
62
63
64
65

1 evolution is not unique, it is very difficult to define which genetic alteration comes first [36].
 2 However, this assumption seems to be disproved by a recent array comparative genomic
 3 hybridization study which revealed that in low-risk MDS, CD34 negative cells harbour genetic
 4 alterations in addition to those present in CD34 positive cells [44]. Thus, disease progression might
 5 be caused by the outgrowth of a clonal CD34 negative cell population already present at clinical
 6 diagnosis and by the type of genetic lesion present in these dysplastic cells. This last suggestion is
 7 also strengthened by our study. In fact, in our series, the majority of secondary defects revealed in
 8 early/low-risk MDS were distinct from those revealed in advanced MDS: 5q- and del(11)(q14q23)
 9 prevailed in patients with early disease, -7, del(7)(q31q34) and del(17p) in those with advanced
 10 disease. Trisomy 8 was equally distributed. Thus, it is not surprising that del(7)(q31q34) was the
 11 only secondary defect significantly effecting the risk of MDS/AML evolution.

12 In conclusion, i) CE occurs in 30.7% of patients, particularly in those with more advanced MDS; ii)
 13 it has a significant influence on OS and PFI independently of other variables with well-known
 14 prognostic relevance; iii) del(7)(q31q34) is the only single secondary chromosomal defect which
 15 significantly effects PFI, whereas trisomy 8 has a moderate influence.

36 References

- 37 1. Heaney ML, Golde DW (1999) Myelodysplasia. *N Engl J Med* 340:1649-1660.
- 38 2. Nimer SD (2008) Myelodysplastic syndromes. *Blood* 11:4841-4851.
- 39 3. Jaffe ES, Harris NL, Stein H, Vardiman JW (Eds.) (2008) Tumours of Haematopoietic and
 40 Lymphoid Tissues. World Health Organization Classification of Tumours. IARC Press,
 41 Lyon.
- 42 4. Germing U, Gattermann N, Strupp C, Aivado M, Aul C (2000) Validation of the WHO
 43 proposal for a new classification of primary myelodysplastic syndromes: a retrospective
 44 analysis of 1600 patients. *Leuk Res* 24:983-992.
- 45 5. Nösslinger T, Reisner R, Koller E, Grüner H, Tüchler H, Nowotny H et al (2001)
 46 Myelodysplastic syndromes, from the French-American-British to World Health
 47 Organization: comparison of classifications on 431 unselected patients from a single
 48 institution. *Blood* 98:2935-2941.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
6. Lee J-H, Lee J-H, Shin J-R, Kim W-K, Chi H-S, Park C-J et al (2003) Application of different prognostic scoring systems and comparison of the FAB and WHO classification in Korean patients with myelodysplastic syndromes. *Leukemia* 17:305-313.
7. Howe B, Porwit-MacDonald A, Wanat R, Tehranchi R, Hellström-Lindberg E (2004) The WHO classification of MDS does make a difference. *Blood* 103:3265-3270.
8. Malcovati L, Della Porta MG, Pascutto C, Invernizzi R, Boni M, Travaglini E et al (2005) Prognostic factors and life expectancy in myelodysplastic syndromes classified according to WHO criteria: a basis for clinical decision making. *J Clin Oncol* 23:7594-7603 .
9. Strupp C, Gattermann N, Giagounidis A, Aul C, Hildebrandt B, Haas R, Germing U (2003) Refractory anemia with excess of blasts in transformation: analysis of reclassification according to the WHO proposals. *Leuk Res* 27:397-404.
10. Germing U, Strupp C, Kuendgen A, Aivado M, Giagounidis A, Hildebrandt B et al (2006a) Refractory anemia with excess of blasts (RAEB): analysis of reclassification according to the WHO proposals. *B J Haematol* 132:162-167.
11. Germing U, Strupp C, Kuendgen A, Isa S, Knipp S, Hildebrandt B et al (2006b) Prospective validation of the WHO proposals for the classification of myelodysplastic syndromes. *Haematologica* 91:1596-1604.
12. Valent P, Horny HP, Bennet JM, Fonatsch C, Germing U, Greenberg P et al (2007) Definitions and standards in the diagnosis and treatments of the myelodysplastic syndromes: Consensus statements and report from a working conference. *Leuk Res* 31:727-736.
13. Mufti GJ, Bennet JM, Goasguen J, Bain BJ, Baumann I, Brunning R et al (2008) Diagnosis and classification of myelodysplastic syndromes: International Working Group on Morphology of Myelodysplastic Syndrome. *Haematologica* 93:1712-1717.
14. Breccia M, Latagliata R, Cannella L, Carmosino I, De Cuia R, Frustaci A et al (2009) Analysis of prognostic factors in patients with refractory anemia with excess of blasts (RAEB) reclassified according to WHO proposal. *Leuk Res* 33:391-394.
15. Jädersten M, Hellström-Lindberg E (2008) Myelodysplastic syndromes: biology and treatment. *J Int Med* 265:307-328.
16. Cazzola M, Malcovati L (2005) Myelodysplastic syndromes – Coping with ineffective hematopoiesis. *N Engl J Med* 352:536-538.
17. Corey SJ, Minden MD, Barber DL, Kantarjian H, Wang JCY, Schimmer AD (2007) Myelodysplastic syndromes: the complexity of stem-cell diseases. *Nature Rev Cancer* 7:118-129.
18. Greenberg P, Cox C, Le Beau MM, Fenaux P, Morel P, Sanz G et al (1987) International system for evaluating prognosis in myelodysplastic syndromes. *Blood* 89:2079-2088.
19. Solè F, Espinet B, Sanz G, Cervera J, Calasanz MJ, Luño E et al (2000) Incidence, characterization and prognostic significance of chromosomal abnormalities in 640 patients with primary myelodysplastic syndromes. *B J Haematol* 108:346-356.

- 1 20. Solè F, Luño E, Sanzo C, Espinet B, Sanz GF, Cervera J et al (2005) Identification of novel
2 cytogenetic markers with prognostic significance in a series of 968 patients with primary
3 myelodysplastic syndromes. *Haematologica* 90:1168-1178.
- 4
5
6 21. Bernasconi P, Klersy C, Boni M, Cavigliano PM, Giardini I, Rocca B et al (2005) Incidence
7 and prognostic significance of karyotype abnormalities in de novo primary myelodysplastic
8 syndromes: a study on 331 patients from a single institution. *Leukemia* 19:1424-1431.
- 9
10
11 22. Bernasconi P, Klersy C, Boni M, Cavigliano PM, Calatroni S, Giardini I et al (2007) World
12 Health Organization classification in combination with cytogenetic markers improves the
13 prognostic stratification of patients with de novo primary myelodysplastic syndromes. *B J*
14 *Haematol* 137:193-205.
- 15
16
17 23. Haase D, Germing U, Schanz J, Pfeilstöcker M, Nösslinger T, Hildebrandt B et al (2007)
18 New insights into the prognostic impact of the karyotype in MDS and correlation with
19 subtypes: evidence from a core data set of 2124 patients. *Blood* 110:4385-4395.
- 20
21
22 24. Malcovati L, Germing U, Kuendgen A, Della Porta MG, Pascutto C, Invernizzi R et al
23 (2007) Time-dependent prognostic scoring system for predicting survival and leukemic
24 evolution in myelodysplastic syndromes. *J Clin Oncol* 25:3503-3510.
- 25
26
27 25. Olney HJ, Le Beau MM (2001) The cytogenetics of myelodysplastic syndromes. *Best Pract*
28 *Res Clin Hematol* 14:479-495.
- 29
30
31 26. Bernasconi P, Boni M, Cavigliano PM, Calatroni S, Giardini I, Rocca B et al (2006) Clinical
32 relevance of cytogenetics in myelodysplastic syndromes. *Ann NY Acad Sci* 1089:395-410.
- 33
34
35 27. Tricot G, Boogaerts MA, de Wolf-Peeters C, Van den Berghe H, Verwilghen RL (1985) The
36 myelodysplastic syndromes: different evolution patterns based on sequential morphological
37 and cytogenetic investigations. *B J Haematol* 59:659-670.
- 38
39
40 28. Benitez J, Carbonell F, Sanchez Fayos J, Heimpel H (1985) Karyotypic evolution in patients
41 with myelodysplastic syndromes. *Cancer Gen Cytogen* 15:157-167.
- 42
43
44 29. Horiike S, Taniwaki M, Misawa S, Abe T (1988) Chromosome abnormalities and karyotype
45 evolution in 83 patients with myelodysplastic syndrome and predictive value for prognosis.
46 *Cancer* 62:1129-1138.
- 47
48
49 30. Suciú S, Kuse R, Weh HJ, Hossfeld DK (1990) Results of chromosome studies and their
50 relation to morphology, course, and prognosis in 120 patients with de novo myelodysplastic
51 syndrome. *Cancer Gen Cytogen* 44:15-26.
- 52
53
54 31. Michalova K, Musilova J, Zemanova Z (1991) Consecutive chromosomal studies in patients
55 with myelodysplastic syndrome (MDS). Czechoslovak MDS Cooperative Group. *Ann Genet*
56 34:212-218.
- 57
58
59 32. White AD, Culligan DJ, Hoy TG, Jacobs A (1992) Extended cytogenetic follow-up of
60 patients with myelodysplastic syndrome (MDS). *B J Haematol* 81:499-502.
- 61
62
63
64
65

- 1 33. Ghaddar HM, Stass SA, Pierce S, Estey EH (1994) Cytogenetic evolution following the
2 transformation of myelodysplastic syndrome to acute myelogenous leukaemia: implications
3 on the overlap between the two diseases. *Leukemia* 8:1649-1653.
- 4 34. White AD, Hoy TG, Jacobs A (1994) Extended cytogenetic follow-up and clinical progress
5 in patients with myelodysplastic syndromes (MDS). *Leuk Lymph* 12:401-412.
- 6 35. Tien HF, Wang CH, Chuang SM, Lee FY, Liu MC, Chen YC et al (1995) Acute leukaemic
7 transformation of myelodysplastic syndrome-immunophenotypic, genotypic, and
8 cytogenetic studies. *Leuk Res* 19:595-603.
- 9 36. De Souza Fernandez T, Ornellas MH, de Carvalho LO, Tabak D, Abdelhay E (2000)
10 Chromosomal alterations associated with evolution from myelodysplastic syndrome to acute
11 myeloid leukaemia. *Leuk Res* 24:839-848.
- 12 37. Mitelman F. (Eds.) *ISCN: An International System for Human Cytogenetic Nomenclature*
13 (1995) S Karger, Basel.
- 14 38. Novotna B, Neuwirtova R, Siskova M, Bagryantseva Y (2008) DNA instability in low-risk
15 myelodysplastic syndromes: refractory anemia with or without ring sideroblasts. *Hum Mol*
16 *Genet* 17:2144-2149.
- 17 39. Bernasconi P. (2008) Molecular pathways in myelodysplastic syndromes and acute myeloid
18 leukaemia: relationship and distinctions-a review. *Brit J Haematol* 142:695-708.
- 19 40. Pomeroy C, Oken MM, Rydell RE, Filice GA (1991) Infection in the myelodysplastic
20 syndromes. *Am J Med* 90:338-344.
- 21 41. Kantarjian H, Giles F, List A, Lyons R, Sekeres MA, Pierce S, et al. (2007) The incidence
22 and impact of thrombocytopenia in myelodysplastic syndromes. *Cancer* 109:1705-1714.
- 23 42. Germing U, Hildebrandt B, Pfeilstöcker M, Nösslinger T, Valent P, Fonatsch C, et al (2005)
24 Refinement of the International Prognostic Scoring System (IPSS) by including LDH as an
25 additional scoring variable to improve risk assessment in patients with primary
26 myelodysplastic syndromes (MDS). *Leukemia* 19:2223-2231.
- 27 43. Kao JM, McMillan A, Greenberg P. (2008) International MDs risk analysis workshop
28 (IMRAW)/IPSS reanalyzed: impact of cytopenias on clinical outcomes in myelodysplastic
29 syndromes. *Am J Hematol* 83:765-770.
- 30 44. Starczynowski DT, Vercauteren S, Telenius A, Sung S, Tohyama K, Brooks-Wilson A et al
31 (2008) High-resolution whole genome tiling path array CGH analysis of CD34+ cells from
32 patients with low-risk myelodysplastic syndromes reveals cryptic copy number alterations
33 and predicts overall and leukemia-free survival. *Blood* 112:3412-3424.

Tab.1: Patients' clinical, hematological and cytogenetic features at clinical diagnosis.

Total of patients	153 (%)
Median age, years (Inter-quartile range)	60.5 (49.8-68.3)
Sex	
Males	80 (52.3)
Females	73 (47.7)
Cytopenias	
None	14 (9.1)
One	74 (48.3)
Two	45 (29.4)
Three	20 (13.0)
Marrow blast cell percentage	
<5	83 (54.2)
5-10	27 (17.6)
11-20	33 (21.5)
21-30	10 (6.5)
WHO classification	
RARS / RA	17 (11.9) / 21 (14.8)
MDS del(5q)	11 (7.7)
RCMDS / RCMD	7 (4.9) / 27 (19.0)
RAEB-1 / RAEB-2	27 (19.0) / 31 (21.8)
MDS-U	1
AML*	11 (7.2)
Karyotype	
Normal	59 (38.5)
Abnormal	94 (61.4)
Chromosomal defects	
Single	75 (49.0)
Double	14 (9.1)
Complex (with ≥ 3 defects)	5 (3.2)
Chromosomal defects in each WHO subtype	
RARS / RA	6 (35.3) / 8 (39.0)
MDS del(5q)	11 (100)
RCMDS / RCMD	5 (71.4) / 14 (51.8)
RAEB-1 / RAEB-2	19 (70.3) / 22 (70.9)
AML*	9 (81.8)
IPSS cytogenetic categories	
Good	85 (55.5)
Intermediate	49 (32.0)
Poor	19 (12.4)

WHO classification, World Health Organization classification; RARS, RA with ringed sideroblasts; RA, refractory anemia; MDS del(5q), 5q- syndrome; RCMD, refractory cytopenia with multilineage dysplasia; RCMDS, RCMD with ringed sideroblasts; RAEB-1, RA with excess of blasts type 1; RAEB-2, RAEB type 2; MDS-U, unclassifiable MDS; * classified as RAEB in transformation by the previously applied FAB classification.

Tab.2: Clinical and cytogenetic features of the forty patients who experienced cytogenetic evolution (CE)

Clinical and cytogenetic characteristics	At clinical diagnosis / At CE*
Cytopenias	
None	2 (5.0) / 1 (2.5)
One	16 (40.0) / 6 (15.0)
Two	15 (37.5) / 16 (40.0)
Three	7 (17.5) / 17 (42.5)
Blast cell percentage	
<5	13 (32.5) / 13 (32.5)
5-10	16 (40.0) / 11 (27.5)
11-20	10 (25.0) / 15 (37.5)
21-30	1 (2.5) / 1 (2.5)
WHO classification	
RARS	1 (2.5) / 1 (2.5)
RA	6 (15.0) / 6 (15.0)
MDS del(5q)	1 (2.5) / 1 (2.5)
RCMDS	- / -
RCMD	5 (12.5) / 5 (12.5)
RAEB-1	7 (17.5) / 7 (17.5)
RAEB-2	15 (37.5) / 15 (37.5)
MDS-U	- / -
AML ^a	5 (12.5) / 5 (12.5)
Karyotype	
Normal	12 (30.0) / -
Abnormal	28 (70.0) / 40 (100.0)

*= number of patients with percentages in parentheses; ^a= classified as RAEB in transformation by the previously applied FAB classification.

Tab.3: Survival and disease progression in relation to CE according to WHO classification, karyotype and IPSS cytogenetic categories

	Survived without CE	Survived with CE	In disease prog. without CE	In disease prog. with CE	Tot. of pts
Pts [‡]	86/113 (76.1)	21/40 (52.5)	34/113 (30.1)	31/40 (77.5)	153
WHO [‡] :					
RARS	16/16 (100)	1/1 (100)	—	—	17
RA	15/15 (100)	5/6 (83.3)	5/15 (33.3)	1/6 (16.7)	21
5q- Syn.	8/10 (80.0)	1/1 (100)	2/10 (20)	1/1 (100)	11
RCMDS	5/7 (71.4)	—	1/7 (14.3)	—	7
RCMD	17/22 (77.3)	2/5 (40.0)	5/22 (22.7)	4/5 (80.0)	27
RAEB-1	11/20 (55.0)	3/7 (42.9)	6/20 (30.0)	6/7 (85.7)	27
RAEB-2	9/16 (56.2)	7/15 (46.7)	10/16 (62.5)	14/15 (93.3)	31
AML*	4/6 (66.7)	2/5 (40.0)	5/6 (83.3)	5/5 (100)	11
Karyotype [‡]					
Normal	41/47 (87.2)	8/12 (66.7)	9/47 (19.1)	9/12 (75.0)	59
Abnormal	45/66 (68.2)	13/28 (46.4)	25/66 (37.8)	22/28 (78.5)	94
IPSS cyto. cat. [‡]					
Good	58/67 (86.6)	10/18 (55.6)	15/67 (22.4)	15/18 (83.3)	85
Intermediate	25/37 (67.6)	8/12 (66.7)	15/37 (40.5)	7/12 (58.3)	49
Poor	3/9 (33.3)	3/10 (30.0)	4/9 (44.4)	9/10 (90.0)	19

[‡] Numbers and percentages in parentheses represent the true numbers and true percentages of patients either surviving or experiencing disease progression over the total number of patients included in each category analysed; WHO, World Health Organization; RARS, refractory anemia with ringed sideroblasts; RA, refractory anemia; MDS del(5q), 5q- syndrome; RCMDS, refractory cytopenia with multilineage dysplasia with ringed sideroblasts; RCMD, RCMD refractory cytopenia with multilineage dysplasia; RAEB-1, RA with excess of blasts type 1; RAEB-2, RAEB type 2; * classified as RAEB in transformation by the previously applied FAB classification.

Table 4

Tab. 4: Cytogenetic evolution (CE) predicts OS and PFI in univariable analysis and maintains its relevance independently of other well-known prognostic parameters in adjusted bivariable analyses.

	Survival						MDS/AML evolution			
	Tot. Pts ^a (%)	Events ^b (%)	2-year surv. (95% CI)	5-year surv. (95% CI)	HR (95% CI)	P-value	Tot. Pts ^a (%)	Events ^b (%)	HR (95% CI)	P-value
Cytogenetic evolution:										
No	113 (100)	27 (24)	93 (86-96)	70 (58-79)	–	–	113 (100)	34 (30)	–	–
Yes	40 (100)	19 (47)	40 (19-60)	10 (0.7-35)	7.1 (3.92-12.86)	<0.001	40 (100)	31 (77)	35.8 (20.68-62.13)	<0.001
Cytogenetic evolution adjusted for	Survival					MDS/AML evolution				
	HR (95% CI)					P-value				
Cytopenias	6.4 (3.54-11.77)					<0.001				
Blast cell percentage	4.4 (2.34-8.60)					<0.001				
WHO classification	4.0 (1.96-8.26)					<0.001				
Chrom. pattern at clin. diagn.	6.6 (3.65-12.06)					<0.001				
IPSS cytogen. categories	6.6(3.65-12.2)					<0.001				

HR= Hazard Ratio; 95% CI= 95% confidence intervals; ^a the percentages are calculated for total number of patients; ^b the percentages are calculated for total number of patients included in each category; WHO, World Health Organization.

Replies to Reviewers' comments

As requested:

1. FAB classification has been completely omitted from the manuscript (text and tables). However, in table 1 it has been underlined that 11 of the 153 patients, classified as RAEB-t according to FAB classification, were classified as AML according to WHO. Furthermore, in table 2 and 3 it is underlined that 5 of the 40 patients who experienced cytogenetic evolution were classified as RAEB-t according to FAB but as AML according to WHO.
2. Peripheral blood counts have been completely omitted from the tables.
3. The WHO classification at diagnosis and cytogenetic evolution with percentage of marrow blasts has been provided (table 2).
4. IPSS has been deleted from the tables and the manuscript has concentrated on cytogenetic evaluation
5. The number of tables has been reduced to four. In particular:
The previous tab.2 has been fused to tab.1; the previous tab.3 has been omitted and substituted with the new tab.2 which reports the clinical and cytogenetic features of the 40 patients who experienced CE; the previous tables 4 and 5 has been fused together in the new table 3 which maintains their relevant information; the previous table 6, now table 4, has been maintained; the previous tables 7 and 8 has been omitted.