



HAL
open science

Percolation in a multiscale Boolean model

Jean-Baptiste Gouéré

► **To cite this version:**

| Jean-Baptiste Gouéré. Percolation in a multiscale Boolean model. 2010. hal-00519310v1

HAL Id: hal-00519310

<https://hal.science/hal-00519310v1>

Preprint submitted on 19 Sep 2010 (v1), last revised 9 Mar 2011 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Percolation in a multiscale Boolean model

Jean-Baptiste Guéré *

Abstract

We consider percolation in a multiscale Boolean model. This model is defined as the union of scaled independent copies of a given Boolean model. The scale factor of the n^{th} copy is ρ^{-n} . Under optimal integrability assumptions, we prove that no percolation occurs in the multiscale Boolean model for large enough ρ if the rate of the Boolean model is below some critical value.

1 Introduction and statement of the main result

1.1 The Boolean model

Let $d \geq 2$. Let μ be a finite measure on $]0, +\infty[$. We assume that the mass of μ is positive. Let ξ be a Poisson point process on $\mathbb{R}^d \times]0, +\infty[$ whose intensity is the product of the Lebesgue measure on \mathbb{R}^d by μ . With ξ we associate a random set $\Sigma(\mu)$ defined as follows:

$$\Sigma(\mu) = \bigcup_{(c,r) \in \xi} B(c,r)$$

where $B(c,r)$ is the open Euclidean ball of radius r centered at c . The random set $\Sigma(\mu)$ is the Boolean model with parameter μ . We shall sometimes write Σ to simplify the notations.

The following description may be more intuitive. Let χ denote the projection of ξ on \mathbb{R}^d . With probability one this projection is one-to-one. We can therefore write:

$$\xi = \{(c, r(c)), c \in \chi\}.$$

Write $\mu = m\nu$ where ν is a probability measure. Then, χ is a Poisson point process on \mathbb{R}^d with density m . Moreover, given χ , the sequence $(r(c))_{c \in \chi}$ is a sequence of independent random variable with common distribution ν . We shall not use this point of view.

1.2 Percolation in the Boolean model

Let C denote the connected component of Σ that contains the origin. We say that Σ percolates if C is unbounded with positive probability. We refer to the book by Meester and Roy [9] for background on continuum percolation. Set:

$$\lambda_c(\mu) = \inf\{\lambda > 0 : \Sigma(\lambda\mu) \text{ percolates}\}.$$

**Postal address:* Université d'Orléans MAPMO B.P. 6759 45067 Orléans Cedex 2 France *E-mail:* jbgouere@univ-orleans.fr

One easily check that $\lambda_c(\mu)$ is finite as soon as μ has a positive mass. In [4] we proved that $\lambda_c(\mu)$ is positive if and only if:

$$\int r^d \mu(dr) < \infty.$$

The only if part had been proved earlier by Hall [7]. For all $A, B \subset \mathbb{R}^d$, we write $A \leftrightarrow_{\Sigma} B$ if there exists a path in Σ from A to B . We denote by $S(c, r)$ the Euclidean sphere of radius r centered at the c :

$$S(c, r) = \{x \in \mathbb{R}^d : \|x - c\|_2 = r\}.$$

We write $S(r)$ when $c = 0$.

The critical parameter $\lambda_c(\mu)$ can also be defined as follows:

$$\lambda_c(\mu) = \sup \{ \lambda > 0 : P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \},$$

We shall need two other critical parameters:

$$\begin{aligned} \widehat{\lambda}_c(\mu) &= \sup \{ \lambda > 0 : P(S(r/2) \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \}, \\ \widetilde{\lambda}_c(\mu) &= \sup \{ \lambda > 0 : r^d P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \}. \end{aligned}$$

We have (see Lemma 10):

$$\widetilde{\lambda}_c(\mu) \leq \widehat{\lambda}_c(\mu) \leq \lambda_c(\mu). \quad (1)$$

When the support of μ is bounded,

$$P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r))$$

decays exponentially fast to 0 as soon as $\lambda < \lambda_c(\mu)$ (see for example [9], Section 12.10 in [6] when $R = 1$ or the papers [10], [13], [18] and [19]). Therefore:

$$\widetilde{\lambda}_c(\mu) = \widehat{\lambda}_c(\mu) = \lambda_c(\mu) \text{ as soon as the support of } \mu \text{ is bounded.} \quad (2)$$

Remark. By results and ideas of [4], we can prove:

- $\widehat{\lambda}_c(\mu)$ is positive if and only if $\int x^d \mu(dx)$ is finite (i.e., if and only if $\lambda_c(\mu)$ is positive).
- $\widetilde{\lambda}_c(\mu)$ is positive if and only if $r^d \int_r^\infty x^d \mu(dx) \rightarrow 0$ as $r \rightarrow \infty$.

1.3 A multiscale Boolean model

Let $\rho > 1$ be a scale factor. Let $(\Sigma_n)_{n \geq 0}$ be a sequence of independent copies of $\Sigma(\mu)$. In this paper, we are interested in percolation properties of the following multiscale Boolean model:

$$\Sigma^\rho(\mu) = \bigcup_{n \geq 0} \rho^{-n} \Sigma_n. \quad (3)$$

We shall sometimes write Σ^ρ to simplify the notations. As before, we say that Σ^ρ percolates if the connected component of Σ^ρ that contains the origin is unbounded with positive probability.

This model seems to have been first introduced as a model of failure in geophysical medias in the 80'. We refer to the paper by Molchanov, Pisarenko and Reznikova [14] for an account of those studies. For more recent results we refer to [2], [9], [10], [11], [12] and [16].

This model is related to a discrete model introduced by Mandelbrot [8]. We refer to the survey by L. Chayes [3] and, for more recent results, to [1], [15] and [17].

In [11], Menshikov, Popov and Vachkovskaia considered the case where the radii of the unscaled process Σ_0 equal 1. They proved the following result.

Theorem 1 ([11]) *If $\lambda < \lambda_c(\delta_1)$ then, for all large enough ρ , $\Sigma^\rho(\lambda\mu)$ does not percolate.*

In [12] the same authors considered the case where the radii are random and can be unbounded. They considered the following sub-autosimilarity assumption on the measure μ :

$$\lim_{a \rightarrow \infty} \sup_{r \geq 1/2} \frac{a^d \mu([ar, +\infty[)}{\mu([r, +\infty[)} = 0 \quad (4)$$

with the convention $0/0 = 0$. They proved the following result.

Theorem 2 ([12]) *Assume that the measure μ satisfies (4). If $\lambda < \tilde{\lambda}_c(\mu)$ then, for all large enough ρ , $\Sigma^\rho(\lambda\mu)$ does not percolate.*

Note that (4) is fulfilled for any measure with bounded support. Because of (2), Theorem 2 is then a generalization of Theorem 1.

In this paper, we prove the following result.

Theorem 3 *Assume:*

$$\int_{[1, +\infty[} \beta^d \ln(\beta) \mu(d\beta) < \infty. \quad (5)$$

If $\lambda < \hat{\lambda}_c(\mu)$, then for all large enough ρ , $\Sigma^\rho(\lambda\mu)$ does not percolate.

Condition (5) is a necessary condition, as shown by the following result from [5].

Theorem 4 *Let $\rho > 1$. There exists $\lambda > 0$ such that $\Sigma^\rho(\lambda\mu)$ does not percolate if and only if (5) holds.*

Theorem 3 is a generalization of Theorem 2 and thus of Theorem 1. Indeed, by (1), one has $\lambda < \hat{\lambda}_c$ as soon as $\lambda < \tilde{\lambda}_c$. Moreover, (5) is implicitly assumed in the statement of Theorem 2. This can be seen as a consequence of Theorem 4. Alternatively, one can check that (5) is a consequence of $\tilde{\lambda}_c(\mu) > 0$ or of (4).

2 Proof

2.1 Some notations

In the whole of Section 2, we make the following assumptions:

- μ satisfies (5).
- $1 < \hat{\lambda}_c(\mu)$.

We aim at proving that, for large enough ρ , $\Sigma^\rho(\mu)$ does not percolate. Theorem 3 follows by applying this result to the measure $\lambda\mu$.

For all $\eta > 0$, we denote by $T_\eta\mu$ the measure defined by $T_\eta\mu(A) = \mu(A - \eta)$. In other words, we can build $\Sigma(T_\eta\mu)$ from $\Sigma(\mu)$ by adding η to each radius.

For all $\rho > 1$, we denote by $H^\rho\mu$ the measure defined by $H^\rho\mu(A) = \rho^d\mu(\rho A)$. With this definition, $\rho^{-1}\Sigma(\mu)$ is a Boolean model driven by the measure $H^\rho\mu$. For all $n \geq 0$, we let:

$$m_n^\rho = \sum_{k=0}^n H^{\rho^k}\mu.$$

With this definition and the notations of (3),

$$\bigcup_{k=0}^n \rho^{-k}\Sigma_k$$

is a Boolean model driven by m_n^ρ . We also let:

$$m_\infty^\rho = \sum_{k \geq 0} H^{\rho^k}\mu.$$

So, $\Sigma^\rho(\mu)$ is a Boolean model driven by the locally finite measure m_∞^ρ .

Let $p(a, \mu)$ denote the probability of existence of a path from $S(a/2)$ to $S(a)$ in $\Sigma(\mu)$:

$$p(a, \mu) = P(S(a/2) \leftrightarrow_{\Sigma(\mu)} S(a)).$$

2.2 Ideas

In this subsection we first sketch the proof of the existence of ρ and a such that $p(a, m_\infty^\rho)$ is small. This gives the main ingredients of the proof of Theorem 3. A full proof is given in Subsection 2.3. We then give the ideas of the proof of Theorems 1 and 2 by Menshikov, Popov and Vachkovskaia. Their basic strategy is similar but the implementation of the proof is different.

Sketch of the proof of Theorem 3

Consider a small $\varepsilon_1 > 0$. Fix a small $\eta > 0$ and a large a such that (see Lemma 5):

$$p(a, T_\eta\mu) \leq \varepsilon_1/2. \tag{6}$$

For all $n \geq 1$, write:

$$m_n^\rho = H^\rho m_{n-1}^\rho + \mu.$$

If the event $S(a/2) \leftrightarrow_{\Sigma(m_n^\rho)} S(a)$ occurs, then either the event $S(a/2) \leftrightarrow_{T_\eta\mu} S(a)$ occurs (with a natural coupling between the Boolean models) either in $\Sigma(H^\rho m_{n-1}^\rho) \cap B(a)$ one can find a component of diameter at least η . We use this observation through its following crude consequence (see Lemma 6):

$$p(a, m_n^\rho) \leq p(a, T_\eta\mu) + C a^d \eta^{-d} p(\eta/2, H^\rho m_{n-1}^\rho).$$

By scaling and by (6), this yields:

$$p(a, m_n^\rho) \leq \varepsilon_1/2 + Ca^d \eta^{-d} p(\rho\eta/2, m_{n-1}^\rho). \quad (7)$$

But for any ε_2 , any small enough ε_1 and any large enough a we can find τ such that (see Lemmas 7 and 8):

$$p(\tau a, m_{n-1}^\rho) \leq \varepsilon_2 \text{ as soon as } p(a, m_{n-1}^\rho) \leq \varepsilon_1. \quad (8)$$

An important fact is that τ does not depend on n nor on ρ , provided $\rho \geq \rho_0$ where ρ_0 is an arbitrary constant strictly larger than 1. Here we use assumption (5) to bound error terms due to the existence of large balls.

We choose ε_2 such that:

$$Ca^d \eta^{-d} \varepsilon_2 = \varepsilon_1/2.$$

We set $\rho = 2\tau a/\eta$. Then, (7) and (8) can be rewritten as follows:

$$p(a, m_n^\rho) \leq \varepsilon_1/2 + Ca^d \eta^{-d} p(\tau a, m_{n-1}^\rho) \quad (9)$$

$$Ca^d \eta^{-d} p(\tau a, m_{n-1}^\rho) \leq \varepsilon_1/2 \text{ as soon as } p(a, m_{n-1}^\rho) \leq \varepsilon_1. \quad (10)$$

As moreover (6) implies $p(a, m_0^\rho) \leq \varepsilon_1$ we get, by induction and then sending n to infinity (see Lemma 9):

$$p(a, m_\infty^\rho) \leq \varepsilon_1.$$

Sketch of the proofs of Theorems 1 and 2 by Menshikov, Popov and Vachkovskaia

Let us quickly describe the ideas of the proofs of Menshikov, Popov and Vachkovskaia. Those ideas are used in their papers [11] and [12] through a discretization of space ; we describe them in a slightly more geometric way. For simplicity we only consider two scales: $\rho^{-1}\Sigma_1$ and Σ_0 . Assume that C is a connected component of $\rho^{-1}\Sigma_1 \cup \Sigma_0$ whose diameter is at least α (it can be much larger) for a small enough constant $\alpha > 0$. Then, C is included in the union of the following kind of sets:

1. connected components of $\rho^{-1}\Sigma_1$ whose diameter is at least α ;
2. balls of Σ_0 enlarged by the factor $1 + \alpha$ (same centers but the radii are $1 + \alpha$ times the radii of Σ_0).

Then, they show that the union of all those sets is stochastically dominated by a Boolean model similar to Σ_0 but with radii enlarged by a factor $1 + \alpha$ and with density of centers $1 + \alpha'$ times the corresponding density for Σ_0 for a suitable $\alpha' > 0$. This part uses $\lambda < \widehat{\lambda}_c$. In some sense, one can therefore control percolation in the union of two models by percolation in one model. Iterating the argument with some care in the constants α and α' , one sees that – for large enough ρ – one can control percolation in the multiscale model by percolation in a subcritical model. This yields the result.

2.3 Proof

As $1 < \widehat{\lambda}_c(\mu)$, we know that $p(a, \mu)$ tends to 0 as a tends to infinity. We need the following slightly stronger consequence.

Lemma 5 *There exists $\eta > 0$ such that $p(a, T_\eta \mu)$ tends to 0.*

Proof. Let $\varepsilon > 0$ and $x > 0$. We have:

$$\begin{aligned} H^{1+\varepsilon} T_{\varepsilon^2} \mu([x, +\infty[) &= (1 + \varepsilon)^d T_{\varepsilon^2} \mu([x(1 + \varepsilon), +\infty[) \\ &= (1 + \varepsilon)^d \mu([x(1 + \varepsilon) - \varepsilon^2, +\infty[) \\ &\leq \kappa(\varepsilon)(1 + \varepsilon)^d \mu([x, +\infty[) \end{aligned} \quad (11)$$

where

$$\kappa(\varepsilon) = \frac{\mu(]0, +\infty[)}{\mu(] \varepsilon, +\infty[)}.$$

The inequality is proven as follows. If $x \geq \varepsilon$, then $[x(1 + \varepsilon) - \varepsilon^2, +\infty[\subset [x, +\infty[$ and the result follows from $\kappa(\varepsilon) \geq 1$. If, on the contrary, $x < \varepsilon$, then the left hand side is bounded above by $(1 + \varepsilon)^d \mu(]0, +\infty[)$ which is itself bounded above by the right hand side.

Note that $\kappa(\varepsilon)(1 + \varepsilon)^d$ tends to 1 as ε tends to 0. Let us say that a measure ν is subcritical if $\widehat{\lambda}_c(\nu) > 1$. As μ is subcritical, we get that $\kappa(\varepsilon)(1 + \varepsilon)^d \mu$ is subcritical for small enough ε . We fix such an ε . By (11) we can couple a Boolean model driven by $H^{1+\varepsilon} T_{\varepsilon^2} \mu$ and a Boolean model driven by $\kappa(\varepsilon)(1 + \varepsilon)^d \mu$ in such a way that the first one is contained in the second one. Therefore the first one is subcritical. By scaling, a Boolean model driven by $T_\eta \mu$ is then subcritical. We take $\eta = \varepsilon^2$. \square

Lemma 6 *Let ν_1 and ν_2 be two finite measures on $]0, +\infty[$. One has, for all $\eta > 0$ and $a \geq 4\eta$:*

$$p(a, \nu_1 + \nu_2) \leq p(a, T_\eta \nu_1) + C_1 a^d \eta^{-d} p(\eta/2, \nu_2)$$

where $C_1 = C_1(d) > 0$ depends only on the dimension d .

Proof. Let $(x_i)_{i \in I}$ be a family of points such that :

- The balls $B(x_i, \eta/4)$, $i \leq I$, cover $B(a)$.
- There are at most $C_1 a^d \eta^{-d}$ points in the family where $C_1 = C_1(d)$ depends only on the dimension d .

We couple the different Boolean model as follows. Let $\Sigma(\nu_1)$ be a Boolean model driven by ν_1 . Let $\Sigma(\nu_2)$ be a Boolean model driven by ν_2 . Assume that $\Sigma(\nu_1)$ and $\Sigma(\nu_2)$ are independent. Then $\Sigma(\nu_1) \cup \Sigma(\nu_2)$ is a Boolean model driven by $\nu_1 + \nu_2$. We set $\Sigma(\nu_1 + \nu_2) = \Sigma(\nu_1) \cup \Sigma(\nu_2)$. We also consider $\Sigma(T_\eta \nu_1)$, the Boolean model obtained by adding η to the radius of each ball of $\Sigma(\nu_1)$. Thus $\Sigma(T_\eta \nu_1)$ is driven by $T_\eta \nu_1$.

Let us prove the following property:

$$\{S(a/2) \leftrightarrow_{\Sigma(\nu_1 + \nu_2)} S(a)\} \subset \{S(a/2) \leftrightarrow_{\Sigma(T_\eta \nu_1)} S(a)\} \cup \bigcup_{i \in I} \{S(x_i, \eta/4) \leftrightarrow_{\Sigma(\nu_2)} S(x_i, \eta/2)\}. \quad (12)$$

Assume that $\Sigma(\nu_1 + \nu_2) = \Sigma(\nu_1) \cup \Sigma(\nu_2)$ connects $S(a/2)$ with $S(a)$. Recall $a \geq 4\eta$. If the diameter of all connected components of $\Sigma(\nu_2) \cap B(a)$ are less or equal to η , then $\Sigma(T_\eta \nu_1)$ connects $S(a/2)$ with $S(a)$. Otherwise, let C be a connected component of $\Sigma(\nu_2) \cap B(a)$ with diameter at least η . Let x, y be two points of C such that $\|x - y\| > \eta$. The point x belongs to a ball $B(x_i, \eta/4)$. As y does not belong to $B(x_i, \eta/2)$, the

component C connects $S(x_i, \eta/4)$ to $S(x_i, \eta/2)$. Therefore, $\Sigma(\nu_2)$ connects $S(x_i, \eta/4)$ to $S(x_i, \eta/2)$. We have proven (12). The lemma follows. \square

The following lemma is essentially the first item of Proposition 3.1 in [4]. For the sake of completeness we nevertheless provide a proof.

Lemma 7 *Let ν be a finite measure on $]0, +\infty[$. There exists a constant $C_2 = C_2(d) > 0$ such that, for all $a > 0$:*

$$p(10a, \nu) \leq C_2 p(a, \nu)^2 + C_2 \int_{[a, +\infty[} r^d \nu(dr).$$

Proof. Let K be a finite subset of $S(5)$ such that $K + B(1/2)$ covers $S(5)$. Let L be a finite subset of $S(10)$ such that $L + B(1/2)$ covers $S(10)$. Let A be the following event: there exists a random ball $B(c, r)$ of $\Sigma(\nu)$ such that $r \geq a$ and $B(c, r) \cap B(10a)$ is non empty. We have:

$$\{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\} \setminus A \subset \{S(5a) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(10a)\}$$

where, in the last event, we ask for a path using only balls of $\Sigma(\nu)$ of radius at most a . Let us prove the following:

$$\begin{aligned} & \{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\} \setminus A \\ & \subset \bigcup_{k \in K, l \in L} \{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\} \cap \{S(al, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(al, a)\}. \end{aligned} \quad (13)$$

Assume that the event on the left hand side occurs. Then, by the previous remark, there exists a path from a point $x \in S(5a)$ to a point $y \in S(10a)$ that is contained in balls of $\Sigma(\nu)$ of radius at most a . As $Ka + B(a/2)$ covers $S(5a)$, there exists $k \in K$ such that x belongs to $B(ka, a/2)$. Using the previous path, one gets that the event

$$\{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\}$$

occurs. By a similar arguments involving y we get (13).

Observe that, for all $k \in K$ and $l \in L$, the events

$$\{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\} \text{ and } \{S(al, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(al, a)\}$$

are independent. Indeed, the first one depends only on balls with centers in $B(ak, 2a)$, the second one depends only on balls with centers in $B(al, 2a)$, and $\|ak - al\| \geq 5a$. Using this independence, stationarity and (13), we then get:

$$P(\{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\}) \leq CP(S(a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(a))^2 + P(A)$$

where C is the product of the cardinality of K by the cardinality of L . The probability $P(A)$ is bounded above by standard computations. \square

From the previous lemma, we deduce the following result.

Lemma 8 *Let $\varepsilon > 0$. There exists $C_3 = C_3(d) > 0$, $a_0 = a_0(d, \mu)$ and $k_0 = k_0(d, \mu, \varepsilon)$ such that, for all N , all $\rho \geq 2$ and all $a \geq a_0$: if $p(a, m_N^\rho) \leq C_3$ then for all $k \geq k_0$, $p(a10^k, m_N^\rho) \leq \varepsilon$.*

Proof. For all $\rho \geq 2$ and all $a \geq 1$ we have:

$$\begin{aligned} \int_{[a, +\infty[} r^d m_\infty^\rho(dr) &= \sum_{k \geq 0} \rho^{kd} \int_{]0, +\infty[} 1_{[a, +\infty[(r\rho^{-k})}(r\rho^{-k})^d \mu(dr) \\ &= \int_{]0, +\infty[} \sum_{k \geq 0} 1_{[a, +\infty[(r\rho^{-k})} r^d \mu(dr) \\ &= \int_{[a, +\infty[} (\lfloor \ln(r/a) \ln(\rho) \rfloor + 1) r^d \mu(dr) \\ &\leq \int_{[a, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr). \end{aligned}$$

Let C_2 be the constant given by Lemma 7. By (5) we can chose $a_0 = a_0(d, \mu) \geq 1$ such that

$$C_2^2 \int_{[a_0, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr) \leq \frac{1}{4}. \quad (14)$$

Let $C_3 = (2C_2)^{-1}$. Let N , ρ and a be as in the statement of the lemma. From Lemma 7 we get:

$$C_2 p(10a, m_N^\rho) \leq (C_2 p(a, m_N^\rho))^2 + C_2^2 \int_{[a, +\infty[} r^d m_N^\rho(dr) \quad (15)$$

$$\leq (C_2 p(a, m_N^\rho))^2 + C_2^2 \int_{[a, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr) \quad (16)$$

Let (u_k) be a sequence defined by $u_0 = 1/2$ and, for all $k \geq 0$:

$$u_{k+1} = u_k^2 + C_2^2 \int_{[a_0 10^k, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr). \quad (17)$$

Note that the sequence (u_k) only depends on d and μ .

Assume that $p(a, m_N^\rho) \leq C_3$. We then have $C_2 p(a, m_N^\rho) \leq u_0$. Using $a \geq a_0$ and (16), we then get $C_2 p(a10^k, m_N^\rho) \leq u_k$ for all k . Therefore, it sufficies to show that the sequence (u_k) tends to 0.

Using (17), (14) and $u_0 = 1/2$ we get $0 \leq u_k \leq 1/2$ for all k . Therefore, $0 \leq \limsup u_k \leq 1/2$. By (17) and by the convergence of the integrale we also get $\limsup u_k \leq (\limsup u_k)^2$. As a consequence, $\limsup u_k = 0$ and the lemma is proven. \square

Lemma 9 *For all $a > 0$ and $\rho > 1$ the following convergence holds:*

$$p(a, m_\infty^\rho) = \lim_{N \rightarrow \infty} p(a, m_N^\rho).$$

Proof. The sequence of events

$$A_N = \{S(a/2) \leftrightarrow_{\Sigma(m_N^\rho)} S(a)\}$$

is increasing (we use the natural coupling between our Boolean models). Therefore, it suffices to show that the union of the previous events is

$$A = \{S(a/2) \leftrightarrow_{\Sigma(m_\infty^\rho)} S(a)\}.$$

If A occurs, then there is a path from $S(a/2)$ to $S(a)$ that is contained in $\Sigma(m_\infty^\rho)$. By a compactness argument, this path is included in a finite union of ball of $\Sigma(m_\infty^\rho)$. Therefore, there exists N such that the path is included in $\Sigma(m_N^\rho)$ and A_N occurs. This proves $A \subset \cup A_N$. The other inclusion is straightforward. \square

Proof of Theorem 3. By Lemma 5, we can fix $\eta_1 > 0$ such that $p(a, T_{10\eta_1}\mu)$ tends to 0 as a tends to ∞ . Let C_1 be given by Lemma 6. Let a_0 and C_3 be as given by Lemma 8. Fix $a_1 \geq \max(40\eta_1, a_0, 1)$ such that $p(a, T_{10\eta_1}\mu) \leq C_3/2$ for all $a \geq a_1$. Let k_0 be given by Lemma 8 with the choice:

$$\varepsilon = C_1^{-1}(10a_1)^{-d}\eta_1^d C_3/2.$$

Therefore, for all $\rho \geq 2$, all N , all $a \in [a_1, 10a_1]$ and all $\eta \in [\eta_1, 10\eta_1]$:

$$C_1 a^d \eta^{-d} p(a10^k, m_N^\rho) \leq \frac{C_3}{2} \text{ for all } k \geq k_0 \text{ as soon as } p(a, m_N^\rho) \leq C_3.$$

Fix $k \geq k_0$, $a \in [a_1, 10a_1]$ and $\eta \in [\eta_1, 10\eta_1]$. Set:

$$\rho = 2a10^k \eta^{-1}.$$

Note $\rho \geq 8 \geq 2$ as $a \geq a_1 \geq 40\eta_1 \geq 4\eta$. By Lemma 6 we have, for all N :

$$\begin{aligned} p(a, m_{N+1}^\rho) &\leq p(a, T_\eta\mu) + C_1 a^d \eta^{-d} p(\eta/2, m_{N+1}^\rho - \mu) \\ &= p(a, T_\eta\mu) + C_1 a^d \eta^{-d} p(\eta/2, H^\rho m_N^\rho). \end{aligned}$$

By definition of a_1 , by $a_1 \leq a$, by $\eta \leq 10\eta_1$, by scaling and by definition of ρ we get, for all N :

$$\begin{aligned} p(a, m_{N+1}^\rho) &\leq \frac{C_3}{2} + C_1 a^d \eta^{-d} p(\rho\eta/2, m_N^\rho) \\ &= \frac{C_3}{2} + C_1 a^d \eta^{-d} p(a10^k, m_N^\rho). \end{aligned}$$

Combining this inequality with the property defining k_0 , we get that $p(a, m_N^\rho) \leq C_3$ implies $p(a, m_{N+1}^\rho) \leq C_3$. As $p(a, m_0^\rho) = p(a, \mu) \leq p(a, T_{10\eta_1}\mu) \leq C_3/2$ we get $p(a, m_N^\rho) \leq C_3$ for all integer N .

Let $\varepsilon > 0$. Using again Lemma 8 we get the existence of an integer k'_0 such that $p(a10^{k'}, m_N^\rho) \leq \varepsilon$ for all $k' \geq k'_0$ as soon as $p(a, m_N^\rho) \leq C_3$. But we have proven the latter property. Therefore $p(a10^{k'}, m_N^\rho) \leq \varepsilon$ for all N and all $k' \geq k'_0$. By Lemma 9, we get $p(a10^{k'}, m_\infty^\rho) \leq \varepsilon$ for all $k' \geq k'_0$. Using the freeness on the choice of $k \geq k_0$ and $\eta \in [\eta_1, 10\eta_1]$, we get that the previous result holds for all $\rho \geq 2a10^{k_0}\eta_1^{-1}$ and then for all $\rho \geq 20a_110^{k_0}\eta_1^{-1}$. Moreover, using the freeness on the choice of $a \in [a_1, 10a_1]$ and $k' \geq k'_0$, we get:

$$p(r, m_\infty^\rho) \leq \varepsilon \text{ for all } r \geq a_110^{k'_0}.$$

Therefore, $p(r, m_\infty^\rho)$ tends to 0 as r tends to infinity. As a consequence, $\Sigma^\rho(\mu)$ does not percolate for any $\rho \geq 20a_110^{k_0}\eta_1^{-1}$. \square

A Critical parameters

Lemma 10

$$\tilde{\lambda}_c(\mu) \leq \hat{\lambda}_c(\mu) \leq \lambda_c(\mu).$$

Proof. The second inequality is a consequence of the following inclusion:

$$\{\{0\} \leftrightarrow_{\Sigma} S(r)\} \subset \{S(r/2) \leftrightarrow_{\Sigma} S(r)\}.$$

The first inequality can be proven as follows. Let $r \geq 1$. By the FKG inequality, we get:

$$\begin{aligned} P(\{0\} \leftrightarrow_{\Sigma} S(r)) &\geq P(B(0,1) \subset \Sigma \text{ and } S(1) \leftrightarrow_{\Sigma} S(r)) \\ &\geq CP(S(1) \leftrightarrow_{\Sigma} S(r)) \end{aligned}$$

where $C = P(B(0,1) \subset \Sigma) > 0$ does not depend on r . For all large enough r , we can cover $S(2r)$ by at most $C'r^d$ balls $B(x_i, 1)$ where C' only depends on the dimension d . If there is a path in Σ from $S(2r)$ to $S(4r)$, then there exists i and a path in Σ from $S(x_i, 1)$ to $S(x_i, r)$. (Consider the ball $B(x_i, 1)$ that contains the initial point of the path.) By stationarity and by the previous inequality we thus get:

$$\begin{aligned} P(S(2r) \leftrightarrow_{\Sigma} S(4r)) &\leq C'r^d P(S(1) \leftrightarrow_{\Sigma} S(r)) \\ &\leq C'C^{-1}r^d P(\{0\} \leftrightarrow_{\Sigma} S(r)). \end{aligned}$$

The first inequality stated in the lemma follows. □

References

- [1] Erik I. Broman and Federico Camia. Large- N limit of crossing probabilities, discontinuity, and asymptotic behavior of threshold values in Mandelbrot's fractal percolation process. *Electron. J. Probab.*, 13:no. 33, 980–999, 2008.
- [2] Erik I. Broman and Federico Camia. Universal behavior of connectivity properties in fractal percolation models. 2009.
- [3] Lincoln Chayes. Aspects of the fractal percolation process. In *Fractal geometry and stochastics (Finsterbergen, 1994)*, volume 37 of *Progr. Probab.*, pages 113–143. Birkhäuser, Basel, 1995.
- [4] Jean-Baptiste Gouéré. Subcritical regimes in the Poisson Boolean model of continuum percolation. *Ann. Probab.*, 36(4):1209–1220, 2008.
- [5] Jean-Baptiste Gouéré. Subcritical regimes in some models of continuum percolation. *Ann. Appl. Probab.*, 19(4):1292–1318, 2009.
- [6] Geoffrey Grimmett. *Percolation*, volume 321 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1999.
- [7] Peter Hall. On continuum percolation. *Ann. Probab.*, 13(4):1250–1266, 1985.
- [8] M. Mandelbrot. Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier. *Journal of Fluid Mechanics*, 62:331–358, 1974.

- [9] Ronald Meester and Rahul Roy. *Continuum percolation*, volume 119 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [10] Ronald Meester, Rahul Roy, and Anish Sarkar. Nonuniversality and continuity of the critical covered volume fraction in continuum percolation. *J. Statist. Phys.*, 75(1-2):123–134, 1994.
- [11] M. V. Menshikov, S. Yu. Popov, and M. Vachkovskaia. On the connectivity properties of the complementary set in fractal percolation models. *Probab. Theory Related Fields*, 119(2):176–186, 2001.
- [12] M. V. Menshikov, S. Yu. Popov, and M. Vachkovskaia. On a multiscale continuous percolation model with unbounded defects. *Bull. Braz. Math. Soc. (N.S.)*, 34(3):417–435, 2003. Sixth Brazilian School in Probability (Ubatuba, 2002).
- [13] M. V. Men'shikov and A. F. Sidorenko. Coincidence of critical points in Poisson percolation models. *Teor. Veroyatnost. i Primenen.*, 32(3):603–606, 1987.
- [14] S. A. Molchanov, V. F. Pisarenko, and A. Ya. Reznikova. Multiscale models of failure and percolation. *Physics of The Earth and Planetary Interiors*, 61(1-2):36–43, 1990.
- [15] M. E. Orzechowski. On the phase transition to sheet percolation in random Cantor sets. *J. Statist. Phys.*, 82(3-4):1081–1098, 1996.
- [16] Serguei Yu. Popov and Marina Vachkovskaia. A note on percolation of Poisson sticks. *Braz. J. Probab. Stat.*, 16(1):59–67, 2002.
- [17] Damien G. White. On the value of the critical point in fractal percolation. *Random Structures Algorithms*, 18(4):332–345, 2001.
- [18] S. A. Zuev and A. F. Sidorenko. Continuous models of percolation theory. I. *Teoret. Mat. Fiz.*, 62(1):76–86, 1985.
- [19] S. A. Zuev and A. F. Sidorenko. Continuous models of percolation theory. II. *Teoret. Mat. Fiz.*, 62(2):253–262, 1985.