



HAL
open science

Percolation in a multiscale Boolean model

Jean-Baptiste Gouéré

► **To cite this version:**

| Jean-Baptiste Gouéré. Percolation in a multiscale Boolean model. 2010. hal-00519310v2

HAL Id: hal-00519310

<https://hal.science/hal-00519310v2>

Preprint submitted on 13 Nov 2010 (v2), last revised 9 Mar 2011 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Percolation in a multiscale Boolean model

Jean-Baptiste Gouéré *

Abstract

We consider percolation in a multiscale Boolean model. This model is defined as the union of scaled independent copies of a given Boolean model. The scale factor of the n^{th} copy is ρ^{-n} . We prove, under optimal integrability assumptions, that no percolation occurs in the multiscale Boolean model for large enough ρ if the rate of the Boolean model is below some critical value.

1 Introduction and statement of the main result

1.1 The Boolean model

Let $d \geq 2$. Let μ be a finite measure on $]0, +\infty[$. We assume that the mass of μ is positive. Let ξ be a Poisson point process on $\mathbb{R}^d \times]0, +\infty[$ whose intensity is the product of the Lebesgue measure on \mathbb{R}^d by μ . With ξ we associate a random set $\Sigma(\mu)$ defined as follows:

$$\Sigma(\mu) = \bigcup_{(c,r) \in \xi} B(c,r)$$

where $B(c,r)$ is the open Euclidean ball of radius r centered at c . The random set $\Sigma(\mu)$ is the Boolean model with parameter μ . We shall sometimes write Σ to simplify the notations.

The following description may be more intuitive. Let χ denote the projection of ξ on \mathbb{R}^d . With probability one this projection is one-to-one. We can therefore write:

$$\xi = \{(c, r(c)), c \in \chi\}.$$

Write $\mu = m\nu$ where ν is a probability measure. Then, χ is a Poisson point process on \mathbb{R}^d with density m . Moreover, given χ , the sequence $(r(c))_{c \in \chi}$ is a sequence of independent random variable with common distribution ν . We shall not use this point of view.

1.2 Percolation in the Boolean model

Let C denote the connected component of Σ that contains the origin. We say that Σ percolates if C is unbounded with positive probability. We refer to the book by Meester and Roy [9] for background on continuum percolation. Set:

$$\lambda_c(\mu) = \inf\{\lambda > 0 : \Sigma(\lambda\mu) \text{ percolates}\}.$$

**Postal address:* Université d'Orléans MAPMO B.P. 6759 45067 Orléans Cedex 2 France *E-mail:* jbgouere@univ-orleans.fr

One easily check that $\lambda_c(\mu)$ is finite as soon as μ has a positive mass. In [4] we proved that $\lambda_c(\mu)$ is positive if and only if:

$$\int r^d \mu(dr) < \infty.$$

The only if part had been proved earlier by Hall [7]. For all $A, B \subset \mathbb{R}^d$, we write $A \leftrightarrow_\Sigma B$ if there exists a path in Σ from A to B . We denote by $S(c, r)$ the Euclidean sphere of radius r centered at c :

$$S(c, r) = \{x \in \mathbb{R}^d : \|x - c\|_2 = r\}.$$

We write $S(r)$ when $c = 0$.

The critical parameter $\lambda_c(\mu)$ can also be defined as follows:

$$\lambda_c(\mu) = \sup \{ \lambda > 0 : P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \},$$

We shall need two other critical parameters:

$$\begin{aligned} \widehat{\lambda}_c(\mu) &= \sup \{ \lambda > 0 : P(S(r/2) \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \}, \\ \widetilde{\lambda}_c(\mu) &= \sup \{ \lambda > 0 : r^d P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \}. \end{aligned}$$

We have (see Lemma 12) :

$$\widetilde{\lambda}_c(\mu) \leq \widehat{\lambda}_c(\mu) \leq \lambda_c(\mu). \quad (1)$$

When the support of μ is bounded,

$$P(\{0\} \leftrightarrow_{\Sigma(\lambda\mu)} S(r))$$

decays exponentially fast to 0 as soon as $\lambda < \lambda_c(\mu)$ (see for example [9], Section 12.10 in [6] in the case of constant radii or the papers [10], [13], [18] and [19]). Therefore:

$$\widetilde{\lambda}_c(\mu) = \widehat{\lambda}_c(\mu) = \lambda_c(\mu) \text{ as soon as the support of } \mu \text{ is bounded.} \quad (2)$$

Remarks.

- The treshold parameter $\widehat{\lambda}_c(\mu)$ is positive if and only if $\int r^d \mu(dr)$ is finite (i.e., if and only if $\lambda_c(\mu)$ is positive). See Lemma 13.
- Using ideas of [4], we can check that $\widetilde{\lambda}_c(\mu)$ is positive if and only if

$$x^d \int_x^\infty r^d \mu(dr) \rightarrow 0 \text{ as } x \rightarrow \infty.$$

If we only use results stated in [4], we can easily get the following weaker statements. Let $D(\lambda\mu)$ denote the Euclidean diameter of the connected component of $\Sigma(\lambda\mu)$ that contains the origin. Note that $\widetilde{\lambda}_c(\mu)$ is positive if and only if there exists λ such that:

$$r^d P(D(\lambda\mu) \geq r) \rightarrow 0, \text{ as } r \rightarrow \infty. \quad (3)$$

If $E(D(\lambda\mu)^d)$ is finite then (3) holds. If (3) holds then $E(D(\lambda\mu)^{d-\varepsilon})$ is finite for any small enough $\varepsilon > 0$. By Theorem 2.2 of [4] we thus get the following implications:

$$\int_0^{+\infty} r^{2d} \mu(dr) < \infty \text{ implies } \widetilde{\lambda}_c(\mu) > 0 \text{ implies } \forall \varepsilon > 0 : \int_0^{+\infty} r^{2d-\varepsilon} \mu(dr) < \infty.$$

1.3 A multiscale Boolean model

Let $\rho > 1$ be a scale factor. Let $(\Sigma_n)_{n \geq 0}$ be a sequence of independent copies of $\Sigma(\mu)$. In this paper, we are interested in percolation properties of the following multiscale Boolean model:

$$\Sigma^\rho(\mu) = \bigcup_{n \geq 0} \rho^{-n} \Sigma_n. \quad (4)$$

We shall sometimes write Σ^ρ to simplify the notations. As before, we say that Σ^ρ percolates if the connected component of Σ^ρ that contains the origin is unbounded with positive probability.

This model seems to have been first introduced as a model of failure in geophysical medias in the 80'. We refer to the paper by Molchanov, Pisarenko and Reznikova [14] for an account of those studies. For more recent results we refer to [2], [9], [10], [11], [12] and [16].

This model is related to a discrete model introduced by Mandelbrot [8]. We refer to the survey by L. Chayes [3] and, for more recent results, to [1], [15] and [17].

In [11], Menshikov, Popov and Vachkovskaia considered the case where the radii of the unscaled process Σ_0 equal 1. They proved the following result.

Theorem 1 ([11]) *If $\lambda < \lambda_c(\delta_1)$ then, for all large enough ρ , $\Sigma^\rho(\lambda\mu)$ does not percolate.*

In [12] the same authors considered the case where the radii are random and can be unbounded. They considered the following sub-autosimilarity assumption on the measure μ :

$$\lim_{a \rightarrow \infty} \sup_{r \geq 1/2} \frac{a^d \mu([ar, +\infty[)}{\mu([r, +\infty[)} = 0 \quad (5)$$

with the convention $0/0 = 0$. They proved the following result.

Theorem 2 ([12]) *Assume that the measure μ satisfies (5). If $\lambda < \tilde{\lambda}_c(\mu)$ then, for all large enough ρ , $\Sigma^\rho(\lambda\mu)$ does not percolate.*

Note that (5) is fulfilled for any measure with bounded support. Because of (2), Theorem 2 is then a generalization of Theorem 1.

In [5] we proved the following related result in which ρ is fixed.

Theorem 3 *Let $\rho > 1$. There exists $\lambda > 0$ such that $\Sigma^\rho(\lambda\mu)$ does not percolate if and only if:*

$$\int_{[1, +\infty[} \beta^d \ln(\beta) \mu(d\beta) < \infty. \quad (6)$$

The main result of this paper is the first item of the following theorem. The second item is easy and already contained in Theorem 3. Recall that, by Lemma 13, $\widehat{\lambda}_c(\mu)$ is positive as soon as $\int r^d \mu(dr)$ is finite and therefore as soon as (6) holds.

Theorem 4

1. Assume (6). Then, for all $\lambda < \widehat{\lambda}_c(\mu)$, there exists $\rho(\lambda) > 1$ such that, for all $\rho \geq \rho(\lambda)$:

$$P(S(r/2) \leftrightarrow_{\Sigma^\rho(\lambda\mu)} S(r)) \rightarrow 0 \text{ as } r \rightarrow \infty \quad (7)$$

and therefore $\Sigma^\rho(\lambda\mu)$ does not percolate.

2. Assume that (6) does not hold. Then, for all $\lambda > 0$ and for all $\rho > 1$, $\Sigma^\rho(\lambda\mu)$ percolates.

Theorem 4 is a generalization of Theorem 2 and thus of Theorem 1. Indeed, by (1), one has $\lambda < \widehat{\lambda}_c$ as soon as $\lambda < \widetilde{\lambda}_c$. Moreover, by the second item of Theorem 4, (6) has to be a consequence of the assumptions of Theorem 2 at least when $\widetilde{\lambda}_c(\mu)$ is positive. For example, one can check that (5) is stronger than (6)¹. Alternatively, one can check that (6) is a consequence of $\widetilde{\lambda}_c(\mu) > 0$ (see the remarks at the end of Section 1.2).

Let us denote by $\lambda_c(m_\infty^\rho)$ and $\widehat{\lambda}_c(m_\infty^\rho)$ the λ_c and $\widehat{\lambda}_c$ critical thresholds for the multiscale model with scale parameter ρ . Theorems 3 and 4 yield the following result:

1. If (6) holds then $\lambda_c(m_\infty^\rho) > 0$ (and actually the proof of Theorem 3 yields $\widehat{\lambda}_c(m_\infty^\rho) > 0$) for all $\rho > 1$ and $\widehat{\lambda}_c(m_\infty^\rho) \rightarrow \widehat{\lambda}_c(\mu) > 0$ as $\rho \rightarrow \infty$.
2. Otherwise, $\widehat{\lambda}_c(m_\infty^\rho) = \lambda_c(m_\infty^\rho) = 0$ for all $\rho > 1$.

Let us denote by $D^\rho(\lambda\mu)$ the diameter of the connected component of $\Sigma^\rho(\lambda\mu)$ that contains the origin. The following result is an easy consequence of Theorem 4 above and Theorems 2.9 and 1.2 in [5].

Theorem 5 *Let $s > 0$, $\lambda > 0$ and $\rho > 1$.*

1. *If $\int_{[1,+\infty[} \beta^{d+s} \mu(d\beta) < \infty$ and (7) holds, then $E(D^\rho(\lambda\mu)^s) < \infty$.*
2. *If $\int_{[1,+\infty[} \beta^{d+s} \mu(d\beta) = \infty$ then $E(D^\rho(\lambda\mu)^s) = \infty$.*

2 Proof of Theorem 4

2.1 Some notations

In the whole of Section 2, we make the following assumptions:

- μ satisfies (6).
- $1 < \widehat{\lambda}_c(\mu)$.

For all $\eta > 0$, we denote by $T_\eta\mu$ the measure defined by $T_\eta\mu(A) = \mu(A - \eta)$. In other words, we can build $\Sigma(T_\eta\mu)$ from $\Sigma(\mu)$ by adding η to each radius.

For all $\rho > 1$, we denote by $H^\rho\mu$ the measure defined by $H^\rho\mu(A) = \rho^d \mu(\rho A)$. With this definition, $\rho^{-1}\Sigma(\mu)$ is a Boolean model driven by the measure $H^\rho\mu$. For all $n \geq 0$, we let:

$$m_n^\rho = \sum_{k=0}^n H^{\rho^k} \mu.$$

With this definition and the notations of (4),

$$\bigcup_{k=0}^n \rho^{-k} \Sigma_k$$

1. From (5) one gets the existence of $a > 1$ such that, for all $r \geq a$, one has $\mu([r, +\infty[) \leq 2^{-1} a^{-d} \mu([r/a, +\infty[)$. By induction and standard computations this yields, for all $r \geq a$, $\mu([r, +\infty[) \leq A r^{-\ln(2)/\ln(a)-d}$. Therefore, for a small enough $\eta > 0$, one has $\int r^{d+\eta} \mu(dr) < \infty$.

is a Boolean model driven by m_n^ρ . We also let:

$$m_\infty^\rho = \sum_{k \geq 0} H^{\rho^k} \mu.$$

So, $\Sigma^\rho(\mu)$ is a Boolean model driven by the locally finite measure m_∞^ρ .

Let $p(a, \mu)$ denote the probability of existence of a path from $S(a/2)$ to $S(a)$ in $\Sigma(\mu)$:

$$p(a, \mu) = P(S(a/2) \leftrightarrow_{\Sigma(\mu)} S(a)).$$

We aim at proving that, for large enough ρ , $p(a, m_\infty^\rho) \rightarrow 0$ as a tends to infinity and $\Sigma^\rho(\mu)$ does not percolate. The first item of Theorem 4 follows by applying this result to the measure $\lambda\mu$. Recall that the second item of Theorem 4 is contained in Theorem 3.

2.2 Ideas

In this subsection we first sketch the proof of the existence of ρ and a such that $p(a, m_\infty^\rho)$ is small. This gives the main ingredients of the proof of the first item of Theorem 4. A full proof is given in Subsection 2.3. We then give the ideas of the proof of Theorems 1 and 2 by Menshikov, Popov and Vachkovskaia. Their basic strategy is similar but the implementation of the proofs are different.

Sketch of the proof of the first item of Theorem 4

Consider a small $\varepsilon_1 > 0$. Fix a small $\eta > 0$ and a large a such that (see Lemma 6):

$$p(a, T_\eta \mu) \leq \varepsilon_1/2. \quad (8)$$

For all $n \geq 1$, write:

$$m_n^\rho = H^\rho m_{n-1}^\rho + \mu.$$

If the event $\{S(a/2) \leftrightarrow_{\Sigma(m_n^\rho)} S(a)\}$ occurs, then either the event $\{S(a/2) \leftrightarrow_{T_\eta \mu} S(a)\}$ occurs (with a natural coupling between the Boolean models) either in $\Sigma(H^\rho m_{n-1}^\rho) \cap B(a)$ one can find a component of diameter at least η . We use this observation through its following crude consequence (see Lemma 7):

$$p(a, m_n^\rho) \leq p(a, T_\eta \mu) + C a^d \eta^{-d} p(\eta/2, H^\rho m_{n-1}^\rho).$$

By scaling and by (8), this yields:

$$p(a, m_n^\rho) \leq \varepsilon_1/2 + C a^d \eta^{-d} p(\rho\eta/2, m_{n-1}^\rho). \quad (9)$$

But for any ε_2 , any small enough ε_1 and any large enough a we can find τ such that (see Lemmas 8 and 9):

$$p(\tau a, m_{n-1}^\rho) \leq \varepsilon_2 \text{ as soon as } p(a, m_{n-1}^\rho) \leq \varepsilon_1. \quad (10)$$

An important fact is that τ does not depends on n nor on ρ , provided $\rho \geq \rho_0$ where ρ_0 is an arbitrary constant strictly larger than 1. Here we use assumption (6) to bound error terms due to the existence of large balls.

We choose ε_2 such that:

$$Ca^d \eta^{-d} \varepsilon_2 = \varepsilon_1/2.$$

We set $\rho = 2\tau a/\eta$. Then, (9) and (10) can be rewritten as follows:

$$p(a, m_n^\rho) \leq \varepsilon_1/2 + Ca^d \eta^{-d} p(\tau a, m_{n-1}^\rho) \quad (11)$$

$$Ca^d \eta^{-d} p(\tau a, m_{n-1}^\rho) \leq \varepsilon_1/2 \text{ as soon as } p(a, m_{n-1}^\rho) \leq \varepsilon_1. \quad (12)$$

As moreover (8) implies $p(a, m_0^\rho) \leq \varepsilon_1$ we get, by induction and then sending n to infinity (see Lemma 10):

$$p(a, m_\infty^\rho) \leq \varepsilon_1.$$

The convergence of $p(a, m_\infty^\rho)$ to 0 is then extracted from the above result for a small enough ε_2 and from arguments behind (10) applied to m_∞^ρ and other ε .

Sketch of the proofs of Theorems 1 and 2 by Menshikov, Popov and Vachkovskaia

Let us quickly describe the ideas of the proofs of Menshikov, Popov and Vachkovskaia. Those ideas are used in their papers [11] and [12] through a discretization of space ; we describe them in a slightly more geometric way. For simplicity we only consider two scales: $\rho^{-1}\Sigma_1$ and Σ_0 . For simplicity, we also assume that the radius is one in the unscaled model ($\mu = \lambda\delta_1$). We assume that the scale factor ρ is large enough. Assume that C is a connected component of $\rho^{-1}\Sigma_1 \cup \Sigma_0$ whose diameter is at least α (it can be much larger) for a small enough constant $\alpha > 0$. Then, C is included in the union of the following kind of sets:

1. connected components of $\rho^{-1}\Sigma_1$ whose diameter is at least α ;
2. balls of Σ_0 enlarged by α (same centers but the radii are $1 + \alpha$ instead of 1).

Then, they show that the union of all those sets is stochastically dominated by a Boolean model similar to Σ_0 but with radii enlarged by a factor α and with density of centers $1 + \alpha'$ times the corresponding density for Σ_0 for a suitable $\alpha' > 0$. This part uses $\lambda < \widehat{\lambda}_c$. In some sense, one can therefore control percolation in the union of two models by percolation in one model. Iterating the argument with some care in the constants α and α' , one sees that – for large enough ρ – one can control percolation in the multiscale model by percolation in a subcritical model. This yields the result.

2.3 Proof of Theorem 4

As $1 < \widehat{\lambda}_c(\mu)$, we know that $p(a, \mu)$ tends to 0 as a tends to infinity. We need the following slightly stronger consequence.

Lemma 6 *There exists $\eta > 0$ such that $p(a, T_\eta \mu)$ tends to 0.*

Proof. Let $\varepsilon > 0$ and $x > 0$. We have:

$$\begin{aligned} H^{1+\varepsilon} T_{\varepsilon^2} \mu([x, +\infty[) &= (1 + \varepsilon)^d T_{\varepsilon^2} \mu([x(1 + \varepsilon), +\infty[) \\ &= (1 + \varepsilon)^d \mu([x(1 + \varepsilon) - \varepsilon^2, +\infty[) \\ &\leq \kappa(\varepsilon)(1 + \varepsilon)^d \mu([x, +\infty[) \end{aligned} \quad (13)$$

where

$$\kappa(\varepsilon) = \frac{\mu(]0, +\infty[)}{\mu([\varepsilon, +\infty[)}.$$

The inequality is proven as follows. If $x \geq \varepsilon$, then $[x(1 + \varepsilon) - \varepsilon^2, +\infty[\subset [x, +\infty[$ and the result follows from $\kappa(\varepsilon) \geq 1$. If, on the contrary, $x < \varepsilon$, then the left hand side is bounded above by $(1 + \varepsilon)^d \mu(]0, +\infty[)$ which is itself bounded above by the right hand side.

Note that $\kappa(\varepsilon)(1 + \varepsilon)^d$ tends to 1 as ε tends to 0. Let us say that a measure ν is subcritical if $\widehat{\lambda}_c(\nu) > 1$. As μ is subcritical, we get that $\kappa(\varepsilon)(1 + \varepsilon)^d \mu$ is subcritical for small enough ε . We fix such an ε . By (13) we can couple a Boolean model driven by $H^{1+\varepsilon}T_{\varepsilon^2}\mu$ and a Boolean model driven by $\kappa(\varepsilon)(1 + \varepsilon)^d \mu$ in such a way that the first one is contained in the second one. Therefore the first one is subcritical. By scaling, a Boolean model driven by $T_{\varepsilon^2}\mu$ is then subcritical. We take $\eta = \varepsilon^2$. \square

Lemma 7 *Let ν_1 and ν_2 be two finite measures on $]0, +\infty[$. One has, for all $\eta > 0$ and $a \geq 4\eta$:*

$$p(a, \nu_1 + \nu_2) \leq p(a, T_\eta \nu_1) + C_1 a^d \eta^{-d} p(\eta/2, \nu_2)$$

where $C_1 = C_1(d) > 0$ depends only on the dimension d .

Proof. Let $(x_i)_{i \in I}$ be a family of points such that :

- The balls $B(x_i, \eta/4)$, $i \leq I$, cover $B(a)$.
- There are at most $C_1 a^d \eta^{-d}$ points in the family where $C_1 = C_1(d)$ depends only on the dimension d .

We couple the different Boolean model as follows. Let $\Sigma(\nu_1)$ be a Boolean model driven by ν_1 . Let $\Sigma(\nu_2)$ be a Boolean model driven by ν_2 . Assume that $\Sigma(\nu_1)$ and $\Sigma(\nu_2)$ are independent. Then $\Sigma(\nu_1) \cup \Sigma(\nu_2)$ is a Boolean model driven by $\nu_1 + \nu_2$. We set $\Sigma(\nu_1 + \nu_2) = \Sigma(\nu_1) \cup \Sigma(\nu_2)$. We also consider $\Sigma(T_\eta \nu_1)$, the Boolean model obtained by adding η to the radius of each ball of $\Sigma(\nu_1)$. Thus $\Sigma(T_\eta \nu_1)$ is driven by $T_\eta \nu_1$.

Let us prove the following property:

$$\{S(a/2) \leftrightarrow_{\Sigma(\nu_1 + \nu_2)} S(a)\} \subset \{S(a/2) \leftrightarrow_{\Sigma(T_\eta \nu_1)} S(a)\} \cup \bigcup_{i \in I} \{S(x_i, \eta/4) \leftrightarrow_{\Sigma(\nu_2)} S(x_i, \eta/2)\}. \quad (14)$$

Assume that $\Sigma(\nu_1 + \nu_2) = \Sigma(\nu_1) \cup \Sigma(\nu_2)$ connects $S(a/2)$ with $S(a)$. Recall $a \geq 4\eta$. If the diameter of all connected components of $\Sigma(\nu_2) \cap B(a)$ are less or equal to η , then $\Sigma(T_\eta \nu_1)$ connects $S(a/2)$ with $S(a)$. Otherwise, let C be a connected component of $\Sigma(\nu_2) \cap B(a)$ with diameter at least η . Let x, y be two points of C such that $\|x - y\| > \eta$. The point x belongs to a ball $B(x_i, \eta/4)$. As y does not belong to $B(x_i, \eta/2)$, the component C connects $S(x_i, \eta/4)$ to $S(x_i, \eta/2)$. Therefore, $\Sigma(\nu_2)$ connects $S(x_i, \eta/4)$ to $S(x_i, \eta/2)$. We have proven (14). The lemma follows. \square

The following lemma is essentially the first item of Proposition 3.1 in [4]. For the sake of completeness we nevertheless provide a proof.

Lemma 8 *Let ν be a finite measure on $]0, +\infty[$. There exists a constant $C_2 = C_2(d) > 0$ such that, for all $a > 0$:*

$$p(10a, \nu) \leq C_2 p(a, \nu)^2 + C_2 \int_{[a, +\infty[} r^d \nu(dr).$$

Proof. Let K be a finite subset of $S(5)$ such that $K + B(1/2)$ covers $S(5)$. Let L be a finite subset of $S(10)$ such that $L + B(1/2)$ covers $S(10)$. Let A be the following event: there exists a random ball $B(c, r)$ of $\Sigma(\nu)$ such that $r \geq a$ and $B(c, r) \cap B(10a)$ is non empty. We have:

$$\{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\} \setminus A \subset \{S(5a) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(10a)\}$$

where, in the last event, we ask for a path using only balls of $\Sigma(\nu)$ of radius at most a . Let us prove the following:

$$\begin{aligned} & \{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\} \setminus A \\ & \subset \bigcup_{k \in K, l \in L} \{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\} \cap \{S(al, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(al, a)\}. \end{aligned} \quad (15)$$

Assume that the event on the left hand side occurs. Then, by the previous remark, there exists a path from a point $x \in S(5a)$ to a point $y \in S(10a)$ that is contained in balls of $\Sigma(\nu)$ of radius at most a . As $Ka + B(a/2)$ covers $S(5a)$, there exists $k \in K$ such that x belongs to $B(ka, a/2)$. Using the previous path, one gets that the event

$$\{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\}$$

occurs. By a similar arguments involving y we get (15).

Observe that, for all $k \in K$ and $l \in L$, the events

$$\{S(ak, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(ak, a)\} \text{ and } \{S(al, a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(al, a)\}$$

are independent. Indeed, the first one depends only on balls with centers in $B(ak, 2a)$, the second one depends only on balls with centers in $B(al, 2a)$, and $\|ak - al\| \geq 5a$. Using this independence, stationarity and (15), we then get:

$$P(\{S(5a) \leftrightarrow_{\Sigma(\nu)} S(10a)\}) \leq CP(S(a/2) \leftrightarrow_{\Sigma(\nu)}^{\leq a} S(a))^2 + P(A)$$

where C is the product of the cardinality of K by the cardinality of L . The probability $P(A)$ is bounded above by standard computations. \square

From the previous lemma, we deduce the following result.

Lemma 9 *Let $\varepsilon > 0$. There exists $C_3 = C_3(d) > 0$, $a_0 = a_0(d, \mu)$ and $k_0 = k_0(d, \mu, \varepsilon)$ such that, for all N , all $\rho \geq 2$ and all $a \geq a_0$: if $p(a, m_N^\rho) \leq C_3$ then for all $k \geq k_0$, $p(a10^k, m_N^\rho) \leq \varepsilon$.*

Proof. For all $\rho \geq 2$ and all $a \geq 1$ we have:

$$\begin{aligned} \int_{[a, +\infty[} r^d m_\infty^\rho(dr) &= \sum_{k \geq 0} \rho^{kd} \int_{]0, +\infty[} 1_{[a, +\infty[}(r\rho^{-k})(r\rho^{-k})^d \mu(dr) \\ &= \int_{]0, +\infty[} \sum_{k \geq 0} 1_{[a, +\infty[}(r\rho^{-k}) r^d \mu(dr) \\ &= \int_{[a, +\infty[} (\lfloor \ln(r/a) \ln(\rho) \rfloor + 1) r^d \mu(dr) \\ &\leq \int_{[a, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr). \end{aligned}$$

Let C_2 be the constant given by Lemma 8. By (6) we can chose $a_0 = a_0(d, \mu) \geq 1$ such that

$$C_2^2 \int_{[a_0, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr) \leq \frac{1}{4}. \quad (16)$$

Let $C_3 = (2C_2)^{-1}$. Let N , ρ and a be as in the statement of the lemma. From Lemma 8 we get:

$$C_2 p(10a, m_N^\rho) \leq (C_2 p(a, m_N^\rho))^2 + C_2^2 \int_{[a, +\infty[} r^d m_N^\rho(dr) \quad (17)$$

$$\leq (C_2 p(a, m_N^\rho))^2 + C_2^2 \int_{[a, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr) \quad (18)$$

Let (u_k) be a sequence defined by $u_0 = 1/2$ and, for all $k \geq 0$:

$$u_{k+1} = u_k^2 + C_2^2 \int_{[a_0 10^k, +\infty[} (\ln(r) \ln(2)^{-1} + 1) r^d \mu(dr). \quad (19)$$

Note that the sequence (u_k) only depends on d and μ .

Assume that $p(a, m_N^\rho) \leq C_3$. We then have $C_2 p(a, m_N^\rho) \leq u_0$. Using $a \geq a_0$ and (18), we then get $C_2 p(a 10^k, m_N^\rho) \leq u_k$ for all k . Therefore, it suffices to show that the sequence (u_k) tends to 0.

Using (19), (16) and $u_0 = 1/2$ we get $0 \leq u_k \leq 1/2$ for all k . Therefore, $0 \leq \limsup u_k \leq 1/2$. By (19) and by the convergence of the integrale we also get $\limsup u_k \leq (\limsup u_k)^2$. As a consequence, $\limsup u_k = 0$ and the lemma is proven. \square

Lemma 10 *For all $a > 0$ and $\rho > 1$ the following convergence holds:*

$$p(a, m_\infty^\rho) = \lim_{N \rightarrow \infty} p(a, m_N^\rho).$$

Proof. The sequence of events

$$A_N = \{S(a/2) \leftrightarrow_{\Sigma(m_N^\rho)} S(a)\}$$

is increasing (we use the natural coupling between our Boolean models). Therefore, it suffices to show that the union of the previous events is

$$A = \{S(a/2) \leftrightarrow_{\Sigma(m_\infty^\rho)} S(a)\}.$$

If A occurs, then there is a path from $S(a/2)$ to $S(a)$ that is contained in $\Sigma(m_\infty^\rho)$. By a compactness argument, this path is included in a finite union of ball of $\Sigma(m_\infty^\rho)$. Therefore, there exists N such that the path is included in $\Sigma(m_N^\rho)$ and A_N occurs. This proves $A \subset \cup A_N$. The other inclusion is straightforward. \square

Proof of the second item of Theorem 4. By Lemma 6, we can fix $\eta_1 > 0$ such that $p(a, T_{10\eta_1} \mu)$ tends to 0 as a tends to ∞ . Let C_1 be given by Lemma 7. Let a_0 and C_3 be as given by Lemma 9. Fix $a_1 \geq \max(40\eta_1, a_0, 1)$ such that $p(a, T_{10\eta_1} \mu) \leq C_3/2$ for all $a \geq a_1$. Let k_0 be given by Lemma 9 with the choice:

$$\varepsilon = C_1^{-1} (10a_1)^{-d} \eta_1^d C_3/2.$$

Therefore, for all $\rho \geq 2$, all N , all $a \in [a_1, 10a_1]$ and all $\eta \in [\eta_1, 10\eta_1]$:

$$C_1 a^d \eta^{-d} p(a10^k, m_N^\rho) \leq \frac{C_3}{2} \text{ for all } k \geq k_0 \text{ as soon as } p(a, m_N^\rho) \leq C_3.$$

Fix $k \geq k_0$, $a \in [a_1, 10a_1]$ and $\eta \in [\eta_1, 10\eta_1]$. Set:

$$\rho = 2a10^k \eta^{-1}.$$

Note $\rho \geq 8 \geq 2$ as $a \geq a_1 \geq 40\eta_1 \geq 4\eta$. By Lemma 7 we have, for all N :

$$\begin{aligned} p(a, m_{N+1}^\rho) &\leq p(a, T_\eta \mu) + C_1 a^d \eta^{-d} p(\eta/2, m_{N+1}^\rho - \mu) \\ &= p(a, T_\eta \mu) + C_1 a^d \eta^{-d} p(\eta/2, H^\rho m_N^\rho). \end{aligned}$$

By definition of a_1 , by $a_1 \leq a$, by $\eta \leq 10\eta_1$, by scaling and by definition of ρ we get, for all N :

$$\begin{aligned} p(a, m_{N+1}^\rho) &\leq \frac{C_3}{2} + C_1 a^d \eta^{-d} p(\rho\eta/2, m_N^\rho) \\ &= \frac{C_3}{2} + C_1 a^d \eta^{-d} p(a10^k, m_N^\rho). \end{aligned}$$

Combining this inequality with the property defining k_0 , we get that $p(a, m_N^\rho) \leq C_3$ implies $p(a, m_{N+1}^\rho) \leq C_3$. As $p(a, m_0^\rho) = p(a, \mu) \leq p(a, T_{10\eta_1} \mu) \leq C_3/2$ we get $p(a, m_N^\rho) \leq C_3$ for all integer N .

Let $\varepsilon > 0$. Using again Lemma 9 we get the existence of an integer k'_0 such that $p(a10^{k'}, m_N^\rho) \leq \varepsilon$ for all $k' \geq k'_0$ as soon as $p(a, m_N^\rho) \leq C_3$. But we have proven the latter property. Therefore $p(a10^{k'}, m_N^\rho) \leq \varepsilon$ for all N and all $k' \geq k'_0$. By Lemma 10, we get $p(a10^{k'}, m_\infty^\rho) \leq \varepsilon$ for all $k' \geq k'_0$. Using the freedom on the choice of $k \geq k_0$ and $\eta \in [\eta_1, 10\eta_1]$, we get that the previous result holds for all $\rho \geq 2a10^{k_0-1}\eta_1^{-1}$ and then for all $\rho \geq 2a_110^{k_0}\eta_1^{-1}$. Moreover, using the freedom on the choice of $a \in [a_1, 10a_1]$ and $k' \geq k'_0$, we get:

$$p(r, m_\infty^\rho) \leq \varepsilon \text{ for all } r \geq a_110^{k'_0} \text{ and all } \rho \geq 2a_110^{k_0}\eta_1^{-1}.$$

Therefore, $p(r, m_\infty^\rho)$ tends to 0 as r tends to infinity. As a consequence, $\Sigma^\rho(\mu)$ does not percolate for any $\rho \geq 2a_110^{k_0}\eta_1^{-1}$. \square

3 Proof of Theorem 5

Lemma 11 *Let $s > 0$ and $\rho > 1$. The following assumptions are equivalent:*

1. $\int_{]0, +\infty[} r^{d+s} \mu(dr) < \infty$.
2. $\int_{]1, +\infty[} r^{d+s} m_\infty^\rho(dr) < \infty$.

Proof. We have:

$$\begin{aligned} \int_{]1, +\infty[} r^{d+s} m_\infty^\rho(dr) &= \sum_{k \geq 0} \rho^{kd} \int_{]0, +\infty[} 1_{]1, +\infty[}(r\rho^{-k})(r\rho^{-k})^{d+s} \mu(dr) \\ &= \int_{]1, +\infty[} \sum_{k \geq 0} 1_{]1, +\infty[}(r\rho^{-k}) \rho^{-ks} r^{d+s} \mu(dr). \end{aligned}$$

Therefore:

$$\int_{[1,+\infty[} r^{d+s} \mu(dr) \leq \int_{[1,+\infty[} r^{d+s} m_\infty^\rho(dr) \leq \frac{1}{1-\rho^{-s}} \int_{[1,+\infty[} r^{d+s} \mu(dr).$$

This yields the result. \square

Proof of the first item of Theorem 5. By the discussion at the beginning of Section 1.5 in [5], $\Sigma^\rho(\lambda\mu)$ is driven by a a Poisson point process whose intensity is the product of the Lebesgue measure by the locally finite measure λm_∞^ρ . Let us check the three items of Theorem 2.9 in [5] with $\rho = 10$ (ρ is not use in the same way in [5]). We refer to Section 2.1 of [5] for definitions.

1. The first item is fulfilled thanks to (7)
2. For all $\beta > 0$ and all $x \in \mathbb{R}^d$, the event $G(x, 0, \beta)$ only depends on balls $B(c, r) \in \Sigma^\rho(\lambda\mu)$ such that c belongs to $B(x, 3\beta)$. By the independance property of Poisson point processes, we then get that $G(0, 0, \beta)$ and $G(x, 0, \beta)$ are independent whenever $\|x\| \geq 10\beta$. Therefore $I(10, 0, \beta) = 0$ and the second item of Theorem 2.9 is fulfilled.
3. The third item (note that μ in [5] is m_∞^ρ in this paper) is fulfilled thanks to Lemma 11

Theorem 2.9 in [5] yields the result. \square

Proof of the second item of Theorem 5. If $\int r^d \mu(dr)$ is infinite then, $\Sigma(\lambda\mu)$ percolates for all $\lambda > 0$ (see the dicussion of Section 1.2). Therefore $\Sigma^\rho(\lambda\mu)$ percolates for all $\rho > 1$ and $\lambda > 0$. Therefore $D^\rho(\lambda\mu) = \infty$ with positive probability for all $\rho > 1$ and $\lambda > 0$.

Now, assume that $\int r^d \mu(dr)$ is finite. Then, by the discussion at the beginning of Section 1.5 in [5], $\Sigma^\rho(\lambda\mu)$ is driven by a a Poisson point process whose intensity is the product of the Lebesgue measure by the locally finite measure λm_∞^ρ . We can therefore apply Theorem 1.2 in [5]. By Lemma 11, assumption (A3) of Theorem 1.2 in [5] is not fulfilled (note that μ in [5] is m_∞^ρ in this paper). Theorem 1.2 in [5] then yields the result. \square

A Critical parameters

Lemma 12

$$\tilde{\lambda}_c(\mu) \leq \hat{\lambda}_c(\mu) \leq \lambda_c(\mu).$$

Proof. The second inequality is a consequence of the following inclusion:

$$\{\{0\} \leftrightarrow_\Sigma S(r)\} \subset \{S(r/2) \leftrightarrow_\Sigma S(r)\}.$$

The first inequality can be proven as follows. Let $r \geq 1$. By the FKG inequality, we get:

$$\begin{aligned} P(\{0\} \leftrightarrow_\Sigma S(r)) &\geq P(B(0, 1) \subset \Sigma \text{ and } S(1) \leftrightarrow_\Sigma S(r)) \\ &\geq CP(S(1) \leftrightarrow_\Sigma S(r)) \end{aligned}$$

where $C = P(B(0, 1) \subset \Sigma) > 0$ does not depend on r . For all large enough r , we can cover $S(2r)$ by at most $C'r^d$ balls $B(x_i, 1)$ where C' only depends on the dimension d . If

there is a path in Σ from $S(2r)$ to $S(4r)$, then there exists i and a path in Σ from $S(x_i, 1)$ to $S(x_i, r)$. (Consider the ball $B(x_i, 1)$ that contains the initial point of the path.) By stationarity and by the previous inequality we thus get:

$$\begin{aligned} P(S(2r) \leftrightarrow_{\Sigma} S(4r)) &\leq C' r^d P(S(1) \leftrightarrow_{\Sigma} S(r)) \\ &\leq C' C^{-1} r^d P(\{0\} \leftrightarrow_{\Sigma} S(r)). \end{aligned}$$

The first inequality stated in the lemma follows. \square

Lemma 13 *The treshold parameter $\widehat{\lambda}_c(\mu)$ is positive if and only if $\int r^d \mu(dr)$ is finite.*

Proof. If $\widehat{\lambda}_c(\mu)$ is positive, then there exists $\lambda > 0$ such that $\Sigma(\lambda\mu)$ does not percolate. By Theorem 2.1 of [4] this implies that $\int r^d \mu(dr)$ is finite.

Let us assume now that $\int r^d \mu(dr)$ is finite. We need to prove the existence of $\lambda > 0$ such that $p(a, \lambda\mu)$ tends to 0. This is proven, as an intermediate result, in the proof of Theorem 1.1 in [5]. As the result is an easy consequence of Lemma 8, we find it more convenient to provide a proof here. Let C_2 be the constant given by Lemma 8. For all $a > 0$ and $\lambda > 0$ we have:

$$C_2 p(10a, \lambda\mu) \leq (C_2 p(a, \lambda\mu))^2 + \lambda C_2^2 \int_{[a, +\infty[} r^d \mu(dr). \quad (20)$$

For all $0 < a \leq 1$ we have, by standard computations:

$$C_2 p(a, \lambda\mu) \leq C_2 P(\text{a ball of } \Sigma(\lambda\mu) \text{ touches } B(a)) \leq C_2 v_d \lambda \int_{]0, +\infty[} (1+r)^d \mu(dr)$$

where v_d is the volume of the unit Euclidean ball. As $\int r^d \mu(dr)$ is finite, we can therefore fix $\lambda > 0$ such that:

$$\lambda C_2^2 \int_{[a, +\infty[} r^d \mu(dr) \leq 1/4 \text{ for all } a > 0 \text{ and } C_2 p(a, \lambda\mu) \leq 1/2 \text{ for all } 0 < a \leq 1. \quad (21)$$

By (20), (21) and by induction we get $C_2 p(a, \lambda\mu) \leq 1/2$ for all $a > 0$. Therefore, we have $0 \leq \limsup C_2 p(a, \lambda\mu) \leq 1/2$. But (20) also yields the inequality $\limsup C_2 p(a, \lambda\mu) \leq (\limsup C_2 p(a, \lambda\mu))^2$. As a consequence we must have $\limsup C_2 p(a, \lambda\mu) = 0$ and then $p(a, \lambda\mu) \rightarrow 0$. \square

References

- [1] Erik I. Broman and Federico Camia. Large- N limit of crossing probabilities, discontinuity, and asymptotic behavior of threshold values in Mandelbrot's fractal percolation process. *Electron. J. Probab.*, 13:no. 33, 980–999, 2008.
- [2] Erik I. Broman and Federico Camia. Universal behavior of connectivity properties in fractal percolation models. 2009.
- [3] Lincoln Chayes. Aspects of the fractal percolation process. In *Fractal geometry and stochastics (Finsterbergen, 1994)*, volume 37 of *Progr. Probab.*, pages 113–143. Birkhäuser, Basel, 1995.

- [4] Jean-Baptiste Gouéré. Subcritical regimes in the Poisson Boolean model of continuum percolation. *Ann. Probab.*, 36(4):1209–1220, 2008.
- [5] Jean-Baptiste Gouéré. Subcritical regimes in some models of continuum percolation. *Ann. Appl. Probab.*, 19(4):1292–1318, 2009.
- [6] Geoffrey Grimmett. *Percolation*, volume 321 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1999.
- [7] Peter Hall. On continuum percolation. *Ann. Probab.*, 13(4):1250–1266, 1985.
- [8] M. Mandelbrot. Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier. *Journal of Fluid Mechanics*, 62:331–358, 1974.
- [9] Ronald Meester and Rahul Roy. *Continuum percolation*, volume 119 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [10] Ronald Meester, Rahul Roy, and Anish Sarkar. Nonuniversality and continuity of the critical covered volume fraction in continuum percolation. *J. Statist. Phys.*, 75(1-2):123–134, 1994.
- [11] M. V. Menshikov, S. Yu. Popov, and M. Vachkovskaia. On the connectivity properties of the complementary set in fractal percolation models. *Probab. Theory Related Fields*, 119(2):176–186, 2001.
- [12] M. V. Menshikov, S. Yu. Popov, and M. Vachkovskaia. On a multiscale continuous percolation model with unbounded defects. *Bull. Braz. Math. Soc. (N.S.)*, 34(3):417–435, 2003. Sixth Brazilian School in Probability (Ubatuba, 2002).
- [13] M. V. Men'shikov and A. F. Sidorenko. Coincidence of critical points in Poisson percolation models. *Teor. Veroyatnost. i Primenen.*, 32(3):603–606, 1987.
- [14] S. A. Molchanov, V. F. Pisarenko, and A. Ya. Reznikova. Multiscale models of failure and percolation. *Physics of The Earth and Planetary Interiors*, 61(1-2):36–43, 1990.
- [15] M. E. Orzechowski. On the phase transition to sheet percolation in random Cantor sets. *J. Statist. Phys.*, 82(3-4):1081–1098, 1996.
- [16] Serguei Yu. Popov and Marina Vachkovskaia. A note on percolation of Poisson sticks. *Braz. J. Probab. Stat.*, 16(1):59–67, 2002.
- [17] Damien G. White. On the value of the critical point in fractal percolation. *Random Structures Algorithms*, 18(4):332–345, 2001.
- [18] S. A. Zuev and A. F. Sidorenko. Continuous models of percolation theory. I. *Teoret. Mat. Fiz.*, 62(1):76–86, 1985.
- [19] S. A. Zuev and A. F. Sidorenko. Continuous models of percolation theory. II. *Teoret. Mat. Fiz.*, 62(2):253–262, 1985.