



HAL
open science

The enhancement of stress-related memory by glucocorticoids depends on synapsin-Ia/Ib

Jean-Michel Revest, Nadia Kaouane, Françoise Rougé-Pont, Anne Le Roux, Monique Vallée, Jacques Barik, Aline Desmedt, François Tronche, Pier-Vincenzo Piazza, Magali Mondin

► **To cite this version:**

Jean-Michel Revest, Nadia Kaouane, Françoise Rougé-Pont, Anne Le Roux, Monique Vallée, et al.. The enhancement of stress-related memory by glucocorticoids depends on synapsin-Ia/Ib. *Molecular Psychiatry*, 2010, 10.1038/mp.2010.40 . hal-00523673

HAL Id: hal-00523673

<https://hal.science/hal-00523673>

Submitted on 6 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The enhancement of stress-related memory by glucocorticoids depends on synapsin-Ia/Ib

J-M Revest^{1,2}, N Kaouane^{3,2}, M Mondin^{4,2}, A Le Roux^{1,2}, F Rougé-Pont^{1,2}, M Vallée^{1,2}, J Barik⁵, F Tronche⁵, A Desmedt^{3,2} and P V Piazza^{1,2}

¹INSERM U862, Neurocentre Magendie, Pathophysiology of Addiction group, Bordeaux-F33077, France.

²Université de Bordeaux, Bordeaux-F33077, France.

³CNRS UMR5228, Cognitive and Integrative Neurosciences, Talence-F33405, France.

⁴CNRS UMR 5091, Cellular Physiology of the Synapse, Bordeaux-F33077, France.

⁵CNRS FRE2401, Molecular Genetics, Neurophysiology and Behavior, Institute of Biology, Paris-F75231, France.

Correspondence should be addressed to PVP (pier-vincenzo.piazza@inserm.fr) and JMR (jean-michel.revest@inserm.fr)

Keywords: glucocorticoids, glucocorticoid receptors (GR), MAPK, Egr-1, synapsin-Ia/Ib, stress-related memory

The activation of glucocorticoid receptors (GR) by glucocorticoids increases stress-related memory through the activation of the MAPK signaling pathway and the downstream transcription factor Egr-1. Here, using converging *in vitro* and *in vivo* approaches, respectively GR expressing cell lines, culture of hippocampal neurons and GR genetically-modified mice (GR^{NesCre}), we identified synapsin-1a/1b as one of the effectors of the glucocorticoid signaling cascade. Stress and glucocorticoid-induced activation of the GR modulate synapsin-1a/1b through two complementary mechanisms. First glucocorticoids driving Egr-1 expression increases the expression of synapsin-1a/1b and second glucocorticoids driving MAPK activation increases its phosphorylation. Finally, we showed that blocking fucosylation of synapsin-1a/1b in the hippocampus inhibits its expression and prevents the glucocorticoid-mediated increase in stress-related memory. In conclusion, our data provide a complete molecular pathway (GR/Egr-1/MAPK/Syn-1a/1b; GEMS) through which stress and glucocorticoids enhance the memory of stress-related events and highlight the role of synapsin-1a/1b as molecular effector of the behavioral effects of stress.

Introduction

Glucocorticoid hormones are released by the adrenal glands during the active phase of the circadian cycle and in response to stress.^{1;1-4} Stress-induced high levels of glucocorticoids have been shown to increase the memory of stress-associated events⁵⁻⁸ and to contribute to the development of stress-related pathologies such as depression, anxiety and drug abuse.^{1;3-6;9;10} Most of these behavioral effects of stress-induced glucocorticoid secretion depend on the activation of glucocorticoid receptors (GR).¹¹ GR are steroid-regulated transcription factors which upon binding to glucocorticoids are translocated to the nucleus where they modify transcription by multiple mechanisms.¹² Consequently, identifying the transcriptional targets of stress-induced activation of GR is paramount in uncovering the molecular basis of stress-related disorders.

In a previous paper we described how activated GR enhance fear memories by inducing a cascade of events that starts with increasing the expression of the transcription factor Egr-1 followed by an increase in expression and activity of the MAPK pathway which further enhances the expression of Egr1⁸. These effects were observed in the hippocampus one of the main brain targets of glucocorticoids that is crucial for the encoding of memory.^{5;6} The GR/Egr-1/MAPK was the first molecular cascade subserving the enhancement of stress-related memory to be identified.⁸ Unfortunately, this cascade starts with a transcriptional factor, GR and ends up with another transcription factor Egr-1.⁸ Consequently the ultimate effector proteins through which stress induces an enhancement of memory still remain unknown.

In this report, we show that, in the hippocampus, synapsin-Ia/Ib isoforms are one of the effector proteins through which the GR/Egr-1/MAPK cascade enhances stress-related memories. Synapsin-Ia/Ib are the most highly expressed hippocampal pre-synaptic vesicle-associated phosphoproteins.^{13;14} The mammalian synapsin family is composed of three genes

(syn1, syn2, and syn3) that are alternatively spliced to give rise to two isoforms a and b regulated by phosphorylation.¹³ Synapsins control the release of neurotransmitters, such as glutamate which plays a crucial role in synaptic plasticity.¹⁵⁻¹⁷ These proteins have also been involved in the memory process in drosophila and in aging-related memory impairment in mammals.¹⁸⁻²⁰ However, the role of the different synapsin isoforms in the formation of specific types of memories remains unknown.

The expression of synapsin-Ia/Ib and their phosphorylation counterparts in response to glucocorticoids was studied using PC12 cell lines that express both endogenous GR and synapsin isoforms^{21;22}, primary culture of hippocampal neurons as well as the hippocampus of GR genetically-modified mice (GR^{NesCre}) in which the expression of GR has been conditionally suppressed.^{8;12;23} The role of synapsin-Ia/Ib on stress-related memory was investigated using 2-Deoxy-D-Galactose (2-dGal) an inhibitor of synapsin-Ia/Ib expression.²⁴

Our results show that the GR/Egr-1/MAPK signaling cascade increases stress-related memories through synapsin-Ia/Ib. GR-activated Egr-1 increases the expression of synapsin-Ia/Ib, the role of which is to enhance the pool of readily releasable synaptic vesicles. GR-activated MAPK then increases synapsin phosphorylation, the role of which is to induce neurotransmitter release. Finally, inhibition of GR-induced increase in hippocampal synapsin-I expression abolishes the enhancement of stress-related memories mediated by the GR/Egr-1/MAPK signaling pathway. Together, our studies identify a complete molecular pathway (GR/Egr-1/MAPK/Syn-Ia/Ib; GEMS) enhancing the consolidation of stress-related memory and highlight the important role of synapsin-Ia/Ib as an effector of the behavioral effects of glucocorticoids.

Materials and methods

Study of the interaction between GR and synapsin in vivo

For all the experiments, 5-6 week-old mice, were fed *ad libitum* and maintained in a temperature ($22 \pm 1^\circ\text{C}$) and humidity ($60 \pm 5\%$)-controlled environment under a 12 hr light/dark cycle (lights on at 8 am). The *gr* gene (also called Nr3c1) was knocked out using the Cre-LoxP recombination system in which the Cre recombinase under the control of the rat nestin promoter and enhancer allows GR^{LoxP} allele excision in the entire brain of mutant mice,²³ but not in other tissues. Control littermates not expressing Cre were $GR^{LoxP/LoxP}$ mice. Experiments carried out in basal conditions compared $GR^{LoxP/LoxP}$ and their littermates, GR^{NesCre} mice. The mice were sacrificed when glucocorticoid levels were low (beginning of the inactive phase of the light cycle (9 am)). For the stress experiments, $GR^{LoxP/LoxP}$ (n = 4) and GR^{NesCre} (n = 4) mice were subjected to a 30 min restraint stress and sacrificed either in basal conditions (t0) or 30 and 120 min after stress onset. Then, the hippocampus and blood were collected respectively and assayed either for protein extraction or for corticosterone assay. All the experiments were conducted in strict compliance with the European Communities Council Directive of 24 November 1986 (86/609/EEC), and approved by the Aquitaine-Poitou Charentes ethical committee.

Chemicals

The corticosterone used in the various experiments was from Sigma (#C-2505, USA). Corticosterone was used at 10 nM in PC12 cells and at 100 nM in primary culture of hippocampal neurons. The latter concentration was chosen on the basis of preliminary experiments showing that 100 nM corresponded to the EC100 for the nuclear translocation of the GR in this cellular model. MEK1/2 inhibitor (UO126, #9903, Cell Signaling Technology, USA) at a final concentration of 10 μM was used to block the MAPK signaling pathway. 2-Deoxy-D-galactose (2-dGal, #259580, Calbiochem, USA) acting as an inhibitor of synapsin fucosylation was used at a final concentration of 400 mM/hemisphere.

Cell culture and transient siRNA transfection

The PC12 cell line (ATCC CRL-1721) derived from a transplantable rat pheochromocytoma was used. The PC12 cells were trypsinized and seeded on poly-D-lysine coated wells to the appropriate concentration (9×10^5 cells/well) in fresh, antibiotic-free medium (15% horse serum + 2.5% FBS). Transfections of siRNA ON-TARGETplus SMARTpool against Egr-1 (10 nmol, #L-100247-01-0010, Dharmacon, USA), siRNA negative control ON-TARGETplus siCONTROL Non-targeting Pool (10 nmol, #D-001810-10-05, Dharmacon, USA) and siRNA-Alexa Fluor 546 (#1027098, Qiagen, USA) were performed with Lipofectamine 2000 (Invitrogen, France) within OPTI-MEM medium (Invitrogen, France) according to the manufacturer's instructions at a final ratio of 1:1. The percentage of transfection efficiency realized with siRNA-Alexa Fluor 546 was determined by flow cytometry (FACS Calibur, BD, USA). Then 6 hr after transfection the OPTI-MEM medium was changed for a steroid-free culture medium. After 65 hr post-transfection, the cells were treated for 1 hr with 10 nM of corticosterone-HBC (Sigma, USA); proteins were extracted with RIPA buffer containing protease and phosphatase inhibitors and subjected to western blot analysis.

Cultures of hippocampal neurons prepared from E18 Sprague-Dawley rats were plated at a density of 200×10^3 cells/ml on poly-lysine pre-coated coverslips.²⁵ Cultures were maintained in serum-free N2 medium²⁶ and kept at 36.5°C in 5.0 % CO₂ for 7 to 9 days *in vitro* (DIV) before transfection. Neuronal Egr-1 down-regulation by RNA interference was performed by co-transfecting EGFP cDNA (1.6 µg) as a reporter and RNA duplexes (2.4 µg) with Lipofectamine 2000 (15 µl, Invitrogen, France) according to the manufacturer's instructions. Neurons were incubated for 1 hr with this solution in serum-free N2 medium. Forty eight hr

after transfection, the neurons were treated for 1 hr with 100 nM of corticosterone-HBC (Sigma, USA) then siRNA efficiency was checked by immunostaining.

Immunostaining

Neurons were fixed with 4% paraformaldehyde (10 min room temperature (RT)), washed and permeabilized using 0.1% Triton-X100 (5 min RT; Sigma, USA). The cells were incubated with primary antibodies for 1 hr then with secondary antibodies for 30 min at RT. After washing, neurons were mounted with Mowiol (Calbiochem, USA) or ProLong Gold Antifade Reagent containing DAPI (Molecular Probes-Invitrogen, UK). Preparations were kept at 4°C until analysis. GR immunoreactivity (IR), Egr-1-IR and synapsin-I-IR were revealed using a rabbit polyclonal anti-GR antibody (#sc-1004-X, 1/500, Santa Cruz, USA), a rabbit polyclonal anti-Egr-1 antibody (#ARP32241_P50, 1/500, Aviva System biology, USA) and a mouse monoclonal anti-synapsin-I (#106001, 1/400, SYSY: Synaptic System, Germany) respectively. Secondary antibodies with fluorescent labels were from Molecular Probes (UK). Microscopic imaging was performed using an upright epi-fluorescence microscope Leica DMR (Leica Microsystems, Wetzlar, Germany) equipped with objective HCX PL APO 63X oil NA 1.32 and a CoolSnapHQ camera (Roper Scientific, Evry, France). Quantification of the fluorescence was realized using imaging tools from Metamorph software (Universal Imaging Corporation, PA, USA).

Protein extraction from cell culture and tissues

Nuclear and cytoplasmic protein extracts from PC12 cell lines and from mice hippocampi containing protease and phosphatase inhibitors (Sigma, USA) were prepared either using a procedure previously described and validated²⁷ or using the homogenizer Precellys 24 which is a benchtop device dedicated to the grinding, lysis and homogenization of biological

samples (Bertin Technologies, France). The homogenizing protocol in RIPA buffer containing protease and phosphatase inhibitors was 5000 rpm 2x30 sec + 10 sec break using ceramic CK14 beads (Bertin Technologies, France).

Immunoblotting analysis

Cytoplasmic and nuclear proteins suspended in Laemmli buffer were separated by SDS-PAGE (10% gels) and transferred onto PVDF membranes (Immobilon P, #IPVH00010, Millipore, USA). The proteins were revealed with relevant antibodies as previously described.

²⁷ The following Santa Cruz polyclonal antibodies were used: anti-GR (#sc-1004-X; 1/50000 dilution), anti-Egr-1 (#sc-189; 1/1000 dilution), anti-Synapsin-Ia/Ib (#sc-8295; 1/2000 dilution), anti-Synapsin-IIa (#sc-8293; 1/2000 dilution), anti-Synapsin-IIIa (#sc-8292; 1/350 dilution). Aviva System biology (CA, USA) supplied the rabbit polyclonal anti-Egr-1 (ARP36841_T100; 1/4000 dilution). BD Biosciences (USA) supplied the monoclonal anti-Synapsin-I (#611392; 1/10000 dilution) and anti-Synapsin-IIa (#610666; 1/2000 dilution). Upstate supplied the polyclonal anti-MAPK (#06-182; 1/100000 dilution). Polyclonal anti-Phospho-MAPK (#9101S; 1/1000 dilution) and Phospho-Synapsin (#2311; 1/1000 dilution), Pan-synapsin (#2312; 1/2000 dilution) were from Cell Signalling Technology (USA). The Neuronal Class III β -Tubulin (TUJ1) monoclonal antibody (#MMS-435P; 1/20000 dilution) was purchased from Eurogentec. The X-Ray films (Kodak, USA) were quantified by densitometry using a GS-800 scanner (in transmission mode) and the associated Quantity One software (Bio-Rad, CA, USA) according to the manufacturer's instructions.

Corticosterone assay

Plasma corticosterone was quantified by radioimmunoassay (RIA) using a specific corticosterone antibody (ICN Pharmaceuticals, CA, USA) as described elsewhere.²⁷

Behavioral experiments.

Subjects and surgery. C57/Bl6 male mice (n=69, 3-4 months old (IFFA Credo, Arbresle, France) were anaesthetized with ketamine/xylazine (Bayer, Germany) (8 ml/kg, ip). Using a Kopf stereotaxic apparatus, stainless-steel guide cannulae (26-gauge, 8 mm length) were implanted bilaterally 1 mm above the dorsal hippocampus and fixed in place with dental cement. Mice were allowed to recover for at least 8 days before the behavioral experiments.

Contextual Fear Conditioning procedure. All experimental procedures took place during the light portion of the light/dark cycle. Mice were handled daily for 5 min three days before the start of the contextual conditioning procedure. This behavioral procedure has been repeatedly used and fully described in previous studies.^{8,28} Briefly, each animal was placed in the conditioning chamber for 4 min during which it received 2 footshocks which never co-occurred with 2 tone (63 db, 1 KHz, 15 sec) delivery. Each animal was then returned to its home cage. Twenty-four hr later, the mice were re-exposed to the conditioning chamber and the behavior of the subjects was continuously recorded on video tape for off-line scoring of freezing. The conditioning procedure consisted of a tone (CS)-shock (US) unpairing known to favor a context-US association.²⁸ Animals received one of two shock regimens: either 0.3 mA, 50 Hz, 3 sec (low shock intensity group) or 0.7 mA, 50 Hz, 3 sec (high shock intensity group). Freezing behavior, used as an index of conditioned fear, was calculated as a percentage (\pm sem) of the total time spent freezing during the first 4 min-period of the retention test.

Microinjections. Immediately after the acquisition of fear conditioning, animals were placed in their home cage and randomly divided into four groups according to the intra-hippocampal infusion they received: 1. Vehicle (1% DMSO in artificial CSF), 2. corticosterone (10 ng/side), 3. 2-dGal (400 mM/side), 4. corticosterone + 2-dGal. Infusions (0.3 μ l/hemisphere)

were performed at a constant rate (0.1 $\mu\text{l}/\text{min}$) through 32-gauge stainless-steel cannulae (9 mm) that were inserted into the guide cannulae and left in place for 2 min after the end of the infusion. Corticosterone (2-Hydroxypropyl-beta-cyclodextrin (HBC) complex; Sigma-Aldrich) and 2-dGal were dissolved in artificial CSF (145 mM NaCl, 2.7 mM KCl, 1 mM MgCl_2 , 1.2 mM CaCl_2 and 0.2 mM $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$ buffered at pH 7.4).

Histology. After completion of the behavioral study, animals were given an overdose of ketamin (24 ml/kg) and transcardially perfused with physiological saline, followed by 10% buffered formalin. Brains were post-fixed in formalin-saccharose 30% solution for 1 week, frozen, cut coronally on a sliding microtome into 50 μm sections that were mounted on a gelatin-coated slide, and stained with thionine in order to evaluate the cannulae placements.

Statistics

Statistical analyses were performed using analysis of variance (ANOVA) followed by the Newman Keuls, Duncan or Scheffe post-hoc test for pairwise comparisons when appropriate. The Student's *t*-test was used for pairwise comparisons.

Results

GR in the hippocampus regulate synapsin-I expression and phosphorylation

Using western blot, we compared the expression of members of the synapsin family in the hippocampus of $\text{GR}^{\text{NesCre}}$ mice, in which the expression of the GR protein is suppressed in the entire brain but not in other tissues using the Cre/LoxP system²³ and in control littermates.

In basal conditions, using pan-synapsin, an antibody that detects all synapsin isoforms, we found a reduced expression of these proteins and of their phosphorylated counterparts in $\text{GR}^{\text{NesCre}}$ mutant mice. However, when more specific antibodies were used we found a selective decrease in the expression of synapsin-Ia/Ib but not of synapsin-IIa and -IIIa (Figure

1a-b). Finally levels of the transcription factor Egr-1 were also reduced in GR^{NesCre} mutant mice.⁸

Since, under physiological conditions, activation of the GR is mainly obtained during stress we also analyzed changes in synapsin levels after an acute restraint stress (30 min). For this experiment blood and hippocampal extracts were collected before stress (t₀) and 30 min or 120 min after stress onset. As previously shown in wild-type animals, stress induced the translocation of the GR and the expression of Egr-1.⁸ In addition, stress also induced an increase in the expression and phosphorylation of pan-synapsin as well of synapsin-Ia/Ib, synapsin-IIa and synapsin-IIIa (Figure 1c). In GR^{NesCre} mutant mice, lacking the GR, stress-induced expression of Egr-1 was profoundly decreased. In addition, stress-induced increase in phosphorylated synapsin was suppressed and the increase in the expression of synapsin-Ia/Ib was greatly reduced. In contrast, expression of synapsin-IIa and -IIIa were not significantly modified by the absence of the GR (Figure 1c).

These modifications of synapsin and Egr-1 levels were not due to a decrease in stress-induced glucocorticoid secretion in mutant animals. Thus, as has been demonstrated several times,^{8;23} basal and stress-induced glucocorticoid levels were higher in GR^{NesCre} mice (Table 1) indicating that the glucocorticoid-negative feedback loop is profoundly impaired in GR mutant mice.

These results indicate that both in basal conditions and after stress GR positively control the phosphorylation of synapsin and the expression of synapsin-Ia/Ib but not of other members of this family. Through this double effect, GR may powerfully increase neurotransmitter release. Indeed, the levels of synapsin-Ia/Ib positively control the pool of readily releasable synaptic vesicles^{17;29;30} and the phosphorylation of synapsin triggers neurotransmitter release.²⁹

Glucocorticoid-induced synapsin-I phosphorylation depends on MAPK activity

In the hippocampus in basal conditions (Figure 1a-b) and after stress (Figure 1c) GR regulate the expression of both Egr-1⁸ and synapsin-I. Since GR are phasically activated by the increase in circulating glucocorticoids that occurs during the active phase of the circadian cycle or in response to stress,^{1;2;5} we analyzed whether the acute administration of glucocorticoids would stimulate the expression of synapsin-I. For this experiment we used the PC12 cell line, which expresses the GR and synapsin-I and -II isoforms.^{21;22} In this *in vitro* model, administration of 10 nM of corticosterone which replicates the concentration of the hormone induced by stress, increases the expression of Egr-1, the phosphorylation of synapsin, the expression levels of synapsin-I but not of synapsin-II (Figure 2a).

Phosphorylation of synapsin plays a major role in controlling pre-synaptic vesicle trafficking by inhibiting the dynamic association between the synaptic vesicles and the actin cytoskeleton.^{13;29-32} We investigated here whether GR-induced activation of the MAPK pathway⁸ was responsible for the GR-induced phosphorylation of synapsin-I. In order to test this hypothesis, we studied glucocorticoid-induced phosphorylation of synapsin-I in PC12 cells in the presence of a selective inhibitor of MAPK phosphorylation, UO126. UO126 was used at 10 μ M, a dose corresponding to the EC100 of this compound (Figure 2b-c).³³

PC12 cells were exposed for 1 hr to 10 nM corticosterone in the presence or absence of UO126, after which the proteins were extracted and analyzed by western blot. In the absence of UO126, we found that 1 hr exposure to corticosterone induced both phosphorylation of the MAPK pathway (P-Erk1/2) and of synapsin-I. Blocking the phosphorylation of the MAPK pathway almost completely blocked the phosphorylation of synapsin but not the expression of synapsin-I (Figure 2b-c).

Glucocorticoid-induced expression of synapsin-I depends on Egr-1

In these experiments we analyzed whether GR increased synapsin-I expression through Egr-1. For this purpose we silenced Egr-1 expression by using a small interfering RNA strategy. PC12 cells were transfected with siRNA (-ctrl or -Egr-1) 65 hr (t_{65}) before exposure to 10 nM corticosterone. After 1 hr of exposure to corticosterone (t_1) the cells were harvested and proteins immediately extracted (Figure 3a). Transfection efficiency was measured by flow cytometry using siRNA-Alexa Fluor 546. With our protocol it was possible to transfect 74% of PC12 cells with siRNA with no significant increase in cell death (Figure 3b). Corticosterone increased both Egr-1 and synapsin-I expressions in cells transfected with control siRNA. In contrast, in cells in which Egr-1 expression was silenced the glucocorticoid-mediated increase in synapsin-I expression was suppressed (Figure 3c-d).

We then analyzed if these effects could also be found in primary culture of hippocampal neurons. We found that also in this cellular context 1 hr of corticosterone treatment promotes the nuclear translocation of the GR proteins (Figure 3e) and a concomitant and proportional increase in both Egr-1 and synapsin-I proteins (Figure 3f-i). Hippocampal neurons were then co-transfected with siRNA (-ctrl or -Egr-1) and EGFP 48 hr (t_{48}) before exposure to 1 hr of 100 nM corticosterone. siRNA effects were checked by immunostaining (Figure 3a, j-m). Silencing Egr-1 expression by RNA interference in hippocampal neurons also suppressed the corticosterone-mediated increase in synapsin-I expression (Figure 3j-m). These results confirm that glucocorticoid-induced expression of synapsin-I also depends on Egr-1 in neuronal cells.

In conclusion these results show that the GR/Egr-1/MAPK cascade controls synapsin-I by complementary mechanisms. Activation of Egr-1 by the GR specifically increases the levels of expression of synapsin-Ia/Ib, in parallel GR-induced MAPK activation induces the phosphorylation of synapsin-I.

Synapsin-I mediates GR-induced enhancement of fear conditioning

The previous results suggest that synapsin-I could be one of the molecular effectors through which activation of the GR/Egr-1/MAPK molecular cascade enhances stress-related memories. In order to test this hypothesis we analyzed the effects of a modification of the levels of expression of synapsin-I on the retention of fear conditioning.

Synapsin-Ia/Ib has recently been shown to be the major Fuc α (1-2)Gal glycoprotein in the hippocampus. Inhibiting synapsin-Ia/Ib fucosylation using 2-dGal critically impacts both its cellular half-life expression and turnover in pre-synaptic nerve terminals.^{24;34}

We therefore examined the effects of an intra-hippocampal infusion of 2-dGal both on the expression of synapsin-I and on the enhancement of hippocampal-dependent contextual fear conditioning^{8;28} induced by corticosterone (Figure 4). Intra-hippocampal infusion of corticosterone immediately after training increased the conditioned fear induced by an electric shock of low intensity (0.3 mA footshock) inducing a response similar to the one obtained with an electric shock of high intensity (0.7 mA footshock) (Figure 4a). However, the enhancing effect of corticosterone on contextual fear conditioning was completely abolished when 2-dGal (400 mM/side) was infused concomitantly with corticosterone (Figure 4a). In order to verify the effect of corticosterone and 2-dGal infusion on synapsin-I, a group of animals was sacrificed 2 hr after the hippocampal infusions. Hippocampal proteins were then extracted and analyzed by western blot. We found that hippocampal infusion of corticosterone stimulates synapsin-Ia/Ib expression and that this increase was largely blocked by 2-dGal infusion (Figure 4b).

In conclusion, these results indicate that synapsin-Ia/Ib is one of the effectors through which the GR/Egr-1/MAPK cascade enhances stress-related memory.

Discussion

The central finding of this work is the identification of synapsin-Ia/Ib as an important molecular effector of glucocorticoid-mediated behavioral effects of stress. In particular, synapsin-Ia/Ib appears as one of the effectors through which the GR/Egr-1/MAPK cascade enhances stress-related memory. Thus, combining *in vitro* and *in vivo* approaches, we found that both in basal conditions and after stress, GR, through the activation of Egr-1 expression and MAPK phosphorylation, increase synapsin-Ia/Ib expression and phosphorylation respectively, and that these events were necessary for the increase in stress-related memories induced by glucocorticoids.

With the results of the present experiment extending those published in a previous paper we can singling out a complete molecular cascade that is involved in the mediation of stress-related memory (Figure 5). We have previously shown that stress-induced glucocorticoids increase stress-related memories by activating the GR which trigger a cascade of events starting with an increase in the expression of the transcription factor Egr-1 and followed by the activation of the MAPK pathway.⁸ We show here that the activation of Egr-1 by glucocorticoids increases the level of synapsin-I and that the activation of the MAPK pathway induces the phosphorylation of this protein. Silencing Egr-1 expression thus represses the increase in synapsin-Ia/Ib expression induced by glucocorticoids both in PC12 cells and in primary culture of hippocampal neurons. This effect is probably a direct one since Egr-1 binds to responsive elements in the promoter region of the syn1 gene and is able to activate its transcription *in vitro*.³⁵⁻³⁷ In parallel, Mitogen-Activated Protein Kinase (Erk1/2) induces the phosphorylation of synapsin-I but does not modify its expression. Thus, UO126-mediated blockade of MAPK phosphorylation largely reduced glucocorticoid-induced synapsin-I phosphorylation but did not change the expression of synapsin-I.

Synapsin-I undergoes several cycles of phosphorylation/dephosphorylation on serine residues. *In vitro* studies have shown that these phosphorylations can involve many different kinases

including cAMP-dependent protein kinase (PKA), Ca²⁺/calmodulin-dependent protein kinase (CaMK) I, CaMK II, CaMK IV, proline-directed kinase, mitogen-activated protein kinase (Erk1/2), cyclin-dependent kinase (cdk) 1 and 5, tyrosine kinase c-Src, protein kinase C (PKC) and p21-activated kinases (PAKs).^{13;14;30;38-40} It is unlikely that *in vivo* all these kinases phosphorylate synapsin-I at the same time, in all probability their actions depend on the type of stimulation.³⁰ Our results indicate that *in vivo* the MAPK pathway plays an important role in stress-induced phosphorylation of synapsin-I.

Our data also show that the regulation of synapsin-I by the GR/Egr-1/MAPK cascade increases stress-related memory. Inhibition of synapsin-I expression by 2-dGal thus abolished glucocorticoid-induced increase in contextual fear conditioning. These results extend previous knowledge and start clarifying the role of different synapsin isoforms in memory formation. Thus it has been previously shown that synapsin-deficient *D. melanogaster* flies display a deficit in learning and memory,^{20;41} and synapsin-I null mice display an increase in age-dependent cognitive impairments.⁴² In parallel, the learning of a spatial memory task increases expression of synapsin-I in the hippocampus^{19;43} and transgenic mice expressing a constitutively active form of H-ras (H-ras^{G12V}), exhibit increased Erk-dependent phosphorylation of synapsin-I and dramatic enhancements in hippocampus-dependent learning that can be reversed by the deletion of synapsin-I.⁴⁴ Our data indicate that synapsin-Ia/Ib play an important role in the formation of hippocampal-mediated stress-related memory. Since all the synapsin isoforms (-I, -II, -III) are expressed within the hippocampus^{14;45} it could be argued that other synapsins could be modified by 2-dGal and consequently be involved in the modulation of fear memory. In fact, synapsin fucosylation sites are localized within the D domain of synapsin, a region that is found only in the synapsin-Ia/Ib isoforms and not in the other isoforms.¹⁴ In addition, western blot analysis of adult hippocampal lysates from synapsin-I-deficient mice revealed a loss of the fucosylated bands, confirming

that synapsins Ia and Ib are the predominant Fuc α (1-2)Gal glycoproteins in the hippocampus.

24

The inhibition of stress-related memory by blocking synapsin-I fucosylation also supports growing evidence indicating that carbohydrate post-translational modifications to *N*- and *O*-linked glycoproteins could participate in cognitive processes such as learning and memory^{46;47} and in synaptic plasticity.^{34;48;49}

The GR/Egr-1/MAPK/Synapsin-I (GEMS) cascade enhances stress-related memory very likely by increasing the release of neurotransmitters. Unphosphorylated synapsins, bind synaptic vesicles (SVs) tightly to actin. These synapsin-actin-bound SVs constitute the readily releasable pool of vesicles that is positively regulated by synapsin-I levels.⁵⁰ Phosphorylation of synapsin-I then releases SVs from the synapsin-actin-bound state³¹ which in turn allows their transport to the sites of exocytosis and the release of neurotransmitters.^{29;30} Consequently, the increase in the level of synapsin-I induced by GR-induced Egr-1 will increase the stock of neurotransmitters readily releasable and its phosphorylation by GR-activated MAPK triggers its actual release.

Several lines of evidences suggest that the enhancement of stress-related memory mediated by the GEMS pathway is mediated by glutamate release. Firstly, in the hippocampus synapsin-I is exclusively expressed in glutamatergic pyramidal neurons, but is absent from GABA interneurons.⁵¹⁻⁵³ Secondly, intraperitoneal administration of corticosterone increases glutamate but not GABA release in the hippocampus.⁵⁴ This effect depends on GR activation since it can be prevented by pre-treatment with the GR antagonist mifepristone.⁵⁵ Finally, synapsin-I has been shown to regulate glutamate release from SVs¹⁷ via a synaptic activity-dependent mechanism⁵⁶ that is impaired in mice lacking the synapsin-I gene.^{16;17;57} An increase in glutamate release triggered by the GEMS pathway can in turn modify synaptic plasticity which is probably involved in the enhancement of stress-related memory.¹⁸

In conclusion our data provide a complete molecular pathway GEMS (GR/Egr-1/MAPK/Synapsin-Ia/Ib) (Figure 5) through which glucocorticoids can enhance the memory of stressful events⁵⁻⁸ and, reveal synapsin-Ia/Ib as an important molecular effector of the behavioral consequences of stress .

Acknowledgments

The authors thank Drs. D. Choquet and O. Thoumine for invaluable help with the experiments using the hippocampal cultures. We acknowledge the technical support of M. Petit, C. Breillat, B. Teissier and D. Bouchet. The microscopy was done in the Bordeaux Imaging Center of the University of Bordeaux II. The help of P. Legros, C. Poujol and L. Malicieux is acknowledged. This work was supported by INSERM, Agence Nationale pour la Recherche (ANR, contract HICOMET), Bordeaux Institute for Neurosciences (IFR8), the University *Victor Segalen-Bordeaux2* and Conseil Régional d'Aquitaine, the 6th Framework European Programme (PHECOMP, contract n°FP6-037669).

References

1. McEwen BS. Effects of adverse experiences for brain structure and function. *Biol.Psychiatry* 2000; **48**: 721-31.
2. De Kloet ER. Hormones, brain and stress. *Endocr.Regul.* 2003; **37**: 51-68.
3. Piazza PV, Le Moal M. Glucocorticoids as a biological substrate of reward: physiological and pathophysiological implications. *Brain Res.Brain Res.Rev.* 1997; **25**: 359-72.
4. Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch.Gen.Psychiatry* 2000; **57**: 925-35.
5. De Kloet ER, Oitzl MS, Joels M. Stress and cognition: are corticosteroids good or bad guys? *Trends Neurosci.* 1999; **22**: 422-6.
6. McGaugh JL, Roozendaal B. Role of adrenal stress hormones in forming lasting memories in the brain. *Curr.Opin.Neurobiol.* 2002; **12**: 205-10.

7. Oitzl MS, Reichardt HM, Joels M, De Kloet ER. Point mutation in the mouse glucocorticoid receptor preventing DNA binding impairs spatial memory. *Proc.Natl.Acad.Sci.U.S.A.* 2001; **98**: 12790-5.
8. Revest JM, Di Blasi F, Kitchener P, Rouge-Pont F, Desmedt A, Turiault M *et al.* The MAPK pathway and Egr-1 mediate stress-related behavioral effects of glucocorticoids. *Nat.Neurosci.* 2005; **8**: 664-72.
9. Holsboer F. The corticosteroid receptor hypothesis of depression. *Neuropsychopharmacology* 2000; **23**: 477-501.
10. Wei Q, Lu XY, Liu L, Schafer G, Shieh KR, Burke S *et al.* Glucocorticoid receptor overexpression in forebrain: A mouse model of increased emotional lability. *Proc.Natl.Acad.Sci.U.S.A* 2004; **101**: 11851-6.
11. Funder JW. Glucocorticoid receptors. *J.Steroid Biochem.Mol.Biol.* 1992; **43**: 389-94.
12. Tronche F, Kellendonk C, Reichardt HM, Schutz G. Genetic dissection of glucocorticoid receptor function in mice. *Curr.Opin.Genet.Dev.* 1998; **8**: 532-8.
13. Evergren E, Benfenati F, Shupliakov O. The synapsin cycle: A view from the synaptic endocytic zone. *J.Neurosci.Res.* 2007; **85**: 2648-56.
14. Valtorta F, Benfenati F, Greengard P. Structure and function of the synapsins. *J.Biol.Chem.* 1992; **267**: 7195-8.
15. Ferreira A, Rapoport M. The synapsins: beyond the regulation of neurotransmitter release. *Cell Mol.Life Sci.* 2002; **59**: 589-95.
16. Jovanovic JN, Czernik AJ, Fienberg AA, Greengard P, Sihra TS. Synapsins as mediators of BDNF-enhanced neurotransmitter release. *Nat.Neurosci.* 2000; **3**: 323-9.
17. Nichols RA, Chilcote TJ, Czernik AJ, Greengard P. Synapsin I regulates glutamate release from rat brain synaptosomes. *J.Neurochem.* 1992; **58**: 783-5.
18. Howland JG, Wang YT. Synaptic plasticity in learning and memory: stress effects in the hippocampus. *Prog.Brain Res.* 2008; **169**: 145-58.
19. John JP, Sunyer B, Hoyer H, Pollak A, Lubec G. Hippocampal synapsin isoform levels are linked to spatial memory enhancement by SGS742. *Hippocampus.* 2009; **19**: 731-8.
20. Godenschwege TA, Reisch D, Diegelmann S, Eberle K, Funk N, Heisenberg M *et al.* Flies lacking all synapsins are unexpectedly healthy but are impaired in complex behaviour. *Eur.J.Neurosci.* 2004; **20**: 611-22.
21. Carpentier R, Sacchetti P, Segard P, Staels B, LeFebvre P. The glucocorticoid receptor is a co-regulator of the orphan nuclear receptor Nurr1. *J.Neurochem.* 2008; **104**: 777-89.
22. Das KP, Freudrich TM, Mundy WR. Assessment of PC12 cell differentiation and neurite growth: a comparison of morphological and neurochemical measures. *Neurotoxicol.Teratol.* 2004; **26**: 397-406.
23. Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban PC *et al.* Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. *Nat.Genet.* 1999; **23**: 99-103.

24. Murrey HE, Gama CI, Kalovidouris SA, Luo WI, Driggers EM, Porton B *et al.* Protein fucosylation regulates synapsin Ia/Ib expression and neuronal morphology in primary hippocampal neurons. *Proc.Natl.Acad.Sci.U.S.A.* 2006; **103**: 21-6.
25. Banker GA, Cowan WM. Rat hippocampal neurons in dispersed cell culture. *Brain Res.* 1977; **126**: 397-42.
26. Bottenstein JE, Sato GH. Growth of a rat neuroblastoma cell line in serum-free supplemented medium. *Proc.Natl.Acad.Sci.U.S.A.* 1979; **76**: 514-7.
27. Kitchener P, Di Blasi F, Borrelli E, Piazza PV. Differences between brain structures in nuclear translocation and DNA binding of the glucocorticoid receptor during stress and the circadian cycle. *Eur.J.Neurosci.* 2004; **19**: 1837-46.
28. Desmedt A, Garcia R, Jaffard R. Differential modulation of changes in hippocampal-septal synaptic excitability by the amygdala as a function of either elemental or contextual fear conditioning in mice. *J.Neurosci.* 1998; **18**: 480-7.
29. Hosaka M, Hammer RE, Sudhof TC. A phospho-switch controls the dynamic association of synapsins with synaptic vesicles. *Neuron.* 1999; **24**: 377-87.
30. Yamagata Y. New aspects of neurotransmitter release and exocytosis: dynamic and differential regulation of synapsin I phosphorylation by acute neuronal excitation in vivo. *J.Pharmacol.Sci.* 2003; **93**: 22-9.
31. Jovanovic JN, Benfenati F, Siow YL, Sihra TS, Sanghera JS, Pelech SL *et al.* Neurotrophins stimulate phosphorylation of synapsin I by MAP kinase and regulate synapsin I-actin interactions. *Proc.Natl.Acad.Sci.U.S.A* 1996; **93**: 3679-83.
32. Turner KM, Burgoyne RD, Morgan A. Protein phosphorylation and the regulation of synaptic membrane traffic. *Trends Neurosci.* 1999; **22**: 459-64.
33. Favata MF, Horiuchi KY, Manos EJ, Daulerio AJ, Stradley DA, Feeser WS *et al.* Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J.Biol.Chem.* 1998; **273**: 18623-32.
34. Murrey HE, Hsieh-Wilson LC. The chemical neurobiology of carbohydrates. *Chem.Rev.* 2008; **108**: 1708-31.
35. Al Sarraj A, Day RM, Thiel G. Specificity of transcriptional regulation by the zinc finger transcription factors Sp1, Sp3, and Egr-1. *J.Cell Biochem.* 2005; **94**: 153-67.
36. James AB, Conway AM, Thiel G, Morris BJ. Egr-1 modulation of synapsin I expression: permissive effect of forskolin via cAMP. *Cell Signal.* 2004; **16**: 1355-62.
37. Thiel G, Schoch S, Petersohn D. Regulation of synapsin I gene expression by the zinc finger transcription factor zif268/egr-1. *J.Biol.Chem.* 1994; **269**: 15294-301.
38. Browning MD, Dudek EM. Activators of protein kinase C increase the phosphorylation of the synapsins at sites phosphorylated by cAMP-dependent and Ca²⁺/calmodulin-dependent protein kinase in the rat hippocampal slice. *Synapse.* 1992; **10**: 62-70.
39. Sakurada K, Kato H, Nagumo H, Hiraoka H, Furuya K, Ikuhara T *et al.* Synapsin I is phosphorylated at Ser603 by p21-activated kinases (PAKs) in vitro and in PC12 cells stimulated with bradykinin. *J.Biol.Chem.* 2002; **277**: 45473-9.

40. Severin SE, Jr., Moskvitina EL, Bykova EV, Lutzenko SV, Shvets VI. Synapsin I from human brain. Phosphorylation by Ca²⁺, phospholipid-dependent protein kinase. *FEBS Lett.* 1989; **258**: 223-6.
41. Michels B, Diegelmann S, Tanimoto H, Schwenkert I, Buchner E, Gerber B. A role for Synapsin in associative learning: the *Drosophila* larva as a study case. *Learn.Mem.* 2005; **12**: 224-31.
42. Corradi A, Zanardi A, Giacomini C, Onofri F, Valtorta F, Zoli M *et al.* Synapsin-I- and synapsin-II-null mice display an increased age-dependent cognitive impairment. *J.Cell Sci.* 2008; **121**: 3042-51.
43. Gomez-Pinilla F, So V, Kesslak JP. Spatial learning induces neurotrophin receptor and synapsin I in the hippocampus. *Brain Res.* 2001; **904**: 13-9.
44. Kushner SA, Elgersma Y, Murphy GG, Jaarsma D, van Woerden GM, Hojjati MR *et al.* Modulation of presynaptic plasticity and learning by the H-ras/extracellular signal-regulated kinase/synapsin I signaling pathway. *J.Neurosci.* 2005; **25**: 9721-34.
45. Kao HT, Porton B, Czernik AJ, Feng J, Yiu G, Haring M *et al.* A third member of the synapsin gene family. *Proc.Natl.Acad.Sci.U.S.A.* 1998; **95**: 4667-72.
46. Lorenzini CG, Baldi E, Bucherelli C, Sacchetti B, Tassoni G. 2-Deoxy-D-galactose effects on passive avoidance memorization in the rat. *Neurobiol.Learn.Mem.* 1997; **68**: 317-24.
47. Rose SP, Jork R. Long-term memory formation in chicks is blocked by 2-deoxygalactose, a fucose analog. *Behav.Neural Biol.* 1987; **48**: 246-58.
48. Tallent MK, Varghis N, Skorobogatko Y, Hernandez-Cuebas L, Whelan K, Vocadlo DJ *et al.* In vivo modulation of O-GlcNAc levels regulates hippocampal synaptic plasticity through interplay with phosphorylation. *J.Biol.Chem.* 2009; **284**: 174-81.
49. Krug M, Jork R, Reymann K, Wagner M, Matthies H. The amnesic substance 2-deoxy-D-galactose suppresses the maintenance of hippocampal LTP. *Brain Res.* 1991; **540**: 237-42.
50. Baldelli P, Fassio A, Valtorta F, Benfenati F. Lack of synapsin I reduces the readily releasable pool of synaptic vesicles at central inhibitory synapses. *J.Neurosci.* 2007; **27**: 13520-31.
51. Bogen IL, Boulland JL, Mariussen E, Wright MS, Fonnum F, Kao HT *et al.* Absence of synapsin I and II is accompanied by decreases in vesicular transport of specific neurotransmitters. *J.Neurochem.* 2006; **96**: 1458-66.
52. Bogen IL, Jensen V, Hvalby O, Walaas SI. Synapsin-dependent development of glutamatergic synaptic vesicles and presynaptic plasticity in postnatal mouse brain. *Neuroscience.* 2009; **158**: 231-41.
53. Melloni RH, Jr., Hemmendinger LM, Hamos JE, DeGennaro LJ. Synapsin I gene expression in the adult rat brain with comparative analysis of mRNA and protein in the hippocampus. *J.Comp Neurol.* 1993; **327**: 507-20.
54. Venero C, Borrell J. Rapid glucocorticoid effects on excitatory amino acid levels in the hippocampus: a microdialysis study in freely moving rats. *Eur.J.Neurosci.* 1999; **11**: 2465-73.
55. Wang CC, Wang SJ. Modulation of presynaptic glucocorticoid receptors on glutamate release from rat hippocampal nerve terminals. *Synapse.* 2009; **63**: 745-51.

56. Chi P, Greengard P, Ryan TA. Synaptic vesicle mobilization is regulated by distinct synapsin I phosphorylation pathways at different frequencies. *Neuron*. 2003; **38**: 69-78.
57. Li L, Chin LS, Shupliakov O, Brodin L, Sihra TS, Hvalby O *et al*. Impairment of synaptic vesicle clustering and of synaptic transmission, and increased seizure propensity, in synapsin I-deficient mice. *Proc.Natl.Acad.Sci.U.S.A.* 1995; **92**: 9235-9.

Legends

Figure 1 Comparison of the expression of the synapsin proteins family and Egr-1 under basal condition and after stress between wild-type (WT) and GR^{NesCre} mice in which the glucocorticoid receptors (GR) were suppressed in the entire brain. (a) Western blot analyses of hippocampus extracts for GR, Egr-1, phosphorylated-synapsin (P-synapsin), total synapsin (Pan-synapsin), synapsin-Ia/Ib, synapsin-IIa, synapsin-IIIa isoforms were performed using specific antibodies. β III-tubulin was used as a loading control. (b) Quantification by densitometry (optical density, OD, mean \pm sem) of the corresponding X-Ray films. Differences between WT ($n = 3$) versus GR^{NesCre} ($n = 3$) mice were all statistically significant, $*P < 0.05$, $**P < 0.005$, $***P < 0.001$. (c) Stress-induced activation of the GR in the hippocampus stimulates synapsin-I expression and phosphorylation. Comparison of the expression of synapsin isoforms in wild-type (WT) and GR^{NesCre} mice lacking the expression of GR in neurons, before (t0) and 30 and 120 min after the onset of 30 min of restraint stress. β III-tubulin was used as a loading control. Nuclear (for GR and Egr-1) and cytoplasmic hippocampal extracts were analyzed by western blot (left panel) and the corresponding X-Ray films were quantified by densitometry (right panel, optical density, OD, mean \pm sem, $n = 6$). $*P < 0.05$; $**P < 0.005$, $***P < 0.001$, WT versus GR^{NesCre} Newman-Keuls and Duncan post-hoc test after ANOVA.

Figure 2 Corticosterone-induced phosphorylation of synapsin-I involves the MAPK pathway. (a) Total extracts from PC12 cells were analyzed by western blot 1 and 3 hr after

treatment with 10 nM corticosterone. The corresponding X-Ray films were quantified by densitometry (optical density, OD, mean \pm sem, $n = 3$). $*P < 0.05$; $***P < 0.001$, in comparison to basal levels. **(b)** The effects of corticosterone administration (10 nM) on Erk1/2, synapsin-I expressions and their phosphorylated counterparts were analyzed by western blot in PC12 cells exposed or not to the inhibitor of the MAPK pathway UO126 (10 μ M). 1 hr after the exposure to 10 nM corticosterone, proteins were extracted and analyzed with relevant antibodies. β III-tubulin was used as a loading control. **(c)** Quantification by densitometry (optical density OD, mean \pm sem, $n = 4$) of the western blot experiments. $*P < 0.05$, $**P < 0.005$, $***P < 0.001$ in comparison with untreated cells (ANOVA).

Figure 3 Corticosterone stimulation of synapsin-I expression depends on Egr-1. **(a)** Schematic illustration of the protocol used to silence Egr-1 expression by using a small interfering RNA strategy. PC12 cells and hippocampal neurons were transfected with siRNA (-Egr-1, -ctrl, -Alexa Fluor 546, or EGFP). 48 to 65 hr later, the cells were exposed to corticosterone (Cort) for 1 hr then proteins were analyzed by western blot (PC12) or immunostaining (neurons). **(b)** The transfection efficiency in PC12 cells was evaluated by flow cytometry after transfecting siRNA-Alexa Fluor 546. **(c)** Western blot analyses of the effects of corticosterone administration (10 nM) on synapsin-I and Egr-1 expressions after silencing Egr-1 expression using a siRNA strategy. β III-tubulin was used as a loading control. **(d)** Quantification by densitometry (optical density OD, means \pm sem, $n = 3$) of the western blot experiments. $*P < 0.05$, $**P < 0.005$, in comparison with untreated cells (ANOVA). **(e)** Immunostaining of hippocampal neurons cultured for 9 DIV showing the sub-localization of the GR proteins (green) in absence or presence of 100 nM of corticosterone for 1 hr. Nuclei were counterstained with DAPI (blue). Merge of the two signals is shown. Scale bar = 25 μ m. **(f)** Co-immunolocalization of Egr-1 (green) and synapsin-I (red) proteins on hippocampal

neurons cultured for 9 DIV in absence or presence of 100 nM of corticosterone during 1 hr. Scale bar = 25 μ m. **(g-h)** Quantification of the fluorescence intensity (AU; Arbitrary Unit) of Egr-1 **(g)** and synapsin-I **(h)** -immunoreactive hippocampal neurons in absence or presence of 100 nM of corticosterone (Cort) during 1 hr. **(i)** Following corticosterone treatment the expression of Egr-1 and synapsin-I proteins in Egr-1-expressing hippocampal neurons increased proportionally. Values shown are means \pm sem. $**P < 0.005$, $***P < 0.001$, in comparison with untreated cells. **(j)** Immunofluorescence analysis of hippocampal neurons treated with corticosterone (100 nM) for 1 hr and previously co-transfected with EGFP cDNA and siRNA control or siRNA silencing Egr-1 protein. Nuclei were counterstained with DAPI (blue). The white arrows show the mapping of the synapsin-I protein in EGFP positive hippocampal neurons transfected either with siRNA-ctrl or siRNA-Egr-1. Scale bar = 25 μ m. **(k-l)** Quantification of the fluorescence intensity (AU; Arbitrary Unit) of Egr-1 **(k)** and synapsin-I **(l)** -immunoreactive hippocampal neurons transfected with siRNA-ctrl or siRNA-Egr-1 in presence of 100 nM of corticosterone (Cort) during 1 hr. **(m)** Proportional decrease between Egr-1 and synapsin-I proteins in neurons in which Egr-1 expression was silenced compared to siRNA-ctrl transfection. Values shown are means \pm sem. $**P < 0.005$, $***P < 0.001$, in comparison with siRNA-ctrl transfected cells.




Figure 4 Synapsin-Ia/Ib mediates the enhancement of contextual fear conditioning induced by glucocorticoids. **(a)** Percentage of conditioned freezing (mean \pm sem) displayed by mice re-exposed for 4 min to the conditioning context 24 hr after conditioning with either high (0.7 mA footshock; $n = 12$) or low shock intensity (0.3 mA footshock; $n = 11$) and receiving a post-training intra-hippocampal infusion of either vehicle ($n = 11$), 2-dGal ($n = 11$; 400 mM/side), corticosterone ($n = 12$; 10 ng/side) or corticosterone + 2-dGal ($n = 12$). **(b)** Western blot analysis and quantification of synapsin-Ia/Ib hippocampal expression in




presence or not of the inhibitor of fucosylation 2-dGal (400 mM/side). β III-tubulin was used as a loading control. 2 hr post hippocampal infusion of 2-dGal proteins were extracted, analysed by western blot and quantified by densitometry (optical density OD, mean \pm sem, $n = 4$). $**P < 0.005$, $***P < 0.0001$, in comparison with all the other groups, Newman-Keuls and Scheffe post-hoc test after ANOVA.


Figure 5 Schematic model of GEMS (GR/Egr-1/MAPK/Synapsin-I) molecular cascade mediating the enhancement of stress-related memories. During stress the increase in glucocorticoid secretion (1) activates the glucocorticoid receptors (GR) inducing its nuclear translocation (2). GR trigger the synthesis of the transcription factor Egr-1 (3) and then the activation of the MAPK pathway (4) which further increases Egr-1 production (5). Egr-1 stimulates the transcription of synapsin-Ia/Ib (6) which increases the number of synaptic vesicles bound to actin and in this way the pool of readily releasable neurotransmitters. Activated MAPK do not modify synapsin-I expression but induce their phosphorylation (7). Phosphorylated synapsin-Ia/Ib releases synaptic vesicles bound to actin (8) allowing their transport to the presynaptic membrane (9) and neurotransmitter release (10). The double action of GR-induced Egr-1 expression and MAPK activation on both synapsin-I synthesis and phosphorylation can guarantee a sustained increase in neurotransmitter release. Specifically, an increase in glutamate release is likely to mediate the enhancement of stress-related memories that can be blocked both by inhibiting MAPK phosphorylation and synapsin-I expression. This model is based on several studies, including ours which are discussed in the main text.



Table 1 Plasma corticosterone concentrations ($\mu\text{g}/100\text{ml}$) induced by a 30 min restraint stress in control littermate (WT) and GR^{NesCre} mice. Blood was collected before the stress (t0), 30


and 120 min after stress onset. Values shown are means \pm sem. * $P < 0.05$, ** $P < 0.001$, compared to respective pre stress levels (t0).











Time (min)	Plasma Corticosterone ($\mu\text{g}/100\text{ ml}$)	
	WT	GR ^{NesCre}
0	3.37 \pm 0.17	25.40 \pm 4.96
30	22.25 \pm 3.22; **	51.77 \pm 6.93; *
120	14.16 \pm 2.73; *	39.89 \pm 0.99