



HAL
open science

The Fujita phenomenon in exterior domains under the dynamical boundary conditions

Jean-François Rault

► **To cite this version:**

Jean-François Rault. The Fujita phenomenon in exterior domains under the dynamical boundary conditions. 2009. hal-00524377

HAL Id: hal-00524377

<https://hal.science/hal-00524377>

Preprint submitted on 8 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Fujita phenomenon in exterior domains under dynamical boundary conditions II*[†]

Jean-François Rault

LMPA Joseph Liouville, FR 2956 CNRS,
Université du Littoral Côte d'Opale
50 rue F. Buisson, B.P. 699, F-62228 Calais Cedex (France)
`jfrault@lmpa.univ-littoral.fr`

Abstract

The Fujita phenomenon for nonlinear parabolic problems $\partial_t u = \Delta u + u^p$ in an exterior domain of \mathbb{R}^N under dissipative dynamical boundary conditions $\sigma \partial_t u + \partial_\nu u = 0$ is investigated in the superlinear case. As in the case of Dirichlet boundary conditions (see Refs. [2] and [9]), it turns out that there exists a critical exponent $p = 1 + \frac{2}{N}$ such that blow-up of positive solutions always occurs for subcritical exponents, whereas in the supercritical case global existence can occur for small non-negative initial data.

Key words: Nonlinear parabolic problems; Dynamical boundary conditions; Global solutions.

AMS Subject Classification: 35B33; 35B40; 35K55; 35K57.

*We improve the previous results from The Fujita phenomenon in exterior domain under dynamical boundary conditions, *Technical Report L.M.P.A. 383*, October 2008.

[†]To appear in *Asymptotic Analysis*.

1 Introduction

Let Ω be an exterior domain of \mathbb{R}^N , that is to say a connected open set Ω such that $\overline{\Omega}^c$ is a bounded domain when $N \geq 2$, and in dimension one, Ω is the complement of a real closed interval. We always suppose that the boundary $\partial\Omega$ is of class \mathcal{C}^2 . The outer normal unit vector field is denoted by $\nu : \partial\Omega \rightarrow \mathbb{R}^N$ and the outer normal derivative by ∂_ν . Let p be a real number with $p > 1$ and φ be a continuous function in $\overline{\Omega}$. Consider the following nonlinear parabolic problem

$$\begin{cases} \partial_t u = \Delta u + u^p & \text{in } \overline{\Omega} \times (0, \infty), \\ \mathcal{B}_\sigma(u) := \sigma \partial_t u + \partial_\nu u = 0 & \text{on } \partial\Omega \times (0, \infty), \\ u(\cdot, 0) = \varphi & \text{in } \overline{\Omega}. \end{cases} \quad (1)$$

The aim of this paper is to show that the well-known Fujita phenomenon in the case of $\Omega = \mathbb{R}^N$ (see Ref. [6]) and in the case of Dirichlet boundary conditions (see Refs. [2] and [9]) still holds for the dynamical boundary conditions. One can notice that dynamical boundary conditions $\mathcal{B}_\sigma(u) = 0$ with $\sigma \equiv 0$ correspond to the Neumann boundary conditions, which case has been discussed by Levine and Zhang [8]. It is already known, by Bandle, von Below and Reichel in [1], that for $p \in (1, 1 + \frac{2}{N})$, also for $p = 1 + \frac{2}{N}$ if $N \geq 3$, and for constant coefficient $\sigma \in [0, \infty)$, all positive solutions of (1) blow up in finite time. In addition, if the complement is star-shaped there exist global positive solutions of class \mathcal{C}^1 for $p > 1 + \frac{2}{N}$ by [1]. Our purpose is to show the existence of global positive solutions of Problem (1) for sufficiently small initial data in the supercritical case ($p > 1 + \frac{2}{N}$) for any exterior domain. Moreover our condition on σ is more general. Throughout, we shall assume the dissipativity condition

$$\sigma \geq 0 \text{ on } \partial\Omega \times (0, \infty) \quad (2)$$

and dealing with classical solutions

$$\sigma \in \mathcal{C}^1(\partial\Omega \times (0, \infty)). \quad (3)$$

The initial data is always supposed to be continuous, non-trivial, bounded, non-negative in $\overline{\Omega}$, and vanishing at infinity:

$$\varphi \in \mathcal{C}(\overline{\Omega}), \quad 0 < \|\varphi\|_\infty < \infty, \quad \varphi \geq 0, \quad \lim_{\|x\|_2 \rightarrow \infty} \varphi(x) = 0. \quad (4)$$

In the case $\Omega = \mathbb{R}^N$, the boundary condition is dropped and the result is well known by the classical paper of Fujita [6]. Thus, we will suppose $\Omega \neq \mathbb{R}^N$.

2 Preliminaries

First, we give the definition of positive solution which is understood along this paper.

Definition 2.1 *A positive solution of Problem (1) is a positive function $u : (x, t) \mapsto u(x, t)$ of class $\mathcal{C}(\overline{\Omega} \times [0, T]) \cap \mathcal{C}^{2,1}(\overline{\Omega} \times (0, T))$, satisfying*

$$\begin{cases} \partial_t u = \Delta u + u^p & \text{in } \overline{\Omega} \times (0, T), \\ \mathcal{B}_\sigma(u) := \sigma \partial_t u + \partial_\nu u = 0 & \text{on } \partial\Omega \times (0, T), \\ u(\cdot, 0) = \varphi & \text{in } \overline{\Omega}, \end{cases}$$

where φ is a function, given in $\mathcal{C}(\overline{\Omega})$. The time $T \in [0, \infty]$ is the maximal existence time of the solution u . If $T = \infty$, the solution u is called global.

From [2], if $T < \infty$, u blows up in finite time, that is to say:

$$\limsup_{\substack{t \nearrow T \\ x \in \overline{\Omega}}} u(x, t) = \infty.$$

Note that for initial data φ of class $\mathcal{C}^2(\overline{\Omega})$, the solution u is $\mathcal{C}^{2,1}(\overline{\Omega} \times [0, T])$, whereas $u \in \mathcal{C}(\overline{\Omega} \times [0, T]) \cap \mathcal{C}^{2,1}(\overline{\Omega} \times (0, T))$ if φ is only continuous in $\overline{\Omega}$. Then, let us recall a standard procedure to construct solutions of Problem (1) in outer domains for uniformly bounded and continuous initial data φ . Let $B(0, R)$ be the ball centered at the origin of radius $R > 0$ such that $\overline{\Omega}^c \subset B(0, R)$. For any $n \in \mathbb{N}$, we set $B_n := B(0, R + n)$ and $\Omega_n := \Omega \cap B_n$. The boundary of Ω_n is

decomposed into two disjoint open sets:

$$\partial\Omega_n = \partial\Omega \dot{\cup} \partial B_n.$$

Define also an increasing sequence of initial data $(\varphi_n)_{n \in \mathbb{N}^*}$ such that

$$\begin{cases} 0 \leq \varphi_n \leq \varphi & \text{in } \overline{\Omega}_n, \\ \varphi_n \equiv 0 & \text{on } \partial B_n, \\ \varphi_n = \varphi & \text{in } \overline{\Omega}_{n-1}, \end{cases} \quad (5)$$

and consider the following problem with mixed boundary conditions

$$\begin{cases} \partial_t u = \Delta u + u^p & \text{in } \overline{\Omega}_n \times (0, \infty), \\ \mathcal{B}_\sigma(u) := \sigma \partial_t u + \partial_\nu u = 0 & \text{on } \partial\Omega \times (0, \infty), \\ u = 0 & \text{on } \partial B_n \times (0, \infty), \\ u(\cdot, 0) = \varphi_n & \text{in } \overline{\Omega}_n. \end{cases} \quad (P(n))$$

Let z be the maximal solution of

$$\begin{cases} \dot{z} = z^p, \\ z(0) = \|\varphi\|_\infty, \end{cases}$$

with maximal existence time $t_0 = \frac{1}{(p-1)\|\varphi\|_\infty^{p-1}}$. It is known from [4] that, for each $n \in \mathbb{N}^*$, Problem $(P(n))$ has a solution $u_n \in \mathcal{C}(\overline{\Omega}_n \times [0, T_n)) \cap \mathcal{C}^{2,1}(\overline{\Omega}_n \times (0, T_n))$, where T_n is the maximal existence time of u_n . Moreover by comparison principle from [3], we have, for any $n \in \mathbb{N}^*$, $0 \leq u_n \leq u_{n+1}$ and $u_n(\cdot, t) \leq z(t)$ in $\overline{\Omega}_n$, so we have also $t_0 \leq T_n$. Hence we obtain a sequence $(u_n)_{n \in \mathbb{N}^*}$ of functions in $\mathcal{C}(\overline{\Omega}_n \times [0, t_0)) \cap \mathcal{C}^{2,1}(\overline{\Omega}_n \times (0, t_0))$. Then, standard arguments based on a priori estimates for the heat equation imply $u_n \rightarrow u$ in the sense of $\mathcal{C}_{loc}^{2,1}(\overline{\Omega} \times (0, t_0))$ as $n \rightarrow \infty$, where u is a positive solution of Problem (1), see Refs. [1] and [7]. Moreover, since u_n vanishes on ∂B_n for each $n \in \mathbb{N}^*$, the solution u vanishes at

infinity:

$$\lim_{\|x\|_2 \rightarrow \infty} u(x, t) = 0, \forall t \in (0, T).$$

Note that t_0 is only a lower bound for the maximal existence time of solutions u_n and u , and it is possible that the times T_n and T are infinite. Indeed, results on blow-up for problems under dynamical boundary conditions from [5] can not be applied to the problems $(P(n))$ with mixed boundary conditions because their solutions u_n vanish on a part of the boundary.

3 Global existence in dimension $N \geq 3$

Throughout this section, we consider supercritical exponent p :

$$p > 1 + \frac{2}{N}.$$

Our technique will be to construct a function that bounds from above each solution u_n of Problem $(P(n))$. This will give us a sequence $(u_n)_{n \in \mathbb{N}^*}$ of global solutions in $\mathcal{C}(\overline{\Omega}_n \times [0, \infty)) \cap \mathcal{C}^{2,1}(\overline{\Omega}_n \times (0, \infty))$, thus the solution u of (1) must be global too. We will proceed by using the solution of the Neumann Problem

$$\begin{cases} \partial_t v = \Delta v + v^p & \text{in } \overline{\Omega} \times (0, \infty), \\ \partial_\nu v = 0 & \text{on } \partial\Omega \times (0, \infty), \\ v(\cdot, 0) = \psi & \text{in } \overline{\Omega}, \end{cases} \quad (6)$$

with ψ verifying (4). In [8], Levine and Zhang proved that Problem (6) admits global positive solutions for sufficiently small initial data. We show that the solution v of Problem (6) bounds from above the solution u of Problem (1) if the initial data are well ordered ($\varphi \leq \psi$) and if ψ satisfies the following hypotheses:

$$\psi \in \mathcal{C}^2(\overline{\Omega}), \quad (7)$$

and for every $n \in \mathbb{N}^*$

$$\Delta\psi_n + \psi_n^p \geq 0 \text{ in } \overline{\Omega}_n, \quad (8)$$

where $(\psi_n)_{n \in \mathbb{N}^*}$ is the sequence of truncated initial data introduced in (5). We need a technical lemma, similar to Lemma 2.1 of [4].

Lemma 3.1 *Let ψ be a function satisfying (4), (7) and (8). For every $n \in \mathbb{N}^*$, the solution v_n of Problem $(P(n))$ under the Neumann boundary conditions and with the truncated initial data ψ_n verifies:*

$$\partial_t v_n \geq 0 \text{ in } \overline{\Omega}_n \times (0, T_{v_n}),$$

where T_{v_n} is the maximal existence time of v_n .

Proof : The function v_n is solution of the following problem

$$\begin{cases} \partial_t v_n = \Delta v_n + v_n^p & \text{in } \overline{\Omega}_n \times (0, T_{v_n}), \\ \partial_\nu v_n = 0 & \text{on } \partial\Omega \times (0, T_{v_n}), \\ v_n = 0 & \text{on } \partial B_n \times (0, T_{v_n}), \\ v_n(\cdot, 0) = \psi_n & \text{in } \overline{\Omega}_n, \end{cases}$$

in the bounded domain $\overline{\Omega}_n$. From (4) and from the strong maximum principle in [3], we claim

$$v_n > 0 \text{ on } (\Omega_n \cup \partial\Omega) \times (0, T_{v_n}).$$

Then, by regularity results from [7], we obtain $v_n \in C^{2,2}(\overline{\Omega}_n \times (0, T_{v_n}))$, and for $y = \partial_t v_n \in C^{2,1}(\overline{\Omega}_n \times (0, T_{v_n}))$ we have

$$\begin{cases} \partial_t y = \Delta y + p v_n^{p-1} y & \text{in } \overline{\Omega}_n \times (0, T_{v_n}), \\ \partial_\nu y = 0 & \text{on } \partial\Omega \times (0, T_{v_n}), \\ y = 0 & \text{on } \partial B_n \times (0, T_{v_n}), \end{cases}$$

and $y(\cdot, 0) \geq 0$ in $\overline{\Omega}_n$ thanks to (8). By the comparison principle in [3], we conclude: $y \geq 0$ in $\overline{\Omega}_n \times (0, T_{v_n})$.

■

Lemma 3.2 *Let a coefficient σ verifying (2) and (3), two functions φ and ψ satisfying (4) and ψ with (7) and (8). If*

$$\varphi \leq \psi \text{ in } \overline{\Omega}, \quad (9)$$

then Problem (1) with initial data φ admits a solution u verifying

$$u \leq v \text{ in } \overline{\Omega} \times (0, T_v),$$

and

$$0 < T_v \leq T \leq \infty,$$

where v is solution of Problem (6) with initial data ψ , of maximal existence time T_v .

Proof : We consider the sequences of truncated solutions $(u_n)_{n \in \mathbb{N}^*}$ and $(v_n)_{n \in \mathbb{N}^*}$ respectively associated to the solutions u and v . Let $n \in \mathbb{N}^*$. First, we show that $v_n \leq v$ in $\overline{\Omega}_n \times [0, T_v)$. By construction (5), we have $\psi_n \leq \psi$ in $\overline{\Omega}_n$. Since v is a positive solution of Problem (6), it satisfies

$$\left\{ \begin{array}{ll} \partial_t v \geq \Delta v + v^p & \text{in } \overline{\Omega}_n \times (0, T_v), \\ \partial_\nu v \geq 0 & \text{on } \partial\Omega \times (0, T_v), \\ v \geq 0 & \text{on } \partial B_n \times (0, T_v), \\ v(\cdot, 0) \geq \psi_n & \text{in } \overline{\Omega}_n. \end{array} \right.$$

As v_n is a positive solution of $(P(n))$ under Neumann boundary conditions, we obtain from the comparison principle in [3]

$$v_n \leq v \text{ in } \overline{\Omega}_n \times [0, \tau), \quad (10)$$

for all $0 < \tau < \min\{ T_{v_n}, T_v \}$. We deduce $T_v \leq T_{v_n}$. Then, we show that $u_n \leq v_n$ in $\overline{\Omega}_n \times [0, T_{v_n})$. The previous lemma ensures that $\partial_t v_n \geq 0$. From (2), we obtain

$$\sigma \partial_t v_n + \partial_\nu v_n \geq 0 \text{ on } \partial\Omega \times (0, T_{v_n}).$$

Next, v_n is a positive solution of $(P(n))$, and thanks to (9), v_n verifies

$$\begin{cases} \partial_t v_n \geq \Delta v_n + v_n^p & \text{in } \overline{\Omega}_n \times (0, T_{v_n}), \\ \sigma \partial_t v_n + \partial_\nu v_n \geq 0 & \text{on } \partial\Omega \times (0, T_{v_n}), \\ v_n \geq 0 & \text{on } \partial B_n \times (0, T_{v_n}), \\ v_n(\cdot, 0) \geq \varphi_n & \text{in } \overline{\Omega}_n. \end{cases}$$

Again, by the comparison principle in [3] and by definition of u_n , we obtain

$$u_n \leq v_n \text{ in } \overline{\Omega}_n \times [0, \tau), \quad (11)$$

for all $0 < \tau < \min\{ T_n, T_{v_n} \}$, and hence $T_{v_n} \leq T_n$. From equations (10) and (11), we have $T_v \leq T_n$ and $u_n \leq v$ in $\overline{\Omega}_n \times [0, T_v)$. Thus the solution u of Problem (1), obtained as the limit of the sequence $(u_n)_{n \in \mathbb{N}^*}$ with the procedure described in section 2, verifies $u \leq v$ in $\overline{\Omega} \times [0, T_v)$ and $T_v \leq T$. ■

Now, we just have to choose an initial data ψ_* sufficiently small such that Problem (6) admits a global positive solution (see Ref. [8]), and satisfying (7) and (8).

Theorem 3.3 *Under conditions (2), (3) and (4), Problem (1) admits global positive solutions for sufficiently small initial data. Moreover, some of these solutions vanish at infinity.*

Proof : An initial data φ verifying (4) and with $\varphi \leq \psi_*$ in $\overline{\Omega}$ allows us to conclude thanks to Lemma 3.2. ■

Remark 3.4 *One can notice that only the dissipativity and the regularity of the coefficient σ are needed. We are not obliged to impose any restriction like σ bounded or $\partial_t \sigma \equiv 0$. Moreover, the hypotheses (7) and (8) on the initial data ψ of Problem (6) are strictly technical and do not concern the initial data φ of Problem (1).*

4 Global existence in lower dimension

In this case, we can not use Levine and Zhang's result because it is proved only for dimension $N \geq 3$: they used some estimates for Green's functions, specific to dimension $N \geq 3$. We need an additional hypothesis on the coefficient σ . There exists a constant $\varsigma \in [0, \infty)$ such that

$$\forall (x, t) \in \partial\Omega \times [0, \infty) : \sigma(x, t) \leq \varsigma . \quad (12)$$

We begin with the case of dimension 2. Until now, Bandle - von Below - Reichel's lemma, concerning star-shaped domains, is the best result:

Lemma 4.1 [1], Lemma 28. *Suppose σ is a positive constant. If Ω^C is star-shaped with respect to the origin and if $\min_{\partial\Omega} |x \cdot \nu(x)| \geq \sigma N$, then there exist positive global solutions of Problem (1), which vanish at infinity, for sufficiently small initial data.*

This allows us to deduce the following result for problems with mixed boundary conditions.

Corollary 4.2 *Suppose conditions (2), (3) and (12). Let $y \in \partial\Omega$. There exists a neighborhood N_y of y relatively open in $\partial\Omega$ such that the following parabolic*

problem with mixed boundary conditions

$$\begin{cases} \partial_t u = \Delta u + u^p & \text{in } \overline{\Omega} \times (0, \infty), \\ \mathcal{B}_\sigma(u) = 0 & \text{on } N_y \times (0, \infty), \\ u = 0 & \text{on } \partial\Omega \setminus N_y \times (0, \infty), \\ u(\cdot, 0) = \varphi & \text{in } \overline{\Omega}. \end{cases}$$

admits global positive solutions which vanish at infinity, for sufficiently small initial data φ satisfying (4).

Proof : Let μ be a vector in \mathbb{R}^N such that the scalar product between the vector $(y + \mu)$ and the outer normal unit vector at y satisfies

$$(y + \mu) \cdot \nu(y) < -\varsigma N. \quad (13)$$

Then, as the mapping $(\partial\Omega \ni z \mapsto (z + \mu) \cdot \nu(z) \in \mathbb{R})$ is continuous, the above inequality remains true on an open neighborhood $N_y \subseteq \partial\Omega$ of y . We obtain the statement of the corollary by using the comparison principle from [3] and the function U defined on $\overline{\Omega} \times [0, \infty)$ by

$$U(x, t) = A(t + 1)^{-\gamma} \exp \frac{-\|x + \mu\|_2^2}{4(t + 1)},$$

with $A = \frac{1}{2} \left(\frac{N}{2} - \frac{1}{p-1} \right)^{\frac{1}{p-1}}$ and $\gamma = \frac{1}{p-1}$. It is clear that $U \geq 0$, belongs to $\mathcal{C}^{2,1}(\overline{\Omega} \times [0, \infty))$ and satisfies:

$$\begin{aligned} \partial_t U &= \left(\frac{-\gamma}{t+1} + \frac{\|x + \mu\|_2^2}{4(t+1)^2} \right) U, \\ \Delta U &= \left(\frac{-N}{2(t+1)} + \frac{\|x + \mu\|_2^2}{4(t+1)^2} \right) U, \\ \partial_\nu U &= \left(\frac{-(x + \mu) \cdot \nu(x)}{2(t+1)} \right) U. \end{aligned}$$

We have:

$$\partial_t U - \Delta U = \left(\frac{-\gamma}{t+1} + \frac{N}{2(t+1)} \right) U = \left(\frac{-2\gamma + N}{2(t+1)} \right) U ,$$

and by definition of the constants A and γ , we obtain

$$\partial_t U - \Delta U - U^p \geq 0 \text{ in } \overline{\Omega} \times [0, \infty) .$$

Then, on $\partial\Omega$, we have

$$\sigma \partial_t U + \partial_\nu U = \left(\frac{-2\sigma\gamma - (x + \mu) \cdot \nu(x)}{2(t+1)} + \frac{\sigma \|x + \mu\|_2^2}{4(t+1)^2} \right) U .$$

Since $p > 1 + \frac{2}{N}$, using (12) and ignoring the non-negative term $\frac{\sigma \|x + \mu\|_2^2}{4(t+1)^2}$, we can reduce the last equation to:

$$\sigma \partial_t U + \partial_\nu U \geq \left(\frac{-\varsigma N - (x + \mu) \cdot \nu(x)}{2(t+1)} \right) U .$$

Thanks to (13) we obtain $\mathcal{B}_\sigma(U) \geq 0$ in $N_y \times [0, \infty)$. And we have $U \geq 0$ on $\partial\Omega \setminus N_y \times (0, \infty)$. An initial data φ with $\varphi \leq U(\cdot, 0)$ in $\overline{\Omega}$ permits to conclude. ■

In the case of dimension one, we use the fact that Ω is not connected. Let us write $\Omega = \mathbb{R} \setminus [a, b]$ with $a < b$ in \mathbb{R} , and let V be the function defined in $\overline{\Omega} \times [0, \infty)$ by:

$$V(x, t) = \begin{cases} A(t+1)^{-\gamma} \exp \frac{-\|x + \mu_1\|_2^2}{4(t+1)} & \text{if } x \leq a \\ A(t+1)^{-\gamma} \exp \frac{-\|x + \mu_2\|_2^2}{4(t+1)} & \text{if } x \geq b , \end{cases}$$

with A and γ like in Corollary 4.2, μ_1 and μ_2 in \mathbb{R} such that

$$-(a + \mu_1) - \varsigma \geq 0 ,$$

and

$$(b + \mu_2) - \varsigma \geq 0 .$$

As $\nu(a) = 1$ and $\nu(b) = -1$, we obtain with (12)

$$\sigma \partial_t V + \partial_\nu V \geq 0 \text{ on } (\{a\} \cup \{b\}) \times [0, \infty) .$$

Following the proof of Corollary 4.2, we obtain this result:

Theorem 4.3 *Under conditions (2), (3), (4), (12), $N = 1$ and $p > 3$, Problem (1) admits global positive solutions vanishing at infinity, for sufficiently small initial data φ .*

Acknowledgment

The author would like to thank Professor Joachim von Below for helpful discussions and advices.

References

- [1] C. Bandle, J. von Below, and W. Reichel, Parabolic problems with dynamical boundary conditions: eigenvalue expansions and blow up, *Rend. Lincei Math. Appl.* **17** (2006), 35–67.
- [2] C. Bandle and H.A. Levine, On the existence and the nonexistence of global solutions of reaction-diffusion equations in sectorial domains, *Trans. Am. Math. Soc.* **316** (1989), 595–622.
- [3] J. von Below and C. De Coster, A Qualitative Theory for Parabolic Problems under Dynamical Boundary Conditions, *Journal of Inequalities and Applications* **5** (2000), 467–486.

- [4] J. von Below and G. Pincet-Mailly, Blow Up for Reaction Diffusion Equations Under Dynamical Boundary Conditions, *Communications in Partial Differential Equations* **28** (2003), 223–247.
- [5] J. von Below and G. Pincet-Mailly, Blow Up for some non linear parabolic problems with convection under dynamical boundary conditions, *Discrete and Continuous Dynamical Systems Supplement Volume 2007* (2007), 1031–1041.
- [6] H. Fujita, On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$, *Journal of the Faculty of Science of the University of Tokyo* **13** (1966), 109–124.
- [7] O.A. Ladyženskaya, V.A. Solonnikov, and N.N. Uraltseva, Linear and quasi-linear equations of parabolic type, *Trans. of Math. Monographs* **23** (1968).
- [8] H.A. Levine and Q.S. Zhang, The critical Fujita number for a semilinear heat equation in exterior domains with homogeneous Neumann boundary values, *Proceedings of the Royal Society of Edinburgh* **130 A** (2000), 591–602.
- [9] K. Mochizuki and R. Suzuki, Critical exponent and critical blow up for quasilinear parabolic equations, *Israel J. Math* **98** (1997), 141–156.