



The feeding ecology of little auks raises questions about winter zooplankton stocks in North Atlantic surface waters

Jérôme Fort, Yves Cherel, Ann M.A. Harding, Carsten Egevang, Harald Steen, Grégoire Kuntz, Warren P Porter, David Grémillet

► To cite this version:

Jérôme Fort, Yves Cherel, Ann M.A. Harding, Carsten Egevang, Harald Steen, et al.. The feeding ecology of little auks raises questions about winter zooplankton stocks in North Atlantic surface waters. *Biology Letters*, 2010, 6, pp.682-684. 10.1098/rsbl.2010.0082 . hal-00527435

HAL Id: hal-00527435

<https://hal.science/hal-00527435>

Submitted on 27 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The feeding ecology of little auks raises questions about winter zooplankton stocks in North Atlantic surface waters

Jérôme Fort^{1,*}, Yves Cherel², Ann M. A. Harding³, Carsten Egevang⁴, Harald Steen⁵, Grégoire Kuntz¹, Warren P. Porter⁶ & David Grémillet^{1,7}

¹ *Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175 du CNRS, 1919 Route de Mende, 34293 Montpellier cedex 5, France*

² *Centre d'Etudes Biologiques de Chizé, UPR 1934 du CNRS, 79360 Villiers-en-Bois, France*

³ *Environmental Science Department, Alaska Pacific University, 4101 University Drive, Anchorage, AK 99508, USA*

⁴ *Greenland Institute of Natural Resources, Postbox 570, Kivioq 2, DK-3900 Nuuk, Greenland*

⁵ *Norwegian Polar Institute, Polar Environmental Centre, N-9296 Tromsø, Norway*

⁶ *Department of Zoology, University of Wisconsin, 250 N. Mills Street, Madison, Wisconsin 53706, USA*

⁷ *Percy FitzPatrick Institute, DST/NRF Centre of Excellence, University of Cape Town, Rondebosch 7701, South Africa*

* Author for correspondence (fort.jerome@gmail.com)

Running head: seabirds and zooplankton stocks

Copepods are essential components of marine food webs worldwide. In the North Atlantic, they are thought to perform vertical migration and to remain at depths >500 m during winter. We challenge this concept through a study of the winter feeding ecology of little auks (*Alle alle*), a highly abundant planktivorous seabird from the North Atlantic. By combining stable isotope and behavioural analyses, we strongly suggest that swarms of copepods are still available to their predators in water surface layers (<50 m) during winter, even during short daylight periods. Using a new bioenergetic model, we estimate that the huge number (20-40 million birds) of little auks wintering off southwest Greenland consume 3600-7200 tonnes of copepods daily, strongly suggesting substantial zooplankton stocks in surface waters of the North Atlantic in the middle of the boreal winter.

Keywords: diving behaviour, energetic modelling, food requirements, marine food web, stable isotope, seabird

1. INTRODUCTION

Copepods are essential components of aquatic food webs and play an important role in marine ecosystems. Their life strategies have been extensively studied (e.g. Falk-Petersen 2009) and their ecology is considered to be well known. At high latitudes and during winter, copepods are thought to perform vertical migration to deeper water to undergo diapause, and to become unavailable to most predators (Falk-Petersen 2009). However, a recent study has shown that in high Arctic waters copepods maintain their synchronized diel vertical migration throughout the polar night (Berge et al. 2009), suggesting that further investigations are necessary to yield a complete view of copepod winter ecology.

Seabirds are powerful indicators of marine food webs (e.g. Piatt et al. 2007). Here we used little auks (*Alle alle*) as biosamplers of copepods, since this small Arctic species is the only seabird in the North Atlantic which feeds almost exclusively on zooplankton, essentially calanoid copepods (Stempniewicz 2001). We used isotopic tracers (Kelly 2000) and Time-Depth-Recorders to investigate little auk diet and foraging depths during winter. It is the first study to record the winter diving behaviour of such a small seabird. We tested the hypothesis that copepods are still available in upper surface layers and consumed by little auks, even during the short winter daylight period.

2. MATERIAL AND METHODS

Little auks breed at high latitudes around the Arctic Ocean. The North-East Atlantic populations winter south of the ice edge, mainly off southwest Greenland and Newfoundland (Stempniewicz 2001). Twenty breeding little auks were caught during summer 2007 at Kap Höegh (East Greenland; 70°44'N 21°35'W). Blood samples (~0.2ml) were collected from the brachial vein, as well as 16 chick meals transported in adult's sublingual pouch. Twenty wintering birds were legally shot at sea off Nuuk (southwest Greenland; 64°10'N 51°45'W) in January 2007 and blood samples collected from the cardiac clot. Stomach contents were removed, but digestion precluded identifying the few prey items. To investigate copepod isotopic signatures, only chick meals containing >90% copepods were analysed. Stable isotope analyses were performed following Cherel et al. (2007), with blood $\delta^{15}\text{N}$ ($^{15}\text{N}/^{14}\text{N}$) values reflecting the trophic position / diet (Kelly 2000). Values are mean \pm S.D.

Eighteen breeding little auks from Longyearbyen (Spitsbergen; N78°13'N 15°20'E) and 22 from Kap Höegh were equipped with a Time-Depth-Recorder archival tag (G5, CEFAS Technology Limited, Lowestoft, UK) during summers 2007 and 2008, respectively. Recorders were implanted into the abdominal cavity following Grémillet et al. (2005). Each batch of loggers was programmed to record pressure during 24 hour periods at a sampling rate of 2 seconds; every 12th day from 6th November 2007 to 22nd February 2008 and every 10th day from 1st October 2008 to 10th May 2009. Five birds from Longyearbyen were resighted in July 2008 and three of those were recaptured. Only one logger successfully recorded pressure. Thirteen birds from Kap Höegh were resighted and recaptured in July 2009. Only two loggers successfully recorded pressure in December-January. Stored data were analysed using MultiTrace-Dive software (Jensen Software Systems). To avoid artefacts due to waves or tag accuracy, only dives ≥ 1.5 m were analyzed. To compare winter results of behavioural and isotopic analyses, only dive data recorded in December and January were used.

Individual daily energy and food requirements during December-January were calculated using the bioenergetics model Niche Mapper™ (Fort et al. 2009). The principles of Niche Mapper™, its validation and all input data are detailed in the electronic supplementary material (ESM-1).

Given the isotopic results (see below), birds were assumed to feed on copepods during winter. We then estimated the daily food intake for the whole little auk population wintering off southwest Greenland. This population is composed to a large extent of birds breeding in northwest Greenland, east Greenland and Svalbard (Stempniewicz 2001). The number of little auks wintering off southwest Greenland is therefore estimated as 20-40 million birds (Barrett et al. 2006, Karnovsky & Hunt 2002).

3. RESULTS

During summer, the $\delta^{15}\text{N}$ value of chick meals (which correspond to copepod prey, Harding et al. 2009) averaged $8.4 \pm 0.5\text{‰}$ (figure 1). Summer and winter bird blood $\delta^{15}\text{N}$ values were almost identical (figure 1), yet statistically different (11.4 ± 0.1 and $11.7 \pm 0.4\text{‰}$, respectively; *t*-test, $t=3.75$, $p<0.001$). During summer, adult bird blood was 3‰ enriched in $\delta^{15}\text{N}$ compared to copepod prey, as expected for an increase of one trophic level (figure 1). During winter, measured bird blood was also 3‰ enriched in $\delta^{15}\text{N}$ compared to copepods (figure 1, Sato et al. 2002).

During winter (December-January), the mean depth of dives was 12 ± 11 m, with a maximum dive depth of 50 m ($n=6056$ dives). The mean dive duration was 45 ± 25 s. On average, birds spent 24% of their time diving, an activity which was equally performed during the daylight and night periods (48% and 52% of time diving, respectively) (figure 2). Dives were mostly concentrated during the daylight period and the first part of the night (figure 2; detailed results in ESM-2).

During January, Niche Mapper™ predicted an individual daily energy expenditure of 438 ± 41 $\text{kJ} \cdot \text{day}^{-1}$, and a food requirement necessary to meet this expenditure of 180 ± 17 g zooplankton $\cdot \text{day}^{-1}$ (wet mass). Consequently, we estimate that the southwest Greenland wintering population of 20-40 millions little auks consumes 3600-7200 tonnes of zooplankton per day (wet mass).

4. DISCUSSION

Using a multidisciplinary-approach combining stable isotope analysis, biotelemetry and energetic modelling, this study presents indirect evidence that during winter daylight periods copepods are still numerous in surface waters <50m off southwest Greenland. Little auks are known to feed almost exclusively on copepods in summer (e.g. Karnovsky et al. 2008), and blood $\delta^{15}\text{N}$ values found here are in agreement with birds preying upon copepods during this period (Karnovsky et al. 2008). Isotopic $\delta^{15}\text{N}$ signatures are also similar for summer and winter copepods (Sato et al. 2002, figure 1), as well as for summer and winter adult blood samples (figure 1). This strongly suggests that little auks feed predominantly on copepods in winter. During this period, birds may also consume other prey, but in a minor proportion since amphipods, euphausiids or fish larvae have higher isotopic signatures than copepods (Tamelander et al. 2008) and would thus lead to higher blood $\delta^{15}\text{N}$ values. Winter blood values slightly higher than those measured in summer might reflect that amphipod/euphausiid effect, as well as the general food web enrichment from summer to winter (Rolff 2000).

Further, we infer from little auk diving behaviour that copepods remain abundant in winter surface waters, even during the daylight period when they are supposed to perform vertical migration to greater depths (figure 2, Fortier et al. 2001, Berge et al. 2009). Even though our dive depth results are consistent with summer recordings (Harding et al. 2009), further studies are now needed to confirm the first winter dive data for little auks. Their performance is impressive, with hundreds of

dives conducted daily, over 1 minute and to depths of up to 50 m (see ESM-2). However, this maximum dive depth is extremely shallow compared to the depths of >500 m at which copepods are assumed to occur in winter.

Copepods are classically thought to enter diapause in the fall and to spend all winter in deeper water (Falk-Petersen 2009). During early spring, and mostly synchronized with the phytoplankton spring bloom, they are supposed to migrate back to surface waters (Falk-Petersen 2009). However, a recent acoustic study has shown that even in winter and during the high-latitude polar night, some copepods perform a synchronized diel vertical migration in the epipelagic layer (Berge et al. 2009), whereby they migrate to the food-rich surface layers during darkness and move to deeper water during daylight where they are out of reach from predators (Fortier et al. 2001). Our study strongly suggests that during winter, swarms of copepods are still present in surface waters (*sensu* Berge et al. 2009), where they are targeted by little auks both during the daylight and the dark period.

Using a bioenergetic model, we estimated that the amount of copepods available to little auks wintering off southwest Greenland is substantial, since the daily intake of the little auk population wintering off southwest Greenland is estimated to amount to 3600-7200 tonnes. This result is based on a diet composed only of copepods and it might be slightly overestimated since little auks probably also consume a few other prey items such as amphipods and euphausiids. The standing stock of copepods is currently unknown and it is impossible to estimate the proportion of the population caught per day by wintering birds. However, a range of 3600-7200 tonnes of copepods caught per day during the winter off southwest Greenland is larger than the daily amount of copepods consumed during summer by the breeding population of the North Water Polynya (i.e. 3450-6900 tonnes of wet food consumed per day, including 2340-4680 tonnes of copepods; Karnovsky & Hunt 2002). Therefore, this suggests that the daily presence of copepods in the upper layer of the water column throughout the winter is much more important than usually assumed, with far-ranging consequences for regional food webs and energy flow.

Acknowledgments

We thank all assistants for their hard work in the field. This project was supported by the Institut Polaire Français Paul Emile Victor (Grant 388 to D.G.) and the National Science Foundation (Grant 0612504 to A.H. & N.Karnovsky). Fieldwork led in Greenland was conducted under permits of the IPEV Ethics Committee (MP/12/24/05) and with the permission of the Greenland Home Rule Government (Danish Polar Center permit 512-240). Fieldwork led in Spitsbergen was conducted under permits 234 Forsøksdyrutvalget, Mattilsynet, Norway. J.F. is supported by a grant from the University Louis Pasteur (Strasbourg, France).

References

- Barrett, R. B., Chapdelaine, G., Anker-Nilssen, T., Mosbech, A., Montevecchi, W. A., Reid, J. B. & Veit, R. R. 2006 Seabird numbers and prey consumption in the North Atlantic. *ICES J. Mar. Sci.* **63**, 1145-1158 (doi:10.1016/j.icesjms.2006.04.004)
- Berge, J., Cottier, F., Last, K. S., Varpe, Ø., Leu, E., Søreide, J., Eiane, K., Falk-Petersen, S., Willis, K., Nygård, H. et al. 2009 Diel vertical migration of Arctic zooplankton during the polar night. *Biol. Lett.* **5**, 69-72. (doi:10.1098/rsbl.2008.0484)
- Cherel, Y., Hobson, K. A. Guinet, C. & Vanpe, C. 2007 Stable isotopes document seasonal changes in trophic niches and winter foraging individual specialization in diving predators from the Southern Ocean. *J. Anim. Ecol.* **76**, 826-836. (doi:10.1111/j.1365-2656.2007.01238.x)
- Falk-Petersen, S., Mayzaud, P., Kattner, G. & Sargent, J. R. 2009 Lipids and life strategy of Arctic *Calanus*. *Mar. Biol. Res.* **5**, 18-39. (doi:10.1080/17451000802512267)
- Fort, J., Porter, W. P. & Grémillet, D. 2009 Thermodynamic modelling predicts energetic bottleneck for seabirds wintering in the northwest Atlantic. *J. Exp. Biol.* **212**, 2483-2490. (doi:10.1242/jeb.032300)
- Fortier, M., Fortier, L., Hattori, H., Saito, H. & Legendre, L. 2001 Visual predators and the diel vertical migration of copepods under Arctic sea ice during the midnight sun. *J. Plankt. Res.* **23**, 1263-1278. (doi:10.1093/plankt/23.11.1263)

- Grémillet, D., Kuntz, G., Gilbert, K., Woakes, A. J., Butler, P. J. & le Maho, Y. 2005 Cormorants dive through the Polar night. *Biol. Lett.* **1**, 469-471. (doi:10.1098/rsbl.2005.0356)
- Harding, A. M. A., Egevang, C., Walkusz, W., Merkel, F., Blanc, S. & Grémillet, D. 2009 Estimating prey capture rates of a planktivorous seabird, the little auk (*Alle alle*), using diet, diving behaviour, and energy consumption. *Polar Biol.* **32**, 785-796. (doi:10.1007/s00300-009-0581-x)
- Karnovsky, N. J. & Hunt, G. L. 2002 Estimation of carbon flux to dovekies (*Alle alle*) in the North Water. *Deep-Sea Res. II* **49**, 5117-5130.
- Karnovsky, N. J., Hobson, K. A., Iverson, S. & Hunt Jr., G. L. 2008 Seasonal changes in diets of seabirds in the North Water Polynya: a multiple-indicator approach. *Mar. Ecol. Prog. Ser.* **357**, 291-299. (doi:10.3354/meps07295)
- Kelly, J. F. 2000 Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Can. J. Zool.* **78**, 1-27.
- Piatt, J. F., Harding, A. M. A., Shultz, M., Speckman, S. G., van Pelt, T. I., Drew, G. S. & Kettle, A. B. 2007 Seabirds as indicators of marine food supplies: Cairns revisited. *Mar. Ecol. Prog. Ser.* **352**, 221-234. (doi:10.3354/meps07295)
- Rolff, C. 2000 Seasonal variation in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of size-fractionated plankton at a coastal station in the northern Baltic proper. *Mar. Ecol. Prog. Ser.* **203**, 47-65.
- Sato, M., Sasaki, H. & Fukuchi, M. 2002 Stable isotopic compositions of overwintering copepods in the arctic and subarctic waters and implications to the feeding history. *J. Mar. Syst.* **38**, 165-174.
- Stempniewicz, L. 2001 Little Auk *Alle alle*. BWP Update. **3**, 175-201.
- Tameler, T., Reigstad, M., Hop, H., Carroll, M. L. & Wassmann, P. 2008 Pelagic and sympagic contribution of organic matter to zooplankton and vertical export in the Barents Sea marginal ice zone. *Deep-Sea Res. II* **55**, 2330-2339. (doi:10.1016/j.dsr2.2008.05.019)




Figure 1. Stable nitrogen isotope values for chick meals (n=9) and for little auks blood samples (n=20) collected at Kap Höegh (east Greenland) during the summer, and for little auks blood samples (n=21) collected at sea off Nuuk (southwest Greenland) during winter. Values are means \pm SD. Values not sharing the same superscript letter are significantly different (see results). Winter value for *Calanus* spp. is from Sato et al. (2002).




Figure 2. Average number of dives (▨) and maximum depth reach for any dive (●) recorded at each hour of day during winter (December-January) (\pm SE). The grey background represents the night/darkness period while the white background represents the daylight period off Nuuk, southwest Greenland ($64^{\circ}10'N$ $51^{\circ}45'W$). Sunrise and sunset hours were converted to GMT-hours. They correspond to the earliest sunrise and latest sunset observed during the study period.