



HAL
open science

Two new species of Antarctic gorgonians (Octocorallia: Primnoidae) with a redescription of Versluys, 1906

Rebeca Zapata-Guardiola, Pablo J. López-González

► To cite this version:

Rebeca Zapata-Guardiola, Pablo J. López-González. Two new species of Antarctic gorgonians (Octocorallia: Primnoidae) with a redescription of Versluys, 1906. *Helgoland Marine Research*, 2009, 64 (3), pp.169-180. 10.1007/s10152-009-0176-5 . hal-00529004

HAL Id: hal-00529004

<https://hal.science/hal-00529004>

Submitted on 24 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Two new species of Antarctic gorgonians (Octocorallia: Primnoidae), with a redescription of *Thouarella laxa* Versluys, 1906

Rebeca Zapata-Guardiola and Pablo J. López-González

Departamento de Fisiología y Zoología, Facultad de Biología, Universidad de Sevilla, Reina Mercedes 6, 41012-Sevilla, Spain. E-mail: rebzapgua@alum.us.es // pjlopez@us.es

Abstract

Two new species of the genus *Thouarella*, from Antarctic waters are described and illustrated from material collected on the *Polarstern* cruises ANT XVII/3 (EASIZ III), ANT XIX/5 (LAMPOS) and ANT XXI/2 (BENDEX). On the one hand, *T. viridis* sp. nov. is placed in the subgenus *Epithouarella* due to the characteristic ornamentation of its marginal scales (the previously most recent species in this group was included by Kükenthal in 1908). On the other hand, *T. minuta* sp. nov. is included in subgenus *Thouarella*, among the 14 species currently recognised, the main distinct feature being tiny polyps. Furthermore a complete redescription is given of *Thouarella laxa* Verluys, 1906. Using the new technology available nowadays, such as images obtained with SEM, we provide accurate images of the polyps and sclerites. In addition, as a result of this study, *T. laxa* and its closest congener *T. tydemani* are maintained as separate species, mainly due to their internal sculpture of body and coenenchymal scales.

Keywords: Cnidaria – Octocorallia – Primnoidae – Antarctica – *Thouarella* – New species.

Introduction

Among the gorgonian families, Primnoidae is one of the richest in genera and species (Bayer, 1980, 1981, 1988, 1998). This family is present in both polar waters, north and south, and in deep waters at temperate latitudes (Bayer, 1996b), dominating, together with the family Isididae, the high latitudes of the Southern Hemisphere (Bayer 1996a, b, 1998; Alderslade 1998; López-González and Gili 2002). According to Cairns and Bayer, 2009, 36 genera with 233 species are currently recognised for Primnoidae, with *Thouarella* Gray, 1870 being one of the most speciose primnoid genera (Kükenthal, 1924; Cairns, 2006; Cairns and Bayer, 2009) represented in Antarctic and SubAntarctic waters. According to the most recent contribution giving a generic revision of, and phylogenetic approach to, the Primnoidae (Cairns and Bayer, 2009), at least 29 species of *Thouarella* could be considered valid, several of them in need of revision. *Thouarella* species are currently organised into four subgenera, *Euthouarella* Kükenthal, 1915, *Epithouarella* Kükenthal, 1915, *Thouarella* Gray, 1870 and *Diplocalyptra* Kinoshita, 1908 (see Kükenthal, 1915; Cairns and Bayer, 2009). The main characters used in the identification of the different subgenera are those related to the ornamentation of marginal scales, the arrangement of polyps and the branching pattern of colonies.

During the Antarctic *Polarstern* cruises ANT XVII/3, ANT XIX/5 and ANT XXI/2, a significant number of Primnoidae specimens were collected, among this material a number of specimens belonged to the genus *Thouarella*. The study of a set of these colonies allows us to propose two new species belonging to two of the four recognized subgenera. With these new species we increase the knowledge of Antarctic diversity and also the variability in colours and polyp sizes previously known in the genus.

An important task in the revision of a large amount of material from many recent international sampling efforts is obviously the identification of the species collected. This work is directly related to the production of reports and information on biodiversity at different geographical scales. One of the main problems in this taxonomic work is the difficulty in recognizing species described in the early history of systematic taxonomy, especially when few characters were described, discussed or illustrated. For this reason, an important effort should be made to redescribe old species under the current descriptive formats, revising the true recognizable distribution and, when possible, relationships with the closest congeners. Recently we have had the opportunity to

consult a fragment of the type material of *Thouarella laxa* Versluys, 1906. This species was also reported by Kükenthal and Grozawsky, 1908, and all subsequent references of this species are based on those old reports. Thus, an additional objective of the present paper is the redescription of *Thouarella laxa*, based on all known material, with special reference to polyp morphology and sclerite variability. The affinities between *T. laxa* and its closest congener *T. tydemani* Versluys, 1906 are also discussed.

Material and Methods

The material of *Thouarella laxa* here examined came from different European museums who kindly send us part of their specimens for SEM examination of polyp morphology and sclerite variability. The holotype is deposited in the Zoological Museum of Amsterdam (ZMA), The Netherlands. The remaining material belonged to the Doflein collection, and is deposited in The Bavarian State Collection of Zoology, Munich (ZSM), Germany. For purposes of comparison, the following material deposited at the Zoological Museum of Amsterdam (ZMA) has also been examined: *Thouarella tydemani* Versluys, 1906. (ZMA, COEL09256) Holotype: Siboga Expedition, stn 297, 10°39'S, 123°40'E, Lesser Sunda Islands, Timor Sea, Indonesia, 520 m depth, 27 January 1900, three slides.

The material studied, concerning the new species, was collected on the *R/V Polarstern* cruises ANT XVII/3 (EASIZ III, Ecology of the Antarctic Sea Ice Zone, 18 March to 11 May 2000), ANT XIX/5 (LAMPOS, Latin American Polarstern Study, 3 Apr to 5 May 2002), ANT XXI/2 (BENDEX, BENthos Disturbance Experiment, 17 Nov 2003 to 18 Jan 2004) sponsored by the Alfred Wegener Institut für Polar- und Meeresforschung (Bremerhaven) under the auspices of the SCAR (Scientific Committee for Antarctic Research). Octocoral colonies were collected using an Agassiz trawl or a Bottom trawl along South Georgia Ridge (Scottia Arc) and Austasen (Eastern Weddell Sea) (Fig. 1). The octocorals were fixed in buffered formalin 10% and then transferred to ethanol 70%. Fragments from different parts of the colonies were prepared for study by SEM employing the usual methodology previously described by various authors (e.g., Bayer and Stefani 1988; Alderslade 1998), and permanent mounts were made for light microscopy observation. All sclerite size measurements and illustrations are from

the holotypes. The colony and sclerite terminology herein mainly follows Bayer *et al.* (1983). The material studied has been deposited at the Zoologisches Institut und Zoologisches Museum, Hamburg, Germany (ZIZMH); the National Museum of Natural History, Smithsonian Institution, Washington, D.C. (USNM), and in the anthozoan reference collection of the research group “Biodiversidad y Ecología de Invertebrados Marinos” of the University of Seville (BEIM).

Results

Family Primnoidae Gray, 1858

Genus *Thouarella* Gray, 1870

Diagnosis

Primnoidae commonly with a bottlebrush colony shape and simple or ramified branchlets, with also fan shaped colonies. Polyps cylindrical to club shaped, upward or almost perpendicular to stem, straight or adaxially incurved, and arranged single or in whorls. Opercular scales usually keeled, eight in number, arranged in two alternate cycles. Marginal scales also eight in number, alternating in two cycles, with a keel folding over operculars in an indented way. Adaxial body scales often reduced.

Subgenus *Eouthouarella* Kükenthal, 1915

Diagnosis

Thouarella with polyps placed in pairs or whorls. Marginal scales bearing thorns.

***Thouarella (Eouthouarella) laxa* Versluys, 1906**

(Figs. 2, 3, 4, 5)

? *Hookerella pulchella* Gray, 1870: 45.

Thouarella laxa Versluys, 1906: 30-32, pl. 1, fig. 5; pl. 3, fig. 8; text figs. 28-33. --

Kükenthal and Grozawsky, 1908: 36-37, pl. 2, fig. 13.

Thouarella (Euthouarella) laxa, Kükenthal, 1915: 150. -- Kükenthal, 1919: 417. -- Kükenthal, 1924: 293-294, text fig. 164. -- Cairns and Bayer, 2009: 28 (in list).

Examined material

Holotype: ZMA (COEL 03576), *Siboga* Expedition, stn 88, 0°34.6'N, 119°8.5'E, Strait of Makassar, Sulawesi, Indonesia, 1301 m depth, 20 June 1899. Additional material: ZSM (20080407), ZSM (20080408) and ZSM (20080409), Steam Vessel "*Zuso Maru*", Sagami Bay, Okinose, 300-700 m depth, 8-15 November 1904, Doflein Collection 1904-1905.

Description of the holotype

Fragment examined (Fig. 2a) 22 cm in length, in one plane, feather-like, with simple branchlets up to 22 mm in length (Fig. 2b). One main lateral branch of 70 mm in length also bearing simple branchlets up to 28 mm in length. Axis ochre coloured, slender and graceful.

Polyps perpendicular to branchlets (Fig. 3b), in pairs, 4 pairs per cm. Polyps singly placed on stem. Polyps (Fig. 3) cylindrical, funnel shaped distally, about 1.2-1.4 mm in height and about 0.8 mm in diameter in their basal portion; polyp diameter reduced at its mid length, wider distally. Operculum small (Fig. 3c), delicate and easily breakable. Distal part of the polyp directed inward toward the branchlet (Fig. 3b, d). Polyps with 7 longitudinal rows of scales, 6-7 transverse rows of scales on each longitudinal abaxial row overlapping one another (Fig. 3a).

Opercular scales, arranged in 2 alternate cycles of 4 scales each, 0.17-0.49 x 0.07-0.27 mm. Inner cycle (Fig. 4a) small, narrow, lancet-shaped; outer cycle (Fig. 4b) large, concave with blunt, square tip. Inner proximal surface tuberculate, inner distal surface smooth, lacking a keel. Outer proximal surface with few granules, distal outer surface smooth. Basal margin with digitate processes, free margin smooth.

Marginal scales (Fig. 4c), 0.53-0.81 x 0.24-0.40 mm, arranged in 2 alternate cycles of 4 scales each, more or less triangular with long ridged spine up to 2/3 of total scale length. Outer surface smooth, 1/3 of inner proximal surface densely tuberculate,

remaining base-lateral surface of spine smooth, granulated or ridged. Basal margin with digitate processes, free margin serrated.

Body scales (Fig 4d) round and more or less oval, 0.26-0.64 mm in diameter. Outer surface smooth or granulated, inner surface tuberculate, with inner face of the distal margin smooth. Basal margin with digitate processes, free margin serrated.

Coenenchyme scales (Fig. 5) similar in shape to body scales but smaller, 0.15-0.45 mm in maximum length.

Variability

Material examined is comprised of small colony fragments of 2.6-4 cm in length, presumable side branches belong to larger colonies. These fragments show a similar branching to the holotype, having small branchlets from 0.75 to 2.8 cm in length. The polyps are 1.2 to 1.7 cm in height, and are arranged in whorls of two or three on branchlets; being singly placed on stems. The number of whorls per centimetre varies from 4 to 7. The sclerites from the polyps are as in the holotype, varying slightly in sizes: opercular scales, 0.16-0.55 x 0.05-0.27 mm; marginal scales, 0.36-0.81 x 0.14-0.4 mm; and body scales, 0.24-0.64 mm in maximum length.

Geographic and bathymetrical distribution

At present, *Thouarella laxa* is known from the West Pacific Ocean, from Indonesia to Japan. The bathymetric distribution of *T. laxa* varies between 400 and 1301 m.

Remarks

Thouarella laxa was described by Versluys (1906) in his primnoid monograph based on the material collected by the Siboga Expedition. In the same work, Versluys also described a similar species, *T. tydemani*. In the words of Versluys “ist sie unschwer zu unterscheiden von der nächst zu beschreibenden Art diese Genus, der *T. tydemani*, vomit ihre Polypen grosse Ähnlichkeit besitzen”. Because of that it is reasonable to discuss here the taxonomic characters of *T. laxa* with those available from the original

description of *T. tydemani*, and the current existing type material (three slides mounted in Canadian balsam, showing two branches, polyps and coenenchymal scales of the stem). No colony or fragments from the type of this latter species are currently present at the ZMA (Dr. R. van Soest, pers. comm.) and the material seems to have been lost.

Both species have very similar polyps but were differentiated by Versluys due to their general appearance, these polyps being wider and more robust in *Thouarella tydemani* than in *T. laxa*, which also shows differences in sclerites. In addition, *T. tydemani* has body scales with the free margin more curved and marginal scales with slightly larger spines (after Versluys, up to 0.45-0.50 mm in total length of the spine, while this measurement could be about 0.37 mm in *T. laxa*). However, in the examined type material we have observed a variation range for this character between 0.26 and 0.54 mm.

After the study of the holotype of *Thouarella laxa* and additional material from Doflein's collection currently held at the ZSM, the initial ranges of variability observed by Versluys have been increased. The examination of these specimens allow us to give a revised diagnostic description of the known material recognized for this species. For *T. tydemani* there is only the original description, and no more reports have been published based on newly collected specimens. Based on the original description of *T. tydemani*, this species does not show any distinctive character to distinguish it from *T. laxa*. The variability ranges concerning the size of branches, the number of whorls per centimetre, the number of polyps per whorls, the size of polyps, the number of sclerites in the longitudinal abaxial row, and also the description of the sclerites are not different from those in *T. laxa* variability. Versluys (1906: 34) mentions the distinct coenenchymal sclerites of the main stem of *T. tydemani*. The sclerites in this area are usually 0.25 to 0.35 mm. However, they can exceptionally reach to more than 0.7 mm. In the material of *T. laxa* here examined, the range of sizes for the coenenchymal sclerites clearly overlap those mentioned by Versluys for *T. tydemani*. In addition, this character, especially when measured at the main stem, is possibly not one of the best to differentiate two primnoid species because it can be influenced by the age of the colony, the presence of symbionts, and the growth conditions. However, after the study of the three slides of the type specimen of *Thouarella tydemani*, the internal ornamentation of the coenenchymal scales as well as some body scales, consist of short marginal ridges (Fig. 6b, d), absent in *T. laxa*, where these scales show a smooth inner surface. Thus, the

inner ornamentation of these scales could be used as a taxonomic character to distinguish both species.

Although *Thouarella laxa* and *T. tydemani* are very close related, the existing type material of *T. tydemani*, cannot be examined (a few slides in Canada balsam) using the comparative techniques (SEM) currently in use for the taxonomy of this family. Because of this, the redescription of this species is highly recommended after the location of recognizable additional material.

Subgenus *Epithouarella* Kükenthal, 1915

Diagnosis

Thouarella with polyps placed singly. Marginal scales pointed, ridged or with spinose projections, without bearing real thorns.

***Thouarella (Epithouarella) viridis* sp. nov.**

(Figs. 7, 8, 9, 10)

Examined material

Holotype: ZIZMH, ANT XIX/5, stn PS61/164-01, 53°23.80'S, 42°42.03'W, west of South Georgia, Antarctica, 312.5-321.6m depth, 9 April 2002. Paratypes: ZIZMH with the same sampling data as the holotype, four fragments, CRO-0038, ANT XIX/5, stn PS61/150-01, 54°30.22'S 56°08.20'W, south east of Falkland Islands, Antarctica, 286.3-291.6m depth, 6 April 2002, 1 fragmented colony; USNM 1128949 and CRO-0039, ANT XIX/5, stn PS61/167-01, 53°23.68'S, 42°42.23'W, west of South Georgia, Antarctica, 306.0-342.7m depth, 9 April 2002, two fragments and one colony and five fragments and one colony respectively+12 fragments; CRO-0040, ANT XIX/5, stn PS61/174-01, 54°24.47'S, 35°36.81'W, north east of South Georgia, Antarctica, 278.3-279.8 m depth, 11 April 2002, 4 fragments; CRO-0041, ANT XIX/5, stn PS61/182-01, 54°27.63'S, 35°41.33'W, north east of South Georgia, Antarctica, 249.3-256.0 m depth, 12 April 2002, 2 fragments.

Description of the holotype

Fragment of a larger colony (Fig. 7a) of 10.2 cm in total height and 5.4 cm in width, with two short lateral branches of 2.3 and 3.5 cm in length. Main stem and lateral branches with simple branchlets (Fig. 7b), occasionally ramified up to two orders, in a bottlebrush branching pattern. Branchlets up to 3 cm in length, from seven to eight branchlets per cm. Living specimens green in colour, preserved material whitish. Axis ochre in colour, stiff and thick, basal axis diameter of 4 mm.

Polyps (Figs. 7b, 8e) straight and directed upward to stem, singly placed, present on main stem, branches and branchlets. Regularly arranged in spirals and more crowded distally, while in the basal portion the polyps are occasionally nearly opposite, with no tendency toward pairs or whorls, 14-15 polyps per cm. Polyps relatively elongated, cylindrical to club-shaped distally, with a conical operculum (Figs. 8a-d); about 2.1-2.5 mm in height and 0.73-0.85 mm in diameter. Polyp body with 8 longitudinal rows of scales, 6-7 scales on each longitudinal abaxial row (Fig. 8b) overlapping one another and slightly reducing in size basally.

Opercular scales (Fig. 9a) large, eight in number, 0.62-0.78 x 0.24-0.42 mm, isosceles-shaped with bilobed base and acute tip. Proximal inner surface tuberculate, covering more than one third of their length. Prominent, strong, apical keel, often bearing additional lateral ridges. Distal inner surface of scale also with longitudinal ridges. Outer surface granulate, with a longitudinal valley corresponding to inner keel. Basal margin with digitate processes, free margin finely serrated.

Eight marginal scales (Fig. 9b) 0.45-0.63 x 0.47-0.64 mm, pentagonal shaped. Proximal inner surface tuberculate, covering up to about 80% of their length of scales with 3-4 short middle ridges and without a spine; other scales with a small blunt spine sculpted by longitudinal ridges, have tubercles covering up to half of their length. Lateral inner distal surface smooth, without granules or ridges. Outer surface granulate. Basal margin with digitate processes, free margin finely serrated.

Body scales (Fig. 9c) like marginal scales in shape with tendency to be fan shape, 0.27-0.51 x 0.34-0.74 mm., without spines and often decreasing in number and size from distal to basal scales. Adaxials with tendency to be reduced. Inner side with

short ridges and up to about 80% tuberculate. Outer surface granulate. Free margin finely serrated, basal margin with digitate processes.

Coenenchyme scales (Fig. 10) round to oval-shaped, 0.13-0.36 mm in maximum length. Inner surface tuberculate, outer surface granulate forming ridges. Margin irregular with warts proximally and the remainder finely serrated.

Variability

The general colonial and polyp structure of the paratypes and additional examined material is quite similar to that of the holotype. The branches and branchlets arise from all around the main stem, from seven to nine branchlets per centimetre in number. Sometimes the colony shows the branchlets bent towards one side of the colony, giving the false impression of being disposed in a plane. The lateral branches can reach up to 9.3 cm in length, and branchlets up to 4 cm, sometimes ramified up to three orders. The number of polyps per centimetre can vary from 10 to 15 on branchlets. The distribution and form of the sclerites from polyps and coenenchyme are as in the holotype.

Geographic and bathymetrical distribution

At present, *Thouarella viridis* sp. nov. is only known from South Georgia Island area and south east of Falkland Islands, Antarctica (Fig. 1), between , 249 and 342 m in depth.

Etymology

The specific epithet *viridis* comes from one of the most distinctive features shown by this species. When sorting the fresh material collected during the catches its olive green colour was very evident, while most of the Antarctic primnoid gorgonians vary in colour from whitish or yellowish to orange.

Remarks

Polyps of *Thouarella viridis* sp. nov. resemble polyps of the genus *Amphilaphis* due to the prominent apical keel on the inner surface of the opercular scales. However, the distinct branching pattern and the slight reduction in size of the basal body scales, suggest that this species could be better placed among the *Thouarella* species, where the presence of keels on the inner surface of operculars has been also documented.

According to previous works and the recent generic revision of the family Primnoidae given by Cairns and Bayer (2009), *Thouarella viridis* sp. nov. is here considered in the subgenus *Epithouarella* because of the lack of a spine on the marginal scales. In this subgenus three species have been previously recognized: *T. crenelata* Kükenthal, 1907, *T. affinis* Wright and Studer, 1889, and *T. chilensis* Kükenthal, 1908. The former two are present in Antarctic and Subantarctic regions.

After the original description for *Thouarella crenelata*, this species has a high number of sclerites in the abaxial rows of the polyp, from nine to ten, while *T. viridis* sp. nov. only presents six or seven scales. Furthermore the ornamentation of the marginal scales also shows differences in both species. In *T. crenelata* the marginal scales (see Kükenthal, 1919: fig. 217; Cairns and Bayer, 2009: fig. 7 k-l) are strongly ornamented on their free margin, with serrated marginal surface and projecting teeth in the middle distal part; the outer surface being ornamented with longitudinal ridges from the nucleus. In *T. viridis* sp. nov. the marginal scales have an almost flat marginal surface, a smooth (Fig. 9b) or finely serrated margin, and three-four prominent ridges on the middle distal part, like short keels; the outer surface being finely granulated, without forming ridges.

Thouarella affinis and *T. chilensis* overlap with *T. viridis* sp. nov. in the number of scales in the longitudinal abaxial row. Kükenthal (1915) distinguished *T. affinis* and *T. chilensis* by the distinct arrangement of polyps on the branches (possibly branchlets), where *T. affinis* shows a regular distribution, while *T. chilensis* shows a dense distribution, which attains a cylindrical shape. Concerning this character *T. viridis* sp. nov. shows a regular distribution of polyps on the branchlets, being close to *T. affinis*.

Thouarella viridis sp. nov. and *T. affinis* are differentiated by the operculars. In *T. affinis* these scales have two lateral teeth and there is no mention of a strong keel on the inner surface (Wright and Studer, 1889: 67-68, pl. 21 fig. 3). In *T. viridis* sp. nov. this keel is distinctly present in the operculars, lateral teeth being completely absent.

In the case of *T. chilensis*, additional differences with *T. viridis* sp. nov. can be observed in the scales of the polyps. The characteristic large operculars with a strong keel on the inner surface of *T. viridis* sp. nov. contrasts with the smaller operculars without keel shown by *T. chilensis* (Kükenthal, 1919: 303, text fig. 7). In the same work, Kükenthal described body scales with the free margin finely serrated, while in *T. viridis* sp. nov. these sclerites show a series of short distal ridges (see Fig. 9c).

Subgenus *Thouarella* Gray, 1870

Diagnosis

Thouarella with polyps placed singly. Marginal scales bearing thorns.

***Thouarella (Thouarella) minuta* sp. nov.**

(Figs. 11, 12, 13)

Examined material

Holotype: ZIZMH, ANT XXI-2, stn PS65-166-01, 70°56.83'S, 10°32.61'W, Austasen, Antarctica, 253.2-338.0 depth, 15 December 2003. Paratypes: ZIZMH, five fragmented colonies and one fragment; USNM 1128948 CRO-0042, with the same sampling data as the holotype, four colonies and 1 colony, one broken colony and two naked stems respectively; CRO-0031, ANT XVII-3, stn 085-01, 71°11.30'S, 12°15.40'W, Austasen, Antarctica, 309-318 m depth, 2 April 2000, two colony fragmented; CRO-0032, ANT XVII-3, stn 119-01, 70°50.40'S, 10°35.20'W, Austasen, Antarctica, 226-266 depth, 7 April 2000, one colony and two fragments; CRO-0033, ANT XXI-2, stn PS65-174-01, 70°56.57'S, 10°31.86'W, Austasen, Antarctica, 253.2-296.0m depth, 16 December 2003, four fragmented colonies and two fragments; CRO-0034, ANT XXI-2, stn PS65-175-01, 70°56.52'S, 10°31.78'W, Austasen, Antarctica, 288.8-337.2 m depth, 16 December 2003, two colonies without holdfast. CRO-0035, ANT XXI-2, stn PS65-265-01, 70°52.74'S, 10°52.72'W, Austasen, Antarctica, 285.6-294.8 m depth, 27 December 2003, four colonies, one of them broken; CRO-0036, ANT XXI-2, stn PS65-274-01, 70°52.16'S, 10°43.69'W, Austasen, Antarctica, 288.0-290.8 m depth, 28 December 2003, one colony and four fragmented colonies; CRO-0037, ANT XXI-2, stn PS65-

292-01, 72°51.43'S, 19°38.62'W, Austasen, Antarctica, 596.4-597.6m depth, 31 December 2003, one colony without holdfast.

Description of the holotype

Bottlebrush colony (Fig. 11a) of 66 cm in height and 5.5 cm in width, main stem without lateral branches. First 9 cm of the main stem without branchlets, on the remain main stem branchlets all around, up to 4.5 cm long, simple at basal portion or after the first 5 mm ramified up to fourth order (usually 2, exceptionally 3 terminal branchlets starting from the same point), terminal branchlets up to 3.5 cm long (Fig. 11b). Axis light brown in colour, stiff and firmly attached to a rock by a greyish, calcareous, discoidal holdfast of 1.3 cm diameter, basal axis diameter of 3 mm.

Polyps appressed to main stem and branchlets (Fig. 12c), directed upward, arranged singly, apparently alternate to loosely spiral, 11-18 polyps per cm. Polyps (Fig. 12) small, cone-shaped; about 0.71-0.96 mm in height and 0.31-0.44 mm in width at marginal scale level. Polyp body with 5 longitudinal rows of scales, 3-4 scales (excluding operculars) on each longitudinal abaxial row (Fig. 12a, b) overlapping one another.

Opercular scales (Fig. 13a) eight in number, small, 0.25-0.43 x 0.07-0.16 mm, isosceles-shaped or spoon-shaped, adaxial scales reduced. Proximal inner surface slightly tuberculate, covering less than half of the length (about 2/3 of the basal part), with an apical serrated spine as a continuation of -well- developed keel. Outer surface quite smooth. Basal margin with digitate processes, with tendency to square-shape, free margin strongly serrated.

Marginal scales (Fig. 13b) in numbers of eight, small, 0.21-0.28 x 0.16-0.23 mm, round to rhomboidal-shaped, adaxials reduced. Proximal inner surface tuberculate, covering up to about 75% of their longitude. Distal inner surface smooth, with prominent keel projecting to form a short spine distally. Outer surface smooth. Basal margin with digitate processes, free margin serrated.

Body scales (Fig. 13c) circular shaped, like marginal scales in size, 0.22-0.28 mm in maximum length, without keel or spine. Adaxials with tendency to be reduced. Inner surface tuberculate, outer surface smooth. Free margin quite entire, basal margin irregular due to presence of tubercles and processes.

Coenenchymal scales (Fig. 13d) round to oval-shaped, 0.12-0.22 mm in maximum diameter. Inner surface sparsely tuberculate, outer surface smooth with some granules or short ridges. Irregular margin due to the presence of tubercles and serrations.

Variability

The general colonial structure of the paratypes and additional examined material is quite similar to that of the holotype. The material studied includes colonies from 11 cm to 66 cm in height. The branchlets arise from all around the main stem; some colonies have bend branchlets, giving a false impression of a flattened or pinnate forms, some branchlets are bent upward, up to 4.5 cm in length, simple or ramified, like those of the holotype. The polyp's proximity to the stem varies from inclined to appressed. There are from 10 to 18 polyps per centimetre on branchlets, and they are up to about 1.1 mm in height. The distribution and shape of the sclerites of the polyps and coenenchyme are as in the holotype.

Geographic and bathymetrical distribution

At present, *Thouarella minuta* sp. nov. is known only from, Austasen, Antarctica (Fig. 1), between 226 and 338 m in depth.

Etymology

In this species, the most interesting character is the minute size of the polyps, one of the smallest in comparison with its congeners. The specific name, *minuta*, comes from Latin, meaning small or minute.

Remarks

We include *Thouarella minuta* sp. nov., in the subgenus *Thouarella* due to the presence of a spine on the marginal scales and the single arrangement of the polyps on the branchlets. Five species in this group are comparable to *T. minuta* sp. nov. because of the presence of polyps with 4-5 scales in the longitudinal abaxial rows: *T. variabilis*

Wright and Studer, 1889, *T. brucei* Thomson and Ritchie, 1906, *T. striata* Kükenthal, 1907, *T. clavata* Kükenthal, 1908, and *T. pendulina* (Roule, 1908).

Thouarella striata differs from all other species listed above due to the distinct radial ridges on the outer surface of the polyp scales (Kükenthal, 1907); quite different from the smooth surface of the sclerites in *T. minuta* sp. nov. In addition, polyps of *T. striata* are 1.5-2 mm in height, while those of *T. minuta* sp. nov. only reach 1.1 mm (usually less).

Thouarella clavata and *Thouarella variabilis* show polyps larger than 1.5 mm in height, the former reaching 2 mm, and the latter usually larger than 2 mm (see Kükenthal, 1907; Wright and Studer, 1889). This feature clearly contrasts with the polyps of *T. minuta* sp. nov. All varieties of *T. variabilis* considered by Wright and Studer (1889:68) have more or less well developed distal spines on the “ultimate and penultimate rows” and those authors mention that var. *gracilis* also has a median keel on the “uppermost” scales. For *T. clavata*, Kükenthal (1907 as *T. aff. antarctica*, 1908) commented that the operculars have keels and the marginals have spines; while in *T. minuta* sp. nov., both the operculars and the marginals show spines and keels, the spine of the marginals being shorter.

Thouarella pendulina and *Thouarella brucei* (taking into account the scarce and sometimes imprecise original description of this latter species) show polyps of about 1 mm (up to 1.5 mm in the former) (see Thomson and Ritchie, 1906; Roule, 1908; Gravier, 1914). As previously described, *T. minuta* sp. nov. has polyps ranging between 0.7 and 1.1 mm in height. *Thouarella pendulina* is clearly distinguishable from the other two species by the dense distribution of polyps in the branchlets, up to 70 polyps per 1 cm. Although *T. pendulina* and *T. minuta* sp. nov. share a characteristic smooth outer surface of the scales, the submarginal scales of *T. pendulina* bear a small spine (after the drawing in Gravier, 1914: fig. 90, these scales apparently have a keel), not present in *T. minuta* sp. nov.

Apart from the few characters mentioned here, and repetitively included in the available key to species of *Thouarella* (Kükenthal, 1924) the original description of *T. brucei* does not offer reliable diagnostic characteristics that could be used here in this species comparison. Kükenthal (1919) commented on the incomplete description of this species, especially on the description of the polyps and designation of distalmost scale

cycles, maintaining this species in the subgenus *Parathouarella* (currently reconsidered as subgenus *Thouarella*). Later on, Kukenthal (1924: 301) included this species as *dubiae atque incertae sedis*. The original material came from shelf depths of three widely separated localities [Burdwood Bank (~54° S), Gough Island (~40° S) and Santa Helena (~16° S, depth not mentioned)], although the type is that from Burwood Bank (Emma Sherlock, pers. comm.). Broch (1965) described the morphology of several specimens collected at Burdwood Bank from about 111 meters depths, and assigned his material to the Thomson and Ritchie' species. However, the material examined by Broch clearly differs from that described by the original authors in the arrangement of polyps: “*The polyps are placed in distinct whorls on the twigs, ... The whorl generally consists of 3 or 4 polyps, but in the distal part of more luxuriantly developed twigs whorls with 5 polyps also are rather common.*” (see Broch, 1965: 27), compared to “*On the twigs they [the polyps] are closely approximated, arising in all directions and without any definite arrangement*” (see Thomson and Ritchie, 1906: 853). Despite the few characters available for *T. brucei*, this species has lateral branches from the main stem, and the branchlets are proportionally shorter and less ramified in comparison with those in *T. minuta* sp. nov. (see Thomson and Ritchie, 1906: pl. I, fig. 1 and Fig. 11b of this paper), while the numerous colonies examined of the new taxon show only elongated colonies without lateral branches (see Fig. 11a).

Acknowledgments

The authors would like to thank Dr. R. van Soest and E.J. Beglinger (Universiteit van Amsterdam), Bernhard Ruthensteiner and Eva Lodde (Zoologische Staatssammlung München) for lending the specimens of *Thouarella laxa* and *T. tydemani* for examination, and Emma Sherlock (NHM in London) for information about the type material of *T. brucei*. We also acknowledge the valuable assistance of the officers and crew of the R/V *Polarstern*, and many colleagues on board during the EASIZ III, LAMPOS and BENDEX cruises. We take this opportunity to extend our thanks to the cruise leaders and steering committee of the cruises especially Wolf E. Arntz (Alfred Wegener Institut, Bremerhaven), who kindly facilitated the work on board and for the opportunity to collaborate in these Antarctic programmes. Special

thanks are addressed to some friends and colleagues for their valuable assistance during EASIZ III (M^a Isabel Alfonso, Josep-Maria Gili), LAMPOS (Neus Vert and Estefanía Rodríguez), and BENDEX (Estefanía Rodríguez, Josep-Maria Gili) cruises. Support for this work was provided by the Spanish CICYT projects ANT98-1739-E and ANT99-1608-E (EASIZ III), REN2001-4929-E/ ANT (LAMPOS), REN2003-04236 (BENDEX), and POL2006-06399/CGL (CLIMANT). Mr. Tony Krupa is thanked for reviewing the English version.

References

- Alderslade P (1998) Revisionary systematics in the gorgonian family Isididae, with description of numerous new taxa (Coelenterata, Octocorallia). *Rec W Aust Mus* 55: 1-359
- Bayer FM (1980) *Armadilloorgia cyathella* and *Ophidiorgia paradoxa*, two new genera and species of primnoid octocorals (Coelenterata: Anthozoa) from South Georgia and South Orkney Islands. *Proc Biol Soc Wash* 93(1): 216-228
- Bayer FM (1981) Key to the genera of Octocorallia exclusive of Pennatulacea (Coelenterata: Anthozoa), with diagnoses of new taxa. *Proc Biol Soc Wash* 94(3): 901-947
- Bayer FM (1988) *Mirostenella articulata*, a remarkable new genus and species of primnoid octocoral with uncalcified axial nodes. *Proc Biol Soc Wash* 101(2): 251-256, figs. 1-3
- Bayer FM (1996a) The Antarctic genus *Callozostron* and its relationship to *Primnoella* (Octocorallia: Gogonacea: Primnoidae). *Proc Biol Soc Wash* 109(1): 150-203
- Bayer FM (1996b) New Primnoid Gorgonians (Coelenterata: Octocorallia) from Antarctic waters. *Bull Mar Sci* 58(2): 511-530
- Bayer FM (1998) A review of the circumaustral gorgonacean genus *Fannyella* Gray, 1870 with description of five new species (Coelenterata: Octocorallia: Primnoidae). *Senckenberg Biol* 77(2): 161-204

- Bayer FM, Stefani J (1988) Primnoidae (Gorgonacea) de Nouvelle-Calédonie. Bull Mus Natn Hist Nat Paris (4) 10(A)3: 449-476
- Bayer FM, Grasshoff M, Verseveldt J (1983) Illustrated trilingual glossary of morphological and anatomical terms applied to Octocorallia. E.J. Brill/Dr. W. Backhuys, Leiden pp 1-75
- Broch H (1965) Some octocorals from Antarctic Waters. Scientific Results of the "Brategg" Expedition, 1947-48. Christensens Hvalfangstmuseum i Sandejford Publikasjon 26: 19-38, 1-7
- Cairns SD (2006) Studies on western Atlantic Octocorallia (Coelenterata: Anthozoa). Part 6: The genera *Primnoella* Gray, 1858; *Thouarella* Gray, 1870; *Dasystenella* Versluys, 1906. Proc Biol Soc Wash 119(2): 161-194
- Cairns SD, Bayer FM (2009) A Generic Revision and Phylogenetic Analysis of the Primnoidae (Cnidaria: Octocorallia). Smithson Contrib Zool 629: 1-79
- Gravier CH (1914) Alcyonaires. Deuxième Expédition Antarctique Française (1908-1910). Sciences Naturelles: Documents scientifiques pp1-118, 1-10
- Gray JE (1858) Synopsis of the Families and Genera of Axiferous Zoophytes or Barked Corals. Proc Zool Soc Lond 1857:278-294
- Gray JE (1870) Catalogue of the lithophytes or stony corals in the collection of the British Museum. London: British Museum pp 1-51
- Kinoshita K (1908) *Diplocalyptra*, eine neue Untergattung von *Thouarella* (Primnoidae). Annot Zool Jap 7 (1): 49-60
- Kükenthal W (1907) Gorgoniden der Deutschen Tiefsee Expedition. Zool Anz 31 (7): 202-212
- Kükenthal W (1908) Diagnosen neuer Gorgoniden (4 Mitteilung). Zool Anz 33 (1): 9-20
- Kükenthal W (1915) System und Stammesgeschichte der Primnoidae. Zool Anz 46 (5): 142-158
- Kükenthal W (1919) Gorgonaria. Wissensch Ergebn deutschen Tiefsee-Exped "Valdivia" 13 (2). 1-946, 30-89

- Kükenthal W (1924) Coelenterata: Gorgonaria. Das Tierreich 47. Walter de Gruyter and Co, Berlin pp 1-478
- Kükenthal W, Gorzawsky H (1908) Diagnosen neuer japanischer Gorgoniden (Reise Doflein 1904-1905). Zool Anz 32: 621-631
- López-González PJ, Gili JM (2002) A new primnoid genus (Anthozoa: Octocorallia) from the Southern Ocean. Sci Mar 66(4): 383-397
- Roule L (1908) Alcyonaires. Expédition Antarctique Française (1903-1905). Sciences Naturelles: Documents scientifiques 15: 1-6, 1
- Thomson JA, Ritchie J (1906) The Alcyonarians of the Scottish National Antarctic Expedition. Trans R Soc Edinburgh 41(3): 851-860, 1-2
- Versluys J (1906) Die Gorgoniden der Siboga-Expedition. II. Die Primnoidae. Siboga-Expedition, 13a: 1-187
- Wright EP, Studer T (1889) Report on the Alcyonaria collected by H.M.S. Challenger during the years 1873-1876. Rep. Sci. Res. Challenger. Zool 31: i-lxxvii, 1-314, 1-43

Figure Captions

Figure 1.- Known distribution of Antarctic *Thouarella* species described in this paper: triangle, *Thouarella viridis* sp. nov.; square, *Thouarella minuta* sp. nov. Solid symbols represent the respective localities where the holotypes were collected.

Figure 2.- *Thouarella laxa* Versluys, 1906, holotype (COEL03576): **a**, whole colony; **b**, detail of branchlets.

Figure 3.- *Thouarella laxa* Versluys, 1906, holotype (COEL03576): **a**, polyp on abaxial view; **b**, polyp on lateral adaxial view; **c**, polyp on oral view; **d**, polyp on lateral view. Abbreviations: **o**, opercular scales; **m**, marginal scales.

Figure 4.- *Thouarella laxa* Versluys, 1906, holotype (COEL03576): Opercular scales from inner (**a**) and outer (**b**) alternate cycles; **c**, marginal scales; **d**, body scales.

Figure 5.- *Thouarella laxa* Versluys, 1906, holotype (COEL03576): Coenenchymal scales.

Figure 6.- *Thouarella tydemani* Versluys, 1906, holotype (COEL09256): **a**, detail of branchlets; **b**, coenenchymal scales of branchlet; **c**, body scales with smooth margin; **d**, body scale with ridged margin; **e-f**, coenenchymal scales of main stem.

Figure 7.- *Thouarella viridis* sp. nov, holotype (_____): **a**, whole colony; **b**, detail of branchlet.

Figure 8.- *Thouarella viridis* sp. nov, holotype (_____): **a**, polyp in oral view, stereo pair; **b**, polyp in abaxial view; **c**, polyp in adaxial view; **d**, polyp in lateral view; **e**, detail of branchlets, stereo pair. Abbreviations: **o**, opercular scales; **m**, marginal scales.

Figure 9.- *Thouarella viridis* sp. nov., holotype (_____): **a**, opercular scales; **b**, marginal scales; **c**, body scales.

Figure 10.- *Thouarella viridis* sp. nov., holotype (_____): Coenenchymal scales.

Figure 11.- *Thouarella minuta* sp. nov, holotype (_____): **a**, whole colony; **b**, detail of branchlets.

Figure 12.- *Thouarella minuta* sp. nov., holotype (_____): **a** polyps in lateral abaxial view, stereo pair; **b**, polyps in abaxial view, stereo pair; **c**, detail of branchlets. Abbreviations: **o**, opercular scales; **m**, marginal scales.

Figure 13.- *Thouarella minuta* sp. nov., holotype (_____): **a**, opercular scales; **b**, marginal scales; **c**, body scales; **d**, coenenchymal scales.































