



HAL
open science

Spectral triples for finitely generated groups.

Sébastien Palcoux

► **To cite this version:**

| Sébastien Palcoux. Spectral triples for finitely generated groups.. 2010. hal-00529553v1

HAL Id: hal-00529553

<https://hal.science/hal-00529553v1>

Preprint submitted on 25 Oct 2010 (v1), last revised 14 Jan 2011 (v8)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spectral triples for finitely generated groups.

Sébastien Palcoux*

Abstract

Using Cayley graphs and Clifford algebras, we are able to give spectral triples for at least a large class of finitely generated groups, such that the phase of the Dirac operator is generically non-trivial.

Contents

1	Basic definitions	2
2	Geometric construction	2
3	Clifford algebra	3
4	Dirac operator	4

*Institut de Mathématiques de Luminy, Marseille, France.
web: <http://iml.univ-mrs.fr/~palcoux> , mail: palcoux@iml.univ-mrs.fr

1 Basic definitions

Definition 1.1. A spectral triple (\mathcal{A}, H, D) is given by a unital \star -algebra \mathcal{A} represented on the Hilbert space H , and an unbounded operator D , called the Dirac operator, such that:

1. D is self-adjoint.
2. $(D^2 + I)^{-1}$ is compact.
3. $\{a \in \mathcal{A} \mid [D, a] \in B(H)\}$ is dense in \mathcal{A} .

See the article [2] of G. Skandalis, dedicated to A. Connes and spectral triple.

Definition 1.2. A group Γ is finitely generated if it exists a finite set $S \subset \Gamma$ generating Γ . We always take S equals to S^{-1} and the identity element $e \notin S$. We can also defined a group by generators and relations: $\Gamma = \langle S \mid R \rangle$.

2 Geometric construction

Definition 2.1. Let $\ell : \Gamma \rightarrow \mathbb{N}$ be the word length related to S . Let \mathcal{G} be the Cayley graph with the distance $d(g, h) = \ell(gh^{-1})$, see [1].

Definition 2.2. Let $p : \Gamma \rightarrow \mathbb{N}$ with $p(g)$ the number of geodesic paths from g to the identity element e , on the Cayley graph \mathcal{G} .

Definition 2.3. Let $p_s : \Gamma \rightarrow \mathbb{N}$ with $p_s(g)$ the number of geodesic paths from g to e , starting by $s \in S$. Then $\sum_{s \in S} p_s = p$.

Remark 2.4. If $p_s(g) \neq 0$ then $p_s(g) = p(gs)$.

Definition 2.5. Let $\ell^2(\Gamma)$ be the canonical Hilbert space of base $(e_g)_{g \in \Gamma}$ and let ∂_s ($s \in S$) be the unbounded operator defined by:

$$\partial_s \cdot e_g = \frac{p_s(g)}{p(g)} \ell(g) e_g.$$

∂_s is a diagonal positive operator, so $\partial_s^* = \partial_s$ and $\partial_s \partial_{s'} = \partial_{s'} \partial_s$.

Example 2.6. Let $\Gamma = \mathbb{Z}^2$, $S = \{a, a^{-1}, b, b^{-1}\}$, $R = \{aba^{-1}b^{-1}\}$.
Let $g = a^n b^m = (n, m)$ with $n, m \in \mathbb{N}$, then $\ell(g) = n + m$, $p(g) = C_{n+m}^n$,
 $p_{a^{-1}}(g) = C_{n+m-1}^{n-1}$, $p_{b^{-1}}(g) = C_{n+m-1}^{m-1}$, and so $\frac{C_{n+m-1}^{n-1}}{C_{n+m}^n}(n + m) = n$.
Let $\partial_1 = \partial_{a^{-1}}$ and $\partial_2 = \partial_{b^{-1}}$, then:

$$\partial_1 e_{(n,m)} = n e_{(n,m)} \quad \text{and} \quad \partial_2 e_{(n,m)} = m e_{(n,m)},$$

as for the canonical derivations !

Example 2.7. Let $\Gamma = \mathbb{F}_2$, $S = \{a, a^{-1}, b, b^{-1}\}$, $R = \emptyset$. Let $g \in \mathbb{F}_2$, $g \neq e$,
then $p(g) = 1$, let $s \in S$ with $p_s(g) = 1$, then $\partial_s . e_g = \ell(g) . e_g$.

3 Clifford algebra

We quickly recall here the notion of Clifford algebra, for a more detailed exposition, see the course of A. Wassermann [3].

Definition 3.1. Let $\Lambda(\mathbb{R}^S)$ be the exterior algebra equals to $\bigoplus_{k=0}^{2d} \Lambda^k(\mathbb{R}^d)$,
with $2d = \text{card}(S)$ and $\Lambda^0(\mathbb{R}^d) = \mathbb{R}\Omega$. We called Ω the vacuum vector.
Recall that $v_1 \wedge v_2 = -v_2 \wedge v_1$ so that $v \wedge v = 0$.

Definition 3.2. Let a_v be the creation operator on $\Lambda(\mathbb{R}^S)$ defined by:

$$a_v(v_1 \wedge \dots \wedge v_r) = v \wedge v_1 \wedge \dots \wedge v_r \quad \text{and} \quad a_v(\Omega) = v$$

Reminder 3.3. The dual a_v^* is called the annihilation operator, then:

$$a_v^*(v_1 \wedge \dots \wedge v_r) = \sum_{i=0}^r (-1)^{i+1} (v, v_i) v_1 \wedge \dots \wedge v_{i-1} \wedge v_{i+1} \wedge \dots \wedge v_r \quad \text{and} \quad a_v^*(\Omega) = 0$$

Reminder 3.4. Let $c_v = a_v + a_v^*$, then $c_v = c_v^*$ and $c_v c_w + c_w c_v = 2(v, w)I$.

Definition 3.5. The operators c_v generate the Clifford algebra $\text{Cliff}(\mathbb{R}^S)$.
Note that the operators c_v are bounded and that $\text{Cliff}(\mathbb{R}^S) . \Omega = \Lambda(\mathbb{R}^S)$.

Remark 3.6. \mathbb{R}^S admits the orthonormal basis $(v_s)_{s \in S}$.
We will write c_s instead of c_{v_s} , so that $[c_s, c_{s'}]_+ = 2\delta_{s,s'}I$.

4 Dirac operator

Definition 4.1. Let H be the Hilbert space $\Lambda(\mathbb{R}^S) \overline{\otimes} \ell^2(\Gamma)$, we define a self-adjoint operator D on a dense domain of H by:

$$D = \sum_{s \in S} c_s \otimes \partial_s.$$

Lemma 4.2. $D^2 = I \otimes (\sum_{s \in S} \partial_s^2)$.

Proof. $D^2 = \sum_{s, s' \in S} c_s c_{s'} \otimes \partial_s \partial_{s'} = \frac{1}{2} \sum (c_s c_{s'} + c_{s'} c_s) \otimes \partial_s \partial_{s'} = I \otimes (\sum \partial_s^2)$ \square

Lemma 4.3. $(D^2 + I)^{-1}$ is compact.

Proof. First of all $\dim(\Lambda(\mathbb{R}^S)) = 2^{2d} < \infty$.

Next $\sum_{s \in S} \frac{p_s(g)}{p(g)} = 1$, then it exists $s_o \in S$ such that $\frac{p_{s_o}(g)}{p(g)} \geq \frac{1}{2d}$.

Now $\sum_{s \in S} [\ell(g) \frac{p_s(g)}{p(g)}]^2 \geq [\ell(g) \frac{p_{s_o}(g)}{p(g)}]^2 \geq [\frac{\ell(g)}{2d}]^2$. \square

Definition 4.4. Let \mathcal{C} be the class of finitely generated group satisfying:
 $\forall g, h \in \Gamma, h \neq e$:

$$\left| \frac{p_s(gh)}{p(gh)} - \frac{p_s(h)}{p(h)} \right| \leq \frac{K_g}{\ell(h)}$$

with $K_g \in \mathbb{R}_+$ depending only on g .

Examples 4.5. The class \mathcal{C} is stable by direct or free product, it contains $\mathbb{Z}^n, \mathbb{F}_n$, the finite groups, and probably every amenable or hyperbolic groups.

Question 4.6. Does \mathcal{C} contains every finitely generated groups ?

From now, the group Γ is in the class \mathcal{C} .

Definition 4.7. Let $\mathcal{A} = C_r^* \Gamma$, acting on H by $I \otimes u_g, g \in \Gamma$.

Proposition 4.8. $\{a \in \mathcal{A} \mid [D, a] \in B(H)\}$ is dense in \mathcal{A} .

Proof. $[D, u_g](\Omega \otimes e_h) = \dots = (\sum_{s \in S} [\ell(gh) \frac{p_s(gh)}{p(gh)} - \ell(h) \frac{p_s(h)}{p(h)}] v_s) \otimes e_{gh}$.

Now $|\ell(gh) \frac{p_s(gh)}{p(gh)} - \ell(h) \frac{p_s(h)}{p(h)}| \leq \ell(gh) \left| \frac{p_s(gh)}{p(gh)} - \frac{p_s(h)}{p(h)} \right| + |\ell(gh) - \ell(h)| \frac{p_s(h)}{p(h)}$
 $\leq K_g \frac{\ell(gh)}{\ell(h)} + \ell(g) \leq K_g(\ell(g) + 1) + \ell(g)$. It's bounded in h . \square

Theorem 4.9. (\mathcal{A}, H, D) is a spectral triple.

The phase of D is generically non-trivial.

References

- [1] P. de la Harpe, *Topics in geometric group theory*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [2] G. Skandalis *Géométrie non commutative d'après Alain Connes: la notion de triplet spectral*. Gaz. Math. No. 94 (2002), 4451.
- [3] A. Wassermann, *Lecture notes on Atiyah-Singer index theorem*, Lent 2010 course, <http://www.dpmms.cam.ac.uk/~ajw/AS10.pdf>