



HAL
open science

Carleman estimates for stratified media

Assia Benabdallah, Yves Dermenjian, Jérôme Le Rousseau

► **To cite this version:**

Assia Benabdallah, Yves Dermenjian, Jérôme Le Rousseau. Carleman estimates for stratified media. 2010. hal-00529924v1

HAL Id: hal-00529924

<https://hal.science/hal-00529924v1>

Preprint submitted on 27 Oct 2010 (v1), last revised 19 Mar 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Carleman estimates for stratified media*

Assia Benabdallah¹, Yves Dermenjian²,
Laboratoire d'Analyse Topologie Probabilités^a
CNRS UMR 6632, Université de Provence,
Aix-Marseille Universités, France

and

Jérôme Le Rousseau³
Laboratoire Mathématiques et Applications, Physique Mathématique d'Orléans^b
CNRS UMR 6628, Université d'Orléans, France

October 27, 2010

Abstract

We consider anisotropic elliptic and parabolic operators in a bounded stratified media in \mathbb{R}^n characterized by discontinuities of the coefficients in one direction. The surfaces of discontinuities cross the boundary of the domain. We prove Carleman estimates for these operators with an arbitrary observation region.

AMS 2010 subject classification: 35J15, 35K10.

Keywords: elliptic operators; parabolic operators; non-smooth coefficients; stratified media; Carleman estimate; observation location;

Contents

1	Introduction, notation and main results	2
2	Spectral properties and weight function	6
3	The Elliptic case: proof of Theorem 1.4	8
4	The parabolic case: proof of Theorem 1.5	12
A	Proof of some intermediate results	19

*The authors were partially supported by l'Agence Nationale de la Recherche under grant ANR-07-JCJC-0139-01.

¹assia@cmi.univ-mrs.fr, ²dermenji@cmi.univ-mrs.fr, ³jlr@univ-orleans.fr

^aLATP – 39, rue F. Joliot-Curie, 13453 Marseille cedex 13, France.

^bMAPMO – Route de Chartres, B.P. 6759, 45067 Orléans cedex 2, France.

1 Introduction, notation and main results

Consider a bounded open set $\Omega \subset \mathbb{R}^n$. For an second-order elliptic operator, say $A = -\Delta_x$, Carleman estimates take the form¹

$$s^3 \|e^{s\varphi} u\|_{L^2(\Omega)}^2 + s \|e^{s\varphi} \nabla_x u\|_{L^2(\Omega)}^2 \lesssim \|e^{s\varphi} Au\|_{L^2(\Omega)}^2, \quad u \in \mathcal{C}_c^\infty(\Omega), \quad s \geq s_0,$$

for a properly chosen weight function $\varphi(x)$ and s_0 sufficiently large (see e.g. [17]). It is common to use a weight function of the form $\varphi(x) = e^{\lambda\beta(x)}$, with β such that $|\beta'| \neq 0$ and λ sufficiently large. Including a second large parameter (see [16]), the Carleman estimate then takes the form

$$s^3 \lambda^4 \|\varphi^{\frac{3}{2}} e^{s\varphi} u\|_{L^2(\Omega)}^2 + s \lambda^2 \|\varphi^{\frac{1}{2}} e^{s\varphi} \nabla_x u\|_{L^2(\Omega)}^2 \lesssim \|e^{s\varphi} Au\|_{L^2(\Omega)}^2, \quad u \in \mathcal{C}_c^\infty(\Omega), \quad s \geq s_0, \quad \lambda \geq \lambda_0.$$

For a parabolic operator, say $P = \partial_t + \Delta_x$ on $Q = (0, T) \times \Omega$, Carleman estimates can be derived [16] in the following form

$$s^3 \lambda^4 \|(a\varphi)^{\frac{3}{2}} e^{s a \eta} u\|_{L^2(Q)}^2 + s \lambda^2 \|(a\varphi)^{\frac{1}{2}} e^{s a \eta} \nabla_x u\|_{L^2(Q)}^2 \lesssim \|e^{s a \eta} Pu\|_{L^2(Q)}^2, \\ u \in \mathcal{C}^\infty(Q), \quad \text{supp}(u(t, \cdot)) \Subset \Omega, \quad s \geq s_0, \quad \lambda \geq \lambda_0,$$

for $a(t) = (t(T-t))^{-1}$, $\varphi(x) = e^{\lambda\beta(x)}$, with β such that $|\beta'| \neq 0$ and $\eta(x) = e^{\lambda\beta(x)} - e^{\lambda\bar{\beta}} < 0$. In this later case the weight function $a(t)\eta(x)$ is singular at time $t = 0$ and $t = T$. For a review of Carleman estimates for elliptic and parabolic operators we refer to [13, 24].

The estimates we have presented are said to be local, as they apply for compactly supported functions in Ω . So-called global Carleman estimates can be derived (see e.g. [16]). They concern functions defined on the whole Ω with prescribed boundary conditions, e.g. homogeneous Dirichlet, Neumann. They are also characterized by the presence of an observation term on $\omega \subset \Omega$ on the r.h.s. of the estimate, e.g., for the elliptic operator $A = -\Delta$,

$$s^3 \lambda^4 \|\varphi^{\frac{3}{2}} e^{s\varphi} u\|_{L^2(\Omega)}^2 + s \lambda^2 \|\varphi^{\frac{1}{2}} e^{s\varphi} \nabla_x u\|_{L^2(\Omega)}^2 \lesssim \|e^{s\varphi} Au\|_{L^2(\Omega)}^2 + s^3 \lambda^4 \|\varphi^{\frac{3}{2}} e^{s\varphi} u\|_{L^2(\omega)}^2, \quad s \geq s_0, \quad \lambda \geq \lambda_0,$$

for $u \in \mathcal{C}^\infty(\bar{\Omega})$, and $u|_{\partial\Omega} = 0$.

Note also that Carleman estimates can be patched together (see e.g. [17, 24]). If local estimates are obtained at the boundary $\partial\Omega$, then one can deduce global estimates from the local ones.

Carleman have many applications. In 1939, T. Carleman introduced these estimates to prove a uniqueness result for some elliptic partial differential equations (PDE) with smooth coefficients in dimension two [10]. This result was later generalized (see e.g. [17, Chapter 8], [18, Chapter 28], [34]). In more recent years, the field of applications of Carleman estimates has gone beyond the original domain. They are also used in the study of inverse problems (see e.g. [9, 20, 19, 22]) and control theory for PDEs. Through unique continuation properties, they are used for the exact controllability of hyperbolic equations [3]. They also yield the null controllability of linear parabolic equations [29] and the null controllability of classes of semi-linear parabolic equations [16, 2, 14].

Difficulties arise for the derivation of Carleman estimates in the case of non-smooth coefficients in the principal part of the operator, i.e., for a regularity lower than Lipschitz. In fact, it is known

¹ $a \lesssim b$ stands for $a \leq Cb$ for some constant $C > 0$.

that unique continuation does not hold in general for a $\mathcal{C}^{0,\alpha}$ Hölder regularity of the coefficient with $0 < \alpha < 1$ [32, 31], which ruins any hope to prove a Carleman estimate.

In the present article, we consider coefficients that are *discontinuous* across a smooth interface, yet regular on each side. This question was first addressed in [12] for a parabolic operator $P = \partial_t - \nabla_x(c(x)\nabla_x)$, with a monotonicity assumption: the observation takes place in the region where the diffusion coefficient c is the ‘lowest’. In the one-dimensional case², the monotonicity assumption was relaxed for general piecewise \mathcal{C}^1 coefficients [6, 7], and for coefficients with bounded variations [23]. The case of an arbitrary dimension without any monotonicity condition in the elliptic case was solved in [5, 27] and in the parabolic case in [28]. In [25, 26] the case of a discontinuous anisotropic matrix coefficients is treated and a sharp condition on the weight function is provided for the Carleman estimate to hold.

The methods used in [5, 27, 28, 25, 26] focus on a neighborhood of a point at the interface where the interface can be given by $\{x_n = 0\}$ for an appropriate choice of coordinates $x = (x', x_n)$, $x' \in \mathbb{R}^{n-1}$, $x_n \in \mathbb{R}$. Then, through microlocal techniques (Calderón projector or first-order factorization), a local Carleman estimate is proven. However, these methods require strong regularity for the coefficients and for the interface. Moreover, they fall short if the interface crosses the boundary. This configuration is typical in bounded stratified media such as those we consider below.

In stratified media, a controllability result for a *linear* parabolic equation in arbitrary dimension was obtained in [8]. The approach was based on the 1D Carleman estimates of [6, 23] and a spectral inequality as in [29, 30, 21] for the transverse elliptic operator, whose coefficients are smooth. The precise definition of such stratified media is given below.

The controllability result obtained in [8] left the question of deriving a Carleman estimate open for stratified media. This result is achieved here. One of the consequences of this result is the null-controllability of classes of semi-linear parabolic equations. We refer to [12] for these developments.

Remark 1.1. The following observation also provides hints that Carleman estimates can be derived for stratified media [33]. As we shall assume below interfaces crosses the boundary transversely. Pick a point at the intersection of an interface and the boundary and choose local coordinates such that the interface is orthogonal to the boundary. Assume that the coefficients associated with the transverse part of the operator are flat at the boundary. Then, by reflection at the boundary, the system under consideration can be turned into a problem with a smooth interface away from any boundary which permits to use the results of [27, 28, 25, 26]. This situation is however not general.

We finish this introductory presentation by pointing out the difficulty that arises when deriving a Carleman estimate for the operator $A = -\nabla_x(c(x)\nabla_x)$ or $P = \partial_t - \nabla_x(c(t, x)\nabla_x)$ in dimension greater than two, in the presence of an interface S . In fact, the standard Carleman derivation method leads to interface terms involving

²Simultaneously to these results, a controllability result for *linear* parabolic equations with $c \in L^\infty$ was proven in [1] in the one-dimensional case *without* Carleman estimate. This controllability result does not cover more general semi-linear equations. An earlier result was that of [15] where the controllability of a *linear* parabolic equations was proven in one dimension with $c \in BV$ through D. Russel’s method.

1. trace of the function $u|_S$. A zero- and first-order operators in the tangent direction acts on $u|_S$.
2. traces of its normal derivative $\partial_{x_n} u|_{S_\pm}$, on both sides of S .

This interface contribution can be interpreted as a quadratic form (see [6]). In [27, 28] the authors show that this quadratic form is only non-negative for low (tangential) frequencies. Here we shall recover this behavior where the tangential Fourier transform is replaced by Fourier series, built on a basis of eigenfunctions of the transverse part of the elliptic operator. For high (tangential) frequencies, the tangential derivative term (i.e., the action of a the first-order operator on $u|_S$) yields a negative contribution, unless a monotonicity assumption on the coefficient c is made as in [12]. In [5, 27, 28] the authors have used microlocal methods in the high frequency regime to solve this difficulty, and more recently in [26, 25]. Here, because of the intersection of the interface with the boundary, and because of the little regularity required for the diffusion coefficients, such methods cannot be used directly. However, the separated-variable assumption made on the diffusion coefficients allows us to use Fourier series and similar ideas can be developed: low frequencies and high frequencies are treated differently. In the parabolic case the separation we make between the two frequency regimes is time dependent. Here, the separated-variable assumption yields explicit computations, which reveals the behavior of the solution in each frequency regime.

In the present article, a particular class of anisotropic coefficients is treated. The question of deriving Carleman estimates for more general coefficients in the neighborhood of the intersection of an interface, where the coefficients jump, with the boundary is left open.

1.1 Setting and notation

We let Ω be an open subset in \mathbb{R}^n , with $\Omega = \Omega' \times (-H, H)$, where Ω' is a nonempty bounded open subset of \mathbb{R}^{n-1} with \mathcal{C}^1 boundary³.

We shall use the notation $x = (x', x_n) \in \Omega' \times (-H, H)$. We set $S = \Omega' \times \{0\}$, that will be understood as an interface where coefficients and functions may jump. For a function $f = f(x)$ we define by $[f]_S$ its jump at S , i.e.,

$$[f]_S(x') = f(x)|_{x_n=0^+} - f(x)|_{x_n=0^-}.$$

For a function u defined on both sides of S , we set

$$u|_{S_\pm} = (u|_{\Omega_\pm})|_S,$$

with $\Omega_+ = \Omega' \times (0, H)$ and $\Omega_- = \Omega' \times (-H, 0)$.

Let $B(t, x)$, $t \in (0, T)$ and $x \in \Omega$, be with values in $M_n(\mathbb{R})$, the space of square matrices with real coefficients of order n . We make the following assumption.

Assumption 1.2. *The matrix diffusion coefficient $B(t, x', x_n)$ has the following block diagonal form*

$$B(t, x', x_n) = \begin{pmatrix} c_1(t, x_n)C_1(x') & 0 \\ 0 & c_2(t, x_n) \end{pmatrix}$$

³Note that the derivation of a Carleman estimate in the case of singular domains can be achieved (see [4]). Addressing the more general case of Lipschitz boundary is an open question to our knowledge.

where the functions c_i , $i = 1, 2$, are⁴ in $\mathcal{C}^1((0, T) \times \overline{\Omega_{\pm}})$ with a possible jump at $x_n = 0$. We assume $C_1 \in W^{1,\infty}(\Omega', M_{n-1}(\mathbb{R}))$ and that $C_1(x')$ is hermitian. We further assume uniform ellipticity

$$\begin{aligned} 0 < c_{\min} \leq c_i(t, x_n) \leq c_{\max} < \infty, \quad (t, x_n) \in (0, T) \times (-H, H) \text{ and } i = 1, 2, \\ 0 < c_{\min} \text{Id}_{n-1} \leq C_1(x') \leq c_{\max} \text{Id}_{n-1}, \quad x' \in \Omega'. \end{aligned}$$

To lighten notation we shall often write $c_{i-} := c_{i|x_n=0^+}$ and $c_{i+} := c_{i|x_n=0^-}$ for $i = 1, 2$.

Remark 1.3. Here, the matrix coefficient B is chosen time dependent in preparation for the Carleman estimate in the parabolic case. We shall also prove such an estimate in the elliptic case: see Theorem 1.4 below and its proof in Section 3. For this theorem we shall *of course* use B independent of time.

For the proof of Theorem 1.4 we shall further assume $c_1 = c_2$. In fact, this simplification allows us to provide a fairly simple proof of the Carleman estimates that shows the different treatment of two frequency regimes. These frequency regimes are connected to the microlocal regions used in [27] and [25, 26]. Note however that the case $c_1 \neq c_2$ can also be treated in the elliptic case. The proof is then closer to that of the parabolic case of Theorem 1.5 in Section 4. We have omitted this proof for the sake of the clarity of the exposition.

Let $T > 0$. For each $t \in [0, T]$, we consider the symmetric bilinear H_0^1 -coercive form

$$a_t(u, v) = \int_{\Omega} (B(t, \cdot) \nabla_x u) \cdot \nabla_x v dx,$$

with domain $D(a_t) = H_0^1(\Omega)$. It defines a selfadjoint operator $A_t = -\nabla_x \cdot (B(t, \cdot) \nabla_x \cdot)$ in $L^2(\Omega)$ with compact resolvent and with domain $D(A_t) = \{u \in H_0^1(\Omega); \nabla_x \cdot (B(t, \cdot) \nabla_x u) \in L^2(\Omega)\}$ (see e.g. [11], p. 1211). In the elliptic case, we shall denote by $\|\cdot\|_{L^2(\Omega)}$ the L^2 norm over Ω and by $|\cdot|_{L^2(S)}$ the L^2 norm over the interface S of codimension 1.

We set $Q_T = (0, T) \times \Omega$, $S_T = (0, T) \times S$. We shall also consider the following parabolic operator $P = \partial_t + A_t$ on Q_T . In the parabolic case, we shall denote by $\|\cdot\|_{L^2(Q_T)}$ the L^2 norm over Q_T and by $|\cdot|_{L^2(S_T)}$ the L^2 norm over the interface S_T of codimension 1.

In this article, when the constant C is used, it refers to a constant that is independent all the parameters. Its value may however change from one line to another. We shall use the notation $a \lesssim b$ if we have $a \leq Cb$ for such a constant. If we want to keep track of the value of a constant we shall use another letter.

1.2 Statements of the main results

We consider ω , a nonempty open subset of Ω . For a function β in $\mathcal{C}^0(\overline{\Omega})$ we set

$$\varphi(x) = e^{\lambda\beta(x)}, \quad \lambda > 0,$$

to be used as weight function. We consider first a matrix coefficient independent of the parameter t . A proper choice of the function β , with respect to the operator A , ω and Ω (see Assumption 2.1 and Assumption 3.2), yields the following Carleman estimate for the elliptic operator A .

⁴Concerning the regularity of the coefficients c_i , an inspection of the proof of the Carleman estimate in the parabolic case shows that the time derivative of the trace of c_2 at $x_n = 0$ needs to make sense (see above (A.9) in Appendix A.5). An alternative regularity is then $W^{1,\infty}(0, T; W^{1,\infty}(\Omega))$.

Theorem 1.4 (Elliptic case). *There exist $C > 0$, λ_0 and $s_0 > 0$ such that*

$$s\lambda^2 \|e^{s\varphi} \varphi^{\frac{1}{2}} \nabla u\|_{L^2(\Omega)}^2 + s^3 \lambda^4 \|e^{s\varphi} \varphi^{\frac{3}{2}} u\|_{L^2(\Omega)}^2 + s\lambda \left(\|e^{s\varphi} \varphi^{\frac{1}{2}} \nabla_\tau u|_S\|_{L^2(S)}^2 + \|e^{s\varphi} \varphi^{\frac{1}{2}} \partial_n u|_S\|_{L^2(S)}^2 \right) + s^3 \lambda^3 \|e^{s\varphi} \varphi^{\frac{3}{2}} u|_S\|_{L^2(S)}^2 \leq C \left(\|e^{s\varphi} Au\|_{L^2(\Omega)}^2 + s^3 \lambda^4 \|e^{s\varphi} \varphi^{\frac{3}{2}} u\|_{L^2(\omega)}^2 \right),$$

for all $u \in D(A)$, $\lambda \geq \lambda_0$, and $s \geq s_0$.

Here, ∇_τ is the tangential gradient on the interface S .

With a function $\tilde{\beta} > 0$ that satisfies Assumption 2.1 below, we introduce $\beta = \tilde{\beta} + m\|\tilde{\beta}\|_\infty$ where $m > 1$. For $\lambda > 0$ we define the following weight functions

$$\varphi(x) = e^{\lambda\beta(x)}, \quad \eta(x) = e^{\lambda\beta(x)} - e^{\lambda\tilde{\beta}}, \quad a(t) = (t(T-t))^{-1},$$

with $\bar{\beta} = 2m\|\tilde{\beta}\|_\infty$ (see [16, 12]). For $\tilde{\beta}$ satisfying some additional requirements, that will be provided in Section 4, we prove the following Carleman estimate for the parabolic operator P .

Theorem 1.5 (Parabolic case). *There exist $C > 0$, λ_0 and $s_0 > 0$ such that*

$$s^{-1} \left(\|e^{san}(a\varphi)^{-\frac{1}{2}} \partial_t u\|_{L^2(Q_T)}^2 + \|e^{san}(a\varphi)^{-\frac{1}{2}} A_t u\|_{L^2(Q_T)}^2 \right) + s\lambda^2 \|e^{san}(a\varphi)^{\frac{1}{2}} \nabla u\|_{L^2(Q_T)}^2 + s^3 \lambda^4 \|e^{san}(a\varphi)^{\frac{3}{2}} u\|_{L^2(Q_T)}^2 + s\lambda \left(\|e^{san}(a\varphi)^{\frac{1}{2}} \nabla_\tau u|_{S_T}\|_{L^2(S_T)}^2 + \|e^{san}(a\varphi)^{\frac{1}{2}} \partial_n u|_{S_T}\|_{L^2(S_T)}^2 \right) + s^3 \lambda^3 \|e^{san}(a\varphi)^{\frac{3}{2}} u|_{S_T}\|_{L^2(S_T)}^2 \leq C \left(\|e^{san} P u\|_{L^2(Q_T)}^2 + s^3 \lambda^4 \|e^{san}(a\varphi)^{\frac{3}{2}} u\|_{L^2((0,T) \times \omega)}^2 \right),$$

for all $u \in C^2((0, T) \times \Omega_\pm)$ such that $u|_{S_T^-} = u|_{S_T^+}$, $(c_2 \partial_{x_n} u)|_{S_T^-} = (c_2 \partial_{x_n} u)|_{S_T^+}$, $\lambda \geq \lambda_0$, and $s \geq s_0(T + T^2)$.

By density argument, we can extend this estimate to functions in $\int_{[0, T]}^\oplus D(A_t) dt \cap H^1(0, T; L^2(\Omega))$.

1.3 Outline

In Section 2, we provide some spectral properties of operator A , which yields a Hilbert direct decomposition of $L^2(\Omega) = \bigoplus_{k \in \mathbb{N}^*} H_k$ that reduces A . We also provide the precise assumptions made on the weight function. In Section 3, we prove the Carleman estimate for the elliptic case. In Section 4 we prove the Carleman estimate for a parabolic case. Some intermediate and technical results are collected in the appendices.

2 Spectral properties and weight function

Similarly to $A_t = -\nabla_x \cdot (B(t, x) \nabla_x)$, one can define the time independent selfadjoint transverse operator on $L^2(\Omega')$

$$A' = -\nabla_{x'} \cdot (C_1 \nabla_{x'}), \quad D(A') = \{u \in H_0^1(\Omega'); \nabla_{x'} \cdot (C_1 \nabla_{x'} u) \in L^2(\Omega')\}.$$

We consider an orthonormal basis of $L^2(\Omega')$, composed of eigenfunctions $(\phi_k)_{k \geq 1}$, associated with the eigenvalues, with finite multiplicities, $0 < \mu_1^2 \leq \mu_2^2 \leq \dots \leq \mu_k^2 \leq \mu_{k+1}^2 \leq \dots$, with $\mu_k \rightarrow \infty$.

With this basis $(\phi_k)_{k \geq 1}$, we build an unitary transform $\mathcal{F} : L^2(\Omega) \mapsto \bigoplus_{k=1}^{\infty} L^2(-H, H)$ defined by

$$(\mathcal{F}u)(k, x_n) := \int_{\Omega'} \phi_k(x') u(x', x_n) dx', \quad (2.1)$$

with the following properties (recall that here $\nabla_{x'} = \nabla_{\tau}$)

$$\begin{aligned} v(x', x_n) &= \sum_{k=1}^{\infty} (v(\cdot, x_n), \phi_k)_{L^2(\Omega')} \phi_k(x') = \sum_{k=1}^{\infty} \hat{v}(k, x_n) \phi_k(x'), \\ \nabla_{x'} v(x', x_n) &= \sum_{k=1}^{\infty} \hat{v}(k, x_n) \nabla_{x'} \phi_k(x'). \end{aligned}$$

We shall often write $\hat{v}_k = \hat{v}(k, \cdot)$.

As the family $(C_1^{1/2} \nabla \phi_k)_k$ is orthogonal in $L^2(\Omega')$ (C_1 is a positive definite matrix) we have

$$\|C_1^{1/2} \nabla_{x'} v(\cdot, x_n)\|_{L^2(\Omega')}^2 = \sum_{k=1}^{\infty} |\hat{v}(k, x_n)|^2 \|C_1^{1/2} \nabla_{x'} \phi_k\|_{L^2(\Omega')}^2 = \sum_{k=1}^{\infty} |\hat{v}(k, x_n)|^2 \mu_k^2,$$

which gives

$$(c_{\max})^{-1} \sum_{k=1}^{\infty} \mu_k^2 |\hat{v}(k, x_n)|^2 \leq \|\nabla_{\tau} v(\cdot, x_n)\|_{L^2(\Omega')}^2 \leq (c_{\min})^{-1} \sum_{k=1}^{\infty} \mu_k^2 |\hat{v}(k, x_n)|^2. \quad (2.2)$$

We choose a weight function β that satisfies the following properties.

Assumption 2.1. *The function $\beta \in \mathcal{C}^0(\Omega)$, and $\beta|_{\Omega_{\pm}} \in \mathcal{C}^2(\overline{\Omega_{\pm}})$ and*

$$\begin{aligned} |\nabla_x \beta| &\geq C > 0 \text{ in } \Omega \setminus \omega, \\ \beta &= \text{Cst on } \Omega' \times \{-H\} \quad \text{and} \quad \beta = \text{Cst on } \Omega' \times \{H\}, \\ \nabla_{x'} \beta &= 0 \text{ on } \partial\Omega' \times (-H, H), \\ \partial_{x_n} \beta &> 0 \text{ on } \Omega' \times \{-H\}, \quad \text{and} \quad \partial_{x_n} \beta < 0 \text{ on } \Omega' \times \{H\}. \end{aligned}$$

There exists a neighborhood V of S in Ω of the form $V = \Omega' \times (-\delta, \delta)$ in which β solely depends on x_n and is a piecewise affine function of x_n .

In particular $\beta|_S$ is constant. As the open set ω can be shrunk if necessary, we further assume that $\omega \cap (\Omega' \times (-\delta, \delta)) = \emptyset$.

Such a weight function β can be obtained by first designing a function that satisfies the proper properties at the boundaries and at the interface and then construct β by means of Morse functions following the method introduced in [16].

Here, in addition we assume that $\beta' > 0$ on S_+ and S_- , which means that the observation region ω is chosen in $\Omega' \times (0, H)$, i.e., where $x_n \geq 0$. There is no loss in generality as we can change x_n into $-x_n$ to treat the case of an observation $\omega \subset \Omega' \times (-H, 0)$.

Note that Assumption 2.1 will be complete below by Assumption 3.2 in the elliptic case and Assumption 4.2 in the parabolic case respectively.

3 The Elliptic case: proof of Theorem 1.4

As mentioned in the introductory section, we have consider only the case $c_1 = c_2 = c$ in this proof. The case $c_1 \neq c_2$ can be treated following the lines of the proof of Theorem 1.5 in Section 4.

Local Carleman estimates can be stitched together to form a global estimate of the form presented in Theorem 1.4 (see e.g. [24, 28]). Such local estimates are classical away from the interface (see [29, 16, 24]). To prove the elliptic Carleman estimate of Theorem 1.4 it thus remains to prove such a local estimate at the interface S , for functions $u \in D(A)$ with support near the interface. We shall thus assume that $\text{supp}(u) \subset \Omega' \times (-\delta, \delta)$, where the weight function β depends only on x_n and is piecewise affine.

Piecewise smooth functions that satisfy the transmission conditions

$$u|_{S_-} = u|_{S_+}, \quad (c\partial_{x_n}u)|_{S_-} = (c\partial_{x_n}u)|_{S_+}, \quad (3.1)$$

are dense in $D(A)$. We may thus restrict our analysis to such functions. Because of these transmission conditions we shall write $u|_S$ and $(c\partial_{x_n}u)|_S$ in place of $u|_{S_\pm}$ and $(c\partial_{x_n}u)|_{S_\pm}$ respectively.

Applying the unitary transform of Section 2, the equation $Au = f$ can be written

$$(-\partial_{x_n}c\partial_{x_n} + c\mu_k^2)\hat{u}_k(x_n) = \hat{f}_k(x_n), \quad x_n \in (-\delta, 0) \cup (0, \delta),$$

with $\text{supp } \hat{u}_k \subset (-\delta, \delta)$.

Our starting point is the following proposition.

Proposition 3.1. *Let $\hat{v}_k = e^{s\varphi}\hat{u}_k$. There exist $C, C', C'' > 0$, $\lambda_0 > 0$, $s_0 > 0$ such that*

$$\begin{aligned} & C(s\lambda^2\|\varphi^{\frac{1}{2}}\partial_{x_n}\hat{v}_k\|_{L^2(-\delta,\delta)}^2 + s\lambda^2\|\varphi^{\frac{1}{2}}\mu_k\hat{v}_k\|_{L^2(-\delta,\delta)}^2 + s^3\lambda^4\|\varphi^{\frac{3}{2}}\hat{v}_k\|_{L^2(-\delta,\delta)}^2) \\ & + s\lambda\varphi|_S \left([c^2\beta'|\partial_{x_n}\hat{v}_k|^2]_S + |s\lambda\varphi\hat{v}_k|_S^2 [c^2\beta'^3]_S - |\mu_k\hat{v}_k|_S^2 [c^2\beta']_S \right) + Z \leq C' \|e^{s\varphi}\hat{f}_k\|_{L^2(-\delta,\delta)}^2, \end{aligned} \quad (3.2)$$

for all $k \in \mathbb{N}^*$, $\lambda \geq \lambda_0$ and $s \geq s_0$, with $Z = C''s\lambda^2\varphi|_S \text{Re}[c^2\beta'^2\partial_{x_n}\hat{v}_k]_S \overline{\hat{v}_k}|_S$.

Such a result can be obtained by adapting the derivations in [12] for instance. We provide a short proof in Appendix A.1. In particular we have

$$Z \leq Cs\lambda^2\varphi|_S \left(|\partial_{x_n}\hat{v}_k|_{S_-}| + |\partial_{x_n}\hat{v}_k|_{S_+}| \right) |\hat{v}_k|_S. \quad (3.3)$$

Moreover, in addition to Assumption 2.1, we shall consider the following particular form of β

Assumption 3.2. *For $K = \frac{c_-}{c_+}$ and some $r \geq 0$, we have*

$$L = \frac{\beta'_{|S_+}}{\beta'_{|S_-}} = \begin{cases} 2 & \text{if } K = 1, \\ K & \text{if } K > 1, \\ (r+1) - rK & \text{if } K < 1. \end{cases} \quad (3.4)$$

Remark 3.3. With this assumption we note that we have $L > 1$ and $L \rightarrow 1$ as $K \rightarrow 1$, $K \neq 1$. Here we choose $L = 2$ if $K = 1$, to preserve interface terms in the Carleman estimates even for this case that corresponds to coefficients with no jump.

The value $r = 3$ is admissible in (3.4) (see Lemma 3.6 and its proof).

We now set $\mathcal{B}(v) = s\lambda\varphi|_S \left([c^2\beta'|\partial_{x_n}v|^2]_S + |s\lambda\varphi v|_S^2 [c^2\beta'^3]_S \right)$.

Lemma 3.4. *We have*

$$\mathcal{B}(\hat{v}_k) = s\lambda\varphi|_S e^{2s\varphi|_S} \left(B_1 |\gamma(\hat{u}_k)|^2 + B_2 |s\lambda\varphi\hat{u}_k|_S^2 \right), \quad \gamma(\hat{u}_k) = c\partial_{x_n}\hat{u}_k|_S + c_+\beta'_{|S_-} \frac{L^2 - K}{L - 1} (s\lambda\varphi\hat{u}_k)|_S,$$

with $B_1 = \beta'_{|S_-} (L - 1) > 0$, and

$$B_2 = c_+^2 (\beta'_{|S_-})^3 \left(2(L^3 - K^2) - \frac{(L^2 - K)^2}{L - 1} \right). \quad (3.5)$$

For a proof see Appendix A.2. Note that $L > 1$ by Assumption 3.2.

Assume that $K > 1$. Then $L = K$ and

$$-[c^2\beta']_S = -c_+^2\beta'_{|S_-} (L - K^2) > 0, \quad B_1 > 0, \quad B_2 = c_+^2 (\beta'_{|S_-})^3 K^2 (K - 1) > 0.$$

The trace terms in (3.2) thus yield a positive contribution. We have

$$\begin{aligned} \mathcal{B}(\hat{v}_k) - s\lambda\varphi|_S |\mu_k \hat{v}_k|_S^2 [c^2\beta']_S &\geq (s\lambda\varphi|_S)^3 e^{2s\varphi} |\hat{u}_k|_S^2 + s\lambda\varphi|_S e^{2s\varphi} \left(|\gamma(\hat{u}_k)|^2 + |\mu_k \hat{u}_k|_S^2 \right) \\ &\geq (s\lambda\varphi|_S)^3 e^{2s\varphi} |\hat{u}_k|_S^2 + s\lambda\varphi|_S e^{2s\varphi} \left(|\partial_{x_n}\hat{u}_k|_S^2 + |\mu_k \hat{u}_k|_S^2 \right). \end{aligned}$$

In particular for s sufficiently large the remainder term Z estimated in (3.3) can be 'absorbed'. We thus obtain

$$\begin{aligned} s\lambda^2 \|\varphi^{\frac{1}{2}} \partial_{x_n} \hat{v}_k\|_{L^2(-\delta, \delta)}^2 + s\lambda^2 \|\varphi^{\frac{1}{2}} \mu_k \hat{v}_k\|_{L^2(-\delta, \delta)}^2 + s^3 \lambda^4 \|\varphi^{\frac{3}{2}} \hat{v}_k\|_{L^2(-\delta, \delta)}^2 \\ + (s\lambda\varphi|_S)^3 e^{2s\varphi} |\hat{u}_k|_S^2 + s\lambda\varphi|_S e^{2s\varphi} \left(|\partial_{x_n}\hat{u}_k|_S^2 + |\mu_k \hat{u}_k|_S^2 \right) \lesssim \|e^{s\varphi} \hat{f}_k\|_{L^2(-\delta, \delta)}^2, \end{aligned} \quad (3.6)$$

for all $k \in \mathbb{N}^*$. Summing over k , using (2.2) we obtain the sought local Carleman estimate in the case $K > 1$

$$\begin{aligned} s\lambda^2 \|\varphi^{\frac{1}{2}} \nabla v\|_{L^2(\Omega' \times (-\delta, \delta))}^2 + s^3 \lambda^4 \|\varphi^{\frac{3}{2}} v\|_{L^2(\Omega' \times (-\delta, \delta))}^2 \\ + (s\lambda\varphi|_S)^3 e^{2s\varphi|_S} |u|_S^2_{L^2(S)} + s\lambda\varphi|_S e^{2s\varphi|_S} |\nabla u|_S^2_{L^2(S)} \lesssim \|e^{s\varphi} f\|_{L^2(\Omega' \times (-\delta, \delta))}^2. \end{aligned} \quad (3.7)$$

The Carleman estimate of Theorem 1.4 can then be deduced classically. This case, $K > 1$ is the case originally covered by [12].

We shall now consider the case $0 < K \leq 1$. Then, either $L = (r + 1) - rK > 1$ or $L = 2$, which gives $B_1 > 0$. The result of Lemma 3.6 below implies that $B_2 > 0$. Hence, for s sufficiently large the remainder term Z estimated in (3.3) can be 'absorbed'. We now aim to estimate the tangential term in (3.2).

Proposition 3.5. *There exists $C > 0$, and $\varepsilon > 0$ such that for all $k \in \mathbb{N}$ we have*

$$\begin{aligned} \left| s\lambda\varphi|_S [c^2\beta']_S |\mu_k \hat{v}_k|_S^2 \right| &\leq \frac{1}{1 + \varepsilon} s\lambda\varphi|_S B_2 |s\lambda(\varphi \hat{v}_k)|_S^2 \\ &\quad + C \left(\|e^{s\varphi} \hat{f}_k\|_{L^2(-\delta, \delta)}^2 + s^2 \lambda^2 \|\hat{v}_k\|_{L^2(-\delta, \delta)}^2 + \|\partial_{x_n} \hat{v}_k\|_{L^2(-\delta, \delta)}^2 \right). \end{aligned} \quad (3.8)$$

Proof. Let $0 < \varepsilon < 1$. The value of ε will be determined below. We treat low and high values of μ_k differently.

Low frequencies. Set k_1 as the largest integer such that $(1 + \varepsilon)[c^2\beta']_S|\mu_k^2 < B_2(s\lambda\varphi)_S^2$, that is

$$(1 + \varepsilon)\mu_k^2 < (\beta'_{|S_-})^2(s\lambda\varphi)_S^2 \frac{1}{|L - K^2|} \left(2(L^3 - K^2) - \frac{(L^2 - K)^2}{L - 1} \right). \quad (3.9)$$

We then have

$$(1 + \varepsilon)s\lambda\varphi_S|[c^2\beta']_S|\mu_k\hat{v}_k|_S|^2 < s\lambda\varphi_S B_2|s\lambda(\varphi\hat{v}_k)_S|^2, \quad k \leq k_1. \quad (3.10)$$

High frequencies. Here we consider frequencies μ_k that satisfy

$$(1 - \varepsilon)\mu_k \geq s|\partial_{x_n}\varphi|_{S_-} = s\lambda\varphi_S\beta'_{|S_-}. \quad (3.11)$$

We denote by k_2 the smallest integer that satisfies (3.11).

We write

$$(\partial_{x_n}^2 - \mu_k^2)\hat{u}_k = -\frac{\hat{f}_k}{c} - \frac{\partial_{x_n}c}{c}\partial_{x_n}\hat{u}_k = -\hat{g}_k.$$

As $\hat{u}_k(-\delta) = \hat{u}_k(\delta) = 0$, with the transmission conditions (3.1), the computations⁵ of Appendix A.3 yield

$$\mu_k\hat{u}_k|_{x_n=0^+} = \frac{1}{(c_+ + c_-)} \int_0^\delta \frac{\sinh(\mu_k(\delta - x_n))}{\cosh(\mu_k\delta)} (c_+\hat{g}_k(x_n) + c_-\hat{g}_k(-x_n)) dx_n. \quad (3.12)$$

We have

$$\frac{\sinh(\mu_k(\delta - x_n))}{\cosh(\mu_k\delta)} = \frac{e^{\mu_k(\delta - x_n)} - e^{-\mu_k(\delta - x_n)}}{e^{\mu_k\delta} + e^{-\mu_k\delta}} \leq e^{-\mu_k x_n}. \quad (3.13)$$

We note that

$$\varphi(0) - \varphi(-x_n) = x_n \int_0^1 \varphi'(-x_n + \sigma x_n) d\sigma = x_n \lambda \beta'_{|S_-} \int_0^1 \varphi(-x_n + \sigma x_n) d\sigma,$$

as the weight function $\beta = \beta(x_n)$ is affine in $(-\delta, 0)$. Since $\beta' > 0$, the function φ increases with x_n and we have $\varphi(0) \leq \varphi(-x_n) + x_n \lambda \varphi(0) \beta'_{|S_-}$, if $x_n > 0$. As we have assumed (3.11) here we obtain

$$s\varphi(0) - \mu_k x_n \leq s\varphi(-x_n) - \varepsilon \mu_k x_n, \quad x_n > 0. \quad (3.14)$$

We also have

$$s\varphi(0) - \mu_k x_n \leq s\varphi(x_n) - \varepsilon \mu_k x_n, \quad x_n > 0. \quad (3.15)$$

From (3.12) we thus obtain

$$\begin{aligned} \mu_k^{\frac{3}{2}} e^{s\varphi|S} |\hat{u}_k|_S &\leq \frac{1}{(c_+ + c_-)} \int_0^\delta \left(e^{s\varphi(-x_n)} |c_-\hat{g}_k(-x_n)| + e^{s\varphi(x_n)} |c_+\hat{g}_k(x_n)| \right) \mu_k^{\frac{1}{2}} e^{-\varepsilon \mu_k x_n} dx_n \\ &\leq \left(\|e^{s\varphi} \hat{g}_k\|_{L^2(-\delta, 0)} + \|e^{s\varphi} \hat{g}_k\|_{L^2(0, \delta)} \right) \left(\int_0^\delta \mu_k e^{-2\varepsilon \mu_k x_n} dx_n \right)^{\frac{1}{2}} \\ &\leq \varepsilon^{-\frac{1}{2}} \|e^{s\varphi} \hat{g}_k\|_{L^2(-\delta, \delta)} \lesssim \varepsilon^{-\frac{1}{2}} \left(\|e^{s\varphi} \hat{f}_k\|_{L^2(-\delta, \delta)} + s\lambda \|\hat{v}_k\|_{L^2(-\delta, \delta)} + \|\partial_{x_n} \hat{v}_k\|_{L^2(-\delta, \delta)} \right), \end{aligned}$$

⁵This is the precise point where $c_1 = c_2$ is used. In the case $c_1 \neq c_2$ the result of Appendix A.3 cannot be used and we have to proceed as in Section 4.

which leads to, for $k_2 \leq k$,

$$\begin{aligned} & s\lambda\varphi|_S |[c^2\beta']_S |\mu_k \hat{v}_k|_S|^2 \\ & \lesssim (1-\varepsilon)|\beta'_-|^{-1} |[c^2\beta']_S |\mu_k^3 \hat{v}_k|_S|^2 \\ & \lesssim (1-\varepsilon)\varepsilon^{-1} |\beta'_-|^{-1} |[c^2\beta']_S |(\|e^{s\varphi} \hat{f}_k\|_{L^2(-\delta,\delta)}^2 + s^2\lambda^2 \|\hat{v}_k\|_{L^2(-\delta,\delta)}^2 + \|\partial_{x_n} \hat{v}_k\|_{L^2(-\delta,\delta)}^2). \end{aligned}$$

We have thus seen that low frequencies in (3.8) are estimated by boundary terms and high frequencies are estimated by the r.h.s. of (3.2) and "absorbable" terms. It remains to prove that we cover the whole spectrum with the two estimates we have obtained. A sufficient condition is then

$$(1-\varepsilon)^{-2}(s\lambda\varphi|_S)^2(\beta'_{|S_-})^2 \leq \frac{1}{1+\varepsilon}(\beta'_{|S_-})^2(s\lambda\varphi|_S)^2 \frac{1}{|L-K^2|} \left(2(L^3-K^2) - \frac{(L^2-K)^2}{L-1}\right),$$

that is

$$P(K, L) := -|L-K^2|(L-1) + \frac{(1-\varepsilon)^2}{1+\varepsilon} (2(L^3-K^2)(L-1) - (L^2-K)^2) \geq 0. \quad (3.16)$$

We recall that $L = (r+1) - rK$ if $0 < K < 1$. The following lemma provides a positive answer (see Appendix 3.6 for a proof).

Lemma 3.6. *There exists $\varepsilon_0 > 0$ such that for $0 < \varepsilon < \varepsilon_0$,*

- $P(K, L) \geq 0$ if $K = 1$,
- *there exists $r \geq 1$ such that $P(K, L) \geq 0$ for $K \in (0, 1)$. In particular the value $r = 3$ is admissible.*

This concludes the proof of Proposition 3.5. ■

Arguing as we did for (3.7) in the case $K > 1$, we now obtain

$$\begin{aligned} & s\lambda^2 \|\varphi^{\frac{1}{2}} \nabla v\|_{L^2(\Omega' \times (-\delta, \delta))}^2 + s^3 \lambda^4 \|\varphi^{\frac{3}{2}} v\|_{L^2(\Omega' \times (-\delta, \delta))}^2 + (s\lambda\varphi|_S)^3 e^{2s\varphi|_S} |u|_S|_{L^2(S)}^2 \\ & + s\lambda\varphi|_S e^{2s\varphi|_S} |\nabla u|_S|_{L^2(S)}^2 \lesssim \|e^{s\varphi} f\|_{L^2(\Omega' \times (-\delta, \delta))}^2 + s^2 \lambda^2 \|v\|_{L^2(\Omega' \times (-\delta, \delta))}^2 + \|\partial_{x_n} v\|_{L^2(\Omega' \times (-\delta, \delta))}^2. \end{aligned} \quad (3.17)$$

The last two terms on the r.h.s. can be "absorbed" by the l.h.s. by choosing s sufficiently large. This concludes the proof of Theorem 1.4.

Remark 3.7. Observe that the local Carleman estimate that we obtain in $\Omega' \times (-\delta, \delta)$ does not require any regularity for the boundary of the open set Ω' . Here, the weight function does not depend on x' . The minimal regularity of the boundary to achieve a Carleman estimate remains an open question to our knowledge.

4 The parabolic case: proof of Theorem 1.5

Here, the matrix coefficient B is assumed to be time dependent as stated in Assumption 1.2. The coefficients $c_1(t, x_n)$ and $c_2(t, x_n)$ can be different.

We choose a function $\tilde{\beta} > 0$ that satisfies the requirements of Assumption 2.1 and we introduce $\beta = \tilde{\beta} + m\|\tilde{\beta}\|_\infty$ where $m > 1$. Observe that β also satisfies Assumption 2.1.

For $T > 0$ and $\lambda > 0$ we define the following weight functions

$$\varphi(x) = e^{\lambda\beta(x)}, \quad \eta(x) = e^{\lambda\beta(x)} - e^{\lambda\bar{\beta}}, \quad x \in \Omega, \quad a(t) = (t(T-t))^{-1}, \quad t \in (0, T), \quad (4.1)$$

with $\bar{\beta} = 2m\|\tilde{\beta}\|_\infty$ (see [12]). As in the previous sections we choose $\beta' > 0$ on S_+ and S_- , which means that the observation region ω is chosen in $\Omega' \times (0, H)$, i.e., where $x_n \geq 0$. It suffices to prove a local Carleman estimate at the interface S , i.e., for functions u with support near the interface, $\text{supp}(u) \subset [0, T] \times \Omega' \times (-\delta, \delta)$, where the weight function β depends only on x_n and is piecewise affine.

We assume moreover that u satisfies the transmission conditions

$$u|_{S_T^-} = u|_{S_T^+}, \quad (c_2\partial_{x_n}u)|_{S_T^-} = (c_2\partial_{x_n}u)|_{S_T^+}. \quad (4.2)$$

Applying the unitary transform of Section 2, the equation $\partial_t u + Au = f$ can be written

$$(\partial_t - \partial_{x_n}c_2\partial_{x_n} + c_1\mu_k^2)\hat{u}_k(t, x_n) = \hat{f}_k(t, x_n), \quad t \in (0, T), \quad x_n \in (-\delta, 0) \cup (0, \delta), \quad k \geq 1,$$

with $\text{supp}(\hat{u}_k) \subset [0, T] \times (-\delta, \delta)$. Setting $q_{T,\delta} = (0, T) \times (-\delta, \delta)$, our starting point is the following proposition.

Proposition 4.1. *Let $T > 0$. Let $\hat{v}_k = e^{s\alpha\eta}\hat{u}_k$. There exist $C, C' > 0$, $\lambda_0 > 0$, $s_0 > 0$ such that*

$$\begin{aligned} & C\left(s^{-1}\|(a\varphi)^{-\frac{1}{2}}\partial_t\hat{v}_k\|_{L^2(q_{T,\delta})}^2 + s\lambda^2\|(a\varphi)^{\frac{1}{2}}\partial_{x_n}\hat{v}_k\|_{L^2(q_{T,\delta})}^2 + s\lambda^2\|(a\varphi)^{\frac{1}{2}}\mu_k\hat{v}_k\|_{L^2(q_{T,\delta})}^2\right. \\ & \quad \left. + s^3\lambda^4\|(a\varphi)^{\frac{3}{2}}\hat{v}_k\|_{L^2(q_{T,\delta})}^2\right) + s\lambda\int_0^T a\varphi|_S \left([c_2^2\beta'|\partial_{x_n}\hat{v}_k|^2]_S + |s\lambda a(\varphi\hat{v}_k)|_S|^2[c_2^2\beta'^3]_S\right) dt \\ & \leq C'\|e^{s\alpha\eta}\hat{f}_k\|_{L^2(q_{T,\delta})}^2 + s\lambda\int_0^T a\varphi|_S |\mu_k\hat{v}_k|_{S_T}^2 [c_1c_2\beta']_S dt + Z, \end{aligned} \quad (4.3)$$

for all $k \in \mathbb{N}^*$, $\lambda \geq \lambda_0$ and $s \geq s_0(T + T^2)$, with

$$|Z| \lesssim s^{\frac{1}{2}}\lambda T \int_0^T a\varphi|_S (|\partial_{x_n}\hat{v}_k|_{S_T^-}^2 + |\partial_{x_n}\hat{v}_k|_{S_T^+}^2) dt + \left(s(T^3 + T^4)\lambda + s^{\frac{3}{2}}T^3\lambda^3\right) \int_0^T a^3\varphi|_S |\hat{v}_k|_{S_T}^2 dt. \quad (4.4)$$

Such a result can be obtained by adapting the derivations in [12] for instance. We provide a short proof in Appendix A.5.

As in Section 3, we set

$$\begin{aligned} \mathcal{B}_p(\hat{v}_k) &= s\lambda a\varphi|_S \left([c_2^2\beta'|\partial_{x_n}\hat{v}_k|^2]_S + |s\lambda a\varphi\hat{v}_k|_{S_T}^2 [c_2^2\beta'^3]_S\right), \\ L &= \frac{\beta'_{|S_+}}{\beta'_{|S_-}}, \quad K_i(t) = \frac{c_{i-}(t)}{c_{i+}(t)}, \quad \underline{K}_i = \inf_{t \in [0, T]} K_i(t), \quad \overline{K}_i = \sup_{t \in [0, T]} K_i(t), \quad i = 1, 2. \end{aligned} \quad (4.5)$$

and

$$\mathbf{B} = \mathbf{B}(L) = \inf_{t \in [0, T]} (c_{2+}^2(t)) (\beta'_{|S_-})^3 \frac{K_2^2 + L^3(L - \underline{L})}{L - 1}, \quad \text{with } \underline{L} = \max\{\bar{K}_2, 2\}, \quad (4.6)$$

and finally

$$\mathbf{D} = \mathbf{D}(L) = \sup_{t \in [0, T]} (c_{1+}c_{2+})(t) \beta'_{|S_-} (L + \bar{K}_1 \bar{K}_2) > 0. \quad (4.7)$$

We make the following assumption on the weight function in addition to Assumption 2.1.

Assumption 4.2. *The weight function β is chosen such that $L \geq \underline{L} = \max\{\bar{K}_2, 2\}$ and*

$$\frac{1}{2} \geq \max \left\{ 2 \sqrt{\frac{\mathbf{D}}{\mathbf{B}}}, \frac{4\beta'_{|S_-}}{\sigma} \sqrt{\frac{\mathbf{D}}{\mathbf{B}}} \right\}, \quad \sigma = \left(\inf_{t, x_n} \frac{c_1(t, x_n)}{c_2(t, x_n)} \right)^{\frac{1}{2}}, \quad (4.8)$$

The coefficients c_1, c_2 being fixed, the forms of the coefficients \mathbf{D} and \mathbf{B} show that this can be achieved by first choosing the value of $\beta'_{|S_-} > 0$ and then picking a sufficiently large value for L .

Lemma 4.3. *We have*

$$\mathcal{B}_p(\hat{v}_k) = s\lambda a\varphi_{|S} e^{2s\alpha\varphi_{|S}} \left(B_1 |\gamma(\hat{u}_k)|^2 + B_2 |s\lambda a(\varphi \hat{u}_k)_{|S}|^2 \right),$$

with $\gamma(\hat{u}_k) = (c_2 \partial_{x_n} \hat{u}_k)_{|S} + c_{2+} \beta'_{|S_-} \frac{L^2 - K_2}{L-1} (s\lambda a \varphi \hat{u}_k)_{|S}$ and where

$$B_1 = \beta'_{|S_-} (L - 1), \quad B_2(t) = c_{2+}^2(t) (\beta'_{|S_-})^3 \left(2(L^3 - K_2^2(t)) - \frac{(L^2 - K_2(t))^2}{L - 1} \right).$$

If β satisfies Assumption 4.2 we have $B_1 > 0$ and $B_2(t) \geq \mathbf{B}$, with \mathbf{B} defined in (4.6).

Proof. The proof of Lemma 3.4 in Appendix A.2 can be directly adapted and gives the first part of the lemma. As $L \geq 1$ we have $B_1 > 0$. A direct computation yields $B_2(t) = c_{2+}^2(t) (\beta'_{|S_-})^3 \frac{P_p(L, K_2(t))}{L-1}$ with

$$\begin{aligned} P_p(L, Y) &= Y^2(1 - 2L) + 2YL^2 + L^4 - 2L^3 \\ &= L^3(L - \underline{L}) + L^3(\underline{L} - 2) + 2LY(L - Y) + Y^2. \end{aligned}$$

As $\underline{L} \geq 2$, and $L \geq \bar{K}_2 \geq K_2(t) \geq \underline{K}_2 > 0$, we thus obtain $P_p(L, K_2(t)) \geq \underline{K}_2^2 + L^3(L - \underline{L})$. \blacksquare

We now prove the following key result, providing an estimate of the tangential derivative of v , i.e., $\mu_k \hat{v}_k$, in the Fourier decomposition.

Proposition 4.4. *For a weight function β that satisfies Assumptions 2.1 and 4.2 there exists $C > 0$ such that for all $k \in \mathbb{N}^*$ we have*

$$\begin{aligned} s\lambda \int_0^T a\varphi_{|S} |c_1 c_2 \beta'|_{|S} |\mu_k \hat{v}_k|_{|S}^2 dt &\leq \frac{\mathbf{B}}{4} (s\lambda)^3 (a\varphi_{|S})^{\frac{3}{2}} \hat{v}_k|_{|S}^2 \Big|_{L^2((0, T))} + C \left(\|e^{s\alpha n} \hat{f}_k\|_{L^2(q_{T, \delta})}^2 \right. \\ &\quad \left. + s^3 \lambda^3 \|(a\varphi)^{\frac{3}{2}} \hat{v}_k\|_{L^2(q_{T, \delta})}^2 + s\lambda \|(a\varphi)^{\frac{1}{2}} \partial_{x_n} \hat{v}_k\|_{L^2(q_{T, \delta})}^2 + s\lambda \|(a\varphi)^{\frac{1}{2}} \mu_k \hat{v}_k\|_{L^2(q_{T, \delta})}^2 \right). \end{aligned} \quad (4.9)$$

for λ and $s/(T + T^2)$ both sufficiently large.




Figure 1: The function Φ . The shaded region is treated in the first step of the proof.

Proof. We fix $k \geq 1$ and we shall keep track of the dependency of the constants on k .

We have

$$|[c_1 c_2 \beta']_S| \leq (c_1 + c_2)(t) \beta'_{|S_-}(L + K_1 K_2(t)) \leq D,$$

with D as defined in (4.7). We set

$$\Phi(t; s, \lambda) := \frac{1}{2} s \lambda a(t) \varphi_{|S} \sqrt{\frac{B}{D}} \quad \text{and} \quad \mu_{s, \lambda} := \Phi\left(\frac{T}{2}; s, \lambda\right) = \min_{t \in (0, T)} \Phi(t; s, \lambda). \quad (4.10)$$

If $\mu_k > \mu_{s, \lambda}$, there exists $t_k := t_k(s, \lambda) \in (0, T/2)$ such that

$$\mu_k = \Phi(t_k; s, \lambda) = \Phi(T - t_k; s, \lambda). \quad (4.11)$$

For $\mu_k \geq \mu_{s, \lambda}$, we set

$$I_k := (0, t_k) \cup (T - t_k, T), \quad J_k := (0, T) \setminus I_k = (t_k, T - t_k), \quad \tilde{J}_k := \left(\frac{t_k}{2}, T - \frac{t_k}{2}\right).$$

For $\mu_k < \mu_{s, \lambda}$, we set

$$I_k := (0, T).$$

We then introduce

$$I(k; s, \lambda) := s \lambda D \int_{I_k} a(t) \varphi_{|S} \mu_k^2 |\hat{v}_{k|S}|^2 dt, \quad (4.12)$$

$$J(k; s, \lambda) := s \lambda D \int_{J_k} a(t) \varphi_{|S} \mu_k^2 |\hat{v}_{k|S}|^2 dt, \quad (4.13)$$

so that the term on the l.h.s. of (4.9) is less than the sum of the two previous quantities.

The first term, $I(k; s, \lambda)$, involving time t close to 0 or T , will be estimated by a trace term. The second term, $J(k; s, \lambda)$, involving time t away from 0 and T , will be estimated by volume terms.

Step 1: $\mu_k \leq \mu_{s,\lambda}$ or $t \in I_k$ In the (t, μ_k) plane presented in Figure 1 this corresponds to the shaded region. We thus treat low (tangential) frequencies here.

Lemma 4.5. *For all $k \geq 1$ we have*

$$D|\mu_k \hat{v}_k|_S \leq \frac{B}{4} |s\lambda a(t)(\varphi \hat{v}_k)|_S^2,$$

with B as defined in (4.6), if either (1) $\mu_k \leq \mu_{s,\lambda}$ or (2) $\mu_k > \mu_{s,\lambda}$ and $t \in I_k$.

Proof. The first point follows from the definition of $\mu_{s,\lambda}$. The second point is a direct consequence of the definition of t_k in (4.11) as the function $t \mapsto a(t)$ decreases on $(0, T/2)$. ■

For all $k \in \mathbb{N}^*$, we thus obtain

$$I(k; s, \lambda) \leq \frac{B}{4} (s\lambda)^3 |(a\varphi)_S|^{\frac{3}{2}} \hat{v}_k|_S^2_{L^2((0,T))}.$$

Step 2: $\mu_k > \mu_{s,\lambda}$ and t in a neighborhood of J_k , preliminary result. In each open set $(0, T) \times (-\delta, 0)$ and $(0, T) \times (0, \delta)$, the function \hat{u}_k satisfies the following equation

$$-\partial_{x_n}^2 \hat{u}_k + \frac{c_1}{c_2} \mu_k^2 \hat{u}_k + \frac{1}{c_2} \partial_t \hat{u}_k = \frac{\hat{f}_k}{c_2} + \frac{\partial_{x_n} c_2}{c_2} \partial_{x_n} \hat{u}_k. \quad (4.14)$$

Because of the form of (4.13) we set

$$p(t; s, \lambda) := s\lambda D a(t) \varphi|_S e^{2sa(t)\eta s}. \quad (4.15)$$

We consider a cutoff function $(0, T) \ni t \rightarrow \chi(t)$, such that

$$\chi \equiv 1 \text{ on } J_k, \quad 0 \leq \chi \leq 1, \quad \text{supp}(\chi) \subset \tilde{J}_k \quad \text{and} \quad \|\chi'\|_\infty \leq C/t_k,$$

and we introduce

$$w = w(t, k, x_n; s, \lambda) = \frac{1}{2} \chi(t) p(t; s, \lambda) |\hat{u}_k(t, x_n)|^2. \quad (4.16)$$

Observe that $w \geq 0$ and that it satisfies the same transmission conditions (4.2) as u . The function w satisfies

$$\partial_{x_n}^2 w - \frac{c_1}{c_2} \left((2 - \gamma) \mu_k^2 - \frac{p'}{c_1 p} \right) w = -g, \quad (4.17)$$

with $0 < \gamma < 1$ and

$$g = -\chi p |\partial_{x_n} \hat{u}_k|^2 - \frac{1}{c_2} \partial_t w + \chi \frac{1}{c_2} p \text{Re} \hat{f}_k \overline{\hat{u}_k} - \frac{c_1}{c_2} \mu_k^2 \frac{\gamma}{2} \chi p |\hat{u}_k|^2 + \chi p \frac{\partial_{x_n} c_2}{c_2} \text{Re} \hat{u}_k \partial_{x_n} \overline{\hat{u}_k} + \frac{\chi'}{2c_2} p |\hat{u}_k|^2.$$

Lemma 4.6. *There exist $s_0 > 0$, $\lambda_0 > 0$, depending on L and γ , such that*

$$(2 - \gamma) \mu_k^2 - \frac{p'}{c_1 p} \geq \mu_k^2 \text{ if } \frac{t_k}{2} < t < T - \frac{t_k}{2}, \quad x_n \in (-\delta, \delta), \quad (4.18)$$

for $s > s_0(T + T^2)$ and $\lambda > \lambda_0$.

See Appendix A.6 for a proof.

Step 3: $\mu_k > \mu_{s,\lambda}$ and t in a neighborhood of J_k , conclusion. For $t \in \tilde{J}_k$ we begin by replacing the time-space dependent coefficient $\frac{c_1}{c_2}((2-\gamma)\mu_k^2 - \frac{p'}{c_1 p})$ by $\sigma^2 \mu_k^2$ on the l.h.s. of (4.17) (the constant σ is introduced in (4.8)). This will allow us to argue as in the elliptic case, viz. solving an ordinary differential equation with constant coefficients.

We set

$$q(t, x_n; k, s, \lambda) := -\sigma^2 \mu_k^2 + \frac{c_1}{c_2} \left((2-\gamma)\mu_k^2 - \frac{p'}{c_1 p} \right).$$

We have

$$\partial_{x_n}^2 w - \sigma^2 \mu_k^2 w = -\tilde{g}, \quad (4.19)$$

with

$$\tilde{g} := -qw - \chi p |\partial_{x_n} \hat{u}_k|^2 - \frac{1}{c_2} \partial_t w + \chi \frac{1}{c_2} p \operatorname{Re} \hat{f}_k \overline{\hat{u}_k} - \frac{c_1}{c_2} \chi \frac{\gamma}{2} p \mu_k^2 |\hat{u}_k|^2 + \chi p \frac{\partial_{x_n} c_2}{c_2} \operatorname{Re} \hat{u}_k \partial_{x_n} \overline{\hat{u}_k} + \frac{\chi'}{2c_2} p |\hat{u}_k|^2.$$

Observe that Lemma 4.6 gives

$$q(t, x_n; k, s, \lambda) \geq 0, \quad \frac{t_k}{2} < t < T - \frac{t_k}{2}, \quad x_n \in (-\delta, \delta), \quad s > s_0, \quad \lambda > \lambda_0.$$

From (4.19) and Appendix A.3 we obtain

$$\begin{aligned} J(k; s, \lambda) &\leq 2 \int_{\tilde{J}_k} \mu_k^2 w|_S dt \\ &= \frac{2\mu_k}{\sigma} \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2+} + c_{2-}) \cosh(\sigma\mu_k\delta)} (c_{2+} \tilde{g}(t, x_n) + c_{2-} \tilde{g}(t, -x_n)) dx_n dt. \end{aligned} \quad (4.20)$$

Note that the introduction of \tilde{J}_k , instead of J_k , is due to the cut-off function χ . Substituting \tilde{g} in (4.20) we obtain seven terms. We shall provide the details for the contribution of $c_{2+} \tilde{g}(t, x_n)$. For the contribution $c_{2-} \tilde{g}(t, -x_n)$ details are given if difference occurs. As in the elliptic case, we shall use that the kernel $e^{-2sa(t)\eta s} \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)}$ be estimated by the weight $e^{2sa\eta}$.

1. We have

$$\int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)} (-qw - \chi p |\partial_{x_n} \hat{u}_k|^2) dx_n \leq 0. \quad (4.21)$$

The negative sign is fortunate as the absolute value of this term cannot be reasonably bounded, i.e., by a term that can be ‘‘absorbed’’ by the l.h.s. of (4.3).

2. (a) **Term $\frac{1}{c_2} \partial_t w$.** Because of the cut-off function χ we have $w|_{t=t_k/2} = w|_{t=T-t_k/2} = 0$ and we get

$$\begin{aligned} &\int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2+} + c_{2-}) \cosh(\sigma\mu_k\delta)} \left(-\frac{c_{2\pm}}{c_2} \partial_t w(t, \pm x_n) \right) dx_n dt \\ &= \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)} \partial_t \left(\frac{c_{2\pm}}{(c_{2+} + c_{2-})c_2} \right) w(t, \pm x_n) dx_n dt, \end{aligned}$$

and, by (3.13), we have

$$\begin{aligned} & \left| \mu_k \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2+} + c_{2-}) \cosh(\sigma\mu_k\delta)} \left(-\frac{c_{2\pm}}{c_2} \partial_t w(t, \pm x_n) \right) dx_n dt \right| \\ & \lesssim s\lambda D \int_{\tilde{J}_k} \int_0^\delta a(t) \varphi|_S e^{-\sigma\mu_k x_n} e^{2sa(t)\eta|_S} \mu_k |\hat{u}_k(t, \pm x_n)|^2 dx_n dt. \end{aligned}$$

We shall thus obtain an estimate of this term by the r.h.s. of (4.9) if we prove

$$-\sigma\mu_k x_n + 2sa(t)\eta|_S \leq 2sa(t)\eta(\pm x_n), \quad \forall (t, x_n) \in \tilde{J}_k \times (0, \delta). \quad (4.22)$$

This is clear for the case + since $\eta|_S \leq \eta(x_n)$.

The argument is different for the case -. Using that β is a piecewise affine, we have

$$\eta(-x_n) - \eta|_S = \varphi(-x_n) - \varphi|_S \geq -x_n \lambda (\beta'_- \varphi)|_S, \quad x_n \in (0, \delta).$$

Therefore, (4.22) will be satisfied if

$$\sigma\mu_k \geq 2sa(t)\lambda(\beta'_- \varphi)|_S, \quad \forall t \in \tilde{J}_k,$$

which, by the definition of Φ in (4.10), can be written as

$$\sigma\mu_k \geq 4\beta'_-|_S \sqrt{\frac{D}{B}} \Phi(t; s, \lambda), \quad \forall t \in \tilde{J}_k.$$

As $\max_{t \in \tilde{J}_k} \Phi(t; s, \lambda) = \Phi\left(\frac{t_k}{2}; s, \lambda\right)$, it suffices to have

$$\sigma\mu_k = \sigma\Phi(t_k; s, \lambda) \geq 4\beta'_-|_S \sqrt{\frac{D}{B}} \Phi\left(\frac{t_k}{2}; s, \lambda\right), \quad \forall t \in \tilde{J}_k.$$

This holds if we have

$$\frac{a(t_k)}{a\left(\frac{t_k}{2}\right)} \geq \frac{4\beta'_-|_S}{\sigma} \sqrt{\frac{D}{B}}. \quad (4.23)$$

Observing that $\frac{a(t_k)}{a\left(\frac{t_k}{2}\right)} \geq \frac{1}{2}$, we find that (4.23) is fulfilled by Assumption 4.2.

(b) **Term $\chi\left(\frac{1}{c_2} p \operatorname{Re} \hat{f}_k \overline{\hat{u}_k} - \frac{c_1}{c_2} \frac{\gamma}{2} p \mu_k^2 |\hat{u}_k|^2\right)$.**

We shall prove that the associated term in (4.20) is estimated by $\|e^{sa\eta} f\|_{L^2(q_{T,\delta})}^2$. Applying the Young inequality, we obtain

$$\frac{\mu_k p \operatorname{Re} \hat{f}_k \overline{\hat{u}_k}}{c_2} \leq \frac{D}{2\gamma \inf_{t \in [0, T]} (c_1 c_2)} |e^{sa\eta|_S} \hat{f}_k|^2 + \frac{c_1}{c_2} \frac{\gamma s \lambda a \varphi|_S p \mu_k^2 |\hat{u}_k|^2}{2}. \quad (4.24)$$

Observe that

$$\mu_k \geq s\lambda a(t)\varphi|_S, \quad t \in \tilde{J}_k \Leftrightarrow \mu_k \geq 2 \sqrt{\frac{D}{B}} \Phi(t; s, \lambda), \quad t \in \tilde{J}_k.$$

Arguing as above this will be fulfilled if

$$\frac{a(t_k)}{a(\frac{t_k}{2})} \geq 2\sqrt{\frac{D}{B}},$$

which holds by Assumption 4.2. We thus find, for $t \in \tilde{J}_k$,

$$\mu_k \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)} \chi \left(\frac{1}{c_2} p \operatorname{Re} \hat{f}_k \overline{\hat{u}_k} - \frac{c_1\gamma}{2c_2} p \mu_k^2 |\hat{u}_k|^2 \right) dx_n \leq \int_0^\delta \frac{D e^{-2(\frac{\sigma\mu_k}{2} x_n - s a \eta s)}}{2\gamma \inf_{t \in [0, T]} (c_1 c_2)} |\hat{f}_k|^2 dx_n,$$

and proceeding as in 2.(a) we find

$$\mu_k \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2-} + c_{2+}) \cosh(\sigma\mu_k\delta)} c_{2+} \chi \left(\frac{1}{c_2} p \operatorname{Re} \hat{f}_k \overline{\hat{u}_k} - \frac{c_1\gamma}{2c_2} p \mu_k^2 |\hat{u}_k|^2 \right) dx_n dt \lesssim \|e^{s a \eta} \hat{f}_k\|_{L^2(q_{T,\delta})}^2.$$

(c) **Term** $\chi p \frac{\partial_{x_n} c_2}{c_2} \operatorname{Re} \hat{u}_k \partial_{x_n} \overline{\hat{u}_k}$.

With the Young inequality we find $\mu_k \chi p \operatorname{Re} \hat{u}_k \partial_{x_n} \overline{\hat{u}_k} \leq \frac{1}{2} p |\partial_{x_n} \hat{u}_k|^2 + \mu_k^2 \frac{1}{2} p |\hat{u}_k|^2$. With (4.22), arguing as above we obtain

$$\begin{aligned} & \mu_k \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2+} + c_{2-}) \cosh(\sigma\mu_k\delta)} c_{2+} \chi p \frac{\partial_{x_n} c_2}{c_2} \operatorname{Re} \hat{u}_k \partial_{x_n} \overline{\hat{u}_k} dx_n dt \\ & \lesssim s \lambda \| (a\varphi)^{\frac{1}{2}} e^{s a \eta} \partial_{x_n} \hat{u}_k \|_{L^2(q_{T,\delta})}^2 + s \lambda \| (a\varphi)^{\frac{1}{2}} \mu_k \hat{v}_k \|_{L^2(q_{T,\delta})}^2 \\ & \lesssim s \lambda \| (a\varphi)^{\frac{1}{2}} \partial_{x_n} \hat{v}_k \|_{L^2(q_{T,\delta})}^2 + s^3 \lambda^3 \| (a\varphi)^{\frac{3}{2}} \hat{v}_k \|_{L^2(q_{T,\delta})}^2 + s \lambda \| (a\varphi)^{\frac{1}{2}} \mu_k \hat{v}_k \|_{L^2(q_{T,\delta})}^2. \end{aligned}$$

(d) **Term** $\frac{\chi'}{2c_2} p |\hat{u}_k|^2$.

As we have $\|\chi'\|_\infty \leq C/t_k$ we get

$$\|\chi'\|_\infty \lesssim T a(t_k) \lesssim \frac{T \Phi(t_k; s, \lambda)}{s \lambda \varphi|_S} \sqrt{\frac{D}{B}} \lesssim \frac{T \mu_k}{s \lambda \varphi|_S} \sqrt{\frac{D}{B}}.$$

We thus find

$$\mu_k \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)} \frac{\chi'}{2c_2} p |\hat{u}_k|^2 dx_n \lesssim \mu_k^2 T a(t) \sqrt{\frac{D^3}{B}} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{\cosh(\sigma\mu_k\delta)} e^{2s a \eta s} |\hat{u}_k|^2 dx_n.$$

Arguing as above with (4.22) we obtain

$$\begin{aligned} \mu_k \int_{\tilde{J}_k} \int_0^\delta \frac{\sinh(\sigma\mu_k(\delta - x_n))}{(c_{2+} + c_{2-}) \cosh(\sigma\mu_k\delta)} c_{2+} \frac{\chi'}{2\sigma c_2} p |\hat{u}_k|^2 dx_n dt & \lesssim T \|a^{\frac{1}{2}} \mu_k \hat{v}_k\|_{L^2(q_{T,\delta})}^2 \\ & \lesssim s \lambda \| (a\varphi)^{\frac{1}{2}} \mu_k \hat{v}_k \|_{L^2(q_{T,\delta})}^2, \end{aligned}$$

if $s \geq s_0 T$, with $s_0 > 0$, and $\lambda \geq \lambda_0 > 0$.

Collecting all the estimates we have obtained we conclude the proof of the Proposition 4.4. \blacksquare

End of the proof of Theorem 1.5.

With Proposition 4.1, estimate (4.4), Lemma 4.3 and Proposition 4.4, for λ and $s/(T + T^2)$ sufficiently large, we obtain, for all $k \in \mathbb{N}^*$,

$$\begin{aligned} & s^{-1} \|(a\varphi)^{-\frac{1}{2}} \partial_t \hat{v}_k\|_{L^2(Q_{T,\delta})}^2 + s\lambda^2 \left(\|(a\varphi)^{\frac{1}{2}} \partial_{x_n} \hat{v}_k\|_{L^2(Q_{T,\delta})}^2 + \|(a\varphi)^{\frac{1}{2}} \mu_k \hat{v}_k\|_{L^2(Q_{T,\delta})}^2 \right) + s^3 \lambda^4 \|(a\varphi)^{\frac{3}{2}} \hat{v}_k\|_{L^2(Q_{T,\delta})}^2 \\ & + s\lambda \left(|(a\varphi|_S)^{\frac{1}{2}} \partial_{x_n} \hat{v}_k\|_{L^2((0,T))}^2 + |(a\varphi|_S)^{\frac{1}{2}} \mu_k \hat{v}_k\|_{L^2((0,T))}^2 \right) + s^3 \lambda^3 |(a\varphi|_S)^{\frac{3}{2}} \hat{v}_k\|_{L^2((0,T))}^2 \lesssim \|e^{s\alpha\eta} \hat{f}_k\|_{L^2(Q_{T,\delta})}^2. \end{aligned} \quad (4.25)$$

Summing over k , using (2.2) we obtain

$$\begin{aligned} & s^{-1} \|(a\varphi)^{-\frac{1}{2}} \partial_t v\|_{L^2(Q_T)}^2 + s\lambda^2 \left(\|(a\varphi)^{\frac{1}{2}} \partial_{x_n} v\|_{L^2(Q_T)}^2 + \|(a\varphi)^{\frac{1}{2}} \mu_k v\|_{L^2(Q_T)}^2 \right) + s^3 \lambda^4 \|(a\varphi)^{\frac{3}{2}} v\|_{L^2(Q_T)}^2 \\ & + s\lambda \left(|(a\varphi|_S)^{\frac{1}{2}} \partial_{x_n} v\|_{L^2(S_T)}^2 + |(a\varphi|_S)^{\frac{1}{2}} \mu_k v\|_{L^2(S_T)}^2 \right) + s^3 \lambda^3 |(a\varphi|_S)^{\frac{3}{2}} v\|_{L^2(S_T)}^2 \lesssim \|e^{s\alpha\eta} f\|_{L^2(Q_T)}^2. \end{aligned} \quad (4.26)$$

The remainder of the proof of the Carleman estimate is now classical (see e.g. [24]).

Remark 4.7. It is important to note that Proposition 4.4 is not a trace result, otherwise we stronger Sobolev norm would appear on the r.h.s. of (4.9). The L^2 -norm of the trace of the tangential derivative is estimated by an $L^2((0, T); H^1(\Omega))$ -norm, but this is valid only for solutions of $Pu = f$. This result appears to us as an expression of the parabolic regularization effect.

Observe that the estimate of Proposition 4.4 is also valid in the case where c_1 and c_2 are smooth if the weight function β is chosen with a discontinuous derivative across S according to Assumptions 2.1 and 4.2.

A Proof of some intermediate results

A.1 Proof of Proposition 3.1

For later use of this proof in Section 4 we consider the case $c_1 \neq c_2$ here. The inequality we prove is uniform w.r.t. k . We shall thus remove the Fourier notation \hat{u}_k and simply write $(-\partial_{x_n} c_2 \partial_{x_n} + \mu^2)u = f$. We introduce $v = e^{s\varphi} u$ and $g = e^{s\varphi} f$ and we obtain

$$(-\partial_{x_n} c_2 \partial_{x_n} - c_2 (s\varphi')^2 + c_1 \mu^2 + 2sc_2 \varphi' \partial_{x_n} + s\partial_{x_n} (c_2 \varphi'))v = g,$$

which, following [16], we write $M_1 v + M_2 v = \tilde{g}$, with

$$\begin{aligned} M_1 &= -\partial_{x_n} c_2 \partial_{x_n} - c_2 (s\varphi')^2 + c_1 \mu^2, & M_2 &= 2sc_2 \varphi' \partial_{x_n} + s\partial_{x_n} (c_2 \varphi''), \\ \tilde{g} &= g + (p-1)sc_2 \varphi'' v - s(\partial_{x_n} c_2) \varphi' v, & 1 &< p < 3. \end{aligned}$$

The introduction of the parameter p is for instance explained in [24]. Following the classical method to prove Carleman estimates we compute

$$\|\tilde{g}\|_{L^2(\mathbb{R}^+)}^2 = \|M_1 v\|_{L^2(\mathbb{R}^+)}^2 + \|M_2 v\|_{L^2(\mathbb{R}^+)}^2 + 2 \operatorname{Re}(M_1 v, M_2 v)_{L^2(\mathbb{R}^+)}, \quad (A.1)$$

considering only the region $\{x_n > 0\}$ for now. We focus on the computation of the third term which we write as sum of 4 terms I_{ij} , $1 \leq i \leq 2$, $1 \leq j \leq 2$, where I_{ij} is the inner product of the i^{th} term in the expression of $M_1 v$ and the j^{th} term in the expression of $M_2 v$ above.

Term I_{11} . With an integration by parts we have

$$\begin{aligned} I_{11} &= -2 \operatorname{Re} \int_{x_n > 0} s\varphi'(\partial_{x_n} c_2 \partial_{x_n} v) c_2 \overline{\partial_{x_n} v} dx_n = - \int_{x_n > 0} s\varphi' \partial_{x_n} |c_2 \partial_{x_n} v|^2 dx_n \\ &= s\varphi' |c_2 \partial_{x_n} v|_{|x_n=0^+}^2 + \int_{x_n > 0} s\varphi'' |c_2 \partial_{x_n} v|^2 dx_n. \end{aligned}$$

Term I_{21} . Similarly we find

$$\begin{aligned} I_{21} &= \operatorname{Re} \int_{x_n > 0} (-c_2 (s\varphi')^3 + s c_1 \mu^2 \varphi') c_2 \partial_{x_n} |v|^2 dx_n \\ &= (c_2 (s\varphi')^3 - s c_1 \mu^2 \varphi') c_2 |v|_{|x_n=0^+}^2 + \int_{x_n > 0} c_2 (3s^3 c_2 (\varphi')^2 \varphi'' - s c_1 \mu^2 \varphi'') |v|^2 dx_n \\ &\quad + \int_{x_n > 0} (2c_2 \partial_{x_n} c_2 (s\varphi')^3 - (c_1 \partial_{x_n} c_2 + c_2 \partial_{x_n} c_1) s \mu^2 \varphi') |v|^2 dx_n. \end{aligned}$$

Term I_{12} . We have

$$\begin{aligned} I_{12} &= -s p \operatorname{Re} \int_{x_n > 0} (\partial_{x_n} c_2 \partial_{x_n} v) c_2 \varphi'' \bar{v} dx_n \\ &= s p \int_{x_n > 0} \varphi'' |c_2 \partial_{x_n} v|^2 dx_n + s p \varphi'' \operatorname{Re}(c_2 \partial_{x_n} v) c_2 \bar{v}|_{x_n=0^+} + s p \operatorname{Re} \int_{x_n > 0} \partial_{x_n} (c_2 \varphi'') (c_2 \partial_{x_n} v) \bar{v} dx_n. \end{aligned}$$

Term I_{22} . We directly find $I_{22} = s p \int_{x_n > 0} c_2 (-c_2 (s\varphi')^2 + c_1 \mu^2) \varphi'' |v|^2 dx_n$.

Collecting together the different terms we have obtained we find

$$\frac{1}{2} \|\tilde{g}\|_{L^2(\mathbb{R}^+)}^2 \geq \int_{x_n > 0} \alpha_0 |v|^2 dx_n + \int_{x_n > 0} \alpha_1 |c_2 \partial_{x_n} v|^2 dx_n + \gamma_0 |v|_{|x_n=0^+}^2 + \gamma_1 |c_2 \partial_{x_n} v|_{|x_n=0^+}^2 + X + Y,$$

with

$$\begin{aligned} \alpha_0 &= s(p-1)c_1 c_2 \mu^2 \varphi'' + (3-p)s^3 (c_2 \varphi')^2 \varphi'', & \alpha_1 &= s(p+1)\varphi'', \\ \gamma_0 &= c_2^2 (s\varphi')_{|x_n=0^+}^3 - c_1 c_2 s \mu^2 \varphi'_{|x_n=0^+}, & \gamma_1 &= s\varphi', \\ X &= s p \operatorname{Re} \int_{x_n > 0} \partial_{x_n} (c_2 \varphi'') (c_2 \partial_{x_n} v) \bar{v} dx_n \\ &\quad + \int_{x_n > 0} (2c_2 \partial_{x_n} c_2 (s\varphi')^3 - (c_1 \partial_{x_n} c_2 + c_2 \partial_{x_n} c_1) \mu^2 s \varphi') |v|^2 dx_n, \\ Y &= s p c^2 \varphi'' \operatorname{Re}(\partial_{x_n} v) \bar{v}|_{x_n=0^+}. \end{aligned}$$

Because of the form of φ , a direct computation shows that

$$\alpha_0 \gtrsim s \lambda^2 \mu^2 \varphi + s^3 \lambda^4 \varphi^3, \quad \alpha_1 \gtrsim C s \lambda^2 \varphi,$$

for λ chosen sufficiently large. Recalling that β is affine in the region we consider we find

$$\begin{aligned} X &= s p \operatorname{Re} \int_{x_n > 0} (c_2^2 \lambda^3 \beta'^3 + c_2 (\partial_{x_n} c_2) \lambda^2 \beta'^2) \varphi (\partial_{x_n} v) \bar{v} dx_n \\ &\quad + \int_{x_n > 0} (2c_2 \partial_{x_n} c_2 (s \lambda \beta' \varphi)^3 - (c_1 \partial_{x_n} c_2 + c_2 \partial_{x_n} c_1) \mu^2 s \lambda \beta' \varphi) |v|^2 dx_n, \end{aligned}$$

and

$$\|\tilde{g}\|_{L^2(\mathbb{R}^+)}^2 \lesssim \|g\|_{L^2(\mathbb{R}^+)}^2 + s^2(\lambda^4 + \lambda^2) \int_{x_n > 0} \varphi^2 |v|^2.$$

Choosing s and λ sufficiently large, with the Young inequality, we obtain

$$\begin{aligned} C\|g\|_{L^2(\mathbb{R}^+)}^2 &\geq C' \int_{x_n > 0} (s\lambda^2\mu^2\varphi + s^3\lambda^4\varphi^3)|v|^2 dx_n + C' \int_{x_n > 0} s\lambda^2\varphi|\partial_{x_n} v|^2 dx_n \\ &\quad + \gamma_0|v|_{x_n=0^+}^2 + \gamma_1|c_2\partial_{x_n} v|_{x_n=0^+}^2 + Y. \end{aligned} \quad (\text{A.2})$$

The same type of estimate can be obtained in the region $\{x_n < 0\}$ with opposite signs for the trace terms. The sum of (A.2) from both sides yields the result. \blacksquare

A.2 Proof of Lemma 3.4

Here we drop the \hat{v}_k notation and simply write v . It follows that

$$c\partial_{x_n} v = e^{s\varphi}(c\partial_{x_n} u + cs(\partial_{x_n} \varphi)u) = e^{s\varphi}(c\partial_{x_n} u + c\beta'(s\lambda\varphi u)).$$

We set $a = c\partial_{x_n} u$ and $b = s\lambda\varphi u$. We then have

$$[|c\partial_{x_n} v|^2 \beta']_S = e^{2s\varphi} \left([\beta']_S |a|^2 + [c^2(\beta')^3]_S |b|^2 + 2[c(\beta')^2] \operatorname{Re} \bar{a}b \right).$$

We thus obtain

$$\mathcal{B}(v) = s\lambda\varphi_S e^{2s\varphi}(Aw, w),$$

with $w = (a, b)^t$ and where A is the following symmetric matrix

$$A = \begin{pmatrix} [\beta']_S & [c(\beta')^2]_S \\ [c(\beta')^2]_S & 2[c^2(\beta')^3]_S \end{pmatrix} = \beta'_- \begin{pmatrix} (L-1) & c_+\beta'_-(L^2-K) \\ c_+\beta'_-(L^2-K) & 2(c_+\beta'_-)^2(L^3-K^2) \end{pmatrix}.$$

We then see that

$$(Aw, w) = \beta'_-(L-1) \left| a + c_+\beta'_- \frac{L^2-K}{L-1} b \right|^2 + \beta'_- \left(2(c_+\beta'_-)^2(L^3-K^2) - (c_+\beta'_-)^2 \frac{(L^2-K)^2}{L-1} \right) |b|^2.$$

which gives the result. \blacksquare

A.3 Traces of the solution

We consider the following ODEs

$$v'' - \mu^2 v = F, \quad s \in (-\delta, 0) \cup (0, \delta), \quad (\text{A.3})$$

$$v(-\delta) = v(\delta) = 0, \quad v|_{s=0^-} = v|_{s=0^+}, \quad cv'|_{s=0^-} = cv'|_{s=0^+}. \quad (\text{A.4})$$

Here $\mu > 0$. The solutions of (A.3) can be written as

$$v(s) = A_{\pm} \cosh(\mu s) + B_{\pm} \sinh(\mu s) + \mu^{-1} \int_0^s \sinh(\mu(s-\sigma)) F(\sigma) d\sigma, \quad s \in (-\delta, 0) \cup (0, \delta).$$

We then have $A_{\pm} = v_{|s=0^{\pm}}$, $\mu B_{\pm} = v_{|s=0^{\pm}}$ and

$$v(\pm\delta) = \mu^{-1} \cosh(\mu\delta) \left(\mu A_{\pm} + \mu B_{\pm} \tanh(\pm\mu\delta) + \int_0^{\pm\delta} \frac{\sinh(\mu(\pm\delta - \sigma))}{\cosh(\mu\delta)} F(\sigma) d\sigma \right).$$

The boundary conditions (A.4) then yield

$$\begin{pmatrix} \mu & \tanh(\mu\delta) \\ \mu & -\frac{c_{\pm}}{c_{-}} \tanh(\mu\delta) \end{pmatrix} \begin{pmatrix} v_{+}(0) \\ v'_{+}(0) \end{pmatrix} = \begin{pmatrix} -\int_0^{\delta} \frac{\sinh(\mu(\delta - \sigma))}{\cosh(\mu\delta)} F(\sigma) d\sigma \\ -\int_0^{-\delta} \frac{\sinh(\mu(-\delta - \sigma))}{\cosh(\mu\delta)} F(\sigma) d\sigma \end{pmatrix}.$$

We observe that the determinant of this system,

$$D = -c_{-}^{-1} \tanh(\mu\delta) \mu (c_{+} + c_{-}),$$

is non zero as $\mu > 0$. It thus follows that

$$v_{-}(0) = v_{+}(0) = -\frac{c_{+}}{\mu} \int_0^{\delta} \frac{\sinh(\mu(\delta - \sigma))}{(c_{+} + c_{-}) \cosh(\mu\delta)} F(\sigma) d\sigma - \frac{c_{-}}{\mu} \int_0^{-\delta} \frac{\sinh(\mu(-\delta - \sigma))}{(c_{+} + c_{-}) \cosh(\mu\delta)} F(\sigma) d\sigma.$$

A.4 Proof of Lemma 3.6

We first consider the case $K = 1$. Then $L = 2$ and $P(K, L) = -(1 + \varepsilon) + 5(1 - \varepsilon)^2$. The result is clear for ε sufficiently small.

We now consider the case $0 < K < 1$. Then $L > 1$; we have $L - K^2 > 0$ and thus

$$P(K, L) = -(1 + \varepsilon)(L - K^2)(L - 1) + (1 - \varepsilon)^2(2(L^3 - K^2)(L - 1) - (L^2 - K^2)^2).$$

For convenience we write $(1 - \varepsilon)^2/(1 + \varepsilon) = 1 - \alpha$ with $0 < \alpha < 1$. We then find

$$Q(K) = P(K, (r + 1) - rK) = -(K - 1)^2 S(K), \quad S(K) = aK^2 + bK + c,$$

with

$$\begin{aligned} a &= -(1 - \alpha)r^4 < 0, & b &= 2(1 - \alpha)(r^4 + r^3 - r^2) - (1 - 2\alpha)r, \\ c &= -(1 - \alpha)(r^4 + 2r^3 - 1) + r^2 + (3 - 2\alpha)r. \end{aligned}$$

It thus remains to prove that $S(1) \leq 0$ and $S'(1) \geq 0$. We find

$$S(1) = (2\alpha - 1)r^2 + 2r + 1 - \alpha, \quad S'(1) = r(2(1 - \alpha)r^2 - 2(1 - \alpha)r - (1 - 2\alpha)).$$

We see that $S(1) < 0$ and $S'(1) > 0$ if $\alpha = 0$ and $r = 3$. It thus remains true for α sufficiently small. ■

A.5 Proof of Proposition 4.1

The inequality we prove is uniform w.r.t. k . We shall thus remove the Fourier notation \hat{u}_k and simply write $(\partial_t - \partial_{x_n} c_2 \partial_{x_n} + c_1 \mu^2)u = f$. We introduce $v = e^{sa\eta}u$ and $g = e^{sa\eta}f$ and we obtain

$$\left(\partial_t - \partial_{x_n} c_2 \partial_{x_n} - c_2 (sa\eta')^2 + c_1 \mu^2 + 2sc_2 a\eta' \partial_{x_n} + sa \partial_{x_n} (c_2 \eta') - sa' \eta\right)v = g,$$

which we write $M_1 v + M_2 v = \tilde{g}$, with

$$\begin{aligned} M_1 &= -\partial_{x_n} c_2 \partial_{x_n} + [-c_2 (sa\eta')^2 + c_1 \mu^2] - sa' \eta, & M_2 &= 2sc_2 a\eta' \partial_{x_n} + spc_2 a\eta'' + \partial_t, \\ \tilde{g} &= g + (p-1)sc_2 a\eta'' v - s(\partial_{x_n} c_2) a\eta' v, & 1 < p < 3. \end{aligned} \quad (\text{A.5})$$

In preparation for what follows we observe that

$$1 \lesssim T^2 a, \quad |a'| \lesssim T a^2, \quad |a''| \lesssim T^2 a^3, \quad |\eta| \lesssim \varphi^2.$$

We have

$$\|\tilde{g}\|_{L^2((0,T)\times\mathbb{R}^+)}^2 \lesssim \|g\|_{L^2((0,T)\times\mathbb{R}^+)}^2 + s^2(\lambda^4 + \lambda^2)T^2 \|a^{\frac{3}{2}}\varphi^{\frac{1}{2}}v\|_{L^2((0,T)\times\mathbb{R}^+)}^2.$$

We compute

$$\|\tilde{g}\|_{L^2((0,T)\times\mathbb{R}^+)}^2 = \|M_1 v\|_{L^2((0,T)\times\mathbb{R}^+)}^2 + \|M_2 v\|_{L^2((0,T)\times\mathbb{R}^+)}^2 + 2 \operatorname{Re}(M_1 v, M_2 v)_{L^2((0,T)\times\mathbb{R}^+)}, \quad (\text{A.6})$$

considering only the region $\{x_n > 0\}$ for now. For the computation of the last term in (A.6), we set I_{ij} , $1 \leq i \leq 3$, $1 \leq j \leq 3$, where I_{ij} is the inner product of the i^{th} term in the expression of $M_1 v$ and the j^{th} term in the expression of $M_2 v$ above. For the computations of I_{11} , I_{12} , I_{21} and I_{22} we refer to the computations performed in Appendix A.1 (simply replacing φ by $a\varphi$ and integrating in time).

Term I_{13} . By integration by parts we find

$$\begin{aligned} I_{13} &= \operatorname{Re} \int_0^T \int_{0, x_n > 0} -(\partial_{x_n} c_2 \partial_{x_n} v) \partial_t \bar{v} \, dx_n dt \\ &= \frac{1}{2} \int_0^T \int_{0, x_n > 0} c_2 \partial_t |\partial_{x_n} v|^2 \, dx_n dt + \operatorname{Re} \int_0^T \left((c_2 \partial_{x_n} v) \partial_t \bar{v} \right)_{|x_n=0^+} dt \\ &= -\frac{1}{2} \int_0^T \int_{0, x_n > 0} (\partial_t c_2) |\partial_{x_n} v|^2 \, dx_n dt + \operatorname{Re} \int_0^T \left((c_2 \partial_{x_n} v) \partial_t \bar{v} \right)_{|x_n=0^+} dt. \end{aligned}$$

We have

$$\left| \frac{1}{2} \int_0^T \int_{0, x_n > 0} (\partial_t c_2) |\partial_{x_n} v|^2 \, dx_n dt \right| \lesssim T^2 \int_0^T \int_{0, x_n > 0} a |\partial_{x_n} v|^2 \, dx_n dt.$$

Term I_{23} . By integration by parts we have

$$\begin{aligned} I_{23} &= \frac{1}{2} \int_0^T \int_{0, x_n > 0} (-c_2 (sa\eta')^2 + c_1 \mu^2) \partial_t |v|^2 \, dx_n dt = s^2 \int_0^T \int_{0, x_n > 0} c_2 a a' \eta'^2 |v|^2 \, dx_n dt \\ &\quad + \frac{1}{2} \int_0^T \int_{0, x_n > 0} [(\partial_t c_2) (sa\eta')^2 - (\partial_t c_1) \mu^2] |v|^2 \, dx_n dt \end{aligned}$$

We thus find

$$|I_{23}| \lesssim (T + T^2)s^2\lambda^2 \int_{0,x_n>0}^T \int a^3\varphi^2|v|^2 dx_n dt + T^2 \int_{0,x_n>0}^T \int a\mu^2|v|^2 dx_n dt.$$

Term I_{33} . By integration by parts we find

$$I_{33} = -\frac{s}{2} \int_{0,x_n>0}^T \int a'\eta\partial_t|v|^2 dx_n dt = \frac{s}{2} \int_{0,x_n>0}^T \int a''\eta|v|^2 dx_n dt.$$

This yields

$$|I_{33}| \lesssim T^2 s \int_{0,x_n>0}^T \int a^3\varphi^2|v|^2 dx_n dt.$$

Terms I_{31} and I_{32} . We have

$$I_{31} = -s^2 \int_{0,x_n>0}^T \int c_2 a a' \eta \eta' \partial_{x_n} |v|^2 dx_n dt = s^2 \int_{0,x_n>0}^T \int a a' \partial_{x_n} (c_2 \eta \eta') |v|^2 dx_n dt.$$

We find directly $I_{32} = -ps^2 \int_0^T \int_{x_n>0} c_2 a a' \eta \eta'' |v|^2 dx_n dt$. We then obtain

$$|I_{31}| + |I_{32}| \lesssim T s^2 (\lambda + \lambda^2) \int_{0,x_n>0}^T \int a^3 \varphi^3 |v|^2 dx_n dt.$$

With the computations of Appendix A.1 we find, for λ and $sa \gtrsim s/T^2$ sufficiently large,

$$\begin{aligned} C\|g\|_{L^2((0,T)\times\mathbb{R}^+)}^2 &\geq C' \int_{0,x_n>0}^T \int (s\lambda^2\mu^2 a\varphi + s^3\lambda^4(a\varphi)^3)|v|^2 dx_n dt + C' \int_{0,x_n>0}^T \int s\lambda^2 a\varphi |\partial_{x_n} v|^2 dx_n dt \\ &+ \frac{1}{2}\|M_2 v\|_{L^2((0,T)\times\mathbb{R}^+)}^2 + \int_0^T (\gamma_0 |v|_{|x_n=0^+}^2 + \gamma_1 |c\partial_{x_n} v|_{|x_n=0^+}^2) dt + X + Y + I_{13} + I_{23} + I_{33} + I_{31} + I_{32}, \end{aligned}$$

with

$$\begin{aligned} \gamma_0 &= c_2^2 (sa\varphi')_{|x_n=0^+}^3 - c_1 c_2 s\mu^2 a\varphi'_{|x_n=0^+}, \quad \gamma_1 = sa\varphi', \\ X &= sp \operatorname{Re} \int_{0,x_n>0}^T \int a \partial_{x_n} (c_2 \varphi'') (c_2 \partial_{x_n} v) \bar{v} dx_n dt \\ &+ \int_{0,x_n>0}^T \int [2c_2 (\partial_{x_n} c_2) (sa\varphi')^3 - \partial_{x_n} (c_1 c_2) \mu^2 sa\varphi'] |v|^2 dx_n dt, \\ Y &= sp \int_0^T c_2^2 a\varphi'' \operatorname{Re}(\partial_{x_n} v) \bar{v}|_{x_n=0^+} dt. \end{aligned}$$

For λ and $s/(T + T^2)$ sufficiently large, the estimations we found above yield,

$$\begin{aligned} C\|g\|_{L^2((0,T)\times\mathbb{R}^+)}^2 &\geq C' \int_{0,x_n>0}^T \int (s\lambda^2\mu^2 a\varphi + s^3\lambda^4(a\varphi)^3)|v|^2 dx_n dt + C' \int_{0,x_n>0}^T \int s\lambda^2 a\varphi |\partial_{x_n} v|^2 dx_n dt \\ &+ \frac{1}{2}\|M_2 v\|_{L^2((0,T)\times\mathbb{R}^+)}^2 + \int_0^T (\gamma_0 |v|_{|x_n=0^+}^2 + \gamma_1 |c_2 \partial_{x_n} v|_{|x_n=0^+}^2) dt + Y', \quad (\text{A.7}) \end{aligned}$$

with

$$Y' = sp \int_0^T c_2^2 a \varphi'' \operatorname{Re}(\partial_{x_n} v) \bar{v}|_{x_n=0^+} dt + \operatorname{Re} \int_0^T ((c_2 \partial_{x_n} v) \partial_t \bar{v})|_{x_n=0^+} dt.$$

The same type of estimate can be obtained in the region $\{x_n < 0\}$ with opposite signs for the trace terms. The sum of (A.7) from both sides yields

$$\begin{aligned} & C(s\lambda^2 \|(a\varphi)^{\frac{1}{2}} \partial_{x_n} v\|_{L^2(Q_{T,\delta})}^2 + s\lambda^2 \|(a\varphi)^{\frac{1}{2}} \mu v\|_{L^2(Q_{T,\delta})}^2 + s^3 \lambda^4 \|(a\varphi)^{\frac{3}{2}} v\|_{L^2(Q_{T,\delta})}^2) \\ & + \frac{1}{2} \|M_2 v\|_{L^2((0,T) \times \mathbb{R}^*)}^2 + s\lambda \int_0^T a \varphi|_S \left([c_2^2 \beta' |\partial_{x_n} v|^2]_S + |s\lambda a \varphi v|_{S_T}|^2 [c_2^2 \beta'^3]_S \right) dt + Y'' \\ & \leq C \|e^{san} f\|_{L^2(Q_{T,\delta})}^2 + s\lambda \int_0^T a \varphi|_S |\mu v|_{S_T}|^2 [c_1 c_2 \beta']_S dt, \quad (\text{A.8}) \end{aligned}$$

with $Y'' = Y_1'' + Y_2''$, where

$$Y_1'' = sp\lambda^2 \int_0^T a \varphi|_S \operatorname{Re} [c_2^2 \beta'^2 \partial_{x_n} v]_S \bar{v}|_{S_T} dt,$$

and

$$Y_2'' = \operatorname{Re} \int_0^T [c_2 \partial_{x_n} v]_S \partial_t \bar{v}|_{S_T} dt.$$

We have

$$\begin{aligned} |Y_1''| & \lesssim s^{\frac{1}{2}} \lambda \int_0^T a^{\frac{1}{2}} \varphi|_S (|\partial_{x_n} v|_{S_T^-}^2 + |\partial_{x_n} v|_{S_T^+}^2) dt + s^{\frac{3}{2}} \lambda^3 \int_0^T a^{\frac{3}{2}} \varphi|_S |v|_{S_T}^2 dt \\ & \lesssim s^{\frac{1}{2}} \lambda T \int_0^T a \varphi|_S (|\partial_{x_n} v|_{S_T^-}^2 + |\partial_{x_n} v|_{S_T^+}^2) dt + s^{\frac{3}{2}} \lambda^3 T^3 \int_0^T a^3 \varphi|_S |v|_{S_T}^2 dt. \end{aligned}$$

As $u = e^{-san} v$ we have $c \partial_{x_n} u = c_2 (\partial_{x_n} v - sa(\eta') v) e^{-san}$ and thus

$$[c_2 \partial_{x_n} v]_S = sa [c_2 \eta']_S v|_{S_T} = s\lambda a [c_2 \beta']_S (v\varphi)|_{S_T}.$$

By integration by parts, we thus have

$$Y_2'' = \frac{1}{2} s\lambda \int_0^T a [c_2 \partial_{x_n} \beta]_S \varphi|_S \partial_t |v|_{S_T}^2 dt = -\frac{1}{2} s\lambda \int_0^T [\partial_t (ac_2) \partial_{x_n} \beta]_S \varphi|_S |v|_{S_T}^2 dt.$$

We thus obtain

$$|Y_2''| \lesssim s(T^3 + T^4) \lambda \int_0^T a^3 \varphi|_S |v|_{S_T}^2 dt. \quad (\text{A.9})$$

Finally, from the form of M_2 in (A.5), we have

$$\begin{aligned} \|(sa\varphi)^{-\frac{1}{2}} \partial_t v\|_{L^2((0,T) \times \mathbb{R}^*)}^2 & \lesssim \|(sa\varphi)^{-\frac{1}{2}} M_2 v\|_{L^2((0,T) \times \mathbb{R}^*)}^2 + \|(sa\varphi)^{\frac{1}{2}} \lambda \partial_{x_n} v\|_{L^2(Q_{T,\delta})}^2 + \|(sa\varphi)^{\frac{1}{2}} \lambda^2 v\|_{L^2(Q_{T,\delta})}^2 \\ & \lesssim \|M_2 v\|_{L^2((0,T) \times \mathbb{R}^*)}^2 + s\lambda^2 \|(a\varphi)^{\frac{1}{2}} \partial_{x_n} v\|_{L^2(Q_{T,\delta})}^2 + s\lambda^4 \|(a\varphi)^{\frac{1}{2}} v\|_{L^2(Q_{T,\delta})}^2 \\ & \lesssim \|M_2 v\|_{L^2((0,T) \times \mathbb{R}^*)}^2 + s\lambda^2 \|(a\varphi)^{\frac{1}{2}} \partial_{x_n} v\|_{L^2(Q_{T,\delta})}^2 + s^3 \lambda^4 \|(a\varphi)^{\frac{3}{2}} v\|_{L^2(Q_{T,\delta})}^2, \end{aligned}$$

as here $sa \gtrsim s/T^2 \geq s_0$, for some $s_0 > 0$ and $\varphi \geq 1$. ■

A.6 Proof of Lemma 4.6

Computing $p'(t) = s\lambda e^{2sa(t)\eta_S} a'(t)\varphi_{|S}[1 + 2s\eta_S a(t)]\mathbb{D}$, we have

$$\frac{p'}{p}(t) = \frac{a'}{a}(t)[1 + 2sa(t)\eta_S],$$

If $\frac{T}{2} \leq t < T - \frac{t_k}{2}$, because of the form of η in (4.1) we find $\frac{p'}{p} \leq 0$, for $sa(t) \gtrsim s/T^2$ and λ both sufficiently large. This implies that inequality (4.18) holds for these values of t .

We now consider the case $\frac{t_k}{2} < t < \frac{T}{2}$. Note that $\frac{p'}{p}$ is nonnegative, for s/T^2 and λ large, as here $a'(t) < 0$. Setting $\tilde{c}_1 = \inf_{t, x_n} c_1(t, x_n)$, and using the definition of t_k in (4.11), it suffices to prove

$$\frac{p'(t)}{\tilde{c}_1 p(t)} \leq (1 - \gamma)\Phi^2(t_k; s, \lambda), \quad \frac{t_k}{2} < t < \frac{T}{2}.$$

For all s, λ , we have

$$\frac{p'(t)}{p(t)} = \frac{2t - T}{t(T - t)}[2s\eta_S a(t) + 1] \leq \frac{2t - T}{t(T - t)} 2s\eta_S a(t) \leq -2T s\eta_S a^2(t).$$

As we have

$$\frac{1}{2} < \frac{\Phi(t_k; s, \lambda)}{\Phi(t_k/2; s, \lambda)} = \frac{a(t_k)}{a(t_k/2)} < \frac{3}{4}, \quad (\text{A.10})$$

it is sufficient to prove

$$-\frac{2T}{\tilde{c}_1} s\eta_S a^2(t) \leq \frac{1}{4}(1 - \gamma)\Phi^2(t_k/2; s, \lambda), \quad \frac{t_k}{2} < t < \frac{T}{2}. \quad (\text{A.11})$$

As the function Φ decreases on $(0, T/2)$, (A.11) holds if we have

$$-\frac{2T}{\tilde{c}_1} s\eta_S a^2(t) \leq \frac{1}{4}(1 - \gamma)\Phi^2(t; s, \lambda), \quad \frac{t_k}{2} < t < \frac{T}{2}.$$

With the definition of Φ , this reads

$$-\frac{\eta_S}{\varphi_{|S}^2} \leq \frac{\tilde{c}_1}{32T}(1 - \gamma)\frac{\mathbb{B}}{\mathbb{D}}s\lambda^2$$

or equivalently

$$e^{\lambda(\bar{\beta} - 2\beta_{|S})} - e^{-\lambda\beta_{|S}} \leq \frac{\tilde{c}_1}{32T}(1 - \gamma)\frac{\mathbb{B}}{\mathbb{D}}s\lambda^2.$$

As $\bar{\beta} < 2\beta_{|S}$ by construction of β (see the beginning of Section 4), this will hold for λ and s/T sufficiently large. \blacksquare

References

- [1] G. Alessandrini and L. Escauriaza, *Null-controllability of one-dimensional parabolic equations*, ESAIM Control Optim. Calc. Var. **14** (2008), no. 2, 284–293.
- [2] V. Barbu, *Exact controllability of the superlinear heat equation*, Appl. Math. Optim. **42** (2000), 73–89.
- [3] C. Bardos, G. Lebeau, and J. Rauch, *Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary*, SIAM J. Control Optim. **30** (1992), no. 5, 1024–1065.
- [4] A. Belghazi, F. Smadhi, N. Zaidi, and O. Zair, *Carleman inequalities for the heat equation in singular domains*, C. R. Acad. Sci. Paris, Ser I. **348** (2010), 277–282.
- [5] M. Bellassoued, *Carleman estimates and distribution of resonances for the transparent obstacle and application to the stabilization*, Asymptotic Anal. **35** (2003), 257–279.
- [6] A. Benabdallah, Y. Dermenjian, and J. Le Rousseau, *Carleman estimates for the one-dimensional heat equation with a discontinuous coefficient and applications*, Comptes Rendus Mécanique **334** (2006), 582–586.
- [7] ———, *Carleman estimates for the one-dimensional heat equation with a discontinuous coefficient and applications to controllability and an inverse problem*, J. Math. Anal. Appl. **336** (2007), 865–887.
- [8] ———, *On the controllability of linear parabolic equations with an arbitrary control location for stratified media*, C. R. Acad. Sci. Paris, Ser I. **344** (2007), 357–362.
- [9] A. L. Bukhgeim and M. V. Klibanov, *Global uniqueness of class of multidimensional inverse problems*, Soviet Math. Dokl. **24** (1981), 244–247.
- [10] T. Carleman, *Sur une problème d’unicité pour les systèmes d’équations aux dérivées partielles à deux variables indépendantes*, Ark. Mat. Astr. Fys. **26B** (1939), no. 17, 1–9.
- [11] R. Dautray and J.-L. Lions, *Analyse Mathématique et Calcul Numérique pour les Sciences et les Techniques*, vol. 4, Masson, Paris, 1988.
- [12] A. Doubova, A. Osses, and J.-P. Puel, *Exact controllability to trajectories for semilinear heat equations with discontinuous diffusion coefficients*, ESAIM Control Optim. Calc. Var. **8** (2002), 621–661.
- [13] E. Fernández-Cara and S. Guerrero, *Global Carleman inequalities for parabolic systems and application to controllability*, SIAM J. Control Optim. **45** (2006), no. 4, 1395–1446.
- [14] E. Fernández-Cara and E. Zuazua, *Null and approximate controllability for weakly blowing up semilinear heat equations*, Ann. Inst. H. Poincaré, Analyse non lin. **17** (2000), 583–616.
- [15] ———, *On the null controllability of the one-dimensional heat equation with BV coefficients*, Comput. Appl. Math. **21** (2002), 167–190.
- [16] A. Fursikov and O. Yu. Imanuvilov, *Controllability of evolution equations*, vol. 34, Seoul National University, Korea, 1996, Lecture notes.
- [17] L. Hörmander, *Linear Partial Differential Operators*, Springer-Verlag, Berlin, 1963.
- [18] ———, *The Analysis of Linear Partial Differential Operators*, vol. IV, Springer-Verlag, 1985.
- [19] O. Yu. Imanuvilov, V. Isakov, and M. Yamamoto, *An inverse problem for the dynamical Lamé system with two sets of boundary data*, Comm. Pure Appl. Math. **56** (2003), 1366–1382.
- [20] V. Isakov, *Inverse problems for partial differential equations*, Springer-Verlag, Berlin, 1998.
- [21] D. Jerison and G. Lebeau, *Harmonic analysis and partial differential equations (Chicago, IL, 1996)*, Chicago Lectures in Mathematics, ch. Nodal sets of sums of eigenfunctions, pp. 223–239, The University of Chicago Press, Chicago, 1999.

- [22] C. E. Kenig, J. Sjöstrand, and G. Uhlmann, *The Calderón problem with partial data*, Ann. of Math. **165** (2007), 567–591.
- [23] J. Le Rousseau, *Carleman estimates and controllability results for the one-dimensional heat equation with BV coefficients*, J. Differential Equations **233** (2007), 417–447.
- [24] J. Le Rousseau and G. Lebeau, *On Carleman estimates for elliptic and parabolic operators. Applications to unique continuation and control of parabolic equations*, Preprint (2009).
- [25] J. Le Rousseau and N. Lerner, *Carleman estimates for elliptic operators with jumps at an interface: Anisotropic case and sharp geometric conditions*, Journées “Équations aux Dérivées Partielles”, 2010, Port d’Albret.
- [26] ———, *Carleman estimates for elliptic operators with jumps at an interface: Anisotropic case and sharp geometric conditions*, Preprint (2010).
- [27] J. Le Rousseau and L. Robbiano, *Carleman estimate for elliptic operators with coefficients with jumps at an interface in arbitrary dimension and application to the null controllability of linear parabolic equations*, Arch. Rational Mech. Anal. **105** (2010), 953–990.
- [28] ———, *Local and global Carleman estimates for parabolic operators with coefficients with jumps at interfaces*, Invent. Math., to appear (2010).
- [29] G. Lebeau and L. Robbiano, *Contrôle exact de l’équation de la chaleur*, Comm. Partial Differential Equations **20** (1995), 335–356.
- [30] G. Lebeau and E. Zuazua, *Null-controllability of a system of linear thermoelasticity*, Arch. Rational Mech. Anal. **141** (1998), 297–329.
- [31] K. Miller, *Nonunique continuation for uniformly parabolic and elliptic equations in self-adjoint divergence form with Hölder continuous coefficients*, Arch. Rational Mech. Anal. **54** (1974), 105–117.
- [32] A. Pliś, *On non-uniqueness in Cauchy problem for an elliptic second order differential equation*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **11** (1963), 95–100.
- [33] O. Poisson, Personal communication, 2010.
- [34] C. Zuily, *Uniqueness and Non Uniqueness in the Cauchy Problem*, Birkhauser, Progress in mathematics, 1983.