



HAL
open science

A counterexample for Improved Sobolev Inequalities over the 2-adic group

Diego Chamorro

► **To cite this version:**

Diego Chamorro. A counterexample for Improved Sobolev Inequalities over the 2-adic group. 2010.
hal-00531555

HAL Id: hal-00531555

<https://hal.science/hal-00531555>

Preprint submitted on 3 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A counterexample for Improved Sobolev Inequalities over the 2-adic group

Diego Chamorro

November 3, 2010

Abstract

On the framework of the 2-adic group \mathbb{Z}_2 , we study a Sobolev-like inequality where we estimate the L^2 norm by a geometric mean of the BV norm and the $\dot{B}_{\infty}^{-1,\infty}$ norm. We first show, using the special topological properties of the p -adic groups, that the set of functions of bounded variations BV can be identified to the Besov space $\dot{B}_1^{1,\infty}$. This identification lead us to the construction of a counterexample to the improved Sobolev inequality.

Keywords: Sobolev inequalities, p -adic groups.

MSC 2010: 22E35, 46E35

1 Introduction

The general improved Sobolev inequalities were initially introduced by P. Gérard, Y. Meyer and F. Oru in [6]. For a function f such that $f \in \dot{W}^{s_1,p}(\mathbb{R}^n)$ and $f \in \dot{B}_{\infty}^{-\beta,\infty}(\mathbb{R}^n)$, these inequalities read as follows:

$$\|f\|_{\dot{W}^{s,q}} \leq C \|f\|_{\dot{W}^{s_1,p}}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta} \quad (1)$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s = \theta s_1 - (1 - \theta)\beta$ and $-\beta < s < s_1$. The method used for proving these estimates relies on the Littlewood-Paley decomposition and on a dyadic bloc manipulation and this explains the fact that the value $p = 1$ is forbidden here.

In order to study the case $p = 1$, it is necessary to develop other techniques. The case when $p = 1$, $s = 0$ and $s_1 = 1$ was treated by M. Ledoux in [9] using a special cut-off function; while the case $s_1 = 1$ and $p = 1$ was studied by A. Cohen, W. Dahmen, I. Daubechies & R. De Vore in [5]. In this last article, the authors give a BV -norm weak estimation using wavelet coefficients and isoperimetric inequalities and obtained, for a function f such that $f \in BV(\mathbb{R}^n)$ and $f \in \dot{B}_{\infty}^{-\beta,\infty}(\mathbb{R}^n)$, the estimation below:

$$\|f\|_{\dot{W}^{s,q}} \leq C \|f\|_{BV}^{1/q} \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-1/q} \quad (2)$$

where $1 < q \leq 2$, $0 \leq s < 1/q$ and $\beta = (1 - sq)/(q - 1)$.

In a previous work (see [3], [4]), we studied the possible generalizations of inequalities of type (1) and (2) to other frameworks than \mathbb{R}^n . In particular, we worked over stratified Lie groups and over polynomial volume growth Lie groups and we obtained some new weak-type estimates.

The aim of this paper is to study inequalities of type (1) and (2) in the setting of the 2-adic group \mathbb{Z}_2 . The main reason for working in the framework of \mathbb{Z}_2 is that this group is completely different from \mathbb{R}^n and from stratified or polynomial Lie groups. Indeed, since the 2-adic group is totally discontinuous, it is not absolutely trivial to give a definition for smoothness measuring spaces. Thus, the first step to do, in order to

study these Sobolev-like inequalities, is to give an adapted characterization of such functional spaces. This will be achieved using the Littlewood-Paley approach and, once this task is done, we will immediately prove -following the classical path exposed in [6]- the inequalities (1) in the setting of the 2-adic group \mathbb{Z}_2 .

For the estimate (2), we introduce the BV space in the following manner: we will say that $f \in BV(\mathbb{Z}_2)$ if there exists a constant $C > 0$ such that

$$\int_{\mathbb{Z}_2} |f(x+y) - f(x)| dx \leq C|y|_2 \quad (\forall y \in \mathbb{Z}_2).$$

As a surprising fact, we obtain the

Theorem 1 *We have the following relationship between the space of functions of bounded variation $BV(\mathbb{Z}_2)$ and the Besov space $\dot{B}_1^{1,\infty}(\mathbb{Z}_2)$:*

$$BV(\mathbb{Z}_2) \simeq \dot{B}_1^{1,\infty}(\mathbb{Z}_2)$$

Of course, this identification is false in \mathbb{R}^n and it is this special relationship in \mathbb{Z}_2 that give us our principal theorem which is the 2-adic counterpart of the inequality (2):

Theorem 2 *The following inequality is false in \mathbb{Z}_2 . There is not an universal constant $C > 0$ such that we have*

$$\|f\|_{L^2}^2 \leq C \|f\|_{BV} \|f\|_{\dot{B}_\infty^{-1,\infty}}$$

for all $f \in BV \cap \dot{B}_\infty^{-1,\infty}(\mathbb{Z}_2)$.

This striking fact says that the improved Sobolev inequalities of type (2) depend on the group's structure and that they are no longer true for the 2-adic group \mathbb{Z}_2 .

The plan of the article is the following: in section 2 we recall some well known properties about p -adic groups, in 3 we define Sobolev and Besov spaces, in 4 we prove theorem 1 and, finally, we prove the theorem 2 in section 5.

2 p -adic groups

We write $a|b$ when a divide b or, equivalently, when b is a multiple of a . Let p be any prime number, for $0 \neq x \in \mathbb{Z}$, we define the p -adic valuation of x by $\gamma(x) = \max\{r : p^r|x\} \geq 0$ and, for any rational number $x = \frac{a}{b} \in \mathbb{Q}$, we write $\gamma(x) = \gamma(a) - \gamma(b)$. Furthermore if $x = 0$, we agree to write $\gamma(0) = +\infty$.

Let $x \in \mathbb{Q}$ and p be any prime number, with the p -adic valuation of x we can construct a norm by writing

$$|x|_p = \begin{cases} p^{-\gamma} & \text{if } x \neq 0 \\ p^{-\infty} = 0 & \text{if } x = 0. \end{cases} \quad (3)$$

This expression satisfy the following properties

- a) $|x|_p \geq 0$, and $|x|_p = 0 \iff x = 0$;
- b) $|xy|_p = |x|_p |y|_p$;
- c) $|x+y|_p \leq \max\{|x|_p, |y|_p\}$, with equality when $|x|_p \neq |y|_p$.

When a norm satisfy c) it is called a non-Archimedean norm and an interesting fact is that over \mathbb{Q} all the possible norms are equivalent to $|\cdot|_p$ for some p : this is the so-called Ostrowski theorem, see [1] for a proof.

Definition 2.1 Let p be a any prime number. We define the field of p -adic numbers \mathbb{Q}_p as the completion of \mathbb{Q} when using the norm $|\cdot|_p$.

We present in the following lines the algebraic structure of the set \mathbb{Q}_p . Every p -adic number $x \neq 0$ can be represented in a unique manner by the formula

$$x = p^\gamma(x_0 + x_1p + x_2p^2 + \dots), \quad (4)$$

where $\gamma = \gamma(x)$ is the p -adic valuation of x and x_j are integers such that $x_0 > 0$ and $0 \leq x_j \leq p - 1$ for $j = 1, 2, \dots$. Remark that this canonical representation implies the identity $|x|_p = p^{-\gamma}$.

Let $x, y \in \mathbb{Q}_p$, using the formula (4) we define the sum of x and y by $x + y = p^{\gamma(x+y)}(c_0 + c_1p + c_2p^2 + \dots)$ with $0 \leq c_j \leq p - 1$ and $c_0 > 0$, where $\gamma(x + y)$ and c_j are the unique solution of the equation

$$p^{\gamma(x)}(x_0 + x_1p + x_2p^2 + \dots) + p^{\gamma(y)}(y_0 + y_1p + y_2p^2 + \dots) = p^{\gamma(x+y)}(c_0 + c_1p + c_2p^2 + \dots).$$

Furthermore, for $a, x \in \mathbb{Q}_p$, the equation $a + x = 0$ has a unique solution in \mathbb{Q}_p given by $x = -a$. In the same way, the equation $ax = 1$ has a unique solution in \mathbb{Q}_p : $x = 1/a$.

We take now a closer look at the topological structure of \mathbb{Q}_p . With the norm $|\cdot|_p$ we construct a distance over \mathbb{Q}_p by writing

$$d(x, y) = |x - y|_p \quad (5)$$

and we define the balls $B_\gamma(x) = \{y \in \mathbb{Q}_p : d(x, y) \leq p^{-\gamma}\}$ with $\gamma \in \mathbb{Z}$. Remark that, from the properties of the p -adic valuation, this distance has the *ultra-metric* property (*i.e.* $d(x, y) \leq \max\{d(x, z), d(z, y)\} \leq |x|_p + |y|_p$).

We gather with the next proposition some important facts concerning the balls in \mathbb{Q}_p .

Proposition 2.1 Let γ be an integer, then we have

- 1) the ball $B_\gamma(x)$ is a open and a closed set for the distance (5).
- 2) every point of $B_\gamma(x)$ is its center.
- 3) \mathbb{Q}_p endowed with this distance is a complete Hausdorff metric space.
- 4) \mathbb{Q}_p is a locally compact set.
- 5) the p -adic group \mathbb{Q}_p is a totally discontinuous space.

For a proof of this proposition and more details see the books [1], [8] or [13].

3 Functional spaces

In this article, we will work with the subset \mathbb{Z}_2 of \mathbb{Q}_2 which is defined by $\mathbb{Z}_2 = \{x \in \mathbb{Q}_2 : |x|_2 \leq 1\}$, and we will focus on real-valued functions over \mathbb{Z}_2 . Since \mathbb{Z}_2 is a locally compact commutative group, there exists a Haar measure dx which is translation invariant *i.e.*: $d(x + a) = dx$, furthermore we have the identity $d(xa) = |a|_2 dx$ for $a \in \mathbb{Z}_2^*$. We will normalize the measure dx by setting

$$\int_{\{|x|_2 \leq 1\}} dx = 1.$$

This measure is then unique and we will note $|E|$ the measure for any subset E of \mathbb{Z}_2 . Lebesgue spaces $L^p(\mathbb{Z}_2)$ are thus defined in a natural way: $\|f\|_{L^p} = \left(\int_{\mathbb{Z}_2} |f(x)|^p dx\right)^{1/p}$ for $1 \leq p < +\infty$, with the usual modifications when $p = +\infty$.

Let us now introduce the Littlewood-Paley decomposition in \mathbb{Z}_2 . We note \mathcal{F}_j the Boole algebra formed by the equivalence classes $E \subset \mathbb{Z}_2$ modulo the sub-group $2^j\mathbb{Z}_2$. Then, for any function $f \in L^1(\mathbb{Z}_2)$, we call $S_j(f)$ the conditionnal expectation of f with respect to \mathcal{F}_j :

$$S_j(f)(x) = \frac{1}{|B_j(x)|} \int_{B_j(x)} f(y) dy.$$

The dyadic blocks are thus defined by the formula $\Delta_j(f) = S_{j+1}(f) - S_j(f)$ and the Littlewood-Paley decomposition of a function $f : \mathbb{Z}_2 \rightarrow \mathbb{R}$ is given by

$$f = S_0(f) + \sum_{j=0}^{+\infty} \Delta_j(f) \quad \text{where } S_0(f) = \int_{\mathbb{Z}_2} f(x) dx. \quad (6)$$

We will need in the sequel some very special sets noted $Q_{j,k}$. Here is the definition and some properties:

Proposition 3.1 *Let $j \in \mathbb{N}$ and $k = \{0, 1, \dots, 2^j - 1\}$. Define the subset $Q_{j,k}$ of \mathbb{Z}_2 by*

$$Q_{j,k} = \{k + 2^j\mathbb{Z}_2\}. \quad (7)$$

Then

- 1) We have the identity $\mathcal{F}_j = \bigcup_{0 \leq k < 2^j} Q_{j,k}$,
- 2) For $k = \{0, 1, \dots, 2^j - 1\}$ the sets $Q_{j,k}$ are mutually disjoint,
- 3) $|Q_{j,k}| = 2^{-j}$ for all k ,
- 4) the 2-adic valuation is constant over $Q_{j,k}$.

The verifications are easy and left to the reader.

With the Littlewood-Paley decomposition given in (6), we obtain the following equivalence for the Lebesgue spaces $L^p(\mathbb{Z}_2)$ with $1 < p < +\infty$:

$$\|f\|_{L^p} \simeq \|S_0(f)\|_{L^p} + \left\| \left(\sum_{j \in \mathbb{N}} |\Delta_j f|^2 \right)^{1/2} \right\|_{L^p}.$$

See the book [10], chapter IV, for a general proof.

Let us turn now to smoothness measuring spaces. As said in the introduction, it is not absolutely trivial to define Sobolev and Besov spaces over \mathbb{Z}_2 since we are working in a totally discontinuous setting. Here is an example of this situation with the Sobolev space $W^{1,2}$: one could try to define the quantity $|\nabla f|$ by the formula

$$|\nabla f| = \lim_{\delta \rightarrow 0} \sup_{d(x,y) < \delta} \frac{|f(x) - f(y)|}{d(x,y)}$$

and define the Sobolev space $W^{1,2}(\mathbb{Z}_2)$ by the norm

$$\|f\|_* = \|f\|_{L^2} + \left(\int_{\mathbb{Z}_2} |\nabla f|^2 dx \right)^{1/2}. \quad (8)$$

Now, using the Littlewood-Paley decomposition we can also write

$$\|f\|_{**} = \|S_0 f\|_{L^2} + \left\| \left(\sum_{j \in \mathbb{N}} 2^{2j} |\Delta_j f|^2 \right)^{1/2} \right\|_2.$$

However, the quantities $\|\cdot\|_*$ and $\|\cdot\|_{**}$ are not equivalent: in the case of (8) consider a function $f = c_k$ constant over each $Q_{j,k} = \{k + 2^j \mathbb{Z}_2\}$ for some fixed j . Then we have $|\nabla f| \equiv 0$ and for these functions the norm $\|\cdot\|_*$ would be equal to the L^2 norm.

This is the reason why we will use in this article the Littlewood-Paley approach to characterize Sobolev spaces:

$$\|f\|_{W^{s,p}} \simeq \|S_0 f\|_{L^p} + \left\| \left(\sum_{j \in \mathbb{N}} 2^{2js} |\Delta_j f|^2 \right)^{1/2} \right\|_{L^p}. \quad (9)$$

with $1 < p < +\infty$ and $s > 0$. For Besov spaces we will define them by the norm

$$\|f\|_{B_p^{s,q}} \simeq \|S_0 f\|_{L^p} + \left(\sum_{j \in \mathbb{N}} 2^{jsq} \|\Delta_j f\|_{L^p}^q \right)^{1/q} \quad (10)$$

where $s \in \mathbb{R}$, $1 \leq p, q < +\infty$ with the necessary modifications when $p, q = +\infty$.

Remark 1 For homogeneous functional spaces $\dot{W}^{s,p}$ and $\dot{B}_p^{s,q}$, we drop out the term $\|S_0 f\|_{L^p}$ in (9) and (10).

Let us give some simple examples of function belonging to these functional spaces.

- 1) The function $f(x) = \log_2 |x|_2$ is in $\dot{B}_1^{1,\infty}(\mathbb{Z}_2)$. First note that $|x|_2 = 2^{-\gamma(x)}$ and thus $f(x) = -\gamma(x)$. Recall (cf. proposition 3.1) that over each set $Q_{j,k}$, the quantity $\gamma(x)$ is constant, so the dyadic bloc $\Delta_j f$ is given by

$$\Delta_j f(x) = \begin{cases} -1 & \text{over } Q_{j+1,0} \\ 0 & \text{elsewhere.} \end{cases}$$

Hence, taking the L^1 norm, we have $\|\Delta_j f\|_{L^1} = \frac{1}{2} 2^{-j}$ and then $f \in \dot{B}_1^{1,\infty}(\mathbb{Z}_2)$.

- 2) Set $h(x) = 1/|x|_2$, we have $h \in \dot{B}_\infty^{-1,\infty}$. For this, we must verify $\sup_{j \geq 0} 2^{-j} \|\Delta_j h\|_{L^\infty} < +\infty$. By definition we obtain $h(x) = 2^{\gamma(x)}$ and then

$$\Delta_j h(x) = \begin{cases} 2^j & \text{over } Q_{j+1,0} \\ 0 & \text{elsewhere.} \end{cases}$$

We finally obtain $\|\Delta_j h\|_{L^\infty} = 2^j$ and hence $2^{-j} \|\Delta_j h\|_{L^\infty} = 1$ for all j , so we write $h \in \dot{B}_\infty^{-1,\infty}$.

With the Littlewood-Paley characterisation of Sobolev spaces and Besov spaces given in (9) and (10) we have the following theorem:

Theorem 3 *In the framework of the 2-adic group \mathbb{Z}_2 we have, for a function f such that $f \in \dot{W}^{s_1,p}(\mathbb{Z}_2)$ and $f \in \dot{B}_\infty^{-\beta,\infty}(\mathbb{Z}_2)$, the inequality*

$$\|f\|_{\dot{W}^{s,q}} \leq C \|f\|_{\dot{W}^{s_1,p}}^\theta \|f\|_{\dot{B}_\infty^{-\beta,\infty}}^{1-\theta}$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s = \theta s_1 - (1 - \theta)\beta$ and $-\beta < s < s_1$.

Proof. We start with an interpolation result: let $(a_j)_{j \in \mathbb{N}}$ be a sequence, let $s = \theta s_1 - (1 - \theta)\beta$ with $\theta = p/q$, then we have for $r, r_1, r_2 \in [1, +\infty]$ the estimate

$$\|2^{js} a_j\|_{\ell^r} \leq C \|2^{js_1} a_j\|_{\ell^{r_1}}^\theta \|2^{-j\beta} a_j\|_{\ell^{r_2}}^{1-\theta}$$

See [2] for a proof. Apply this estimate to the dyadic blocks $\Delta_j f$ to obtain

$$\left(\sum_{j \in \mathbb{Z}} 2^{2js} |\Delta_j f(x)|^2 \right)^{1/2} \leq C \left(\sum_{j \in \mathbb{Z}} 2^{2js_1} |\Delta_j f(x)|^2 \right)^{\theta/2} \left(\sup_{j \in \mathbb{Z}} 2^{-j\beta} |\Delta_j f(x)| \right)^{1-\theta}$$

To finish, compute the L^q norm of the preceding quantities. ■

4 The $BV(\mathbb{Z}_2)$ space and the proof of theorem 1

We study in this section the space of functions of bounded variation BV and we will prove some surprising facts in the framework of 2-adic group \mathbb{Z}_2 . Let us start recalling the definition of this space:

Definition 4.1 *If f is a real-valued function over \mathbb{Z}_2 , we will say that $f \in BV(\mathbb{Z}_2)$ if there exists a constant $C > 0$ such that*

$$\int_{\mathbb{Z}_2} |f(x+y) - f(x)| dx \leq C |y|_2, \quad (\forall y \in \mathbb{Z}_2). \quad (11)$$

We prove now the theorem 1 which asserts that in \mathbb{Z}_2 , the BV space can be identified to the Besov space $\dot{B}_1^{1,\infty}$. For this, we will use two steps given by the propositions 4.1 and 4.2 below.

Proposition 4.1 *If f is a real-valued function over \mathbb{Z}_2 belonging to the Besov space $\dot{B}_1^{1,\infty}$, then $f \in BV$ and we have the inclusion $\dot{B}_1^{1,\infty} \subseteq BV$.*

Proof. Let $f \in \dot{B}_1^{1,\infty}(\mathbb{Z}_2)$ and let us fix $|y|_2 = 2^{-m}$. We have to prove the following estimation for all $m > 0$

$$I = \int_{\mathbb{Z}_2} |f(x+y) - f(x)| dx \leq C 2^{-m}.$$

Using the Littlewood-Paley decomposition given in (6), we will work on the formula below

$$I = \left\| \left(S_0 f(x+y) + \sum_{j \geq 0} \Delta_j f(x+y) \right) - \left(S_0 f(x) + \sum_{j \geq 0} \Delta_j f(x) \right) \right\|_{L^1}$$

Then, by the dyadic block's properties we have to study

$$I \leq \|S_m f(x+y) - S_m f(x)\|_{L^1} + \sum_{j=m+1}^{+\infty} \|\Delta_j f(x+y) - \Delta_j f(x)\|_{L^1}. \quad (12)$$

We estimate this inequality with the two following lemmas.

Lemma 4.1 *The first term in (12) is identically zero.*

Proof. Since we have fixed $|y|_2 = 2^{-m}$, then for $x \in Q_{m,k}$, we have $x+y \in Q_{m,k}$ with $k = \{0, \dots, 2^m - 1\}$. Applying the operators S_m to the functions $f(x+y)$ and $f(x)$ we get the desired result. ■

The second term in (12) is treated by the next lemma.

Lemma 4.2 *Under the hypothesis of proposition 4.1 and for $|y|_2 = 2^{-m}$ we have*

$$\sum_{j=m+1}^{+\infty} \|\Delta_j f(x+y) - \Delta_j f(x)\|_{L^1} \leq C 2^{-m}.$$

Proof. Indeed,

$$\sum_{j=m+1}^{+\infty} \|\Delta_j f(x+y) - \Delta_j f(x)\|_{L^1} \leq 2 \sum_{j=m+1}^{+\infty} \|\Delta_j f\|_{L^1}.$$

We use now the fact $\|\Delta_j f\|_{L^1} \leq C 2^{-j}$ for all j , since $f \in \dot{B}_1^{1,\infty}$, to get

$$\sum_{j=m+1}^{+\infty} \|\Delta_j f(x+y) - \Delta_j f(x)\|_{L^1} \leq C 2^{-m}.$$

With these two lemmas, and getting back to (12), we deduce the following inequality for all $y \in \mathbb{Z}_2$:

$$\int_{\mathbb{Z}_2} |f(x+y) - f(x)| dx \leq C |y|_2$$

and this concludes the proof of proposition 4.1. ■

Our second step in order to prove theorem 1 is the next result.

Proposition 4.2 *In \mathbb{Z}_2 we have the inclusion $BV(\mathbb{Z}_2) \subseteq \dot{B}_1^{1,\infty}(\mathbb{Z}_2)$.*

Proof. Observe that we can characterize the Besov space $\dot{B}_1^{1,\infty}(\mathbb{Z}_2)$ by the condition

$$\|f(\cdot + y) + f(\cdot - y) - 2f(\cdot)\|_{L^1} \leq C |y|_2, \quad \forall y \neq 0.$$

Let f be a function in $BV(\mathbb{Z}_2)$, then we have

$$\|f(\cdot + y) - f(\cdot)\|_{L^1} \leq C |y|_2.$$

Summing $\|f(\cdot - y) - f(\cdot)\|_{L^1}$ in both sides of the previous inequality we obtain

$$\|f(\cdot + y) - f(\cdot)\|_{L^1} + \|f(\cdot - y) - f(\cdot)\|_{L^1} \leq C |y|_2 + \|f(\cdot - y) - f(\cdot)\|_{L^1}$$

and by the triangular inequality we have

$$\|f(\cdot + y) + f(\cdot - y) - 2f(\cdot)\|_{L^1} \leq C |y|_2 + \|f(\cdot - y) - f(\cdot)\|_{L^1}$$

We thus obtain

$$\|f(\cdot + y) + f(\cdot - y) - 2f(\cdot)\|_{L^1} \leq 2C |y|_2.$$

We have proved, in the setting of the 2-adic group \mathbb{Z}_2 , the inequalities ■

$$C_1 \|f\|_{\dot{B}_1^{1,\infty}} \leq \|f\|_{BV} \leq C_2 \|f\|_{\dot{B}_1^{1,\infty}},$$

so the theorem 1 follows. ■

5 Improved Sobolev inequalities, BV space and proof of theorem 2

We do not give here a global treatment of the family of inequalities of type (2); instead we focus on the next inequality

$$\|f\|_{L^2}^2 \leq C \|f\|_{BV} \|f\|_{\dot{B}_\infty^{-1,\infty}} \quad (13)$$

and we want to know if this estimation is true in a 2-adic framework. Since in the \mathbb{Z}_2 setting we have the identification $\|f\|_{BV} \simeq \|f\|_{\dot{B}_\infty^{1,\infty}}$, the estimation (13) becomes

$$\|f\|_{L^2}^2 \leq C \|f\|_{\dot{B}_1^{1,\infty}} \|f\|_{\dot{B}_\infty^{-1,\infty}}. \quad (14)$$

This remark lead us to the theorem 2 which states that the previous inequalities are false.

Proof. We will construct a counterexample by means of the Littlewood-Paley decomposition, so it is worth to recall very briefly the dyadic bloc characterization of the norms involved in inequality (14). For the L^2 norm we have $\|f\|_{L^2}^2 = \sum_{j \in \mathbb{N}} \|\Delta_j f\|_{L^2}^2$, while for the Besov spaces $\dot{B}_1^{1,\infty}$ and $\dot{B}_\infty^{-1,\infty}$ we have

$$\begin{aligned} \|f\|_{\dot{B}_1^{1,\infty}} &= \sup_{j \in \mathbb{N}} 2^j \|\Delta_j f\|_{L^1} \quad \text{and} \\ \|f\|_{\dot{B}_\infty^{-1,\infty}} &= \sup_{j \in \mathbb{N}} 2^{-j} \|\Delta_j f\|_{L^\infty}. \end{aligned}$$

We construct a function $f : \mathbb{Z}_2 \rightarrow \mathbb{R}$ by considering his values over the dyadic blocs and we will use for this the sets $Q_{j,k}$ defined in (7). First fix α and β two non negative real numbers and j_0, j_1 two integers such that $0 \leq j_0 \leq j_1$ with the condition

$$2^{2j_0} \leq \frac{\beta}{\alpha}.$$

Now define N_j as a function of α and β :

$$N_j = 2^j \quad \text{if } 0 \leq j \leq j_0 \quad \text{and} \quad N_j = \frac{\beta}{\alpha} 2^{-j} \leq 2^j \quad \text{if } j_0 < j \leq j_1. \quad (15)$$

and write

$$\Delta_j f(x) = \begin{cases} \alpha 2^j & \text{over } Q_{j+1,0}, \\ -\alpha 2^j & \text{over } Q_{j+1,1}, \\ \alpha 2^j & \text{over } Q_{j+1,2}, \\ -\alpha 2^j & \text{over } Q_{j+1,3}, \\ \vdots & \\ \alpha 2^j & \text{over } Q_{j+1,2N_j-2}, \\ -\alpha 2^j & \text{over } Q_{j+1,2N_j-1}, \\ 0 & \text{elsewhere.} \end{cases}$$

Once this function is fixed, we compute the following norms

- $\|\Delta_j f\|_{L^1} = \sum_{k=0}^{N_j} \alpha 2^j 2^{-j} = \alpha N_j,$
- $\|\Delta_j f\|_{L^\infty} = \alpha 2^j,$
- $\|\Delta_j f\|_{L^2}^2 = \sum_{k=0}^{N_j} \alpha^2 2^{2j} 2^{-j} = \alpha^2 2^j N_j,$

and we build from these quantities the Besov and Lebesgue norms in the following manner:

- 1) For the Besov space $\dot{B}_\infty^{-1,\infty}$:

$$\|f\|_{\dot{B}_\infty^{-1,\infty}} = \sup_{0 \leq j \leq j_1} 2^{-j} \alpha 2^j = \alpha,$$

2) For the Besov space $\dot{B}_1^{1,\infty}$:

By the definition (15) of N_j we have $2^j \|\Delta_j f\|_{L^1} = 2^j \alpha N_j = 2^{2j} \alpha$ if $0 \leq j \leq j_0$ and $2^j \|\Delta_j f\|_{L^1} = \beta$ if $j_0 < j \leq j_1$. Since $2^{2j_0} \leq \frac{\beta}{\alpha}$ we have:

$$\|f\|_{\dot{B}_1^{1,\infty}} = \beta.$$

3) For the Lebesgue space L^2 :

$$\begin{aligned} \|f\|_{L^2}^2 &= \sum_{j=0}^{j_1} \alpha^2 2^j N_j = \sum_{j=0}^{j_0} \alpha^2 2^{2j} + \sum_{j>j_0}^{j_1} \alpha^2 2^j \frac{\beta}{\alpha} 2^{-j} = \sum_{j=0}^{j_0} \alpha^2 2^{2j} + (j_1 - j_0) \alpha \beta \\ &= \alpha \beta \left(\frac{\alpha}{\beta} \sum_{j=0}^{j_0} 2^{2j} + (j_1 - j_0) \right). \end{aligned}$$

With the condition $2^{2j_0} \leq \frac{\beta}{\alpha}$, we obtain from the previous formula that

$$\|f\|_{L^2}^2 \simeq \alpha \beta (j_1 - j_0) = \|f\|_{\dot{B}_1^{1,\infty}} \|f\|_{\dot{B}_\infty^{-1,\infty}} (j_1 - j_0).$$

Thus, getting back to (14) and therefore to (13), we have for an universal constant C the inequality

$$\|f\|_{\dot{B}_1^{1,\infty}} \|f\|_{\dot{B}_\infty^{-1,\infty}} (j_1 - j_0) \leq C \|f\|_{\dot{B}_1^{1,\infty}} \|f\|_{\dot{B}_\infty^{-1,\infty}}$$

$$\iff (j_1 - j_0) \leq C,$$

which is false since we can freely choose the values of j_1 and j_0 . The theorem 2 is proved. ■

References

- [1] Yvette AMICE. *Les nombres p-adiques*. PUF, 1975.
- [2] J. BERGH & J. LÖFSTRÖM. *Interpolation Spaces*. Grundlehren der mathematischen Wissenschaften, 223. Springer Verlag (1976).
- [3] D. CHAMORRO. *Improved Sobolev Inequalities and Muckenhoupt weights on stratified Lie groups*, <http://arxiv.org/abs/1007.4086>, (2010).
- [4] D. CHAMORRO. *Some functional inequalities on polynomial volume growth Lie groups*, <http://arxiv.org/abs/1009.3106>, (2010).
- [5] A. COHEN, W. DAHMEN, I. DAUBECHIES & R. DE VORE. *Harmonic Analysis of the space BV*. Rev. Mat. Iberoamericana 19, n°1, 235-263 (2003).
- [6] P. GÉRARD, Y. MEYER & F. ORU. *Inégalités de Sobolev Précisées*. Equations aux Dérivées Partielles, Séminaire de l'Ecole Polytechnique, exposé n° IV (1996-1997).
- [7] L. GRAFAKOS. *Classical and Modern Fourier Analysis*. Prentice Hall (2004).
- [8] N. KOBLITZ, *p-adic Numbers, p-adic Analysis and Zeta-functions*. GTM 58. Springer Verlag, 1977.
- [9] M. LEDOUX. *On improved Sobolev embedding theorems*. Math. Res. Letters 10, 659-669 (2003).
- [10] E. M. STEIN. *Topics in Harmonic analysis*. Annals of mathematics studies, 63. Princeton University Press (1970).

- [11] E. M. STEIN. *Harmonic Analysis*. Princeton University Press (1993).
- [12] H. TRIEBEL. *Theory of function spaces II*, Birkhäuser (1992).
- [13] V. S. VLADIMIROV, I. V. VOLOVICH and E. I. ZELENOV. *p-Adic Analysis and Mathematical Physics*, World Scientific, Singapore (1994).

Diego CHAMORRO

Laboratoire d'Analyse et de Probabilités
Université d'Evry Val d'Essonne & ENSIIE

1 square de la résistance,
91025 Evry Cedex

diego.chamorro@m4x.org