



HAL
open science

Locally tame plane polynomial automorphisms

Joost Berson, Adrien Dubouloz, Jean-Philippe Furter, Stefan Maubach

► **To cite this version:**

Joost Berson, Adrien Dubouloz, Jean-Philippe Furter, Stefan Maubach. Locally tame plane polynomial automorphisms. 2010. hal-00531776

HAL Id: hal-00531776

<https://hal.science/hal-00531776v1>

Preprint submitted on 3 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LOCALLY TAME PLANE POLYNOMIAL AUTOMORPHISMS

JOOST BERSON, ADRIEN DUBOULOZ, JEAN-PHILIPPE FURTER, AND STEFAN MAUBACH

ABSTRACT. For automorphisms of a polynomial ring in two variables over a domain R , we show that local tameness implies global tameness provided that every 2-generated locally free R -module of rank 1 is free. We give many examples illustrating this property.

INTRODUCTION

A natural problem in commutative algebra and algebraic geometry is to understand the group $\mathrm{GA}_n(R)$ of automorphisms of a polynomial ring $R[X_1, \dots, X_n]$ over a ring R . Although much progress has been made in this direction during the last decades, one can state that only the case $n = 2$ and R is a field is fully understood. A central and fruitful notion in the study of polynomial automorphisms is the notion of tameness: an automorphism is called *tame* if it can be written as a composition of affine and triangular ones, where by a triangular automorphism, we mean an automorphism $F = (F_1, \dots, F_n) \in \mathrm{GA}_n(R)$ such that $F_i \in R[X_i, \dots, X_n]$ for every $i = 1, \dots, n$. Tame automorphisms form a subgroup $\mathrm{TA}_n(R)$ of $\mathrm{GA}_n(R)$ and a classical theorem due to Jung in characteristic zero [8] and van der Kulk in the general case [9] asserts that if R is a field k then $\mathrm{GA}_2(k) = \mathrm{TA}_2(k)$. The result is even more precise: $\mathrm{GA}_2(k)$ is the free product of the subgroups of affine and triangular automorphisms amalgamated over their intersection. In contrast, even the equality $\mathrm{TA}_2(R) = \mathrm{GA}_2(R)$ is no longer true for a general domain R , as illustrated by a famous example due to Nagata : for an element $z \in R \setminus \{0\}$ the endomorphism

$$F = (X - 2Y(zx + Y^2) - z(zX + Y^2)^2, Y + z(zX + Y^2))$$

of $R[X, Y]$ is in $\mathrm{GA}_2(R)$ and can be decomposed as $F = (X - z^{-1}Y^2, Y)(X, z^2X + Y)(X + z^{-1}Y^2, Y)$ in $\mathrm{GA}_2(K(R)) = \mathrm{TA}_2(K(R))$. Such a decomposition being essentially unique, this implies in particular that if z is not invertible in R , then F cannot be tame over R . Note that more generally, given a prime ideal $\mathfrak{p} \in \mathrm{Spec}(R)$, $F \in \mathrm{TA}_2(R_{\mathfrak{p}})$ if and only if $z \notin \mathfrak{p}$.

Automorphisms $F \in \mathrm{GA}_n(R)$ such that $F \in \mathrm{TA}_n(R_{\mathfrak{p}})$ for every $\mathfrak{p} \in \mathrm{Spec}(R)$ are said to be *locally tame*. Of course, every tame automorphism is locally tame. In contrast, the Nagata automorphism is neither tame nor locally tame. This could suggest that, at least for plane polynomial automorphisms, tameness is a property that can be checked locally on the base ring. In particular, one could hope that the only reason why an automorphism $F \in \mathrm{GA}_2(R)$ is not tame is because there exists a prime $\mathfrak{p} \in \mathrm{Spec}(R)$ such that F is already non tame over $R_{\mathfrak{p}}$. It turns out that this hope is too optimistic, and that in general, some “global” properties of R have to be taken into account to be able to infer tameness directly from local tameness. The main result of this article is the following characterization of rings for which global tameness can be checked locally :

Theorem. *For a domain R , the following assertions are equivalent :*

- 1) $\mathrm{TA}_2(R) = \bigcap_{\mathfrak{p} \in \mathrm{Spec}(R)} \mathrm{TA}_2(R_{\mathfrak{p}})$,
- 2) *Every 2-generated locally free R -module of rank 1 is free.*

In particular, it follows that over a unique factorization domain R , tameness is a local property of automorphisms.

The article is organized as follows. Section one is devoted to the proof of the above characterization, that we essentially derive from the fact that tame automorphisms of a polynomial ring in two variables can be recognized algorithmically. In section two, we consider many examples that illustrate condition 2) in the Theorem above.

Funded by a free competition grant of the Netherlands Organisation for scientific research (NWO).

Partially supported by FABER grant 07-512-AA-010-S-179 and PHC Grant Van Gogh 18153NA.

Partially supported by PHC Grant Van Gogh 18153NA.

Funded by Veni-grant of council for the physical sciences, Netherlands Organisation for scientific research (NWO).

1. FROM LOCAL TAMENESS TO GLOBAL TAMENESS

In this section, we characterize domains R with the property that an automorphism $F = (F_1, F_2) \in \text{GA}_2(R)$ is tame if and only if it is locally tame.

1.1. Notations. For an automorphism $F = (F_1, F_2) \in \text{GA}_2(R)$, we let $\deg F = (\deg F_1, \deg F_2) \in (\mathbb{N}^*)^2$ considered as equipped with the product order. We denote by \overline{F}_i the homogeneous component of F_i of degree $\deg F_i$, $i = 1, 2$. An automorphism with $\deg F = (1, 1)$ is affine, and we denote by $\text{Aff}_2(R)$ the corresponding subgroup of $\text{GA}_2(R)$.

1.1. Properties of automorphisms.

Even if the equality $\text{GA}_2(R) = \text{TA}_2(R)$ is no longer true for a general domain R , tame automorphisms of a polynomial ring in two variables can be recognized algorithmically. Indeed, the following result quoted from [5, Prop. 1] (see also [4, Cor. 5.1.6]) says in essence that for every $F \in \text{TA}_2(R)$ with $\deg F > (1, 1)$ there exists a linear or a triangular automorphism φ such that $\deg \varphi F < \deg F$.

Proposition 1. *Let $F = (F_1, F_2) \in \text{TA}_2(R)$ and let $(d_1, d_2) = \deg F$. Then the following holds:*

a) $d_1 \mid d_2$ or $d_2 \mid d_1$.

b) If $\max(d_1, d_2) > 1$ then we have:

(i) If $d_1 < d_2$ then $\overline{F}_2 = c\overline{F}_1^{d_2/d_1}$ for some $c \in R$,

(ii) If $d_2 < d_1$ then $\overline{F}_1 = c\overline{F}_2^{d_1/d_2}$ for some $c \in R$,

(iii) If $d_1 = d_2$ then there exists $\varphi \in \text{Aff}_2(R)$ such that $\varphi F = (F'_1, F'_2)$ satisfies $\deg F'_1 = d_1$ and $\deg F'_2 < d_1$.

1.2. In contrast to the tame case, for an arbitrary automorphism $F = (F_1, F_2) \in \text{GA}_2(R)$ with $\deg F_1 = \deg F_2$ there is no guarantee in general that there exists $\varphi \in \text{Aff}_2(R)$ such that $\deg \varphi F < \deg F$. Indeed, such a φ exists if and only there exists a unimodular vector $(\alpha_1, \alpha_2) \in R^2$ such that $\alpha_1 \overline{F}_1 + \alpha_2 \overline{F}_2 = 0$, which is the case if and only if the R -module $R\overline{F}_1 + R\overline{F}_2$ is free of rank 1. Combined with [4, Ex. 6 p. 94], this observation leads to a natural procedure to construct families of locally tame but not (globally) tame automorphisms, namely:

Proposition 2. *If $z, w \in R$ and $q(T) \in R[T]$ is a polynomial of degree at least 2, then*

$$F := (X + wq(zX + wY), Y - zq(zX + wY))$$

is an element of $\text{GA}_2(R)$. Furthermore F is tame if and only if (z, w) is a principal ideal of R .

In particular, if (z, w) is a locally principal but not principal ideal, then F is a locally tame but not globally tame automorphism.

Proof. A straightforward verification shows that $G = (X - wq(zX + wY), Y + zq(zX + wY))$ is an inverse for F . Suppose that $(z, w) = aR$ for some $a \in R$. Replacing $q(T)$, z and w by $aq(aT)$, $a^{-1}z$ and $a^{-1}w$ respectively, we may assume that $(z, w) = R$. But then if we take any $\varphi \in \text{SL}_2(R)$ having $zX + wY$ as its first component, one checks that $F = \varphi^{-1}(X, Y - q(X))\varphi \in \text{TA}_2(R)$. Conversely, if $F \in \text{TA}_2(R)$, then, since $\deg F_1 = \deg F_2 = \deg Q > 1$, it follows from Proposition 1 and the above discussion that the R -module generated by $\overline{F}_1 = wq(zX + wY)$ and $\overline{F}_2 = -zq(zX + wY)$ is free of rank 1. Simplifying by $\overline{q}(zX + wY)$, we get that the R -module generated by w and $-z$ is free a rank 1, i.e., $(w, -z)$ is a principal ideal. \square

1.3. It follows that locally tame but not globally tame automorphisms abound : for instance, in the proposition above, one can take for R the coordinate ring of a smooth non rational affine curve C and for z, w a pair of generators of the defining ideal of a non principal Weil divisor on C (see also section 2 below for more examples).

1.2. A criterion.

It turns out that the examples discussed above illustrate the only global obstruction to infer global tameness from local tameness, namely, the existence of 2-generated locally free but not globally free modules of rank 1. Indeed, we have the following criterion.

Theorem 3. *For a domain R , the following assertions are equivalent:*

1) $\text{TA}_2(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \text{TA}_2(R_{\mathfrak{p}})$,

2) Every 2-generated locally free R -module of rank 1 is free.

Proof. 1) \Rightarrow 2). Since R is a domain, every locally free R -module of rank 1 is isomorphic to an R -submodule of the field of fractions $K(R)$ of R (see e.g. [7, Prop. 6.15]). In turn, every such submodule is isomorphic to an ideal of R . In particular, if there exists a locally free but non free 2-generated R -module of rank 1, then there exists locally principal but not principal ideal (z, w) of R . But then any $F \in \text{GA}_2(R)$ as in Proposition 2 above is locally tame but not tame.

2) \Rightarrow 1). Conversely, for any domain R , it is clear that

$$\text{TA}_2(R) \subseteq \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \text{TA}_2(R_{\mathfrak{p}}) \subseteq \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \text{GA}_2(R_{\mathfrak{p}}) = \text{GA}_2(R).$$

Let $F = (F_1, F_2) \in \text{GA}_2(R)$ be a locally tame automorphism and let $d_i = \deg F_i$, $i = 1, 2$. We may assume that $d_1 \leq d_2$. If $d_1 = d_2 = 1$ then F is affine, whence tame. We now proceed by induction on (d_1, d_2) , assuming that every locally tame automorphism of degree $(d'_1, d'_2) < (d_1, d_2)$ is globally tame.

• Case 1 : $d_1 < d_2$. Since $F \in \text{TA}_2(R_{(0)}) = \text{TA}_2(K(R))$, it follows from Proposition 1 that $e = d_2/d_1 \in \mathbb{N}^*$ and that there exists $\alpha \in K(R)$ such that $\overline{F_2} = \alpha \overline{F_1}^e$. But since $F \in \text{TA}_2(R_{\mathfrak{p}})$ for every $\mathfrak{p} \in \text{Spec}(R)$, it follows that

$$\alpha \in \bigcap_{\mathfrak{p} \in \text{Spec}(R)} R_{\mathfrak{p}} = R.$$

Now, the automorphism $(X, Y - \alpha X^e)F$ satisfies the induction hypothesis and we are done with case.

• Case 2 : $d_1 = d_2$. Since for any $\mathfrak{p} \in \text{Spec}(R)$, we have $F \in \text{TA}_2(R_{\mathfrak{p}})$, it follows from Proposition 1 and the discussion 1.2 that for every $\mathfrak{p} \in \text{Spec}(R)$, the $R_{\mathfrak{p}}$ module generated by $\overline{F_1}$ and $\overline{F_2}$ is free of rank 1. This means exactly that the R -module generated by $\overline{F_1}$ and $\overline{F_2}$ is locally free of rank 1. Our assumption implies that it is globally free, and so, we deduce from 1.2 that there exist $\varphi \in \text{Aff}_2(R)$ such that $\deg \varphi F < \deg F$. \square

1.4. Recall that the *Picard group* of a ring R is the group $\text{Pic}(R)$ of isomorphism classes of locally free R -modules of rank 1. In view of the above criterion, it is natural to introduce the subgroup $\text{Pic}_2(R)$ of $\text{Pic}(R)$ generated by isomorphism classes of locally free R -modules of rank 1 that can be generated by 2 elements. With this definition, property 2) in Theorem 3 is equivalent to the triviality of $\text{Pic}_2(R)$. In particular, we obtain:

Corollary 4. *If $\text{Pic}_2(R) = \{1\}$ and F belongs to $\text{GA}_2(R)$, then F is tame if and only if it is locally tame.*

Example 5. The class of rings with $\text{Pic}_2(R) = \{1\}$ contains in particular unique factorization domains since for these domains the Picard group itself is trivial. This also holds for *Bézout rings*, that is, domains in which every finitely generated ideal is principal (see e.g. [2]).

1.3. Minimal overring for tameness.

Recall that $\text{GA}_2(R) = \text{TA}_2(R)$ if and only if R is a field [4, Proposition 5.1.9]. If $F \in \text{GA}_2(R)$, then F is tame over the field of fractions K of R , but, in general, there does not exist a smallest ring S between R and K such that F is tame over S . Indeed, letting $R = \mathbb{C}[z, w]$ every automorphism F as in Proposition 2 is tame over $R[z^{-1}]$ and $R[w^{-1}]$ but not over $R = R[z^{-1}] \cap R[w^{-1}]$. However, if we further assume that R is a Bézout domain, we have the following result.

Proposition 6. *Let R be a Bézout domain and let $(R_j)_{j \in J}$ be a family of rings between R and K such that $R = \bigcap_{j \in J} R_j$. Then $\text{TA}_2(R) = \bigcap_{j \in J} \text{TA}_2(R_j)$.*

Proof. Similarly as in the proof of Theorem 3, we proceed by induction on the degree of $F = (F_1, F_2) \in \text{GA}_2(R) \cap \bigcap_{j \in J} \text{TA}_2(R_j)$, the case $\deg F = (1, 1)$ being obvious. Letting $d_i = \deg F_i$, we may assume that

$d_1 \leq d_2$.

• Case 1 : $d_1 < d_2$. Then $e = d_2/d_1 \in \mathbb{N}^*$ and there exists $\alpha \in K$ such that $\overline{F_2} = \alpha \overline{F_1}^e$. Since $F \in \text{TA}_2(R_j)$, we have $\alpha \in R_j$ for every $j \in J$, and so $\alpha \in R = \bigcap_{j \in J} R_j$. Now the automorphism

$(X, Y - \alpha X^e)F$ satisfies the induction hypothesis.

• Case 2 : $d_1 = d_2$. Since $\overline{F_1}$ and $\overline{F_2}$ are K -linearly dependent, the R -module $R\overline{F_1} + R\overline{F_2}$ is isomorphic to a proper ideal of R . As R is a Bézout domain, the latter is free of rank 1, and so, we conclude from 1.2 above that there exists $\varphi \in \text{Aff}_2(R)$ such that $\deg \varphi F < \deg F$. \square

Proposition 7. *If R is a Bézout domain and $F \in \text{GA}_2(R)$ then there exists a smallest ring S between R and $K(R)$ such that $F \in \text{TA}_2(S)$. Furthermore, S is a finitely generated R -algebra.*

If we assume further that R is a principal ideal domain, then there exists $r \in R \setminus \{0\}$ such that $S = R[r^{-1}]$.

Proof. Any ring between R and $K(R)$ is again a Bézout domain [2, Theorem 1.3]. Therefore, the existence of S is a consequence of the previous proposition. The fact that S is finitely generated follows from Proposition 1 by easy induction. For the last assertion, since S is finitely generated over R , there exists a finitely generated ideal $I \subset R$ and an element $r \in R \setminus \{0\}$ such that $S = R[I/r] = \{a/r^k \in K(R), a \in I^k, k = 0, 1 \dots\}$. Since R is a p.i.d, I is a principal ideal, say generated by an element $g \in R$. After eliminating common factors if any, we may assume that r and g are relatively prime and that $S = R[g/r] \subset R[r^{-1}]$. But by Bézout identity, there exists $u, v \in R$ such that $ur + vg = 1$ and so, $S = R[r^{-1}]$. \square

Example 8. If $F \in \text{GA}_2(\mathbb{C}[z])$, then there exists a smallest ring S between $\mathbb{C}[z]$ and $\mathbb{C}(z)$ of the form $\mathbb{C}[z][r^{-1}]$ such that $F \in \text{GA}_2(S)$.

2. EXAMPLES AND COMPLEMENTS

Here we discuss examples of domains R which illustrate the property $\text{Pic}_2(R) = \{1\}$.

2.1. The condition $\text{Pic}_2(\mathbf{R}) = \{1\}$ for 1-dimensional noetherian domains.

If R is a noetherian domain of Krull dimension 1, every locally free R -module of rank j is generated by at most $j + 1$ elements (see e.g. [10, Th. 5.7]). In particular, we have $\text{Pic}(R) = \text{Pic}_2(R)$ for every noetherian domain of dimension 1. As a consequence, we get:

Example 9. If R is a Dedekind domain, the following are equivalent :

- (1) $\text{Pic}_2(R) = \{1\}$; (2) $\text{Pic}(R) = \{1\}$; (3) R is a UFD; (4) R is a p.i.d.

For the coordinate ring R of an affine curve C defined over an algebraically closed field, we have the following classical result:

Proposition 10. *The Picard group of R is trivial if and only if C is a nonsingular rational curve.*

Proof. Let $\tilde{C} = \text{Spec}(\tilde{R})$ be the normalization of C . By virtue of [13, Theorem 3.2], the natural surjection $\text{Pic}(C) \rightarrow \text{Pic}(\tilde{C})$ is an isomorphism if and only if $R = \tilde{R}$. Therefore, if $\text{Pic}(C)$ is trivial, then C is necessarily a nonsingular curve. Now it is well known that a nonsingular curve has trivial Picard group if and only if it is rational (see e.g. [3, §11.4 p. 261]). \square

Corollary 11. *Let R be the coordinate ring of a rational affine curve and let \tilde{R} be its integral closure in $K(R)$. If $F \in \text{GA}_2(R)$ is locally tame, then $F \in \text{TA}_2(\tilde{R})$.*

Proof. Indeed, with the notation of the previous proof, one has $F \in \text{TA}_2(\mathcal{O}_p)$ for every $p \in C = \text{Spec}(R)$ and so $F \in \text{TA}_2(\tilde{\mathcal{O}}_p)$ for every $p \in C$. Since $\tilde{R} = \bigcap_{p \in C} \tilde{\mathcal{O}}_p$, it follows that F is locally tame over \tilde{R} , whence tame by virtue of Proposition 10. \square

Example 12. Let $R = \mathbb{C}[u, v] / (v^2 - u^3)$ be the coordinate ring of a cuspidal rational curve C . Via the homomorphism $\mathbb{C}[u, v] \rightarrow \mathbb{C}[t], (u, v) \mapsto (t^2, t^3)$ we may identify R with the subring $\mathbb{C}[t^2, t^3]$ of $\mathbb{C}[t]$ and the integral closure \tilde{R} of R with $\mathbb{C}[t]$. For every $a \in \mathbb{C}^*$, we let $I_a = (t^2 - a^2, t^3 - a^3)$ be the maximal ideal of the smooth point (a^2, a^3) of C . In particular, I_a is locally principal but one checks easily that it is not principal. So for $(z, w) = (t^2 - a^2, t^3 - a^3)$, any automorphism F as in Proposition 2 is locally tame but not tame. On the other hand, $I_a \tilde{R}$ is principal, generated by $t - a$, and so, $F \in \text{TA}_2(\mathbb{C}[t])$.

2.2. Examples of rings with $\text{Pic}_2(\mathbf{R}) = \{1\}$ but $\text{Pic}(\mathbf{R}) \neq \{1\}$.

As observed above, for 1-dimensional domains R , the triviality of $\text{Pic}_2(R)$ is equivalent to the one of $\text{Pic}(R)$. Here we give examples of domains with $\text{Pic}_2(R) = \{1\}$ and $\text{Pic}(R) \neq \{1\}$ which are coordinate rings of smooth affine algebraic varieties.

2.1. Let Q be a smooth quadric in the complex projective space $\mathbb{P}^n = \mathbb{P}_{\mathbb{C}}^n$, $n \geq 2$, and let $U = \mathbb{P}^n \setminus Q$. As is well-known, U is smooth affine variety with Picard group isomorphic to \mathbb{Z}_2 , generated by the restriction to U of the invertible sheaf $\mathcal{O}_{\mathbb{P}^n}(1)$ on \mathbb{P}^n . Letting $R_n = \Gamma(U, \mathcal{O}_U)$ and $M_n = \Gamma(U, \mathcal{O}_{\mathbb{P}^1}(1))$, which is a locally free R_n -module of rank 1, we have the following result.

Proposition 13. *The minimal number of generators of M_n as an R_n -module is $\lfloor n/2 \rfloor + 1$. In particular, if $n \geq 4$ then $\text{Pic}_2(R_n) = \{1\}$ whereas $\text{Pic}(R_n) \simeq \mathbb{Z}_2$.*

Proof. Up to the action of $\text{PGL}_{n+1}(\mathbb{C})$, we may assume that $Q \subset \mathbb{P}^n = \text{Proj}(\mathbb{C}[x_0, \dots, x_n])$ is the hypersurface $q = 0$, where $q = x_0^2 + \dots + x_n^2 \in \mathbb{C}[x_0, \dots, x_n]$. Letting $\mathcal{Q} \subset \mathbb{A}^{n+1} = \text{Spec}(\mathbb{C}[x_0, \dots, x_n])$ be the quadric defined by the equation $q = 1$, the natural map $\mathbb{A}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}^n$ restricts to an étale double cover $\mathcal{Q} \rightarrow \mathbb{P}^n \setminus Q$ expressing the coordinate ring R_n of $\mathbb{P}^n \setminus Q$ as the ring of invariant functions of $A = \mathbb{C}[x_0, \dots, x_n]/(q-1)$ for the \mathbb{Z}_2 -action induced by $-\text{id}$ on \mathbb{A}^{n+1} . With this description, $\mathcal{O}_{\mathbb{P}^n}(1)|_U$ coincides with the trivial line bundle $\mathcal{Q} \times \mathbb{A}^1$ equipped with the nontrivial \mathbb{Z}_2 -linearization $\mathcal{Q} \times \mathbb{A}^1 \ni (x, u) \mapsto (-x, -u) \in \mathcal{Q} \times \mathbb{A}^1$ (see e.g. [11, §1.3]). It follows that we may identify regular functions on U and global sections of $\mathcal{O}_{\mathbb{P}^n}(1)|_U$ with cosets in A of even and odd polynomial functions on \mathbb{A}^{n+1} respectively.

- Case 1 : $n = 2m$ is even. Clearly, the $m + 1$ odd polynomials $p_j = x_{2j} + ix_{2j+1}$ for $0 \leq j \leq m - 1$ and $p_m = x_{2m}$ have no common zero on \mathcal{Q} . Therefore, the corresponding sections of $\mathcal{O}_{\mathbb{P}^n}(1)|_U$ generate M_n as an R_n -module. Let us show that M_n cannot be generated by less than $m + 1$ elements. Otherwise, we could find in particular m odd polynomial functions s_1, \dots, s_m on \mathbb{A}^{n+1} with no common zero on \mathcal{Q} . Writing $s_j = a_j + ib_j$ for suitable odd polynomials $a_j, b_j \in \mathbb{R}[x_0, \dots, x_n]$, this would imply in particular that the n odd real polynomials a_1, \dots, a_m and b_1, \dots, b_m have no common zero on $\mathcal{Q} \cap \mathbb{R}^{n+1}$. This is impossible. Indeed, since $\mathcal{Q} \cap \mathbb{R}^{n+1}$ is the real n -sphere \mathbb{S}^n , it follows from Borsuk-Ulam theorem that the map $\phi = (a_1, \dots, a_m, b_1, \dots, b_m) : \mathbb{S}^n \rightarrow \mathbb{R}^n$ takes the same value on a pair of antipodal points, hence, being odd, vanishes on a pair of antipodal points.

- Case 2 : $n = 2m + 1$ is odd. One checks in a similar way as above that the $m + 1$ global sections of $\mathcal{O}_{\mathbb{P}^n}(1)|_U$ corresponding the odd polynomials $p_j = x_{2j} + ix_{2j+1}$, $0 \leq j \leq m$ generate M_n as an R_n -module. Now if M_n was generated by m elements, then there would exist m odd polynomials $s_j = a_j + ib_j$ as above for which the polynomials $a_j, b_j \in \mathbb{R}[x_0, \dots, x_n]$, $j = 1, \dots, m$ have no common zero on the real n -sphere \mathbb{S}^n . But then the continuous map $\phi = (a_1, \dots, a_m, b_1, \dots, b_m, 0) : \mathbb{S}^n = \mathcal{Q} \cap \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ would contradict the Borsuk-Ulam theorem. \square

Remark 14. An argument very similar to the one used in the proof above shows that over the subring \tilde{R}_n of $\mathbb{R}[x_0, \dots, x_n]/(x_0^2 + \dots + x_n^2 - 1)$ consisting of cosets of even polynomials, the module \tilde{M}_n consisting of cosets of odd polynomials cannot be generated by less than $n + 1$ elements. This property seems to have been first observed by Chase (unpublished). Our proof is deeply inspired by an argument due to Gilmer [6] on a slightly different example.

2.3. Further research.

One may wonder if there exists a complete characterization of obstructions to infer global tameness from local tameness for higher dimensional polynomial automorphisms similar to Theorem 3. A good starting point would be to have in general an effective algorithmic way to recognize tame automorphisms. Unfortunately, at the present time, such a higher dimensional algorithm only exists for automorphisms of a polynomial ring in 3 variables over a field [12].

REFERENCES

1. J. Berson, A. van den Essen, D. Wright, *Stable tameness of two-dimensional polynomial automorphisms over a regular ring*, preprint: arXiv:0707.3151v8 [math.AC], submitted to J. Amer. Math. Soc. April 2009.
2. P. M. Cohn, *Bezout rings and their subrings*, Proc. Cambridge Philos. Soc. 64 (1968), 251-264.
3. D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.
4. A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics 190 (329 pp), Birkhäuser Verlag, Basel, 2000.
5. J.-P. Furter, *On the variety of automorphisms of the affine plane*, J. Algebra 195 (1997), 604-623.
6. R. Gilmer, *A note on generating sets for invertible ideals*, Proc. Amer. Math. Soc. 22 (1969), 426-427.
7. R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics 52, Springer-Verlag, New York-Heidelberg, 1977.
8. H. Jung, *Über ganze birationale Transformationen der Ebene*, (German) J. Reine Angew. Math. 184, (1942). 161-174.
9. W. van der Kulk, *On polynomial rings in two variables*, Nieuw Arch. Wiskunde (3) 1, (1953). 33-41.
10. H. Matsumura, *Commutative ring theory*, translated from the Japanese by M. Reid, Cambridge Studies in Advanced Mathematics 8, Cambridge University Press (1986).
11. D. Mumford, J. Fogarty, F. Kirwan, *Geometric invariant theory. Third edition* (292 pp), Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], 34. Springer-Verlag, Berlin, 1994.
12. Ivan P. Shestakov and Ualbai U. Umirbaev, *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. 17 (2004), no. 1, 197-227
13. R. Wiegand, *Cancellation over commutative rings of dimension one and two*, J. Algebra 88 (1984), no. 2, 438-459.

RADBOUD UNIVERSITY NIJMEGEN, POSTBUS 9010, 6500 GL NIJMEGEN, THE NETHERLANDS
E-mail address: j.berson@science.ru.nl

INSTITUT DE MATHÉMATIQUES DE BOURGOGNE, UNIVERSITÉ DE BOURGOGNE, 9 AVENUE ALAIN SAVARY, BP 47870, 21078
DIJON CEDEX, FRANCE
E-mail address: Adrien.Dubouloz@u-bourgogne.fr

UNIVERSITÉ DE LA ROCHELLE, AVENUE MICHEL CRÉPEAU, 17000 LA ROCHELLE, FRANCE
E-mail address: jpfurter@univ-lr.fr

JACOBS UNIVERSITY BREMEN, CAMPUS RING 1, 28759 BREMEN, GERMANY
E-mail address: s.maubach@jacobs-university.de