



**HAL**  
open science

## **Viral causes of Influenza-Like Illness: Insights from a study during the winters 2004-2007**

Simona Puzelli, Catia Valdarchi, Marco Ciotti, Maria Dorrucchi, Francesca Farchi, Muhammed Babakir-Mina, Carlo Federico Perno, Isabella Donatelli, Giovanni Rezza

► **To cite this version:**

Simona Puzelli, Catia Valdarchi, Marco Ciotti, Maria Dorrucchi, Francesca Farchi, et al.. Viral causes of Influenza-Like Illness: Insights from a study during the winters 2004-2007. *Journal of Medical Virology*, 2009, 81 (12), pp.2066. 10.1002/jmv.21610 . hal-00531823

**HAL Id: hal-00531823**

**<https://hal.science/hal-00531823>**

Submitted on 4 Nov 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Viral causes of Influenza-Like Illness: Insights from a study during the winters 2004-2007

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-08-1151.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	23-Jun-2009
Complete List of Authors:	<p>Puzelli, Simona; Istituto Superiore di Sanità, Department of Infectious, Parasite and Immune-mediated Diseases          Valdarchi, Catia; Istituto Superiore di Sanità, Department of Infectious, Parasite and Immune-mediated Diseases          Ciotti, Marco; Policlinico Università Tor Vergata, Department of Molecular Virology          Dorrucchi, Maria; Istituto Superiore di Sanità,, Department of Infectious, Parasite and Immune-mediated Diseases          Farchi, Francesca; Istituto Superiore di Sanità,, Department of Infectious, Parasite and Immune-mediated Diseases          Babakir-Mina, Muhammed; Policlinico Università Tor Vergata, Department of Molecular Virology          Perno, Carlo Federico; Policlinico Università Tor Vergata, Department of Molecular Virology          Donatelli, Isabella; Istituto Superiore di Sanità,, Department of Infectious, Parasite and Immune-mediated Diseases          Rezza, Giovanni; Istituto Superiore di Sanità, Department of Infectious, Parasite and Immune-mediated Diseases</p>
Keywords:	ILI, Respiratory viruses, Influenza, Adenovirus, Italy



**Table 1.**

**Number of positive samples, by specific virus and month of diagnosis (the percentage is calculated from total number in a column)**

	<b>November (n=28)</b>	<b>December<sup>a</sup> (n=151)</b>	<b>January<sup>b</sup> (n=123)</b>	<b>February<sup>c</sup> (n=208)</b>	<b>March<sup>d</sup> (n=61)</b>	<b>April (n=9)</b>
	n. (%)	n. (%)	n. (%)	n. (%)	n. (%)	n. (%)
<b>Influenza virus</b>	1 (4)	12 (8)	37 (30)	65 (31)	13 (21)	4 (44)
<b>Adenovirus</b>	0 (0)	22 (15)	11 (9)	26 (13)	4 (6)	0 (0)
<b>PIV</b>	5 (18)	9 (6)	7 (6)	12 (6)	2 (3)	0 (0)
<b>RSV</b>	0 (0)	5 (3)	2 (2)	3 (1)	0 (0)	0 (0)
<b>Other/nd</b>	22 (79)	106 (70)	70 (57)	112 (55)	45 (74)	5 (56)

NOTE. <sup>a</sup> 3 coinfections (1 RSV + Adenovirus; 2 Adenovirus + PIV)

<sup>b</sup> 4 coinfections (1 Influenzavirus + Adenovirus; 3 Influenzavirus + PIV)

<sup>c</sup> 10 coinfections (8 Influenzavirus + Adenovirus; 1 RSV + Adenovirus; 1 Influenzavirus + PIV)

<sup>d</sup> 3 coinfections (3 Influenzavirus + Adenovirus)

**Table 2.**

**Number of positive samples, by specific virus and age class (the percentage is calculated from total number in a row)**

	<b>FLU</b>	<b>ADENO</b>	<b>PIV</b>	<b>RSV</b>	<b>OTHER/ND</b>
<b>Age (years)</b>	N (%)	N (%)	N (%)	N (%)	N (%)
<b>0 – 2 (63)</b>	20 (32)	13 (21)	3 (5)	3 (5)	29 (46)
<b>3 – 13 (161)</b>	42 (26)	19 (12)	10 (6)	3 (2)	94 (58)
<b>14 – 64 (319)</b>	69 (22)	27 (8)	20 (6)	4 (1)	207 (65)
<b>≥65 (37)</b>	1 (3)	4 (11)	2 (5)	0 (0)	30 (81)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60**Table 3.****Number of samples, by specific virus and sign and symptom.**

Sign/symptom	Influenza (n.116)		Adenovirus (n.47)		PIV (n.29)		RSV (n.8)		Other/nd (n.360)	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Sore throat	76	(66)	26	(55)	20	(69)	3	(38)	237	(66)
Nasal congestion	70	(60)	21	(45)	14	(48)	4	(50)	179	(48)
Muscle pain	63	(54)	10	(21)	5	(17)	1	(12)	134	(37)
Headache	53	(46)	15	(32)	13	(45)	1	(12)	141	(39)
Dry cough	57	(50)	21	(45)	12	(41)	2	(25)	183	(51)
Productive cough	41	(36)	17	(36)	6	(21)	7	(87)	110	(31)
Shivers	41	(36)	10	(21)	8	(28)	1	(12)	130	(36)
Joint pain	42	(36)	10	(21)	3	(10)	1	(12)	91	(25)
Retrosternal pain	28	(24)	7	(15)	5	(17)	2	(25)	65	(18)
Sweating	23	(20)	4	(8)	5	(17)	3	(37)	72	(20)
Abdominal pain	14	(12)	9	(19)	3	(10)	0	(0)	31	(9)
Dyspnea	12	(10)	4	(8)	1	(3)	3	(37)	46	(13)
Nausea	12	(10)	3	(6)	4	(14)	0	(0)	43	(12)
Vomiting	8	(7)	5	(11)	3	(10)	0	(0)	34	(9)
Diarrhoea	7	(6)	7	(15)	3	(10)	1	(12)	22	(6)
Shortness of breath	0	(0)	1	(2)	0	(0)	0	(0)	1	(0)

**Table 4.**

**Crude and adjusted OR (and 95% CI) of Influenza, according to gender, age, month of diagnosis and symptoms.**

Variables	OR	95% CI	Adj for variables *	95% CI
females vs. males	0.82	0.56 1.21	-----	
age x 10 years	0.89	0.81 0.98	0.83	0.74 0.93
Month of diagnosis				
December (ref.)	1.00	---	1.00	---
Jan	6.39	3.23 12.64	7.11	3.53 14.33
Feb	4.74	2.42 9.27	5.32	2.65 10.64
March	3.39	1.45 7.97	3.75	1.55 9.06
April	9.40	2.22 39.71	8.25	1.90 35.78
Nov	0.33	0.04 2.59	0.38	0.05 3.08
Symptoms				
Dry cough	0.86	0.58 1.26	---	
Productive cough	1.24	0.83 1.87	---	
Dyspnea	0.80	0.42 1.51	---	
Sore throat	1.07	0.71 1.62	---	
Headache	1.18	0.80 1.75	---	
Shivers	1.10	0.73 1.65	---	
Sweating	1.01	0.62 1.66	---	
Retrosternal pain	1.29	0.80 2.09	---	
Nasal congestion	1.48	1.00 2.20	1.50	0.97 2.31
Muscle pain	1.97	1.33 2.91	2.00	1.25 3.20
Joint pain	1.82	1.20 2.76	1.81	1.09 3.01
Nausea	0.94	0.50 1.77	---	
Vomit	0.71	0.33 1.49	---	
Abdominal pain	1.30	0.71 2.39	---	
Diarrhoea	0.87	0.39 1.94	---	
Shortness of breath	---		---	
Earache	0.77	0.36 1.63	---	

with a p-value < 0.10 at univariate level.

**Table 5.**  
**Crude and adjusted OR (and 95% CI) of Adenovirus, according to gender, age, month of diagnosis and symptoms**

Variables	OR	95% CI	Adj for variables *	95% CI
females vs. males	0.84	0.50 1.41	----	
age x 10 years	0.92	0.81 1.04	----	
<b>Month of diagnosis</b>				
December (ref.)	1.00	---		
Jan	0.69	0.32 1.49		
Feb	1.20	0.63 2.30		
March	0.56	0.18 1.72		
April	---			
Nov	0.91	0.29 2.87		
<b>Symptoms</b>				
Dry cough	0.78	0.46 1.32	---	
Productive cough	1.12	0.65 1.95	---	
Dyspnea	0.63	0.24 1.64	---	
Sore throat	0.65	0.38 1.10	---	
Headache	0.64	0.36 1.12	---	
Shivers	0.63	0.35 1.15	---	
Sweating	0.34	0.13 0.87	0.41	0.16 1.06
Retrosternal pain	0.70	0.33 1.45	---	
Nasal congestion	0.78	0.46 1.32		
Muscle pain	0.40	0.21 0.75	0.43	0.22 0.82
Joint pain	1.01	0.56 1.83		
Nausea	0.52	0.18 1.47	---	
Vomit	0.88	0.34 2.31	---	
Abdominal pain	2.07	1.01 4.22	2.18	1.05 4.52
Diarrhoea	1.54	0.62 3.84	---	
Shortness of breath	8.32	0.51 134.73	---	
Earache	0.95	0.36 2.49	---	

\* with a p-value < 0.10 at univariate level.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Viral causes of Influenza-Like Illness: Insights from a study during the winters 2004-2007

Simona Puzelli<sup>1\*</sup>, Catia Valdarchi<sup>1\*</sup>, Marco Ciotti<sup>2\*</sup>, Maria Dorrucchi<sup>1</sup>, Francesca Farchi<sup>1</sup>,  
Muhammed Babakir-Mina<sup>2</sup>, Carlo Federico Perno<sup>2</sup>, Isabella Donatelli<sup>1</sup>, Giovanni Rezza<sup>1</sup> for the  
Italian ARI Study Group

\*The first three authors are equal contributors

<sup>1</sup>*Department of Infectious, Parasitic and Immunomediated Diseases, Istituto Superiore di Sanità,  
Rome, Italy*

<sup>2</sup>*Department of Molecular Virology Policlinico Tor Vergata, Rome, Italy*

**Key words:** Influenza-like illness, Respiratory viruses, Influenza, Adenovirus, Italy

**Short running headline:** Clinical predictors of influenza and Adenovirus associated influenza-like  
illness

*Correspondence to:*

Giovanni Rezza

Epidemiology Unit

Department of Infectious Diseases

Istituto Superiore di Sanità

Viale Regina Elena, 299

00161 Roma, Italy

[g.rezza@iss.it](mailto:g.rezza@iss.it)

**ABSTRACT**

Limited information is available on the viral etiology of influenza-like illness in southern European countries, and it is still a matter of debate whether certain symptoms can be used to distinguish the specific viruses that cause influenza-like illness. The main objective of the present study was to identify the demographic and clinical predictors of influenza-like illness due to specific viruses. The study, which was observational in design, was conducted in Rome and Naples, Italy. Cases of influenza-like illness were defined as individuals with fever  $>37.5^{\circ}\text{C}$  and at least one systemic and one respiratory symptom, recruited during the winters of 2004-2005, 2005-2006 and 2006-2007. Influenza and other respiratory viruses were identified using the polymerase chain reaction (PCR), carried out on throat swabs. Basic individual information was collected using a standard form. A total of 580 persons were included in the analysis. Viral pathogens were identified in fewer than 50% of the cases. Overall, 240 viruses were detected: 22.8% were positive for influenza viruses, 10.9% for adenoviruses, 6% for parainfluenza viruses, and 1.7% for respiratory syncytial virus. The month of diagnosis, and muscle and joint pain were associated with influenza virus, although the positive predictive value was low. Abdominal pain was associated with adenovirus infection. Although the positive predictive value of symptoms of influenza virus infection was low, especially during low activity periods, these findings may help diagnosis by clinicians.

## INTRODUCTION

Respiratory infections are common in both adults and children, although the incidence of acute respiratory illness is highest in young children and decreases with increasing age [Badger et al.,1953]. Most acute respiratory infections are fairly mild, self-limiting, and confined to the upper respiratory tract, but severe illness may occur.

In industrialised countries, most of the respiratory infections which occur in winter have been attributed to viruses [Monto, 1995; Gwaltney, 2002]. However, the frequency of detection of specific viruses varies among studies, depending on the case-definition and diagnostic technique used, the type of specimen collected, the year of the study and the season [Badger et al., 1953]. The viruses detected most commonly have been rhinoviruses and influenza viruses, followed by parainfluenza viruses (PIV), respiratory syncytial virus (RSV), and adenoviruses [Badger et al.,1953; Heijnen et al., 1999]. The case-definition may affect the results: for example, for influenza-like illness, influenza viruses are detected most commonly, whereas rhinoviruses may rank first when a more generic definition of acute respiratory infection is used [van Gageldong-Lafeber et al., 2005]. Although high detection rates of RSV in influenza-like illness have also been reported [Lina et al.,1996], RSV infection is more likely to cause severe morbidity and higher mortality than influenza; for this reason, it is reported more commonly among hospitalized patients with severe acute lower respiratory infections, compared to persons in community-based studies [Fleming et al., 2005].

It is still a matter of debate whether certain symptoms can be used to distinguish among specific infections, although it is a common opinion that there are no symptoms specific for any viral infection. Although a few studies have attempted to identify signs or symptoms associated specifically with influenza virus [Peltola et al., 2005; Bolvin et al. 2000], no definitive conclusions have been drawn, and to the best of our knowledge, few studies have focused on influenza-like illness due to viruses other than influenza.

1  
2  
3 An observational study was undertaken with the following objectives: (i) to identify viruses  
4 responsible for influenza-like illness, (ii) to determine the proportion of cases attributable to each  
5 virus, and (iii) to identify the demographic and clinical predictors of influenza-like illness due to  
6 specific viruses.  
7  
8  
9  
10  
11

## 12 **MATERIALS AND METHODS**

### 13 *Study population and sample collection*

14  
15  
16  
17  
18 The study design and methods have been described in detail elsewhere [Rezza et al. 2006].  
19 Briefly, the study was conducted in Rome (winter seasons 2004-05, 2005-06 and 2006-07) and  
20 Naples (2005-06 and 2006-07), Italy, and in some peripheral areas of these cities. Eighteen general  
21 practitioners (GPs) from these areas participated, eight of whom were paediatricians. The  
22 physicians were asked to obtain a throat swab for all consecutive patients who, during the period  
23 November-April, presented with influenza-like illness within three days of the onset of symptoms;  
24 at the beginning of the winter season, each practitioner was provided with “Virocult swabs”  
25 (Medical Wire and Equipment, Corsham, UK). Influenza-like illness was defined as the presence of  
26 fever  $>37.5^{\circ}\text{C}$  (axillary temperature) and at least another systemic symptom (i.e., headache,  
27 malaise, myalgia, chills or sweating, retrosternal pain, or asthenia) and one respiratory symptom  
28 (i.e., cough, sore throat, nasal congestion or runny nose). The minimum temperature for defining  
29 fever ( $>37.5^{\circ}\text{C}$ ) was slightly lower than that used by the Italian Ministry of Health for influenza-  
30 like illness surveillance ( $38^{\circ}\text{C}$ ) [Gabutti et al., 2004], so as to include persons with a milder fever.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Information on gender, age, clinical symptoms, and vaccination history was obtained for  
49 each participant. The clinical samples were sent to the laboratory for examination. Verbal informed  
50 consent was obtained from all of the participants. Because this was an observational study, ethical  
51 committee approval was not required, although the study was conducted in accordance with the  
52 ethical requirements of the Italian Ministry of Health.  
53  
54  
55  
56  
57  
58  
59  
60

### *RNA and DNA Extraction and PCR amplification*

At the laboratory, two separate aliquots of each clinical sample were prepared. To identify influenza A and B viruses and RSV types A and B, a multiplex RT-PCR assay was carried out. Viral RNAs were extracted either directly from clinical samples or from virus-infected MDCK culture fluid using an RNA extraction kit (RNeasy; Qiagen, Santa Clara, CA). cDNA synthesis and amplification were carried out as described elsewhere [Puzelli et al., 2004]. PCR amplification was performed using primers targeting specific regions within the following genes: (i) influenza A nucleoprotein and influenza A/H1- and A/H3-subtype haemagglutinins; and (ii) influenza B haemagglutinin and neuraminidase. The primers used in the PCR reactions are available from the authors upon request.

To identify other respiratory viruses, total DNA and RNA was extracted from a separate aliquot of the clinical sample using Ultrasens kit (Qiagen, Hilden, Germany), following the manufacturer's instructions. To verify the acid nucleic extraction (DNA and RNA), the nucleic acid was amplified with the  $\beta$ -actin gene [Rezza et al., 2006; Strassburg et al., 1999]: all of the samples were positive. The samples were tested for the presence of influenza virus, adenovirus, RSV, and PIV types 1, 2, 3 and 4 [Heim et al., 2003; Templeton et al., 2004].

### *Statistical analysis*

The distribution of the specific viral agents was evaluated first by month of diagnosis and age of the participant. The distribution of signs and symptoms was determined according to the etiology (i.e., influenza virus, adenovirus, RSV, PIV and other/unknown). Odds ratios (OR) and their 95% confidence intervals (95% CI) were calculated to determine the association between each variable (i.e., age, month of diagnosis, symptoms) and the most common viral infections at the univariate level. The variables which were found to be associated statistically at the univariate level were included in a multivariate model to test their independent association with the specific viral etiology (i.e., the model was used only for influenza, since only one variable was found to have

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

been associated with adenovirus infection at the univariate level). The positive predictive value (PPV) of specific symptoms for specific viral etiology was also calculated.

## RESULTS

During the three winters considered in this study, 580 persons were recruited: 173 (29.8%) in 2004-05, 190 (32.8%) in 2005-06, and 217 (37.4%) in 2006-07. Of the participants, 299 (51.6%) were females and 281 (48.4%) males. The median age was 24 years (range: 0.5-88 years); 224 (38.6%) participants were of paediatric age ( $\leq 13$  years), of whom 89 were less than 3 years old. Of the study participants, 561 (96.7%) were Italian. With regard to the study area, 409 (70.5%) participants were recruited in the province of Rome and 171 (29.5%) in Naples.

Of the samples collected, 220 were positive for at least one of the four viruses; of these, 20 were positive for two viruses. In the case of 360 samples, virus was not detected or the virus was not identified. The virus detected most frequently was influenza virus (n=132 samples), followed by adenovirus (n=63), PIV (n=35), and RSV (n=10).

One hundred and eighteen isolates were identified as influenza A and 14 as influenza B. Of the influenza A viruses, H3-subtype strains prevailed in the 2004-05 season, whereas H1-subtype strains prevailed in the 2005-06 season; during the 2006/07 season, H3 strains were again prevalent (57%). Of the parainfluenza viruses isolates, 27 were parainfluenza viruses type 3; 5 were type 2; 2 were type 4; and one was type 1.

The number of samples, by specific virus and month of diagnosis, is shown in Table 1. Influenza virus tended to be the most common virus from January to April, whereas PIV and adenovirus were the viruses detected most frequently in November and December, respectively. Of the 20 coinfecting participants, most (n=12) were positive for both influenza virus and adenovirus; four were positive for influenza virus and PIV, two for adenovirus and PIV, and two for adenovirus and RSV.

1  
2  
3 Influenza virus was detected most frequently during two winter seasons (i.e., 2004-05 and  
4  
5 2006-07), whereas adenovirus was detected more frequently than influenza virus in 2005-06 (i.e.,  
6  
7 21% vs. 18%) (data not shown).  
8  
9

10 The number of samples, by specific virus and age class of the participant, is shown in Table  
11  
12 2. With regard to influenza viruses, the proportion of positive samples was highest in the youngest  
13  
14 age group (0-2 years) and tended to decrease with increasing age (chi square for trend,  $p < 0.01$ ).  
15  
16 Children ( $\leq 13$  years; the two youngest age groups) were more than 2 times more likely than  
17  
18 adolescents and adults/elderly (the two oldest age groups) to be infected with influenza viruses (OR:  
19  
20 1.56, 95% CI: 1.04-2.36). Only one of the 13 participants  $\geq 65$  years of age was positive for  
21  
22 influenza virus.  
23  
24  
25

26  
27 With regard to influenza vaccination, 118 (20.3%) of the 580 participants had been  
28  
29 vaccinated; of these individuals, 23 (19.5%) were infected by influenza viruses, compared to 109  
30  
31 (23.6%) of the 462 individuals who had not been vaccinated (difference not statistically significant;  
32  
33 OR: 0.78, 95% CI: 0.46-1.33).  
34  
35

36 The number of samples, by specific virus and sign and symptom, is shown in Table 3.  
37  
38 Muscle pain ( $p = 0.028$ ) and joint pain ( $p = 0.046$ ) were reported more likely in persons infected with  
39  
40 influenza viruses, compared to those infected with other viruses, and nausea ( $p = 0.045$ ) was reported  
41  
42 less likely. The crude and adjusted ORs for variables potentially associated with positivity for  
43  
44 influenza viruses are reported in Table 4. At the multivariate analysis, influenza viruses were more  
45  
46 likely to be detected between January and April (compared to December), in younger age groups  
47  
48 (i.e.,  $< 10$  years, compared to older persons), and in participants with muscle or joint pain (compared  
49  
50 to participants with other symptoms). However, none of the symptoms associated with influenza  
51  
52 positivity had a high PPV: muscle pain had the highest PPV (42.9%) during the months of flu  
53  
54 activity (i.e., January and February), whereas the combination of symptoms with the highest PPV in  
55  
56 these two months was nasal congestion plus muscle pain (48.3%); the PPV of this combination was  
57  
58 12.9% in low influenza activity months (i.e., November and December). Given that some symptoms  
59  
60

1  
2  
3 were reported less frequently by participants younger than 10 years, such as muscle pain (18%  
4 among those less than 10 years old vs. 46% among those 10 years of age or older) and joint pain  
5 (9% vs. 35%), which could be due in part to the inability of very young children to describe  
6 subjective symptoms, the logistic analyses were repeated excluding persons younger than 10 years  
7 of age, and the results were confirmed (data not shown). Similarly, sweating was reported less  
8 frequently by persons younger than 10 years of age (9% vs. 23%).  
9  
10  
11  
12  
13  
14  
15  
16

17 The crude and adjusted ORs for variables associated potentially with positivity for  
18 adenovirus are reported in Table 5. Adenovirus positivity was more likely to be detected among  
19 those reporting abdominal pain, whereas there was no association with either the month of  
20 diagnosis or age (Table 5). When excluding participants younger than 10 years of age, the adjusted  
21 OR for abdominal pain remained of the same magnitude yet was no longer significant statistically  
22 (results not shown).  
23  
24  
25  
26  
27  
28  
29  
30  
31

## 32 DISCUSSION

33  
34 In this study, about 38% of the samples were positive for at least one virus, which is  
35 consistent with the results of other studies in which from 36-38% [Lina et al., 1999; Vabret et al.,  
36 2003] to 58% [van Gageldong-Lafeber et al., 2005] of the samples were positive. In a study of  
37 community-acquired respiratory infections, including *Mycoplasma pneumoniae* and *Chlamydomphila*  
38 *pneumoniae*, in addition to viruses, at least one potentially pathogenic microorganism was detected  
39 in 52% of the swabs [Heijnen et al., 1999].  
40  
41  
42  
43  
44  
45  
46  
47

48 As was found in other studies [Heijnen et al., 1999; Lina et al., 1996; Vabret et al., 2003],  
49 influenza was the virus detected most commonly, which could in part be due to the large epidemic  
50 of infection with influenza virus in 2004-05. The finding that influenza vaccination did not have a  
51 protective effect was probably related to viral drift [Ansaldi et al., 2005]. As expected, the overall  
52 results showed that the proportion of samples positive for influenza viruses was higher in young  
53 children than in adults; this is likely to be due either to higher vaccination coverage among older  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 adults or to the duration of viral shedding in young children, which is usually longer than in adults  
4  
5 and characterized by larger virus load.  
6  
7

8 In the present study, RSV was detected rarely, which differs from the results of a study, in  
9  
10 which it was almost as common as influenza virus, with the highest impact in the youngest age  
11  
12 groups [Lina et al., 1996]. The rarity of RSV detection could be due to not only the low proportion  
13  
14 of children but also to the fact that children admitted to hospital were not included.  
15  
16

17 In this study, an attempt was made to identify symptoms associated with influenza and  
18  
19 adenovirus infections. However, it is known that clinical diagnosis of influenza is not easy. In a  
20  
21 prospective study of children  $\leq 13$  years old, the overall sensitivity of clinical diagnosis of influenza  
22  
23 was 38% and the PPV was 32% [Peltola et al., 2005]. The analysis of a study involving 5-12 year-  
24  
25 old children enrolled in two trials showed that cough, headache, and fever  $\geq 38.2^{\circ}\text{C}$  predicted  
26  
27 independently positivity to influenza virus; the PPV of combined fever and cough was 83%,  
28  
29 whereas sore throat and myalgia were negative predictors. Myalgia was found to be associated  
30  
31 independently with influenza only among younger children (1-4 year-olds) [Ohmit et al., 2006]. In  
32  
33 another study of children ( $>6$  years of age) and adults, cough and fever  $>38$  were the only factors  
34  
35 associated with a positive PCR for influenza [Bolvin et al., 2000].  
36  
37  
38  
39  
40

41 In the present study, muscle and joint pain were associated with influenza, whereas nasal  
42  
43 congestion was associated only marginally with the infection; the month of diagnosis (from January  
44  
45 to April) and older age were also associated with influenza virus. However, the PPV of these  
46  
47 symptoms remained low, even in the months with high influenza activity. Only abdominal pain was  
48  
49 associated with adenovirus infection, whereas there was no association with either age or month of  
50  
51 diagnosis.  
52  
53

54 Before drawing conclusions, the limits and bias of this study should be mentioned. Firstly,  
55  
56 some viruses (rhinoviruses and metapneumoviruses) and bacteria (*M. pneumoniae* and *C.*  
57  
58 *pneumoniae*) were not studied. In particular, the inclusion of rhinoviruses might increase greatly the  
59  
60 frequency of virus detection, as indicated by studies reporting a higher proportion of these viruses

1  
2  
3 compared to influenza viruses [Badger et al., 1953]; it could also explain the relatively high  
4  
5 proportion of unidentified causes. Nevertheless, the results of this study do not differ significantly  
6  
7 from those of other studies. Secondly, the specific influenza type may produce some differences in  
8  
9 clinical features. Rhinorrhea and sick appearance have been reported more frequently by children  
10  
11 with influenza A, whereas myalgia is more common among those with influenza B [Peltola et al.,  
12  
13 2003]. Symptoms such as muscle ache and headache, which may be characteristic of influenza in  
14  
15 adults, are reported by a low proportion of children and cannot be communicated reliably by those  
16  
17 <3 years of age. Thirdly, the potential occurrence of false-negative results due to the variable  
18  
19 sensitivity of the laboratory techniques and to the type of clinical samples cannot be ruled out  
20  
21 completely. Finally, the timing of collection of samples may have decreased the rate of detection,  
22  
23 since some swabs were taken up to four days after the onset of symptoms (when viruses may have  
24  
25 been cleared, at least in part, by the immune response). For this reason, the maximum sample delay,  
26  
27 which was set at four days in the first year of the study, was reduced to three days in the last two  
28  
29 years.  
30  
31  
32  
33  
34

35  
36 In conclusion, influenza virus was the virus identified most commonly, followed by  
37  
38 adenoviruses, whereas cases attributable to other viruses were uncommon. The comparison between  
39  
40 persons affected by the two different viral infections permitted the identification of some predictors  
41  
42 of a specific viral disease. Although the surveillance of respiratory viruses associated with  
43  
44 influenza-like illness is not sustainable, because of the high costs and lack of preventive measures,  
45  
46 limited etiological surveys may provide useful information on the effect of specific viruses affecting  
47  
48 human populations in the winter season and may possibly allow predictors of specific viral  
49  
50 infections to be identified. The findings of this study may help clinicians to discriminate between  
51  
52 influenza virus, adenovirus, and other infections; due to the low PPV of symptoms and other  
53  
54 correlates of infection for influenza virus, the decision to use specific anti-influenza drugs should  
55  
56 not be based only on the clinical diagnosis of influenza. Rapid laboratory tests might be used to  
57  
58  
59  
60

1  
2  
3 confirm influenza virus infection and facilitate decision-making when the use of antiviral drugs is  
4  
5 considered.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## ACUTE RESPIRATORY INFECTION (ARI) STUDY GROUP

Members of the acute respiratory infection study group include Laura Calzoletti, Angela Di Martino, Concetta Fabiani, Marzia Facchini, Maria Interisano, (Istituto Superiore di Sanità); Maria Corongiu, Antonio Palma, Alberto Chiriatti, Giuseppe Grasso, Paolo Marin, Giulio Nati, Mauro Mocchi, Virgilio Calzini, Andrea Filabozzi, Antonio Verginelli, Tiziana Sisti, Silvana Pescosolido (General Practitioner of Rome); Antonio Parlato, Rosaria Granata, Anna Farro, Teresa Coppola, (ASL NA/2, Naples); Carmine Barile, Salvatore Romani, Luigi Longobardi, Romualdo Mauro, Cimino Margherita, Cecere Aniello, Maisto Angelo, Marrazzo Giuseppe, Auciello Aurora (General Practitioner of Naples)

## ACKNOWLEDGMENTS

The study was funded by the CCM (Centro Controllo Malattie) of the Italian Ministry of Health.

Footnote: The authors declare that they do not have any commercial or other associations that might pose a conflict of interest.

For Peer Review

## REFERENCES

1  
2  
3  
4  
5  
6 Ansaldi F, Icardi G, Gasparini R, Campello C, Puzelli S, Bella A, Donatelli I, Salmaso S,  
7  
8 Crovari P. 2005. New A/H3N2 influenza variant: a small genetic evolution but a heavy burden  
9  
10 on the Italian population during the 2004-2005 season. *J Clin Microbiol* 43: 3027-3029.  
11  
12

13  
14  
15 Badger GF, Dingle JH, Feller AE, Hodges RD, Jordan WS, Rammelkamp CH Jr. 1953. A study  
16  
17 of illness in a group of Cleveland families. II. Incidence of common respiratory disease. *Am J*  
18  
19 *Hyg* 58:31-40.  
20  
21

22  
23  
24  
25 Bolvin G, Hardy I, Tellier G, Maziade J. 2000. Predicting influenza infections during epidemics  
26  
27 with use of a clinical case definition. *Clin Infect Dis* 31: 1166-1169.  
28  
29

30  
31  
32 Division of Health Promotion and Disease Prevention IoMWU. 1985. New vaccine  
33  
34 development: establishing priorities. *New Vaccine Dev* 1:397-410.  
35  
36

37  
38  
39 Fleming DM, Pannell RS, Cross KW. 2005. Mortality in children from influenza and respiratory  
40  
41 syncytial virus. *J Epidemiol Community Health* 59:586-590.  
42  
43

44  
45  
46 Gabutti G, Guido M, Quattrocchi M, Zizza A, De Donno A, Gasparini R, Donatelli I, Prato R,  
47  
48 Germinario C, Crovari P; Collaborative Group for the Study of Infectious Diseases;  
49  
50 Collaborative group for Influenza Surveillance. 2004. "Surveillance of influenza in Apulia, Italy,  
51  
52 1999-2000, 2000-2001, 2001-2002, and 2002-2003 seasons". *Med Mal Infect.* Oct;34(10):469-  
53  
54 76.  
55  
56

57  
58  
59 Gwaltney JM. 2002. Clinical significance and pathogenesis of viral respiratory infections. *Am J*  
60  
*Med* 112 (Suppl 6A):13S-18S.

1  
2  
3  
4  
5  
6 Heijnen MLA, Dorigo-Zetsma JW, Bartelds AIM, Wilbrink B, Sprenger MJW. 1999.  
7  
8 Surveillance of respiratory pathogens and influenza-like illness in general practices – The  
9  
10 Netherlands, winter 1997-98. *Eurosurv* 4:81-84.  
11

12  
13  
14  
15 Heim A, Ebnet C, Harste G, Pring-Akerblom. 2003. Rapid and quantitative detection of human  
16  
17 adenovirus DNA by real-time PCR. *J Med Virol* 70:228-239.  
18

19  
20  
21  
22 Leader S, Kohlhase K. 2003. Recent trends in severe respiratory syncytial virus (RSV) among  
23  
24 US infants, 1997 to 2000. *J Pediatr* 143:S127-S132.  
25

26  
27  
28  
29 Lina B, Valette M, Foray S, Luciani J, Stagnara J, See DM, Aymard M. 1996. Surveillance of  
30  
31 community-acquired viral infections due to respiratory viruses in Rhone-Alpes (France) during  
32  
33 winter 1994 to 1995. *J Clin Microbiol* 34:3007-3011.  
34

35  
36  
37  
38 Monto AS. 1995. Viral respiratory infections in the community: epidemiology, agents, and  
39  
40 interventions. *Am J Med* 99:24S-27S.  
41

42  
43  
44  
45 Ohmit SE, Monto AS. 2006. Symptomatic predictors of influenza virus positivity in children  
46  
47 during the influenza season. *Clin Infect Dis* 43:564-568.  
48

49  
50  
51  
52 Peltola V, Ruuskanen O. 2003. Influenza A and B virus infections in children. *Clin Infect Dis*  
53  
54 36:299-305.  
55

56  
57  
58  
59 Peltola V, Reunanen T, Ziegler T, Silveinonen H, Heikkinen T. 2005. Accuracy of clinical  
60  
diagnosis of influenza in outpatients children. *Clin Infect Dis* 41:1198-1200.

1  
2  
3  
4  
5  
6 Puzelli S, Frezza F, Fabiani C, Ansaldi F, Campitelli L, Lin YP, Gregory V, Bennett M, D'Agaro  
7  
8 P, Campello C, Crovari P, Hay A, Donatelli I. 2004. Changes in the hemagglutinins and  
9  
10 neuraminidases of human influenza B viruses isolated in Italy during the 2001-02, 2002-03, and  
11  
12 2003-04 seasons. *J Med Virol* 74(4):629-40.  
13  
14

15  
16  
17 Rezza G, Valdarchi C, Puzelli S, Ciotti M, Farchi F, Fabiani C, Calzoletti L, Donatelli I, Perno  
18  
19 CF. 2006. Respiratory viruses and influenza-like illness: a survey in the area of Rome, winter  
20  
21 2004-2005. *Eurosurv* 11 (10): 251-3.  
22  
23

24  
25  
26 Strassburg CP, Strassburg A, Nguyen N, Li Q, Manns MP, Tukey RH. 1999. Regulation and  
27  
28 function of family 1 and family 2 UDP-glucuronosyltransferase genes (*UGT1A*, *UGT2B*) in  
29  
30 human oesophagus. *Biochem J* 338:489-498.  
31  
32

33  
34  
35 Templeton KE, Scheltinga SA, Beersma MFC, Kroes ACM, and Claas ECJ. 2004. Rapid and  
36  
37 sensitive method using multiplex real-time PCR for diagnosis of infections by Influenza A and  
38  
39 Influenza B viruses, Respiratory Syncytial virus and Parainfluenza viruses 1, 2, 3, and 4. *J Clin*  
40  
41 *Microbiol* 42:1564-1569.  
42  
43  
44

45  
46  
47 Vabret A, Mourez T, Gouarin S, Petitjean J, Freymuth F. 2003. A outbreak of coronavirus OC43  
48  
49 respiratory infection in Normandy, France. *Clin Infect Dis* 36: 985-989.  
50  
51  
52

53  
54  
55 van Gageldong-Lafeber AB, Haijnen M-LA, Bartelds AIM, Peters MF, van der Plas SM,  
56  
57 Wilbrink B. 2005. A case-control study of acute respiratory tract infection in general practice  
58  
59 patients in The Netherlands. *Clin Infect Dis* 41:490-497.  
60