



HAL
open science

Adenosine receptor agonists for promotion of dermal wound healing

María D. Valls, Bruce N. Cronstein, M. Carmen Montesinos

► **To cite this version:**

María D. Valls, Bruce N. Cronstein, M. Carmen Montesinos. Adenosine receptor agonists for promotion of dermal wound healing. *Biochemical Pharmacology*, 2009, 77 (7), pp.1117. 10.1016/j.bcp.2008.11.002 . hal-00531838

HAL Id: hal-00531838

<https://hal.science/hal-00531838>

Submitted on 4 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Adenosine receptor agonists for promotion of dermal wound healing

Authors: María D. Valls, Bruce N. Cronstein, M. Carmen Montesinos



PII: S0006-2952(08)00789-2
DOI: doi:10.1016/j.bcp.2008.11.002
Reference: BCP 10004

To appear in: *BCP*

Received date: 16-9-2008
Revised date: 31-10-2008
Accepted date: 3-11-2008

Please cite this article as: Valls MD, Cronstein BN, Montesinos MC, Adenosine receptor agonists for promotion of dermal wound healing, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2008.11.002

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1
2
3
4
5 **Adenosine receptor agonists for promotion of dermal wound healing**
6

7 María D. Valls¹, Bruce N. Cronstein², M. Carmen Montesinos^{1,2}
8
9

10
11
12 **¹Department de Farmacologia**
13

14 **Universitat de València**
15

16 **Ave. Vicent Andrés Estellès s/n**
17

18 **46100 Burjassot, Valencia, Spain**
19
20
21

22
23
24 **²Department of Medicine**
25

26 **New York University School of Medicine**
27

28 **550 First Ave.**
29

30 **New York, NY 10016**
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 **Running title:** Adenosine in wound healing
4
5
6
7
8
9

10
11
12 **Corresponding author:**
13

14 M. Carmen Montesinos,
15

16
17 Departament de Farmacologia / Universitat de València,
18

19 Avenida Vicent Andrès Estellès s/n
20

21 46100 Burjassot, Spain
22

23
24 Tel : +(34) 96 354 4946
25

26 Fax : +(34) 96 354 4946
27

28
29 m.carmen.montesinos@uv.es
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 **ABSTRACT**
4
5
6

7 Wound healing is a dynamic and complex process that involves a well coordinated, highly
8 regulated series of events including inflammation, tissue formation, revascularization and tissue
9 remodeling. However, this orderly sequence is impaired in certain pathophysiological conditions
10 such as diabetes mellitus, venous insufficiency, chronic glucocorticoid use, aging and
11 malnutrition. Together with proper wound care, promotion of the healing process is the primary
12 objective in the management of chronic poorly healing wounds. Recent studies have
13 demonstrated that A_{2A} adenosine receptor agonists promote wound healing in normal and
14 diabetic animals and one such agonist, Sonedenoson, is currently being evaluated as a prospective
15 new therapy of diabetic foot ulcers. We will review the mechanisms by which adenosine receptor
16 activation affects the function of the cells and tissues that participate in wound healing,
17 emphasizing the potential beneficial impact of adenosine receptor agonists in diabetic impaired
18 healing.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Keywords:** Adenosine receptors; Impaired healing; Inflammation; Diabetic foot ulcer;
39 angiogenesis; Granulation tissue
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

The mechanisms underlying the normal repair process, cell migration and proliferation, and extracellular matrix deposition and remodelling, have been extensively studied [1-3]. Cellular responses to inflammatory mediators, growth factors, cytokines, and to mechanical forces must be appropriate and precise in order to obtain optimum healing of a cutaneous wound. However, even during the normal process of wound healing complications can occur, including infection, thrombosis, and ischemia. [4, 5]. More importantly, the orderly progression of the healing process is impaired in chronic wounds, including those due to diabetes.

Impaired wound healing is a major concern for diabetic patients because their wounds do not heal properly and are a source of major suffering and cost. Only two-thirds of diabetic foot ulcers eventually heal and up to 28% may result in amputation [6]. The pathogenesis of diabetic foot ulcers is complex and it is well recognized that a number of contributory factors working together ultimately lead to impaired healing. Several intrinsic factors, such as peripheral neuropathy, foot deformity, peripheral vascular disease and peripheral oedema have been identified as the commonest factors responsible of impaired healing after trauma. In addition, extrinsic factors, such as wound infection, callus formation, and excessive pressure to the site, further aggravate the healing process [7].

Recent studies suggest that nerves play a central role in tissue homeostasis and can orchestrate complex reparative as well as destructive processes. First, an intact nociceptor system of primary afferent sensory nerves is important for the initiation of the inflammatory process and successful tissue repair [8]. Apart from the loss of pain perception, which is a key factor in the development

1
2
3 of neuropathic foot ulcers, loss of autonomic function and small fibre neuropathy can result in
4
5 impaired neurogenic control of local microcirculatory blood flow, impaired fluid homeostasis,
6
7 diminished energy metabolism, oxygen delivery, and inflammatory responses. These processes
8
9 could render the feet of diabetic patients with neuropathy more susceptible to tissue damage and
10
11 infection [9].
12
13

14
15
16 Given the complexity of the pathogenesis of diabetic foot ulcers, many different interventions
17
18 have been proposed to accelerate the healing process, but few have been subjected to formal
19
20 evaluation. Despite the relatively large number of studies of growth factors like PDGF
21
22 (becaplermin), EGF, basic FGF and other agents modulating aspects of wound physiology, there
23
24 is currently little evidence to suggest that any of the reported interventions should be adopted in
25
26 routine practice. Diabetic foot ulcer management is based on the simple principles of eliminating
27
28 infection, debridement, cleansing and the use of dressings to maintain a moist wound bed, and
29
30 lastly, becaplermin is the only promoting agent approved for use of those ulcers resistant to
31
32 simpler interventions [6].
33
34
35
36
37
38
39

40
41 Based in studies carried out in vitro and in experimental animal models, we are proposing a new
42
43 strategy for the promotion of impaired wound healing, the use of adenosine receptor agonists.

44
45 We will summarized the biology of adenosine and review its actions on different tissues and cells
46
47 implicated in the healing of cutaneous wounds, as well as its effect on experimental wounds in
48
49 animals.
50
51
52
53
54
55
56
57
58
59

Purine metabolism and biology

Adenosine is a ubiquitous purine nucleoside produced by stepwise dephosphorylation of ATP by the coordinated action of ecto-apyrase (CD39) and ecto-5'-nucleotidase (CD73) (Figure 1).

While extracellular ATP and other nucleotides (ADP, UTP and UDP) have many biological effects through direct activation of cell surface receptors for adenine nucleotides (seven P2X ionotropic and eight P2Y metabotropic receptor subtypes), adenosine modulates cellular and organ function via occupancy of four specific cell surface receptors, (A_1 , A_{2A} , A_{2B} and A_3), all members of the large family of 7-transmembrane spanning, heterotrimeric G protein-associated receptors [10, 11]. The A_1 and A_3 adenosine receptors coupled with G_i proteins are associated with two effector systems, namely, adenylate cyclase and phospholipase C. The binding of adenosine or its agonists to A_1 and A_3 adenosine receptors either induce inhibition of adenylate cyclase leading to a decrease in intracellular cAMP levels or stimulate phospholipase C and the release of intracellular Ca^{2+} . A_{2A} and A_{2B} receptors are associated with G_s proteins and their activation stimulates an increase in intracellular cAMP. In addition, they couple to mitogen-activated protein kinases (MAPK), which may give them a role in cell growth, survival, death and differentiation [12]. The affinity of selected agonists at the different adenosine receptor subtypes is summarized on Table 1.

Extracellular actions of purines in non-neuronal cells, including fast signalling roles in exocrine and endocrine secretion, platelet aggregation, cardiovascular effects and kidney function, have been known for a long time. More recently, slow purinergic signalling has been implicated in embryological development, wound healing, restenosis, atherosclerosis, ischaemia, cell turnover of epithelial cells in skin and visceral organs, inflammation, neuroprotection and cancer [13, 14].

1
2
3 More interestingly, purines and pyrimidines have major roles in the activities of neurons. This
4 includes nociceptive mechanosensory transduction, as well as acting as a cotransmitter and
5 neuromodulator in most, if not all, nerve types in the peripheral and central nervous systems [13].
6
7 This raises the innovative hypothesis that diabetic patients suffering from peripheral neuropathy
8 will have altered purinergic neurotransmission.
9
10
11
12
13

14 15 16 17 **Could adenosine play a role in normal wound healing?** 18

19
20 Under basal conditions, the extracellular adenosine concentration is rather constant (30–300 nM),
21 and held in tight check by the equilibrium between adenosine production/release into the
22 extracellular space and adenosine uptake by cells or catabolism to inosine (Figure 1). In contrast
23 its concentration can increase dramatically to micromolar or even higher ranges when there is an
24 imbalance between energy use and energy supply, such as in oxygen depletion, or under
25 conditions of cellular or tissue necrosis or stress as a result of ATP catabolism [12].
26
27
28
29
30
31
32
33
34
35
36

37 All cell subtypes involved in wound healing, macrophages, epidermal cells, fibroblasts and
38 microvascular endothelial cells, differentially express functional adenosine receptors, although
39 the receptor expression patterns vary between cellular types and have not been fully established.
40 Moreover, even the same cellular type such as endothelial cells express different adenosine
41 receptor subtypes depending on the vascular bed of origin [15]. Adenosine A_{2A} receptors, in
42 particular, are expressed on most cell types involved in wound healing, including macrophages,
43 fibroblasts and microvascular endothelial cells [15-17]. We have reported that A_{2A} receptor-
44 deficient mice suffer from disordered wound healing with poor matrix formation and diminished
45 blood vessel formation in the granulation tissue of excisional wounds and mechanically injured
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 skin, an observation that indicates a role for adenosine, acting at A_{2A} receptors, in normal wound
4
5 healing [18]. We have further observed that cytokines released during the inflammatory phase of
6
7 wound healing, TNF- α and IL-1, up-regulate adenosine A_{2A} receptor expression in human
8
9 monocytoïd cells (THP-1) and microvascular endothelial cells [16, 19, 20]. Similarly, A_{2B}
10
11 receptor expression in murine bone marrow-derived macrophages is also up-regulated by IFN-
12
13 gamma [21]. The differential adenosine receptor expression in cells from diabetic patients has not
14
15 been established yet. In this regard, streptozotocin-induced diabetes in rats altered adenosine
16
17 receptors expression in liver, heart and kidney and administration of insulin returned their levels
18
19 to normality [22]
20
21
22
23
24
25

26 **Adenosine in inflammation**

27
28
29
30
31 Hemostasis and inflammation constitute the first phase of tissue repair. The formation of a blood
32
33 clot re-establishes tissue hemostasis and provides a provisional matrix for cell migration.
34

35
36 Released cellular mediators initiate inflammatory leukocyte recruitment necessary for removing
37
38 necrotic tissue and preventing infection [1]. In addition, inflammatory cells release a variety of
39
40 cytokines and chemokines that play an important role in the evolution of granulation tissue
41
42 through stimulation of fibroblasts and epithelial cells [4]. However, despite its importance,
43
44 persistent or exaggerated inflammation could be detrimental for the healing process in some
45
46 pathological settings. In this respect, the concept that diabetes is a low-level chronic
47
48 inflammatory disease is commonly accepted [23, 24].
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 Numerous studies indicate that adenosine, through the activation of its different receptor
4 subtypes, is a potent regulator of inflammation and innate immunity [25, 26]. In fact, genetic
5 deficiency in adenosine deaminase (ADA), the enzyme responsible of the deamination of
6 adenosine to its less potent derivative inosine (Figure 1), is characterized by a severely
7 compromised immune system [27]. Moreover, several studies have established that adenosine
8 mediates the anti-inflammatory effect of mainstay anti-rheumatic drugs such as methotrexate and
9 sulfasalazine, and also salicylates, in different animal models of acute and chronic inflammation
10 [28-30].
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Adenosine A₁ receptor activation has been associated with pro-inflammatory properties in most
25 inflammatory cell types [31, 32]. Nevertheless, studies *in vivo* have demonstrated the anti-
26 inflammatory effect of selective A₁ agonists acting in the Central Nervous System by increasing
27 adenosine concentration at the inflamed site [33, 34].
28
29
30
31
32
33
34
35

36 Adenosine A_{2A} receptors are generally regarded as the receptor subtype most relevant for the
37 anti-inflammatory effect of adenosine. Their activation inhibits neutrophil and monocyte
38 oxidative burst, degranulation and release of cytokines and chemokines [35, 36]. Activation of
39 A_{2B} receptors selectively inhibits collagenase mRNA accumulation in synovial fibroblasts,
40 mediates neutrophil-stimulated intestinal epithelial leakiness and prevents vascular leakage and
41 edema formation [37-39]
42
43
44
45
46
47
48
49
50
51

52 The role of adenosine A₃ receptors in inflammation has been more controversial, maybe due to
53 the observed difference in agonist affinity between species. have also been described as anti-
54 inflammatory in human blood leukocytes and in murine models of inflammation [40, 41]. We
55
56
57
58
59
60
61
62
63
64
65

1
2
3 have confirmed the anti-inflammatory effects of adenosine acting at A₃ receptors in experimental
4
5 animals, since animals deficient in this receptor show an exacerbated response to an
6
7 inflammatory insult when compared to their wild type littermates [42].
8
9

10
11
12 It has been firmly established that adenosine modulates the production of both inflammatory and
13
14 anti-inflammatory cytokines including TNF α , IL-10, and IL-12 [16, 43]. The anti-inflammatory
15
16 effect of adenosine could be beneficial since some reports depict a deleterious effect of TNF in
17
18 wound healing [44]
19
20
21
22
23

24 **Adenosine in tissue formation**

25
26
27
28
29 Driven by growth factors synthesized by local and migratory cells, fibroblasts migrate on the
30
31 provisional fibrin scaffold into the wound where they proliferate and construct a more robust
32
33 collagen rich extracellular matrix. Wound fibroblasts acquire a distinctive contractile and
34
35 secretory phenotype, known as myofibroblasts, responsible for wound contraction, a very
36
37 important event in full thickness wounds. In response to many of the same growth factors,
38
39 epidermal cells migrate from the edge of the wound over the surface of the injured area and
40
41 proliferate until there is complete wound closure [1, 5].
42
43
44
45
46
47

48 Many studies have shown that purinoceptors are involved in the regulation of proliferation and
49
50 differentiation of most target cells. Thus, activation of adenosine A_{2B} receptor and P2Y₂
51
52 receptors have mitogenic effects in murine keratinocytes [45], contrasting with earlier reports
53
54 showing that adenosine and its related nucleotides (ATP, ADP, AMP) were antiproliferative for
55
56
57
58
59
60
61
62
63
64
65

1
2
3 normal human epidermal keratinocytes cultured in the absence or presence of exogenous
4 epidermal growth factor[46]. Similarly, the result of adenosine receptor activation in fibroblast
5 proliferation remains unclear [47, 48]. Studies in our laboratory indicated that adenosine A₂
6 receptor occupancy, both A_{2A} and A_{2B}, contributes to enhanced fibroblast and endothelial cell
7 migration.[49].
8
9

10
11
12
13
14
15
16 We have recently reported that activation of adenosine A_{2A} receptors promotes collagen synthesis
17 by human dermal fibroblasts and that blockade or deletion of this receptor in mice protects
18 against bleomycin-induced dermal fibrosis, a murine model of scleroderma [17]. The stimulation
19 of collagen synthesis in human dermal fibroblasts occurs through an A_{2A}R/mitogen-activated
20 protein kinase kinase-1/mitogen-activated protein kinase-mediated pathway [17] Adenosine
21 deaminase (ADA), the principal catabolic enzyme for adenosine in vivo, and its deficiency leads
22 to the spontaneous development of pulmonary and skin fibrosis in mice, in which increased
23 collagen deposition is accompanied by increased levels of key mediators of fibrosis, including
24 transforming growth factor beta1, connective tissue growth factor, and interleukin-13.
25
26

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Pharmacological treatment of ADA-deficient mice with the A_{2A} receptor antagonist ZM-241385
prevented the development of dermal fibrosis in this model of elevated tissue adenosine, by
reducing dermal collagen content and expression of profibrotic cytokines and growth factors.

These data confirm a fibrogenic role for adenosine in the skin [50]. We also found an increased
number of myofibroblasts associated with elevated skin adenosine concentration, a phenomenon
that was prevented by pharmacological blockade of A_{2A} receptors [51]. Although these results are
consistent with a fibrogenic role for adenosine A_{2A} receptor activation, a possible role for
adenosine A_{2B} receptors cannot be ruled out, since the concentration of the antagonist ZM-

1
2
3 241385 used in these studies is high enough to also antagonize rodent A₂B receptors [52].
4

5 Wound healing models have not been studied in ADA- deficient mice.
6
7

8 9 **Adenosine in neovascularization**

10
11
12
13
14 Revascularization of the wound bed is essential to supply oxygen, nutrients, and inflammatory
15 cells to the newly growing tissue. Two mechanisms contribute to the development of new vessels
16 in the adult: angiogenesis, the formation of new vessels from pre-existing ones; and
17 vasculogenesis, the initial series of events in vascular growth in which endothelial cell precursors
18 (angioblasts) differentiate in situ and assemble into solid endothelial cords [53] This multistep
19 process is highly regulated by a variety of soluble angiogenic growth factors, proteolytic
20 enzymes, which allow endothelial cell detachment and extracellular matrix invasion, and a close
21 interaction between adhesive proteins of the extracellular matrix and their integrin receptors [54,
22 55]. Among the growth factors implicated in the angiogenic process, bFGF and vascular
23 endothelial growth factor (VEGF) are known to be potent angiogenic molecules that induce the
24 growth of new blood vessels during wound healing and embryonic development. [56, 57].
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Diabetic patients frequently suffer from macro and microvascular complications characterized by
42 an early dysfunction of vascular endothelium that could further aggravate their impaired healing.
43
44
45

46
47
48 Many studies have shown that the administration of adenosine or adenosine agonists as well as
49 the upregulation of endogenous adenosine can increase the expression of vascular endothelial
50 growth factor (VEGF) in a variety of different cells studied in vitro [58-62] and after intravenous
51 infusion of adenosine in humans [63]. Adenosine A_{2A} receptors activation stimulates macrophage
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 production of VEGF as well [60], meanwhile adenosine A_{2B} receptors induce VEGF release by
4
5 retinal endothelial cells [59, 61].
6

7
8
9 The elevated expression of VEGF might account for the angiogenic effects of adenosine;
10
11 however, it is more likely that adenosine also stimulates angiogenesis via other secondary
12
13 angiogenic and antiangiogenic mediators or by way of an intracellular action [64]. Thus, the
14
15 adenosine analog NECA also increased the expression of the proangiogenic factors insulin-like
16
17 growth factor-I (IGF-I) and basic fibroblast growth factor (bFGF) in human retinal endothelial
18
19 cells as well as the expression of the proangiogenic factors interleukin-8 (IL-8) and angiopoietin-
20
21 2 in human mast cells. A_{2B} receptors mediated the IL-8 and bFGF responses and A₃ receptors
22
23 may have mediated the angiopoietin-2 response [65]. We have shown that adenosine and A_{2A}
24
25 agonists can promote in vitro angiogenesis by inhibiting endothelial secretion of the anti-
26
27 angiogenic factor thrombospondin-1 [66].
28
29
30
31
32
33
34
35

36 **Adenosine in remodeling**

37
38
39
40
41 The reparative process culminates with the remodelling of the newly formed granulation tissue in
42
43 order to restore complete functionality. During this extended phase, the provisional extracellular
44
45 matrix rich in type 3 collagen is degraded by serine proteases and metalloproteases and
46
47 sequentially substituted for the definitive matrix rich in type 1 collagen [1]
48
49
50

51
52
53 Activation of plasminogen plays a role in proteolytic degradation of extracellular matrices in
54
55 tissue remodelling events and it is required for normal repair of skin wounds in mice [67]. Earlier
56
57 reports showed that the non selective adenosine receptor agonist NECA increased plasminogen
58
59

1
2
3 activator release in rabbit alveolar macrophages due to intracellular elevations of cAMP [68].

4
5 Similarly, cAMP elevating agents increased the acute release of tissue plasminogen activator in
6
7 human umbilical endothelial cells [69, 70] and intraarterial infusions of ATP, an adenosine
8
9 precursor, to healthy volunteers induce tPA release [71].
10

11 12 13 **Adenosine agonist promotion of wound healing in animal models**

14
15
16
17
18
19 Much of the knowledge of the normal healing process of cutaneous wounds and the mediators
20
21 involved has evolved from information derived from experimental wounds in animals, especially
22
23 genetically modified mice. Unfortunately, a valid model of chronic wounds in animals has not yet
24
25 been developed. From the in vitro experiment data it is difficult to predict which adenosine
26
27 agonists, if any, will be useful for promoting impaired wound healing in humans.
28
29
30

31
32
33
34 There are few reports of the use of adenosine agonists for the enhancement of wound healing. We
35
36 have demonstrated pharmacologically and by the use of mice lacking A_{2A} receptors that topical
37
38 application of adenosine A_{2A} receptor agonists accelerate healing of dermal wounds in both
39
40 healthy animals and in streptozotocin-induced diabetic rats with impaired wound healing [18, 49,
41
42 72]. Histological analysis showed faster re-epithelialization and increased matrix deposition,
43
44 fibroblast density and vascularity in the granulation tissue of the agonist treated wounds as soon
45
46 as 3 days after injury [18]. In a model of pressure ulcer formation as a result of recurrent
47
48 ischemia-reperfusion of the skin, an adenosine A_{2A} receptor agonist infused via osmotic
49
50 minipumps reduced leukocyte infiltration and protected from ischemia/reperfusion injury in skin
51
52
53 [73]. No signs of fibrosis were described in these experimental animal models.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 When studying the dose response effect of the selective A_{2A} receptor agonist CGS-21680, a
4 narrow therapeutic window was observed. Both low (0.5 µg /wound) and high concentrations (10
5 µg /wound) did not affect the rate of wound closure; while intermediate doses were equally
6
7
8
9
10 effective. Loss of efficacy may be caused by the loss of specificity of CGS-21680 at higher
11
12 concentrations. In contrast Sonedenoson did not lose its accelerating effect at the highest
13
14 concentration studied. Moreover, it stimulated more rapid wound closure than CGS-21680,
15
16 suggesting the possibility that a higher selectivity of MRE0094 for A_{2A} receptors may be
17
18 important [72]. It is worth noting that the pharmacokinetics of these compounds administered
19
20 topically in a hydrophilic gel have not been determined and could have important repercussions
21
22 in the final effect, in terms of penetration or systemic absorption and elimination rate. In the same
23
24 study 0.01% Becaplermin gel (rhPDGF-BB), the only growth factor approved for clinical use, did
25
26 not accelerate the rate of wound closure as compared to the control, and both CGS-21680 and
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

36 Topical application of the non selective adenosine receptor agonist 5'-N-ethyl-
37
38 carboxamidoadenosine (NECA) also normalized the impaired healing induced by subcutaneous
39
40 injection of dexamethasone- in mice. This effect was only partially abrogated by a selective A_{2B}
41
42 antagonist, but the contribution of other receptors was not determined [45]. Topical application of
43
44 the adenosine A₁ receptor agonist, N(6)-Cyclopentyladenosine (CPA), also promoted healing of
45
46 incisional and excisional wounds on the dorsum of both diabetic (db/db) and normal (db/+) mice,
47
48 although the concentration used was above the selectivity threshold and the participation of other
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 Finally, using an incisional skin-wound model produced on the back of female diabetic mice,
6
7 intraperitoneal administration of Polydeoxyribonucleotide, a compound mixture of
8
9 deoxyribonucleotide polymers, improved the impaired wound healing and increased the wound-
10
11 breaking strength in diabetic mice through adenosine A_2 receptors activation. The promoting
12
13 effect of this compound was blocked by an antagonist, DMPX (3,7-dimethyl-1-
14
15 propargilxanthine), which blocks both A_{2A} and A_{2B} receptors [74].
16
17
18
19
20

21
22 As inferred from the in vitro studies previously described, many factors contribute to the effect of
23
24 adenosine receptor activation in promoting wound healing. The synergistic interaction between
25
26 Toll-like receptor and adenosine A_{2A} receptor signaling switches macrophages towards an
27
28 angiogenic phenotype [60, 75], and plays a role in an excisional wound healing model. Mice
29
30 lacking MyD88, an adapter protein in the signal transduction pathway of Toll-like receptors, heal
31
32 at a markedly slower rate than wounds in wild-type mice, showing delayed contraction, decreased
33
34 and delayed granulation tissue formation, and reduced new blood vessel density. CGS21680, an
35
36 A_{2A} receptor agonist, promoted wound closure and angiogenesis in wild type mice, but had no
37
38 significant effect on healing of MyD88(-/-) mice, suggesting the synergistic interaction between
39
40 Toll-like receptors and adenosine A_{2A} receptors signaling in wound healing in vivo [76]
41
42
43
44
45
46

47 **Conclusion**

48
49
50
51
52 Nowadays there is a high awareness of the problem that diabetic foot ulcers represent in terms of
53
54 costs and quality of life for those suffering them. Studies to better understand the normal healing
55
56 process and the pathology of impaired healing have been extremely useful to improve wound
57
58
59
60

1
2
3 management and care. Randomized controlled trials have shown that topical application of
4
5 becaplermin gel is effective in increasing healing rates for diabetic neuropathic foot ulcers with
6
7 adequate blood supply. However, this efficacy has not translated to positive clinical experience,
8
9 and the drug is not widely used. Moreover, it is an expensive medication of relative short storage
10
11 stability and an increased risk of mortality secondary to malignancy was observed in patients.
12
13 Adenosine receptor agonists represent an attractive and novel alternative to growth factors for
14
15 promotion of diabetic foot chronic ulcers. They are small synthetic molecules of longer stability
16
17 and lower cost than growth factors. Given the complexity of the pathogenesis of foot ulcers and
18
19 the differential adenosine receptor expression in the cells involved in the impaired tissue repair
20
21 process, it is difficult to predict which agonists, selective or not, could be most useful. To add
22
23 more complexity, all four receptor adenosine subtypes are G protein-coupled receptors, but also
24
25 exert cAMP-independent actions. In addition, they couple to mitogen-activated protein kinases by
26
27 mechanisms that appear to differ substantially, both between receptor subtypes in the same cell
28
29 type and between the same receptor in different cell types. From our studies, selective adenosine
30
31 A_{2A} receptor agonists present the advantages of reducing inflammation and increasing
32
33 vascularization and extracellular matrix deposition. Nevertheless, the fibrogenic potential and
34
35 increased risk of infection should also be monitored. Other important factors to be considered
36
37 are the pharmaceutical formulation and the pharmacokinetics of the agonist used, in order to
38
39 minimize its systemic effects. As any new strategy in the management of chronic wounds, the use
40
41 of adenosine agonists has to be evaluated and validated before taking it into practice.
42
43
44
45
46
47
48
49
50
51

52 **Acknowledgments**

53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 MCM is beneficiary of the Ramón y Cajal program from the Spanish Government (Ministerio de
4
5 Educación y Ciencia) and a grant from the Instituto de Salud Carlos III (FIS 05/1659) of the
6
7 Ministry of Science and Innovation, Government of Spain. B.N.C. is a consultant for King
8
9 Pharmaceuticals, Can-Fite Biopharma, Inc., Bristol-Myers, Squibb, and Tap Pharmaceuticals. He
10
11 is funded by grants from King Pharmaceuticals and the National Institutes of Health (AA13336,
12
13 AR41911, GM56268, and HL70952) and is the recipient of honoraria for speaking from Merck
14
15 Pharmaceuticals, Tap Pharmaceuticals, and Amgen.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. Pierce GF and Mustoe TA, Pharmacologic enhancement of wound healing. *Annual Review of Medicine* **46**: 467-481, 1995.
2. Romer J, Bugge TH, Pyke C, Lund LR, Flick MJ, Degen JL and Dano K, Plasminogen and wound healing. *Nat Med* **2**(7): 725, 1996.
3. Cha J and Falanga V, Stem cells in cutaneous wound healing. *Clin Dermatol* **25**(1): 73-8, 2007.
4. Werner S and Grose R, Regulation of wound healing by growth factors and cytokines. *Physiol Rev* **83**(3): 835-70, 2003.
5. Falanga V, Wound healing and its impairment in the diabetic foot. *Lancet* **366**(9498): 1736-43, 2005.
6. Hinchliffe RJ, Valk GD, Apelqvist J, Armstrong DG, Bakker K, Game FL, Hartemann-Heurtier A, Londahl M, Price PE, van Houtum WH and Jeffcoate WJ, A systematic review of the effectiveness of interventions to enhance the healing of chronic ulcers of the foot in diabetes. *Diabetes Metab Res Rev* **24 Suppl 1**: S119-44, 2008.
7. Gary Sibbald R and Woo KY, The biology of chronic foot ulcers in persons with diabetes. *Diabetes Metab Res Rev* **24 Suppl 1**: S25-30, 2008.
8. Khalil Z and Helme R, Sensory peptides as neuromodulators of wound healing in aged rats. *Journals of Gerontology Series A, Biological*(5): B354-61, 1996.
9. Schaper NC, Huijberts M and Pickwell K, Neurovascular control and neurogenic inflammation in diabetes. *Diabetes Metab Res Rev* **24 Suppl 1**: S40-4, 2008.
10. Khakh BS, Burnstock G, Kennedy C, King BF, North RA, Seguela P, Voigt M and Humphrey PPA, International Union of Pharmacology. XXIV. Current Status of the

- 1
2
3 Nomenclature and Properties of P2X Receptors and Their Subunits. *Pharmacol Rev*
4
5 **53(1)**: 107-118, 2001.
6
7
8 11. Fredholm BB, AP IJ, Jacobson KA, Klotz KN and Linden J, International Union of
9
10 Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol*
11
12 *Rev* **53(4)**: 527-52, 2001.
13
14
15 12. Schulte G and Fredholm BB, Signalling from adenosine receptors to mitogen-activated
16
17 protein kinases. *Cell Signal* **15(9)**: 813-27, 2003.
18
19
20 13. Burnstock G, Purinergic signalling. *Br J Pharmacol* **147 Suppl 1**: S172-81, 2006.
21
22
23 14. Jacobson KA and Gao Z-G, Adenosine receptors as therapeutic targets. **5(3)**: 247-264,
24
25 2006.
26
27
28 15. Feoktistov I, Goldstein AE, Ryzhov S, Zeng D, Belardinelli L, Voyno-Yasenetskaya T
29
30 and Biaggioni I, Differential expression of adenosine receptors in human endothelial
31
32 cells: role of A2B receptors in angiogenic factor regulation. *Circ Res* **90(5)**: 531-8, 2002.
33
34
35 16. Khoa ND, Montesinos MC, Reiss AB, Delano D, Awadallah N and Cronstein BN,
36
37 Inflammatory cytokines regulate function and expression of adenosine A(2A) receptors in
38
39 human monocytic THP-1 cells. *J Immunol* **167(7)**: 4026-32, 2001.
40
41
42 17. Chan ES, Fernandez P, Merchant AA, Montesinos MC, Trzaska S, Desai A, Tung CF,
43
44 Khoa DN, Pillinger MH, Reiss AB, Tomic-Canic M, Chen JF, Schwarzschild MA and
45
46 Cronstein BN, Adenosine A2A receptors in diffuse dermal fibrosis: pathogenic role in
47
48 human dermal fibroblasts and in a murine model of scleroderma. *Arthritis Rheum* **54(8)**:
49
50 2632-42, 2006.
51
52
53 18. Montesinos MC, Desai A, Chen JF, Yee H, Schwarzschild MA, Fink JS and Cronstein
54
55 BN, Adenosine promotes wound healing and mediates angiogenesis in response to tissue
56
57 injury via occupancy of A(2A) receptors. *Am J Pathol* **160(6)**: 2009-18, 2002.
58
59
60
61
62
63
64
65

- 1
2
3 19. Nguyen DK, Montesinos MC, Williams AJ, Kelly M and Cronstein BN, Th1 cytokines
4 regulate adenosine receptors and their downstream signaling elements in human
5 microvascular endothelial cells. *J Immunol* **171**(8): 3991-8, 2003.
6
7
8
9
10 20. Khoa ND, Postow M, Danielsson J and Cronstein BN, Tumor necrosis factor-alpha
11 prevents desensitization of Galphas-coupled receptors by regulating GRK2 association
12 with the plasma membrane. *Mol Pharmacol* **69**(4): 1311-9, 2006.
13
14
15
16
17 21. Xaus J, Mirabet M, Lloberas J, Soler C, Lluís C, Franco R and Celada A, IFN-gamma up-
18 regulates the A2B adenosine receptor expression in macrophages: a mechanism of
19 macrophage deactivation. *J Immunol* **162**(6): 3607-14, 1999.
20
21
22
23
24 22. Grden M, Podgorska M, Szutowicz A and Pawelczyk T, Diabetes-induced alterations of
25 adenosine receptors expression level in rat liver. *Exp Mol Pathol* **83**(3): 392-8, 2007.
26
27
28
29 23. Brownlee M, The pathobiology of diabetic complications: a unifying mechanism.
30
31 *Diabetes* **54**(6): 1615-25, 2005.
32
33
34 24. Lafuente N, Matesanz N, Azcutia V, Romacho T, Nevado J, Rodriguez-Manas L,
35 Moncada S, Peiro C and Sanchez-Ferrer CF, The deleterious effect of high concentrations
36 of D-glucose requires pro-inflammatory preconditioning. *J Hypertens* **26**(3): 478-85,
37
38
39
40
41
42
43 25. Montesinos M and Cronstein B, Role of P1 receptors in inflammation. In: *Handbook of*
44
45 *Experimental Pharmacology, Vol 151/II: Purinergic and Pyrimidinergic Signalling II*
46
47 *Cardiovascular, Respiratory, Immune, Metabolic and Gastrointestinal Tract Function*
48
49 (Eds. Abbrachio M and Williams M), pp. 303-321. Springer-Verlag, Berlin, 2001.
50
51
52
53 26. Hasko G, Linden J, Cronstein B and Pacher P, Adenosine receptors: therapeutic aspects
54
55 for inflammatory and immune diseases. *7*(9): 759-770, 2008.
56
57
58
59
60
61
62
63
64
65

- 1
2
3 27. Sitkovsky MV, Extracellular purines and their receptors in immunoregulation. Review of
4 recent advances. *Nippon Ika Daigaku Zasshi* **65**(5): 351-7, 1998.
5
6
7 28. Cronstein BN, Naime D and Ostad E, The antiinflammatory mechanism of methotrexate:
8 increased adenosine release at inflamed sites diminishes leukocyte accumulation in an in
9 vivo model of inflammation. *Journal of Clinical Investigation* **92**: 2675-2682, 1993.
10
11
12 29. Cronstein BN, Montesinos MC and Weissmann G, Salicylates and sulfasalazine, but not
13 glucocorticoids, inhibit leukocyte accumulation by an adenosine-dependent mechanism
14 that is independent of inhibition of prostaglandin synthesis and p105 of NFkappaB. *Proc*
15 *Natl Acad Sci U S A* **96**(11): 6377-81, 1999.
16
17
18 30. Morabito L, Montesinos MC, Schreiber DM, Balter L, Thompson LF, Resta R, Carlin
19 G, Huie MA and Cronstein BN, Methotrexate and Sulfasalazine Promote Adenosine
20 Release by a Mechanism that Requires Ecto-5'-nucleotidase-mediated Conversion of
21 Adenine Nucleotides. *J Clin Invest* **101**(2): 295-300, 1998.
22
23
24 31. Cronstein BN, Duguma L, Nicholls D, Hutchison A and Williams M, The
25 adenosine/neutrophil paradox resolved. Human neutrophils possess both A1 and A2
26 receptors which promote chemotaxis and inhibit O₂- generation, respectively. *Journal of*
27 *Clinical Investigation* **85**: 1150-1157, 1990.
28
29
30 32. Salmon JE, Brogle N, Brownlie C, Edberg JC, Kimberly RP, Chen BX and Erlanger BF,
31 Human mononuclear phagocytes express adenosine A1 receptors. A novel mechanism for
32 differential regulation of Fc gamma receptor function. *Journal of Immunology* **151**(5):
33 2775-85, 1993.
34
35
36 33. Lesch ME, Ferin MA, Wright CD and Schrier DJ, The effects of (R)-N-(1-methyl-2-
37 phenylethyl) adenosine (L-PIA), a standard A1-selective adenosine agonist on rat acute
38 models of inflammation and neutrophil function. *Agents & Actions* **34**: 25-27, 1991.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3 34. Bong GW, Rosengren S and Firestein GS, Spinal cord adenosine receptor stimulation in
4 rats inhibits peripheral neutrophil accumulation. The role of N-methyl-D-aspartate
5 receptors. *Journal of Clinical Investigation* **98**: 2779-2785, 1996.
6
7
8
9
10 35. Cronstein BN, Kramer SB, Weissmann G and Hirschhorn R, Adenosine: a physiological
11 modulator of superoxide anion generation by human neutrophils. *Journal of Experimental*
12 *Medicine* **158**: 1160-1177, 1983.
13
14
15
16
17 36. Bouma MG, Stad RK, van den Wildenberg FA and Buurman WA, Differential regulatory
18 effects of adenosine on cytokine release by activated human monocytes. *Journal of*
19 *Immunology* **153**(9): 4159-68, 1994.
20
21
22
23
24 37. Firestein GS, Paine MM and Boyle DL, Mechanisms of methotrexate action in
25 rheumatoid arthritis. Selective decrease in synovial collagenase gene expression. *Arthritis*
26 *and Rheumatism* **37**: 193-200, 1994.
27
28
29
30
31 38. Lennon PF, Taylor CT, Stahl GL and Colgan SP, Neutrophil-derived 5'-adenosine
32 monophosphate promotes endothelial barrier function via CD73-mediated conversion to
33 adenosine and endothelial A2B receptor activation. *J Exp Med* **188**(8): 1433-43, 1998.
34
35
36
37
38 39. Thompson LF, Eltzschig HK, Ibla JC, Van De Wiele CJ, Resta R, Morote-Garcia JC and
39 Colgan SP, Crucial role for ecto-5'-nucleotidase (CD73) in vascular leakage during
40 hypoxia. *J Exp Med* **200**(11): 1395-405, 2004.
41
42
43
44
45 40. Salvatore CA, Tilley SL, Latour AM, Fletcher DS, Koller BH and Jacobson MA,
46 Disruption of the A(3) adenosine receptor gene in mice and its effect on stimulated
47 inflammatory cells. *J Biol Chem* **275**(6): 4429-34, 2000.
48
49
50
51
52 41. Baharav E, Bar-Yehuda S, Madi L, Silberman D, Rath-Wolfson L, Halpren M, Ochaion
53 A, Weinberger A and Fishman P, Antiinflammatory effect of A3 adenosine receptor
54 agonists in murine autoimmune arthritis models. *J Rheumatol* **32**(3): 469-76, 2005.
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3 42. Montesinos MC, Desai A and Cronstein BN, Suppression of inflammation by low-dose
4 methotrexate is mediated by adenosine A2A receptor but not A3 receptor activation in
5 thioglycollate-induced peritonitis. *Arthritis Res Ther* **8(2)**: R53, 2006.
6
7
8
9
10 43. Hasko G, Kuhel DG, Chen JF, Schwarzschild MA, Deitch EA, Mabley JG, Marton A and
11 Szabo C, Adenosine inhibits IL-12 and TNF-alpha production via adenosine A2a
12 receptor-dependent and independent mechanisms [In Process Citation]. *Faseb J* **14(13)**:
13 2065-74, 2000.
14
15
16
17
18
19 44. Mi Q, Riviere B, Clermont G, Steed DL and Vodovotz Y, Agent-based model of
20 inflammation and wound healing: insights into diabetic foot ulcer pathology and the role
21 of transforming growth factor-beta1. *Wound Repair Regen* **15(5)**: 671-82, 2007.
22
23
24
25
26 45. Braun M, Lelieur K and Kietzmann M, Purinergic substances promote murine
27 keratinocyte proliferation and enhance impaired wound healing in mice. *Wound Repair*
28 *Regen* **14(2)**: 152-61, 2006.
29
30
31
32
33 46. Cook PW, Ashton NM and Pittelkow MR, Adenosine and adenine nucleotides inhibit the
34 autonomous and epidermal growth factor-mediated proliferation of cultured human
35 keratinocytes. *Journal of Investigative Dermatology* **104(6)**: 976-981, 1995.
36
37
38
39
40 47. Thellung S, Florio T, Maragliano A, Cattarini G and Schettini G,
41 Polydeoxyribonucleotides enhance the proliferation of human skin fibroblasts:
42 Involvement of A2 purinergic receptor subtypes. *Life Sciences* **64(18)**: 1661-1674, 1999.
43
44
45
46
47 48. Sun LL, Xu LL, Nielsen TB, Rhee P and Burris D, Cyclopentyladenosine improves cell
48 proliferation, wound healing, and hair growth. *J Surg Res* **87(1)**: 14-24., 1999.
49
50
51
52 49. Montesinos MC, Gadangi P, Longaker M, Sung J, Levine J, Nilsen D, Reibman J, Li M,
53 Jiang CK, Hirschhorn R, Recht PA, Ostad E, Levin RI and Cronstein BN, Wound healing
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3 is accelerated by agonists of adenosine A₂ (G alpha s- linked) receptors. *J Exp Med*
4
5 **186(9)**: 1615-20, 1997.
6
7
8 50. Fernandez P, Trzaska S, Wilder T, Chiriboga L, Blackburn MR, Cronstein BN and Chan
9
10 ES, Pharmacological blockade of A_{2A} receptors prevents dermal fibrosis in a model of
11
12 elevated tissue adenosine. *Am J Pathol* **172(6)**: 1675-82, 2008.
13
14 51. Katebi M, Fernandez P, Chan ES and Cronstein BN, Adenosine A_{2A} Receptor
15
16 Blockade or Deletion Diminishes Fibrocyte Accumulation in the Skin in a Murine Model
17
18 of Scleroderma, Bleomycin-induced Fibrosis. *Inflammation*, 2008.
19
20 52. Kreckler LM, Wan TC, Ge ZD and Auchampach JA, Adenosine inhibits tumor necrosis
21
22 factor-alpha release from mouse peritoneal macrophages via A_{2A} and A_{2B} but not the A₃
23
24 adenosine receptor. *J Pharmacol Exp Ther* **317(1)**: 172-80, 2006.
25
26 53. Asahara T, Murohara T, Sullivan A, Silver M, van der Zee R, Li T, Witzenbichler B,
27
28 Schatteman G and Isner JM, Isolation of putative progenitor endothelial cells for
29
30 angiogenesis. *Science* **275(5302)**: 964-7, 1997.
31
32 54. Naik MU, Mousa SA, Parkos CA and Naik UP, Signaling through JAM-1 and
33
34 $\{\alpha\}\nu\{\beta\}3$ is required for the angiogenic action of bFGF: dissociation of the JAM-1
35
36 and $\{\alpha\}\nu\{\beta\}3$ complex. *Blood* **102(6)**: 2108-2114, 2003.
37
38 55. Pepper MS, Role of the Matrix Metalloproteinase and Plasminogen Activator-Plasmin
39
40 Systems in Angiogenesis. *Arterioscler Thromb Vasc Biol* **21(7)**: 1104-1117, 2001.
41
42 56. Pepper MS, Rosnoblet C, Di Sanza C and Kruithof EK, Synergistic induction of t-PA by
43
44 vascular endothelial growth factor and basic fibroblast growth factor and localization of t-
45
46 PA to Weibel-Palade bodies in bovine microvascular endothelial cells. *Thromb Haemost*
47
48 **86(2)**: 702-9, 2001.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3 57. Ratel D, Mihoubi S, Beaulieu E, Durocher Y, Rivard G-E, Gingras D and Béliveau R,
4
5 VEGF increases the fibrinolytic activity of endothelial cells within fibrin matrices:
6
7 Involvement of VEGFR-2, tissue type plasminogen activator and matrix
8
9 metalloproteinases. *Thrombosis Research* **121**(2): 203-212, 2007.
10
11
12 58. Lutty GA, Mathews MK, Merges C and McLeod DS, Adenosine stimulates canine retinal
13
14 microvascular endothelial cell migration and tube formation. *Curr Eye Res* **17**(6): 594-
15
16 607, 1998.
17
18
19 59. Grant MB, Tarnuzzer RW, Caballero S, Ozeck MJ, Davis MI, Spoerri PE, Feoktistov I,
20
21 Biaggioni I, Shryock JC and Belardinelli L, Adenosine receptor activation induces
22
23 vascular endothelial growth factor in human retinal endothelial cells. *Circ Res* **85**(8): 699-
24
25 706., 1999.
26
27
28
29 60. Leibovich SJ, Chen JF, Pinhal-Enfield G, Belem PC, Elson G, Rosania A, Ramanathan
30
31 M, Montesinos C, Jacobson M, Schwarzschild MA, Fink JS and Cronstein B, Synergistic
32
33 up-regulation of vascular endothelial growth factor expression in murine macrophages by
34
35 adenosine A(2A) receptor agonists and endotoxin. *Am J Pathol* **160**(6): 2231-44, 2002.
36
37
38
39 61. Feoktistov I, Ryzhov S, Zhong H, Goldstein AE, Matafonov A, Zeng D and Biaggioni I,
40
41 Hypoxia modulates adenosine receptors in human endothelial and smooth muscle cells
42
43 toward an A2B angiogenic phenotype. *Hypertension* **44**(5): 649-54, 2004.
44
45
46
47 62. Clark AN, Youkey R, Liu X, Jia L, Blatt R, Day Y-J, Sullivan GW, Linden J and Tucker
48
49 AL, A1 Adenosine Receptor Activation Promotes Angiogenesis and Release of VEGF
50
51 From Monocytes. *Circ Res* **101**(11): 1130-1138, 2007.
52
53
54
55 63. Adair TH, Cotten R, Gu JW, Pryor JS, Bennett KR, McMullan MR, McDonnell P and
56
57 Montani JP, Adenosine infusion increases plasma levels of VEGF in humans. *BMC*
58
59 *Physiol* **5**: 10, 2005.
60
61
62
63
64
65

- 1
2
3 64. Adair TH, Growth regulation of the vascular system: an emerging role for adenosine. *Am*
4
5 *J Physiol Regul Integr Comp Physiol* **289**(2): R283-R296, 2005.
6
7
8 65. Feoktistov I, Ryzhov S, Goldstein AE and Biaggioni I, Mast cell-mediated stimulation of
9
10 angiogenesis: cooperative interaction between A2B and A3 adenosine receptors. *Circ Res*
11
12 **92**(5): 485-92, 2003.
13
14 66. Desai A, Victor-Vega C, Gadangi S, Montesinos MC, Chu CC and Cronstein BN,
15
16 Adenosine A2A receptor stimulation increases angiogenesis by down-regulating
17
18 production of the antiangiogenic matrix protein thrombospondin 1. *Mol Pharmacol* **67**(5):
19
20 1406-13, 2005.
21
22
23 67. Romer J, Bugge TH, Pyke C, Lund LR, Flick MJ, Degen JL and Dano K, Impaired
24
25 wound healing in mice with a disrupted plasminogen gene [see comments]. *Nature*
26
27 *Medicine* **2**(3): 287-292, 1996.
28
29
30 68. Hasday JD and Sitrin RG, Adenosine receptors on rabbit alveolar macrophages: binding
31
32 characteristics and effects on cellular function. *J.Lab.Clin.Med.* **110**: 264-273, 1987.
33
34
35 69. Hegeman RJ, van den Eijnden-Schrauwen Y and Emeis JJ, Adenosine 3':5'-cyclic
36
37 monophosphate induces regulated secretion of tissue-type plasminogen activator and von
38
39 Willebrand factor from cultured human endothelial cells. *Thromb Haemost* **79**(4): 853-8,
40
41 1998.
42
43
44 70. Santell L and Levin EG, Cyclic AMP potentiates phorbol ester stimulation of tissue
45
46 plasminogen activator release and inhibits secretion of plasminogen activator inhibitor-1
47
48 from human endothelial cells. *J Biol Chem* **263**(32): 16802-8, 1988.
49
50
51 71. Hrafnkelsdottir T, Erlinge D and Jern S, Extracellular nucleotides ATP and UTP induce a
52
53 marked acute release of tissue-type plasminogen activator in vivo in man. *Thromb*
54
55 *Haemost* **85**(5): 875-81, 2001.
56
57
58
59
60
61
62
63
64
65

- 1
2
3 72. Victor-Vega C, Desai A, Montesinos MC and Cronstein BN, Adenosine A2A receptor
4 agonists promote more rapid wound healing than recombinant human platelet-derived
5 growth factor (Becaplermin gel). *Inflammation* **26**(1): 19-24, 2002.
6
7
8
9
10 73. Peirce SM, Skalak TC, Rieger JM, Macdonald TL and Linden J, Selective A(2A)
11 adenosine receptor activation reduces skin pressure ulcer formation and inflammation. *Am*
12 *J Physiol Heart Circ Physiol* **281**(1): H67-74., 2001.
13
14
15
16 74. Galeano M, Bitto A, Altavilla D, Minutoli L, Polito F, Calo M, Lo Cascio P, Stagno
17 d'Alcontres F and Squadrito F, Polydeoxyribonucleotide stimulates angiogenesis and
18 wound healing in the genetically diabetic mouse. *Wound Repair Regen* **16**(2): 208-17,
19 2008.
20
21
22
23
24
25
26 75. Ramakers BP, Riksen NP, Rongen GA, van der Hoeven JG, Smits P and Pickkers P, The
27 effect of adenosine receptor agonists on cytokine release by human mononuclear cells
28 depends on the specific Toll-like receptor subtype used for stimulation. *Cytokine*, 2006.
29
30
31
32
33 76. Macedo L, Pinhal-Enfield G, Alshits V, Elson G, Cronstein BN and Leibovich SJ, Wound
34 Healing Is Impaired in MyD88-Deficient Mice: A Role for MyD88 in the Regulation of
35 Wound Healing by Adenosine A2A Receptors. *Am J Pathol* **171**(6): 1774-88, 2007.
36
37
38
39
40 77. Fredholm BB, Irenius E, Kull B and Schulte G, Comparison of the potency of adenosine
41 as an agonist at human adenosine receptors expressed in Chinese hamster ovary cells.
42 *Biochem Pharmacol* **61**(4): 443-8, 2001.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 **FIGURE LEGEND**
4

5 **Figure 1. Adenosine metabolism.** AICAR, 5-aminoimidazole-4-carboxamide ribonucleotide;
6
7 AICART, 5-aminoimidazole-4-carboxamide ribonucleotide transformylase; FAICAR, formyl 5-
8
9 aminoimidazole-4-carboxamide ribonucleotide; ADA Adenosine deaminase; EctoADA
10
11 Ectoadenosineaminase; CD73 Ecto-5'-nucleotidase; CD39 ecto-nucleoside triphosphate
12
13 diphosphohydrolase (NTPDase); ENT1 equilibrative nucleoside transporter 1.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Binding affinity of selected adenosine receptor agonists at the four receptor subtypes

Ki (nM)	A₁	A_{2A}	A_{2B}	A₃
Adenosine* [77]	310 (EC ₅₀)	700 (EC ₅₀)	24,000 (EC ₅₀)	290 (EC ₅₀)
2-Cl-adenosine [11]	1.39	180	N.D.	19
NECA[11]	14	20	330	6.2
CPA[11]	2.3	790	21,000	43
CGS21680[11]	290	27	361,000	67
ATL-146e [14]	77	0.5	N.D.	45
Sonedenoson (MRE0094)** [66, 72]	> 10,000 (IC ₅₀)	490 ± 50 (IC ₅₀)	> 10,000 (IC ₅₀)	N.D.
IB-MECA[77]	3.7	2,500	54,000	1.2

Ki, Dissociation constant of unlabeled compounds in radioligand competition experiments at recombinant human A₁, A_{2A}, A_{2B} and A₃ adenosine receptors in Chinese hamster ovary (CHO) cells.

* Adenosine data from a cyclic AMP functional assay in CHO cells stably transfected with recombinant human A₁, A_{2A}, A_{2B} and A₃ adenosine receptors.

** Data provided by King Pharmaceuticals at a variety of receptor systems by radioligand-binding studies.

N.D. Not determined or Not Disclosed

NECA 5'-N-ethyl-carboxamidoadenosine

CPA N6-cyclopentyladenosine

CGS 21680 2-[p-(2-carbonyl-ethyl)-phenylethylamino]-5'-N-ethylcarboxamidoadenosine

ATL146e 4-{3-[6-amino-9-(5-ethylcarbamoyl-3,4-dihydroxy-tetrahydro-furan-2-yl)-9H-purin-2-yl]-prop-2-ynyl}-cyclohexanecarboxylic acid methyl ester

Sonedenoson (MRE0094) 2-[2-(4-chlorophenyl)ethoxy] adenosine

IB-MECA N6-(3-iodobenzyl)adenosine-5'-N-methyluronamide

Accepted Manuscript

