



**HAL**  
open science

## Basin tectonics during the Early Cretaceous in the Levant margin, Lebanon

C. Homberg, E. Barrier, M. Mroueh, W. Hamdan, F. Higazi

► **To cite this version:**

C. Homberg, E. Barrier, M. Mroueh, W. Hamdan, F. Higazi. Basin tectonics during the Early Cretaceous in the Levant margin, Lebanon. *Journal of Geodynamics*, 2009, 47 (4), pp.218. 10.1016/j.jog.2008.09.002 . hal-00531893

**HAL Id: hal-00531893**

**<https://hal.science/hal-00531893>**

Submitted on 4 Nov 2010


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript

Title: Basin tectonics during the Early Cretaceous in the Levant margin, Lebanon

Authors: C. Homberg, E. Barrier, M. Mroueh, W. Hamdan, F. Higazi



PII: S0264-3707(08)00079-3  
DOI: doi:10.1016/j.jog.2008.09.002  
Reference: GEOD 866

To appear in: *Journal of Geodynamics*

Received date: 13-2-2008  
Revised date: 5-9-2008  
Accepted date: 5-9-2008

Please cite this article as: Homberg, C., Barrier, E., Mroueh, M., Hamdan, W., Higazi, F., Basin tectonics during the Early Cretaceous in the Levant margin, Lebanon, *Journal of Geodynamics* (2008), doi:10.1016/j.jog.2008.09.002

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Basin tectonics during the Early Cretaceous in the Levant margin, Lebanon.**

2

3 C. Homberg <sup>1\*</sup>

4 E. Barrier <sup>1</sup>

5 M. Mroueh <sup>2</sup>

6 W. Hamdan <sup>2</sup>

7 F. Higazi <sup>2</sup>

8 1 : Université Pierre et Marie Curie, Laboratoire de Tectonique, UMR7072, Case 129, 4 place jussieu,  
9 75252 Paris Cedex 05, France

10 2 : Université libanaise, Faculté d'Agronomie, B.P. 13-5368 Chourane, Beyrouth 1102-2040 Lebanon.

11 \* corresponding author

12 **ABSTRACT**

13 We present new brittle tectonic data constraining the onset of formation of the eastern passive  
14 margin of the Levant basin (Eastern Mediterranean basin) in Lebanon. From the identification of syn-  
15 tectonic growth faults, we infer an extensional tectonic regime starting in the Early Cretaceous and  
16 ceasing during the Cenomanian. The related stress field had a NNE-SSW direction of extension. It  
17 produced WSW-ENE to WNW-ESE normal faults with offsets as large as several hundred meters. Late  
18 Jurassic volcanic activity preceded this rifting event and continued until the late Aptian. Thickness and  
19 facies variations of the Upper Cretaceous sequence indicate that this rifting event led to the development  
20 of an E-W basin in Lebanon. This basin deepens westward, with a possible offshore continuation. The  
21 significant obliquity between the ~NE-SW Early Mesozoic faults in southeastern corner of the Levant  
22 basin and ~E-W Early Cretaceous faults recognized in Lebanon indicates that the mechanisms driving the  
23 development of the Eastern Mediterranean basin drastically changed during the Mesozoic.

24

25 **KEY WORDS** : Levant basin, Eastern Mediterranean Basin, Neotethys, Early Cretaceous extension, rift  
26 tectonics, Lebanon.

27

## 28 **1. Introduction**

29 The Levant basin (LB), the easternmost part of the Eastern Mediterranean basin (EMB), is  
30 generally regarded as a basin that resulted from rifting. This is supported by crustal thinning from 30-35  
31 km on the Africa and Arabia continents (Makris et al., 1988) to ~8km below the LB overlain by a 10-14  
32 km thick sedimentary pile of probably Jurassic to Present age (Makris et al., 1983; Ginzburd and Ben-  
33 Avraham, 1987; Vidal et al., 2000; Ben-Avraham et al., 2002). However, several aspects of the history of  
34 the LB remain unsolved. First, the affinity of the crust below the LB is regarded either as highly stretched  
35 continental (Woddsides et al., 1977; Hirsch et al., 1995; Robertson et al., 1996; Vidal et al., 2000) or as  
36 oceanic (Ginzburd and Ben-Avraham, 1987 and Garfunkel, 1998). Second, various ages for the opening of  
37 the basin have been proposed: from Triassic or Late Permian (Freund, 1975; Garfunkel, 1998 and Stampfli  
38 et al., 2002), Jurassic (Ginzburg and Guitzman, 1979) to Cretaceous (Dercourt et al., 1986). Third, some  
39 difficulties arise in reconciling the kinematic models that predicts a N-S opening of the basin (Dercourt et  
40 al., 1986; Stampfli et al., 2002) with tectonic structures such as Mesozoic NE-SW faults recognized in the  
41 LB and along its margins (Vidal et al., 2000).

42 These unsolved issues permit a variety of plate tectonic models of the western Neotethys. A major  
43 difficulty in getting relevant information on the LB is attenuation of seismic signals by Messinian  
44 evaporates, thus precluding good imaging of the underlying Mesozoic strata and structures and enhancing  
45 the uncertainties on the reflectors. This paper presents new observations in the Mesozoic sedimentary  
46 succession of the eastern passive margin of the LB, in Lebanon. After a review of the regional structures,  
47 we present arguments for an Early Cretaceous extensional tectonic event in Lebanon. Comparing  
48 published data and those of this paper, we then discuss the tectonic history and setting of the LB.

49

## 50 **2. Geological frame of Lebanon**

### 51 *2. 1. Tectonic structures of Lebanon*

52 The Levant margin is now the active left-lateral transform boundary between the Nubia and Arabia  
53 plates, namely the Dead Sea Fault System (DSFS). The DSFS developed during Late Cenozoic times with  
54 a roughly N-S direction and now connects the Red Sea Rift basin in the south to the Arabia-Eurasia-Nubia  
55 triple junction in the north (fig. 1). About 100 km of left-lateral slip have been suggested for the southern  
56 DSFS (Quennel, 1958; Freund et al., 1970). The main fault of the DSFS in Lebanon is the ~150 km-long  
57 left-lateral NNE-SSW Yammouneh fault. The N-S Roum fault and the NE-SW Rachaya and Sergaya  
58 faults are secondary faults, with more moderate offsets (Butler et al., 1988). The Meso-Cenozoic sequence  
59 is folded in three wide NNE-SSW folds that are, from west to east, the Mount Lebanon Anticline, the  
60 Bekaa syncline, and the Anti-Lebanon anticline (fig. 1). These folds are thought to have accommodated  
61 the shortening imposed by the obliquity of the Nubia-Arabia plate motion (e. g., DeMets, 1990; Jestin et  
62 al., 1994) relative to the strike of the Yammouneh fault (Freund et al., 1970; Garfunkel, 1981; Butler et al.,  
63 1988). Local NE-SW and NW-SE faults also exist. They are thought to be minor dextral and sinistral  
64 faults, respectively, related to the transform tectonics (Hancock and Atiya, 1979). The Cenozoic tectonics  
65 exhumed and translated the Mesozoic structures away from their initial paleogeographic positions.  
66 Although strike-slip movements occurred on several faults in Lebanon, a large part of the northward  
67 translation was absorbed along the Yammouneh fault (Walley, 1998). Because most of our observations  
68 were done west of this fault, and thus in the ‘Nubia attached’ domain, integration of the Mesozoic  
69 structures in Lebanon in the geodynamic context of the opening of the EMB does not necessitate taking  
70 into account the Cenozoic movements.

71

### 72 *2. 2. The Mesozoic sequence*

73 The Mesozoic sequence in Lebanon crops out in the cores of the Mount Lebanon and Anti-  
74 Lebanon anticlines (fig. 1). The Jurassic and Lower Cretaceous sequences consist of shallow marine

75 carbonates or continental sandstones. The oldest outcropping levels are Lower to Middle Jurassic  
76 limestones or dolomites. They are overlain by a thick sandy sequence, the Chouf sandstones (called Grès  
77 de Base by previous authors). These fluvial deposits are Neocomian to Barremian in age (Dubertret,  
78 1975); the first levels postdate the early Valanginian (Ferry, personal communication). The overlying  
79 Aptian and Albian formations are shallow marine carbonates, or locally sandstones and marls. They  
80 include a remarkable marker-bed that is the lagoonal Jezine Formation (previously known as Barre de  
81 Blanche), uppermost early Aptian in age. The alternation of shallow water carbonates and deeper marls of  
82 the Cenomanian and Turonian sequence precedes widespread marine flooding at Senonian time and  
83 deposition of chalky limestones. The pioneer Lebanese workers (e. g., Saint Marc, 1974; Dubertret, 1975)  
84 recognized that the Mesozoic sequence exhibits, at the Levant basin scale, a westward thickening and  
85 facies evolution from shallow marine sediments onshore to deep pelagic sediments offshore. This led  
86 some authors to propose that the present-day western limb of the Mount Lebanon anticline was the eastern  
87 margin of the Levant basin during the Mesozoic, trending, therefore, NNE-SSW (e. g., Walley, 1998).  
88 However, the Lower Cretaceous sequence also exhibits a N-S thickness variation. It is particularly  
89 spectacular for the Chouf sandstones, whose thickness reach 300m in Central Lebanon (Chouf area) and is  
90 reduced to a few tens of meters in northern Lebanon. The geometry and the opening direction of the  
91 Lebanese sector of the LB are thus difficult to establish solely on the basis of the sedimentological  
92 architecture. We thus sought out direct arguments for tectonic activity during Mesozoic times.

### 93 94 **3. Early Cretaceous extensional tectonics**

#### 95 *3. 1. Dating arguments and faulting*

96 In order to constrain the Mesozoic tectonics in Lebanon, we examined the faults that cut the  
97 Mesozoic formations. Complete or partial sections of the Middle Jurassic to Cenomanian sequence are  
98 visible along the large valleys that incise the core of the Mount Lebanon anticline. Most faults in this area  
99 trend WSW-ENE to WNW-ESE as mapped in the 1/200 000 geological map of Dubertret (1955). They

100 are particularly well-developed in the Chouf area and are a few tens of kilometers long. These faults dip  
101 either to the North or to the South, with a regular  $\sim 60^\circ$  angle (Fig. 2). They offset vertically the Upper  
102 Jurassic to Lower Cretaceous beds of several tens to hundreds of meters. Because the strata are sub-  
103 horizontal, their offsets cannot result from a pure strike-slip fault movement. When the fault plane is  
104 visible ( $\sim 50\%$  of the faults), the striae inclination is systematically close to  $90^\circ$ , indicating that faults are  
105 dip-slip normal faults (fig. 3). It follows that most WSW-ENE to WNW-ESE faults cutting the Mesozoic  
106 series were produced by an extensional event the age of which is now discussed.

107 When outcrop conditions allowed observation of the whole Lower to Middle Cretaceous sequence,  
108 the Cenomanian beds sealed most of the normal faults, as illustrated in figure 2. A few faults enter into the  
109 basal Cenomanian, which generally forms the top of the cross-sections. According to the geological maps  
110 of Dubertert, these faults do not extend more than a few hundred meters from the Albian-Cenomanian  
111 boundary. St-Marc (1970) showed that the first Cenomanian levels are in fact Late Albian in age.  
112 Therefore, the extensional tectonics that produced the WSW-ENE to WNW-ESE normal faults ended just  
113 prior to the Cenomanian. Our observations in the younger series confirm tectonic quiescence during the  
114 Late Cretaceous. Some NE-SW and NW-SE faults, well developed in southern Lebanon, cut the Upper  
115 Cretaceous and Eocene succession (fig. 1). Because our paper focuses on Mesozoic structures, we do not  
116 further discuss these Cenozoic features.

117 In order to define the time span of this extensional tectonic event, the whole Mesozoic sedimentary  
118 succession was examined for normal growth faults. The oldest such faults were observed in the Chouf  
119 sandstones. Basaltic lenses interlayered in these fluvial deposits indicate that volcanic activity  
120 accompanied the extensional tectonics. The normal fault shown in figure 4 illustrates that normal faulting  
121 continued during Aptian and Albian times. No growth faults were recognized in the Middle to Late  
122 Jurassic levels. However, such structures may be difficult to document in this poorly bedded sequence.  
123 We thus do not exclude normal faulting during Late Jurassic time, but believe that significant vertical  
124 movements did not start not before Early Cretaceous. The widespread Late Oxfordian to Early (Mid?)

125 Kimmeridgian basalts and tuffs attest to a regional volcanic event that immediately pre-dates the Early  
126 Cretaceous phase of extensional tectonics. Extensional tectonics documented in Lebanon thus started in  
127 the Early Cretaceous and ended at in the late Albian (or early Cenomanian). The WSW-ENE to WNW-  
128 ESE orientation of the several kilometer long normal faults suggests a rough N-S direction of extension.

129

### 130 3.2. *Stress field*

131 Because the sole direction of normal faults does not necessarily indicate an accurate direction for  
132 the driving stresses, fault-slip data on meso-scale faults cutting the Lower Cretaceous and older formations  
133 at 29 sites were also measured. Fault-slip data were collected on faults with offsets of 1-100 mm and,  
134 when possible, on Early Cretaceous growth faults. The local stress states were inferred from the inversion  
135 of faults using the method of Angelier et al. (1990), which is based upon the Wallace–Bott hypothesis that  
136 faults slip in the direction of the resolved shear traction. This method allows the determination of the  
137 orientation of the three principal stresses,  $\sigma_1$ ,  $\sigma_2$  and  $\sigma_3$ , as well as a shape ratio between principal stress  
138 differences, without any assumption on material strength. The uncertainty on the stress axis orientation  
139 depends on the 3-D distribution of the fault-slip data. When fault-slip data are of various attitudes and  
140 include conjugate sets, the accuracy on the direction and plunge of the principal axes is  $\sim 10^\circ$ .

141 The fault population includes strike-slip faults at all the visited sites and includes normal faults at  
142 18 sites. Strike-slip faults were disregarded because their compatibility with Dead Sea Fault System  
143 tectonics suggests that they formed during the Cenozoic. The normal faults found at 18 sites typically  
144 strike between N060°E and N140°E (azimuthal range described in clockwise sense). Their dips generally  
145 exceed 55°, with 77% of them dipping between 60° and 85°. Fault inversion yields normal stress tensors  
146 in which the minimum and intermediate principal stresses are sub-horizontal and the maximum stress is  
147 sub-vertical. The rose diagram in figure 5 illustrates the azimuthal distribution of the minimum horizontal  
148 stress axis ( $\sigma_3$ ). The azimuth of  $\sigma_3$  spreads out the N029°E arithmetic mean and lies between N152°E and  
149 N064°E. This range is rather large, but 71% of the stress states have a  $\sigma_3$  direction between N160°E and



150 N040°E. The data are therefore quite consistent and indicate a NNE-SSW direction of extension. Data out  
151 of this range may reflect local stress deflections.

152 At five sites, we could establish that the calculated stress states are Early Cretaceous in age. At two  
153 of these, the fault population includes a majority of fault-slip data on Early Cretaceous growth faults with  
154 offsets in the range of meters. At the other three, faults were formed in the close vicinity of large-scale  
155 Early Cretaceous normal growth faults showing offsets of several or tens of meters. Because faulting  
156 generally implies a variety of fracture scales, we are confident that slip on these meso-scale faults reflects  
157 the same mechanisms as the large-scale faulting. For the remaining 13 sites, we could not establish with  
158 confidence the absolute age of the normal fault-slips. However, for three of them situated in highly  
159 dipping strata, the calculated stress states clearly predated the Neogene folding (fig. 5). Indeed,  
160 measurement of present-day stresses (e.g., Cornet et Burlet, 1992; Brudy et al., 1997) and paleostress  
161 reconstructions (e. g., Homberg et al., 2002) support the hypothesis that stresses follow the Anderson  
162 model in which two of the principal stresses are horizontal, the third one being vertical. For the three sites  
163 discussed above, this criterion is fulfilled when performing the stress inversion on faults with their  
164 backtilted attitude (faults rotated around the local strike of bedding by the amount of tilting). We attributed  
165 the calculated stress states in the remaining 13 sites to the Early Cretaceous extension, although we cannot  
166 firmly exclude that some of them may reflect a later tectonic event. However, because these 13 stress  
167 states together show a NNE-SSW direction of extension that is compatible with the strike of the large-  
168 scale WSW-ENE to WNW-ESE normal growth faults, we believe that they also reflect the driving  
169 mechanism of Early Cretaceous tectonics.

#### 171 **4. Discussion and conclusion**

172 Our investigation of the Mesozoic structures indicates that a phase of extensional tectonics  
173 occurred in Lebanon during the Early Cretaceous. It started during the deposition of the Chouf sandstones,  
174 and thus during late Valanginian or Hauterivian times, and ceased during the Cenomanian. This NNE-

175 SSW extension produced WSW-ENE to WNW-ESE normal faults with length of several kilometers to  
176 several tens of kilometers and with offsets as large as several hundred meters. In view of these fault data,  
177 we interpret the regional northward thinning (and facies evolution) of the Lower Cretaceous sedimentary  
178 sequence as the result of the development of a roughly E-W striking basin in Lebanon. This interpretation  
179 differs from that of Walley (1998) who interpreted the westward thickening of the Mesozoic sequence  
180 near the coast line as a NNE-SSW structural grain, with no reference to specific faults. This interpretation  
181 seems no longer valid considering that the observed abrupt changes of the sequence thickness are  
182 controlled by E-W faults (fig. 4) and that the regional extensional direction trends NNE-SSW (fig. 5). The  
183 westward thickening implies rather that the E-W Lebanese basin continues and deepens westward, that is  
184 offshore through the Levant basin. The axis of the Lebanese basin was situated somewhere in central (or  
185 southern) Lebanon, most probably in the Chouf region. Here, the Lower Cretaceous succession reaches its  
186 maximum thickness of about 700 m and is offset by numerous normal growth faults. The northern margin  
187 of the basin was situated ~ 50km to the North, south of Tripoli, where fluvial and shallow marine  
188 sedimentation was strongly reduced and sometimes replaced by lavas flows. The southern margin is not  
189 visible due to later Paleogene deposits that cover southern Lebanon. Late Jurassic volcanic activity  
190 preceded the development of the basin and continued until the late Aptian. Quite remarkable is that Early  
191 Cretaceous tectonics produced a combination of WSW-ENE and WNW-ESE faults in Lebanon (fig. 1).  
192 Considering the mean NNE-SSW direction of extension inferred from the fault analysis, the WSW-ENE  
193 faults may reflect an earlier structural fabric. Such a fabric is well-developed eastward in Syria in the  
194 WNW-ESE Permian to Early Mesozoic Palmyride Trough (e. g., Brew et al., 2001).

195 As discussed above, the westward increase of the thickness of the Lower Cretaceous succession  
196 described by Walley (1998) suggests that the E-W Lebanese basin extends further offshore into the deep  
197 Levant basin (LB), although its structural continuity may have been slightly disrupted by offshore  
198 Cenozoic faulting. Extrapolating our investigation in Lebanon, the LB and more generally the Eastern  
199 Mediterranean Basin (EMB) registered a major extensional event during the Early Cretaceous. Although

200 the Early Cretaceous is marked by intense volcanism in Lebanon, our data do not allow us to decipher  
201 whether oceanic spreading occurred or not in the easternmost part of the EMB. On the other hand, they  
202 exclude that the EMB had already attained its present configuration during the Early Cretaceous time as  
203 claimed by Garfunkel (1998). Taking a full 0.5 cm/yr opening rate during the about 40 Myr of extension  
204 recorded in Lebanon, the EMB widened by 200km during the Early Cretaceous. This value is hypothetical  
205 and can be much smaller, especially if the basin did not reach the level of sea-floor spreading. Our data  
206 also suggest that the mechanisms driving the development of the EMB drastically changed during the  
207 Mesozoic (fig. 6). A WNW-ESE direction of opening during the Triassic and Jurassic was proposed by  
208 Garfunkel (1998) and Walley (1998), in the light of the NNE-SSW pre-Late Permian, Early to Middle  
209 Triassic, and Early to Middle Jurassic grabens onshore and offshore of Israel (see Cohen et al., 1990 for a  
210 review). This direction is almost perpendicular to the Early Cretaceous NNE-SSW extension documented  
211 in Lebanon, also recognized but with a N-S strike in Egypt (Hantar, 1990; Fawzy and Dahi, 1992). This  
212 change in the direction of opening, from a mean WNW-ESE to a NNE-SSW direction, occurred  
213 somewhere in the Late Jurassic. In addition to this polyphase development, the absence of significant  
214 Early Cretaceous faulting in the southern Levant margin (Israel-Jordan) could indicate that the locus of  
215 extension in the LB migrated northward.

216

### 217 **Acknowledgements**

218 This work was supported by the MEBE (Middle East Basins Evolution) Program and partially by the  
219 French and Lebanese CNRS, and the Lebanese University.

220

### 221 **References**

222 Angelier, J., 1990, Inversion of field data in fault tectonics to obtain the regional stress. III: A new rapid  
223 direct inversion method by analytical means: *Geophysical Journal International*, v. 103, 363-376, 1990.

- 224 Ben Avraham Z., Ginzburh, A., Makris, J., Eppelbaum, 2002, Crustal structure of the Levant Basin,  
225 eastern Mediterranean: Tectonophysics, v. 346, 23-43.
- 226 Brew, G., Barazangi, M., Al-Maleh, A., K., Sawaf, T., 2000. Tectonic map and geologic evolution of  
227 Syria: GeoArabia, 6, 573-616.
- 228 Brudy, M., Zoback, M., D., Fuchs, K., Rumel, F., Baumgärtner, J., 1997, Estimation of the complete stress  
229 tensors to 8km depth in the KTB scientific drill holes: implications for crustal strength: Journal of  
230 Geophysical Research, v. 102, 18453-18475.
- 231 Buttler, R., W., H., Spencer, S., Griffiths, H., M., 1998, The structural response to evolving plate  
232 kinematics during transpression: evolution of the Lebanese restraining bend of the Dead Sea Transform: *in*  
233 Holdsworth, R. E., Strachan, R. A., Dewey, J.- F., eds., Continental Transpressional and Transtensional  
234 Tectonics: Geological Society (London) Special Publication 135, 81-106.
- 235 Cohen, Z., Kaptan, V., Flexer, A., 1990, The tectonic mosaic of the southern Levant: implications for  
236 hydrocarbon prospects: Journal of Petroleum Geology, v. 13, 437-462.
- 237 Cornet F., H., Buret, D., 1992, Stress field determinations in France by hydraulic tests in boreholes:  
238 Journal of Geophysical Research, v. 97, 11829-11849.
- 239 Dercourt, J., Zonenshai, L., P., Ricou, L., E., Kazmin, V., G., Le Pchon, X., Knipper, A., L.,  
240 Grandjacquet, C., Sbertshikov, D., Westphal, M., Bazhenov, M., L., Laurer, J., P., Bijou-Duval, B., 1986,  
241 Geological evolution of the Tethys belt from the Atlantic to the Pamirs since the Lias: Tectonophysics, v.  
242 123, 241-315.
- 243 Dubertret, L. and collaborators, 1955. Carte géologique du Liban au 1/200 000. République libanaise,  
244 Ministère des Travaux publics, Institut géographique national.
- 245 Dubertret, L., 1975, Introduction à la carte géologique a 1/50 000e du Liban : Notes et Mémoires sur le  
246 Moyent-Orient ; v. 13, 345-403.
- 247 Freund, R., Garfunkel, Z., Zak, I, Goldberg, M., Weissbrod, T., Derin, B., 1970, The shear along the Dead  
248 Sea Rift: Philosophical Transactions of the Royal Society of London, v. 267, 107-130.

- 249 Freund, R., Goldberg, M., Weissbrod, T., Druckman, Y., Derin, B., 1975, The Triassic-Jurassic structure of  
250 Israel and its relation to the origin of the eastern Mediterranean: Geological Survey of Israel, v. 65, 1-26.
- 251 Garfunkel, Z., 1998, Constrains on the origin and history of the Eastern Mediterranean basin:  
252 Tectonophysics, v. 298, 5-35.
- 253 Garfunkel, Z., 1981, Internal structure of the Dead Sea leaky transform (rift) in relation to plate  
254 kinematics: Tectonophysics, v. 80, 81-108.
- 255 Ginzburg, A., and Gvirtzman, G., 1979, Changes in the crust and in the sedimentary cover across the  
256 transition from the Arabia platform to the Mediterranean basin: evidence from seismic refraction and  
257 sedimentary studies in Israel and Sinai: Sedimentary Geology, v. 23, 19-36.
- 258 Ginzburg, A., and Ben-Avraham, Z., 1987, The deep structure of the central and southern Levant  
259 continental margin: Annales Tectonicae, v. 1, 105-115.
- 260 Hancock, P., L., Atiya, M., S., 1979, Tectonic significance of mesofracture systems associated with the  
261 Lebanese segment of the Dead Sea transform fault: Journal of Structural Geology, v. 1, 143-153.
- 262 Hirsch, F., Flexer, A., Rosenfeld, A., Yellin-Dror, A., 1995, Palaeozoic crustal setting of the Eastern  
263 Mediterranean: Journal of Petroleum Geology, v. 18, 149-170.
- 264 Homberg, C., Bergerat, F., Philippe, Y., Lacombe, O., and Angelier, J., 2002, Structural inheritance and  
265 Cenozoic stress fields in the Jura fold-and-thrust belt (France): Tectonophysics, v. 327, 137-158, 2002.
- 266 Jestin, F., Huchon, P., Gaulier, J. M., 1994, The Somalia plate and the eastern Africa Rift System, present-  
267 day kinematics: Geophysical Journal International, v. 116, 637-654.
- 268 Makris, J., Ben-Avraham, Z., Behle, A., Ginzburg, A., Giese, A., Steinmetz, L., Whithmarsh, R., B.,  
269 Eleftheriou, S., 1983, Seismic refraction profiles between Cyprus and Israel and their interpretations:  
270 Geophysical Journal of the Royal Astronomical Society, v. 75, 575-591.
- 271 Makris, J., Rihm, R., Allam, A., 1988, Some geophysical aspects of the evolution and structure of the  
272 crust in Egypt: in El-Gaby, S., Greiling, R., O., eds., The Pan-African belt of the Northeast Africa and

- 273 adjacent areas: International monograph series on interdisciplinary earth science and applications, 345-  
274 369.
- 275 Quennell, A., M., 1958, The structure and the evolution of the Dead Sea Rift: Quarterly Journal of the  
276 Geological Society of London, v. 64, 1-24.
- 277 Robertson, A.H.F., Dixon, J., E., Brown, S., Collins, A., Morris, A., Pickett, E., Sharp, I., and Ustaomer,  
278 T., 1996, Alternative tectonic models for the Late paleozoic-Early Tertiary development of the Tethys in  
279 the Eastern Mediterranean region: *in* Morris, A., and Tarling, D., H., eds., Paleomagnetism and tectonics of  
280 the Mediterranean region: Geological Society (London) Special Publication 105, 239-263.
- 281 Saint-Marc, P., 1974, Etude stratigraphique et micropaléontologique de l'Albien, du Cénomaniens et du  
282 Turonien du Liban : Notes Mém. Moyen-Orient, v. 8, 8-342.
- 283 Saint-Marc, P., 1970, Le Crétacé inférieur et moyen du bord occidental du Jabl Sannine (Liban) : Notes et  
284 Mém. Moyen Orient, v. 12, 217-226.
- 285 Stampfli, G., M., and Borel, G., D., 2002, A plate tectonic model for the Paleozoic and Mesozoic  
286 constrained by dynamic plate boundaries and restored synthetic oceanic isochrones: Earth and Planetary  
287 Science Letters, v. 196, 17-33.
- 288 Vidal, N., Alvarez-Marron, J., Klaeschen, D., 2000, Internal configuration of the Levantine Basin from  
289 seismic reflection data (Eastern Mediterranean). Earth and Planetary Science Letters, v. 180, 77-89.
- 290 Walley, C., 1998, Some outstanding issues in the geology of Lebanon and their importance in the tectonic  
291 evolution of the Levantine region: Tectonophysics, v. 298, 1, 37-62.
- 292 Woodside, J., M., 1977, Tectonic elements and crust of the eastern Mediterranean Sea. Marine  
293 Geophysical Researches, v. 3, 317-354.

294

295 Fig. 1. Kinematic frame of Lebanon A: Simplified tectonic map of the Eastern Mediterranean domains Ar,  
 296 Eu, Nu: Arabia, Eurasia, Nubia plates. DSF; Dead Sea Fault. RS and CZ: Red Sea rift and Arabia-  
 297 Eurasia collision zone. LV: Levant Basin. B: Main structures of Lebanon. YF, RoF, RaF, SF: Yamouneh,  
 298 Roum, Rachaya, Sergaya fault. Be, Ba, Tr: cities of Beirut, Balbeck, Tripoli.

299

300 Fig. 2. Normal faults cutting the Jurassic to Lower Cretaceous sequences. See fig. 1 for location. The  
 301 uppermost Albian beds seal the normal faults. Jf: Jezine formation.

302

303 Fig. 3. Early Cretaceous normal fault plane cutting the Jezine formation. Site of Kafer Hachno (see fig. 1  
 304 for location). The striae or slip vector (thin arrows) is aligned with the maximum dip. View is to the  
 305 N°30°E. Meso-scale faults collected in this site and corresponding stress state are shown. Continuous  
 306 lines: fault planes. Slickenside lineations in dots with double arrows for strike-slip motion and outward-  
 307 directed single arrow for normal motion. Gray stars with 5, 4, 3 arms:  $\sigma_1$ ,  $\sigma_2$ , and  $\sigma_3$ , respectively.  
 308 Divergent large black arrows show directions of  $\sigma_3$ .

309

310 Fig. 4. Aptian-Albian growth fault. Inset shows how the growth fault split upward into two branches, the  
 311 southern making of clockwise angle with the single deep fault. Jf: Jezine formation. See fig. 1 for location.

312

313 Fig. 5. Stress states during the Early Cretaceous extension in Lebanon.

314 Arrows indicate the direction of extension ( $\sigma_3$ ) obtained from secondary fault inversion. Examples of  
 315 fault-slip data and stress calculation are shown. In highly dipping strata, age of stress states relative to  
 316 folding was obtained using the Anderson model in which two one of the principal stresses is vertical.  
 317 Stress sates predate folding if this criterion is fulfilled when performing the stress inversion on faults with


318 their backtilted attitude. The rose diagram of the  $\sigma_3$  direction is shown. CA: Chouf area. Same legend as in  
319 fig. 3.


320

321 Fig. 6. Main extensional tectonic event in the polyphase Levant basin.

Accepted Manuscript













Figure 4











Figure 6

	Time		Faulting	Stresses	Regional event	Remarks
<b>Cretaceous</b>	Late				Second rifting event	WNW-ESE to WSW-ENE faults
	Early	Cenomanian Albian Aptian Barremian Hauterivian Valanginian Berriasian				
<b>Jurassic</b>	Malm				Development of the Levant Basin	NE-SW faults
	Dogger					
	Lias					
<b>Triassic</b>	Late					
	Middle					
	Early					
Normal faulting event: 			Direction of extension 