



HAL
open science

Rolling Manifolds: Intrinsic Formulation and Controllability

Yacine Chitour, Petri Kokkonen

► **To cite this version:**

Yacine Chitour, Petri Kokkonen. Rolling Manifolds: Intrinsic Formulation and Controllability. 2010. hal-00535711v1

HAL Id: hal-00535711

<https://hal.science/hal-00535711v1>

Preprint submitted on 12 Nov 2010 (v1), last revised 13 May 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rolling Manifolds: Intrinsic Formulation and Controllability*

Yacine Chitour[†] Petri Kokkonen[‡]

November 12, 2010

Abstract

In this paper, we consider two cases of rolling of one smooth connected complete Riemannian manifold (M, g) onto another one (\hat{M}, \hat{g}) of equal dimension $n \geq 2$. The rolling problem (NS) corresponds to the situation where there is no relative spin (or twist) of one manifold with respect to the other one. As for the rolling problem (R) , there is no relative spin and also no relative slip. Since the manifolds are not assumed to be embedded into an Euclidean space, we provide an intrinsic description of the two constraints "without spinning" and "without slipping" in terms of the Levi-Civita connections ∇^g and $\nabla^{\hat{g}}$. For that purpose, we recast the two rolling problems within the framework of geometric control and associate to each of them a distribution and a control system. We then investigate the relationships between the two control systems and we address for both of them the issue of complete controllability. For the rolling (NS) , the reachable set (from any point) can be described exactly in terms of the holonomy groups of (M, g) and (\hat{M}, \hat{g}) respectively, and thus we achieve a complete understanding of the controllability properties of the corresponding control system. As for the rolling (R) , the problem turns out to be more delicate. We first provide general properties for the reachable set and determine the associated Lie bracket structure. Regarding the controllability issue, we only have partial results, for instance dealing with the situation where one of the manifold is a space form. Finally, we extend the two types of rolling to the case where the manifolds have different dimensions.

*The work of the first author is supported by the ANR project GCM, program "Blanche", (project number NT09_504490) and the DIGITEO-Région Ile-de-France project CONGEO. The work of the second author is supported by Finnish Academy of Science and Letters and Saastamoinen Foundation.

[†]yacine.chitour@lss.supelec.fr, L2S, Université Paris-Sud XI, CNRS and Supélec, Gif-sur-Yvette, 91192, France.

[‡]petri.kokkonen@lss.supelec.fr, L2S, Université Paris-Sud XI, CNRS and Supélec, Gif-sur-Yvette, 91192, France and University of Eastern Finland, Department of Physics and Mathematics, 70211, Kuopio, Finland.

Contents

1	Introduction	3
2	Notations	7
3	Characterization of the Rolling Problems as Control Problems	12
3.1	Definition of the State Space	12
3.2	The Bundle Structure of Q	13
3.3	The State Space as a Quotient	16
3.4	The Control Problem	18
4	Study of the Rolling problem (NS)	20
4.1	The No-Spinning Distribution \mathcal{D}_{NS}	20
4.2	The Control System Associated to the No-Spinning Problem	24
4.3	The Analysis of the Control System $(\Sigma)_{NS}$	24
4.3.1	Description of the Orbits of $(\Sigma)_{NS}$	25
4.3.2	Consequences for Controllability	29
4.4	Computations of Lie Brackets	33
5	Study of the Rolling problem (R)	38
5.1	The Rolling Distribution \mathcal{D}_R	38
5.2	The Bundle Structure of a \mathcal{D}_R -orbit	42
5.3	Computations of some Lie Brackets	46
5.4	Controllability Properties of \mathcal{D}_R	51
5.4.1	First Results	51
5.4.2	A "Rolling Along Loops" Characterization of Isometry	54
5.4.3	The Ambrose's Theorem Revisited	62
6	Rolling Against a Space Form	67
6.1	Rolling Against an Euclidean Space	67
6.2	Rolling Against a Non-Flat Space Form	73
6.2.1	Standard Results on Space Forms	73
6.2.2	Orbit Structure	75
6.2.3	The Rolling Connection	77
6.3	Controllability Results for Rolling Against a Non-Flat Space Form	82
7	Rolling of Spaces of Different Dimensions	98
7.1	Definitions of the State Space and the Rolling Distributions	98
7.2	Controllability Results	101
7.2.1	Rolling Problem (NS)	101
7.2.2	Rolling Problem (R)	101
A	Fiber and Local Coordinates Point of View	107
B	Sasaki-metric on $T^*M \otimes T\hat{M}$ and Q	109
C	The Rolling Problem Embedded in \mathbb{R}^N	114

1 Introduction

In this paper, we study the rolling of a manifold over another one. Unless otherwise precised, manifolds are smooth, connected, oriented, of finite dimension $n \geq 2$, endowed with a complete Riemannian metric. The rolling is assumed to be either without spinning (NS) or without spinning nor slipping (R). When both manifolds are isometrically embedded into an Euclidean space, the rolling problem is classical in differential geometry (see [27]), through the notions of "development of a manifold" and "rolling maps". To get an intuitive grasp of the problem, consider the rolling problem (R) of a 2D convex surface S_1 onto another one S_2 in the euclidean space \mathbb{R}^3 , for instance the plate-ball problem, i.e., a sphere rolling onto a plane in \mathbb{R}^3 , (cf. [12] and [20]). The two surfaces are in contact i.e., they have a common tangent plane at the contact point and, equivalently, their exterior normal vectors are opposite at the contact point. If $\gamma : [0, T] \rightarrow S_1$ is a C^1 regular curve on S_1 , one says that S_1 rolls onto S_2 along γ without spinning nor slipping if the following holds. The curve traced on S_1 by the contact point is equal to γ and let $\hat{\gamma} : [0, T] \rightarrow S_2$ be the curve traced on S_2 by the contact point. At time $t \in [0, T]$, the relative orientation of S_2 with respect to S_1 is measured by the angle $\theta(t)$ between $\dot{\gamma}(t)$ and $\dot{\hat{\gamma}}(t)$ in the common tangent plane at the contact point. The state space Q of the rolling problem is therefore five dimensional since a point in Q is defined by fixing a point on S_1 , a point on S_2 and an angle in S^1 , the unit circle. Requiring that there is no instantaneous relative speed nor twist of S_1 with respect to S_2 defines uniquely the curves $\hat{\gamma}$ and θ once a point on S_2 and an angle are chosen at time $t = 0$. For the rolling (NS), one must choose two C^1 regular curves γ and $\hat{\gamma}$ on S_1 and S_2 respectively, and an angle θ_0 so that one says that S_1 rolls onto S_2 along γ and $\hat{\gamma}$ without spinning if (a) the curves traced on S_1 and S_2 by the contact point are equal to γ and $\hat{\gamma}$ respectively; (b) the no-spin constraint and the initial condition θ_0 determine a unique curve θ which measures the relative orientation of S_2 with respect to S_1 along the rolling. The most basic issue linked to the rolling problems is that of *controllability* i.e., to determine, for two given points q_{init} and q_{final} in the state space Q , if there exists a curve γ so that the rolling of S_1 onto S_2 along γ steers the system from q_{init} to q_{final} . If this is the case for every points q_{init} and q_{final} in Q , then the rolling of S_1 onto S_2 is said to be *completely controllable*.

If the manifolds rolling on each other are two-dimensional, then the controllability issue is well-understood thanks to the work of [2], [5] and [16] especially. For instance, in the simply connected case, the rolling (R) is completely controllable if and only if the manifolds are not isometric. In the case where the manifolds are isometric, [2] also provides a description of the reachable sets in terms of isometries between the manifolds.

In particular, these reachable sets are immersed submanifolds of Q of dimension either 2 or 5. In case the manifolds rolling on each other are isometric convex surfaces, [16] provides a beautiful description of a two dimensional reachable set: consider the initial configuration given by two (isometric) surfaces in contact so that one is the image of the other one by the symmetry with respect to the (common) tangent plane at the contact point. Then, this symmetry property (chirality) is preserved along the rolling (R). Note that if the (isometric) convex surfaces are not spheres nor planes, the reachable set starting at a contact point where the Gaussian curvatures are distinct, is open (and thus of dimension 5).

From a robotics point of view, once the controllability is well-understood, the next issue to address is that of *motion planning*, i.e., defining an effective procedure that produces, for every pair of points $(q_{\text{init}}, q_{\text{final}})$ in the state space Q , a curve $\gamma_{q_{\text{init}}, q_{\text{final}}}$ so that the rolling of S_1 onto S_2 along $\gamma_{q_{\text{init}}, q_{\text{final}}}$ steers the system from q_{init} to q_{final} . In [7], an algorithm based on the continuation method was proposed to tackle the rolling problem (R) of a strictly convex compact surface onto an Euclidean plane. That algorithm was also proved in [7] to be convergent and it was numerically implemented in [1] (see also [17] for another algorithm).

To the best of our knowledge, only the rolling (R) was considered in the literature, eventhough it is the more delicate, as explained below. The rolling problem (R) is traditionally presented by isometrically embedding the rolling manifolds M and \hat{M} in an Euclidean space (cf. [27], [10]) since it is the most intuitive way to provide a rigorous meaning to the notions of relative spin (or twist) and relative slip of one manifold with respect to the other one. However, the rolling model will depend in general on the embedding. For instance, rolling two 2D spheres of different radii on each other can be isometrically embedded in (at least) two ways in \mathbb{R}^3 : the smaller sphere can roll onto the bigger one either inside of it or outside. Then one should be able to define rolling without having to resort to any isometric embedding into an Euclidean space. To be satisfactory, that *intrinsic* formulation of the rolling should also allow one to address at least the controllability issue.

The first step towards an intrinsic formulation of the rolling starts with an intrinsic definition of the state space Q . It is of dimension $2n + n(n - 1)/2$ since it is locally diffeomorphic to neighborhoods of $M \times \hat{M} \times \text{SO}(n)$. There are two main approaches for an intrinsic formulation of the rolling problem (R) , first considered by [2] and [5] respectively. Note that the two references only deal with the two dimensional case but it is not hard to generalize them to higher dimensions. In [2], the state space Q is given by

$$Q = \{A : T|_x M \rightarrow T|_{\hat{x}} \hat{M} \mid A \text{ o-isometry, } x \in M, \hat{x} \in \hat{M}\},$$

where "o-isometry" means positively oriented isometry, (see Definition 3.1 below) while in [5], one has equivalently

$$Q = (F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})) / \Delta,$$

where $F_{\text{OON}}(M)$, $F_{\text{OON}}(\hat{M})$ be the oriented orthonormal frame bundles of (M, g) , (\hat{M}, \hat{g}) respectively, and Δ is the diagonal right $\text{SO}(n)$ -action (see Proposition 3.9 below).

The next step towards an intrinsic formulation consists of using either the parallel transports with respect to ∇^g and $\nabla^{\hat{g}}$ (Agrachev-Sachkov's approach) or alternatively, orthonormal moving frames and the structure equations (Bryant-Hsu's approach) to translate the constraints of no-spinning and no-slipping and derive the admissible curves, i.e., the curves of Q describing the rolling (R) , cf. Eq. (14). Finally, one defines either a distribution or a codistribution depending which approach is chosen. In the present paper, we adopt the Agrachev-Sachkov's approach and we construct an n -dimensional distribution \mathcal{D}_R on Q so that the locally absolutely continuous curves tangent to \mathcal{D}_R are exactly the admissible curves for the rolling problem, cf. Definition 5.2. The construction of \mathcal{D}_R comes along with the construction of (local) basis of vector fields, which allow one to compute the Lie algebraic structure associated to \mathcal{D}_R .

One should mention the recent work [8] dealing with an intrinsic formulation of the rolling problem (R) (see Definition 4 page 18 in the reference therein). However, that definition does not allow one to parameterize the admissible curves using a control system and a fortiori to construct a distribution (or a codistribution) associated to the rolling. Therefore, the computations in that paper related to controllability issues are all performed by embedding the rolling into an Euclidean space.

We now describe precisely the results of the present paper. In Section 2, are gathered the notations used throughout the paper. After that, the control systems associated to the rolling problems (NS) and (R) are introduced in Section 3. Besides the state space Q , one must define the set of admissible controls. For (NS), it is the set of locally absolutely continuous (l.a.c.) curves on $M \times \hat{M}$ while, for (R), it is the set of locally absolutely continuous (l.a.c.) curves on M only. As control systems, we obtain two driftless control systems affine in the control $(\Sigma)_{NS}$ and $(\Sigma)_R$ for (NS) and (R) respectively. We also provide, in Appendix A, expressions in local coordinates for these control systems.

The study of the rolling problem (NS) is the objet of Section 4. We first construct the distribution \mathcal{D}_{NS} of rank $2n$ in Q so that its tangent curves coincide with the admissible curves of $(\Sigma)_{NS}$ and we provide (local) basis of vector fields for \mathcal{D}_{NS} . The controllability issue is completely addressed since we can describe exactly the reachable sets of $(\Sigma)_{NS}$ in terms of H^{∇^g} and $\hat{H}^{\nabla^{\hat{g}}}$, the holonomy groups of ∇^g and $\nabla^{\hat{g}}$ respectively. We thus derive a necessary and sufficient condition for complete controllability of (NS) in terms of the Lie algebras of H^{∇^g} and $\hat{H}^{\nabla^{\hat{g}}}$. For instance, if both manifolds M and \hat{M} are simply connected and non symmetric, then the rolling problem (NS) is completely controllable in dimension $n \neq 8$ if and only if H^{∇^g} or $\hat{H}^{\nabla^{\hat{g}}}$ is equal to $SO(n)$. We conclude that section by computing Lie brackets of vector fields tangent to \mathcal{D}_{NS} .

In Section 5, we start the study of the rolling problem (R). As done for (NS), we construct the rolling distribution \mathcal{D}_R as a sub-distribution of rank n of \mathcal{D}_{NS} so that its tangent curves coincide with the admissible curves of $(\Sigma)_R$ and we provide (local) basis of vector fields for \mathcal{D}_R . We show that the rolling (R) of M over \hat{M} is symmetric to that of \hat{M} over M i.e., the reachable sets are diffeomorphic. Already from these computations, one can see why we considered the rolling problem (NS): from a technical point of view, it is much easier to perform Lie brackets computations first with vector fields spanning \mathcal{D}_{NS} and then specify these computations to vector fields spanning \mathcal{D}_R . Moreover, the complete controllability of (NS) being a necessary condition for the complete controllability of (R), one can derive at once that, for simply connected and non symmetric rolling manifolds, if the rolling problem (R) is completely controllable in dimension $n \neq 8$ then H^{∇^g} or $\hat{H}^{\nabla^{\hat{g}}}$ must be equal to $SO(n)$.

The controllability issue for (R) turns out to be much more delicate than that for (NS). One reason is that, in general, there is no "natural" principal bundle structure on $\pi_{Q,M} : Q \rightarrow M$ which leaves invariant the rolling distribution \mathcal{D}_R . Indeed, if it were the case, then all the reachable sets would be diffeomorphic and this is not true in general (cf. the description of reachable sets of the rolling problem (R) for two-dimensional isometric manifolds). Despite this fact, we prove that the reachable sets are smooth bundles over M (cf. Proposition 5.11).

We also have an equivariance property of the reachable sets of \mathcal{D}_R with respect to the (global) isometries of the manifolds M and \hat{M} , as well as an interesting result

linking the rolling problem (R) for a pair of manifolds M and \hat{M} and the rolling problem (R) associated to Riemannian coverings of M and \hat{M} respectively. As a consequence, we have that the complete controllability for the rolling problem (R) associated to a pair of manifolds M and \hat{M} is equivalent to that of the rolling problem (R) associated to their universal Riemannian coverings. This implies that, as far as complete controllability is concerned, one can assume without loss of generality that M and \hat{M} are simply connected. We then compute the first order Lie brackets of the vector fields generating \mathcal{D}_R and find that they are (essentially) equal to the vector fields given by the vertical lifts of

$$\text{Rol}(X, Y)(A) := AR(X, Y) - \hat{R}(AX, AY)A, \quad (1)$$

where X, Y are smooth vector fields of M , $q = (x, \hat{x}; A) \in Q$ and $R(\cdot, \cdot)$, $\hat{R}(\cdot, \cdot)$ are the curvature tensors of g and \hat{g} respectively. We call the vertical vector field defined in Eq. (1) the *Rolling Curvature*, cf Definition 5.18 below. Higher order Lie brackets can now be expressed as linear combinations of covariant derivatives of the Rolling Curvature for the vertical part and evaluations on \hat{M} of the images of the Rolling Curvature and its covariant derivatives.

In dimension two, the Rolling Curvature is (essentially) equal to $K^M(x) - K^{\hat{M}}(\hat{x})$, where $K^M(\cdot)$, $K^{\hat{M}}(\cdot)$ are the Gaussian curvatures of M and \hat{M} respectively. At some point $q \in Q$ where $K^M(x) - K^{\hat{M}}(\hat{x}) \neq 0$, one immediately deduces that the dimension of the evaluation at q of the Lie algebra of the vector fields spanning \mathcal{D}_R is equal to five, (the dimension of Q) and thus the reachable set from q is open in Q . From that fact, one has the following alternative: (a) there exists $q_0 \in Q$ so that $K^M - K^{\hat{M}} \equiv 0$ over the reachable set from q_0 , yielding easily that M and \hat{M} have the same Riemannian covering space (cf. [2] and [5]); (b) all the reachable sets are open and then the rolling problem (R) is completely controllable. In dimension $n \geq 3$, the Rolling Curvature cannot be reduced to a scalar and it seems difficult to compute in general the rank of the evaluations of the Lie algebra of the vector fields spanning \mathcal{D}_R .

We however propose several characterizations of isometry between two Riemannian manifolds based on the rolling perspective. The first one refers to a "rolling against loops" property which assumes that there is a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that for every loop γ on M based at x_0 , the corresponding rolling curve $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)$ on \hat{M} starting from q_0 is a loop based \hat{x}_0 . Then we prove that, under the previous condition (M, g) and (\hat{M}, \hat{g}) have the same universal Riemannian covering, cf. Theorem 5.32.

The second characterization consists of revisiting the classical Ambrose theorem (see [25] Theorem III.5.1) and showing how the standard argument actually gets simplified when recast in the rolling context. We also prove a version of the Cartan-Ambrose-Hicks theorem, Proposition 7.10, by using the rolling model. In this version, we also include a condition for certain submersions to exist, not only (local) geodesic embeddings. Our proofs are in parallel to those presented in [4], [23].

In Section 6, we present controllability results when one of the manifolds, let say (\hat{M}, \hat{g}) , is a space form i.e., a simply connected complete Riemannian manifold of constant curvature. Our results are actually preliminary and we hope to complete them in a future version of the present draft. Let us summarize them. The main feature of this particular case is that there is a principal bundle structure on the

bundle $\pi_{Q,M} : Q \rightarrow M$, which is compatible with the rolling distribution \mathcal{D}_R . In the case \hat{M} has non-zero constant curvature, this allows us to reduce the problem to a study of a vector bundle connection ∇^{Rol} of the vector bundle $\pi_{TM \oplus \mathbb{R}} : TM \oplus \mathbb{R} \rightarrow M$ and its holonomy group, which is a subgroup of $\text{SO}(n+1)$ or $\text{SO}(n,1)$ depending whether the curvature of \hat{M} is positive or negative, respectively. If \hat{M} has zero curvature i.e., it is the Euclidean plane, the problem reduces to the study of an affine connection and its holonomy group, a subgroup of $\text{SE}(n)$, in the sense of [13]. In all the cases, the fibers over M of the \mathcal{D}_R -orbits are all diffeomorphic to the holonomy group of the connection in question.

In the zero curvature case, we prove that the rolling (R) is completely controllable if and only if the (Riemannian) holonomy group of ∇^g is equal to $\text{SO}(n)$. This result is actually similar to Theorem IV.7.1, p. 193 and Theorem IV.7.2, p. 194 in [13]. In the non-zero curvature case, we only study the rolling onto an n -dimensional sphere. We prove that if the holonomy group of the rolling connection ∇^{Rol} is reducible, then the sphere endowed with the metric induced by the Euclidean metric of \mathbb{R}^{n+1} must be a Riemannian covering space of (M, g) .

In Section 7, we show how to extend the formalism developed previously to the case where the rolling manifolds have different dimensions. In that case, we show that the rolling of M over \hat{M} is not anymore symmetric with that of \hat{M} over M , which is reasonable. We also provide basic controllability results.

We finally gather in a series of appendices several results either used in the text or directly related to it. In particular, we show how the \mathcal{D}_{NS} relates to the Sasaki-metric on the tensor space $T^*(M) \otimes T(\hat{M})$. In the final appendix, we provide, for the sake of completeness, the classical formulation of the rolling problem (R) as embedded in an Euclidean space.

Acknowledgements. The authors want to thank P. Pansu and E. Falbel for helpful comments as well as L. Rifford for having organized the conference "New Trends in Sub-Riemannian Geometry" in Nice and where this work was first presented in April 2010.

2 Notations

For any sets A, B, C and $U \subset A \times B$ and any map $F : U \rightarrow C$, we write U_a and U^b for the sets defined by $\{b \in B \mid (a, b) \in U\}$ and $\{a \in A \mid (a, b) \in U\}$ respectively. Similarly, let $F_a : U_a \rightarrow C$ and $F^b : U^b \rightarrow C$ be defined by $F_a(b) := F(a, b)$ and $F^b(a) := F(a, b)$ respectively. For any sets V_1, \dots, V_n the map $\text{pr}_i : V_1 \times \dots \times V_n \rightarrow V_i$ denotes the projection onto the i -th factor.

For a real matrix A , we use A_j^i to denote the real number on the i -th row and j -th column and the matrix A can then be denoted by $[A_j^i]$. If, for example, one has $A_j^i = a_{ij}$ for all i, j , then one uses the notation $A_j^i = (a_{ij})_j^i$ and thus $A = [(a_{ij})_j^i]$. The matrix multiplication of $A = [A_j^i]$ and $B = [B_j^i]$ is therefore given by $AB = [(\sum_k A_k^i B_j^k)_j^i]$.

Suppose V, W are finite dimensional \mathbb{R} -linear spaces, $L : V \rightarrow W$ is an \mathbb{R} -linear map and $F = (v_i)_{i=1}^{\dim V}$, $G = (w_i)_{i=1}^{\dim W}$ are bases of V, W respectively. The $\dim W \times \dim V$ -real matrix corresponding to L w.r.t the bases F and G is denoted by $\mathcal{M}_{F,G}(L)$. In other words, $L(v_i) = \sum_j \mathcal{M}_{F,G}(L)_i^j w_j$ (corresponding to the right multiplication by a matrix of a row vector). Notice that, if $K : W \rightarrow U$ is yet another

\mathbb{R} -linear map to a finite dimensional linear space U with basis $H = (u_i)_{i=1}^{\dim U}$, then

$$\mathcal{M}_{F,H}(K \circ L) = \mathcal{M}_{G,H}(K)\mathcal{M}_{F,G}(L).$$

If (V, g) , (W, h) are inner product spaces with inner products g and h , one defines $L^{T_{g,h}} : W \rightarrow V$ as the transpose (adjoint) of A w.r.t g and h i.e., $g(L^{T_{g,h}}w, v) = h(w, Lv)$. With bases F and G as above, one has $\mathcal{M}_{F,G}(L)^T = \mathcal{M}_{G,F}(L^{T_{g,h}})$, where T on the left is the usual transpose of a real matrix i.e., the transpose w.r.t standard Euclidean inner products in \mathbb{R}^N , $N \in \mathbb{N}$.

In this paper, by a smooth manifold, one means a smooth finite-dimensional, second countable, Hausdorff manifold (see e.g. [15]). A smooth manifold $N \subset M$ is an immersed submanifold of M if the inclusion map $i : N \rightarrow M$ is a smooth immersion. We call N embedded submanifold if the topology on N induced by the inclusion i coincides with the manifold topology of N . By a smooth submanifold of M , we always mean a smooth embedded submanifold.

A smooth bundle (over M) is a smooth map $\pi : E \rightarrow M$ between two smooth manifolds E and M together with a prescribed smooth manifold F (unique up to diffeomorphism), called the typical fiber of π , such that, for each $x \in M$, there is a neighbourhood U of x in M and a smooth diffeomorphism $\tau : \pi^{-1}(U) \rightarrow U \times F$ with the property that $\text{pr}_1 \circ \tau = \pi|_{\pi^{-1}(U)}$. Such maps τ are called (smooth) local trivializations of π .

For any smooth map $\pi : E \rightarrow M$ between smooth manifolds E and M , the set $\pi^{-1}(\{x\}) =: \pi^{-1}(x)$ is called the π -fiber over x and it is sometimes denoted by $E|_x$, when π is clear from the context. A smooth section of a smooth map $\pi : E \rightarrow M$ is a smooth map $s : M \rightarrow E$ such that $\pi \circ s = \text{id}_M$. The set of smooth sections of π is denoted by $\Gamma(\pi)$. Local sections of π are sections defined only on open (possibly proper) subsets of M . The value $s(x)$ of a section s at x is usually denoted by $s|_x$.

A smooth manifold M is oriented if there exists a smooth (or continuous) section, defined on all of M , of the bundle of n -forms $\pi_{\wedge^n(M)} : \wedge^n(M) \rightarrow M$ where $n = \dim M$. Otherwise mentioned, the smooth manifolds considered in this paper are connected and oriented.

A smooth vector bundle is a smooth bundle where the typical fiber F is a finite dimensional \mathbb{R} -linear space together with a collection of local trivializations so that there is a well defined vector space structure on each π -fiber (see [15] for the precise definition). Some important vector bundles for us over a manifold M are the tangent bundle $\pi_{T(M)} : T(M) \rightarrow M$ and different (k, m) -tensor bundles $\pi_{T_m^k(N)} : T_m^k(N) \rightarrow M$. We will many times write $TM := T(M)$ etc. to ease the notation.

If G is a smooth Lie-group, a smooth bundle $\pi : E \rightarrow M$ is called a right principal G -bundle if there exists a smooth right action $\mu : E \times G \rightarrow E$ of G on E (i.e., $\mu(\mu(y, g), h) = \mu(y, gh)$ where the product gh is computed in G) such that $\pi \circ \mu = \pi \circ \text{pr}_1$ and μ is free (i.e., $\mu(y, g) = y$ for a $y \in E$ implies $g = e$ the identity of G) and transitive on π -fibers (i.e., for every $y, z \in \pi^{-1}(x)$ there is a $g \in G$ such that $\mu(y, g) = z$). It follows from the definition that this bundle has G as the typical fiber. Similarly, using a left action one defines a left principal G -bundle. For short, by a principal bundle we mean a left or right principal bundle (the side of the action being clear from context). There is no difference between left and right principal bundles since a right principal bundle π_E with action $\mu : E \times G \rightarrow E$ can be identified with a left principal bundle π_E with action $\lambda : G \times E \rightarrow E$; $\lambda(g, y) = \mu(y, g^{-1})$ and vice versa.

For a smooth map $\pi : E \rightarrow M$ and $y \in E$, let $V|_y(\pi)$ be the set of all $Y \in T|_y E$ such that $\pi_*(Y) = 0$. If π is a smooth bundle, the collection of spaces $V|_y(\pi)$, $y \in E$, defines a smooth submanifold $V(\pi)$ of $T(E)$ and the restriction $\pi_{T(E)} : T(E) \rightarrow E$ to $V(\pi)$ is denoted by $\pi_{V(\pi)}$. In this case $\pi_{V(\pi)}$ is a vector subbundle of $\pi_{T(E)}$ over E .

For a smooth manifold M , one uses $\text{VF}(M)$ to denote the set of smooth vector fields on M i.e., the set of smooth sections of the tangent bundle $\pi_{T(M)} : T(M) \rightarrow M$. The flow of a vector field $Y \in \text{VF}(M)$ is a smooth onto map $\Phi_Y : D \rightarrow M$ defined on an open subset D of $\mathbb{R} \times M$ containing $\{0\} \times M$ such that $\frac{\partial}{\partial t} \Phi_Y(t, y) = Y|_{\Phi_Y(t, y)}$ for $(t, y) \in D$ and $\Phi_Y(0, y) = y$ for all $y \in M$. As a default, we will take D to be the maximal flow domain of Y .

A subset $\mathcal{D} \subset T(M)$ of the tangent bundle of M is called a smooth distribution on M if $\pi_{T(M)}|_{\mathcal{D}}$ is a smooth vector subbundle of $\pi_{T(M)}$ over M . For $x \in M$, the fiber $\pi_{T(M)}|_{\mathcal{D}}^{-1}(\{x\})$ is denoted by $\mathcal{D}|_x$ and the common dimension of the spaces $\mathcal{D}|_x$, $x \in M$, is called the rank of the distribution \mathcal{D} .

For any distribution \mathcal{D} on a manifold M , we use $\text{VF}_{\mathcal{D}}$ to denote the set of vector fields $X \in \text{VF}(M)$ tangent to \mathcal{D} (i.e., $X|_x \in \mathcal{D}|_x$ for all $x \in M$) and we define inductively for $k \geq 2$

$$\text{VF}_{\mathcal{D}}^k = \text{VF}_{\mathcal{D}}^{k-1} + [\text{VF}_{\mathcal{D}}, \text{VF}_{\mathcal{D}}^{k-1}],$$

where $\text{VF}_{\mathcal{D}}^1 := \text{VF}_{\mathcal{D}}$. The Lie algebra generated by $\text{VF}_{\mathcal{D}}$ is denoted by $\text{Lie}(\mathcal{D})$ and it equals $\bigcup_k \text{VF}_{\mathcal{D}}^k$.

For any maps $\gamma : [a, b] \rightarrow X$, $\omega : [c, d] \rightarrow X$ into a set X such that $\gamma(b) = \omega(c)$ we define

$$\omega \sqcup \gamma : [a, b + d - c] \rightarrow X; \quad (\omega \sqcup \gamma)(t) = \begin{cases} \gamma(t), & t \in [a, b] \\ \omega(t - b + c), & t \in [b, b + d - c]. \end{cases}$$

A map $\gamma : [a, b] \rightarrow X$ is a loop in X based at $x_0 \in X$ if $\gamma(a) = \gamma(b) = x_0$. In the space of loops $[0, 1] \rightarrow X$ based at some given point x_0 , one defines a group operation "·", concatenation, by

$$\omega \cdot \gamma := (t \mapsto \gamma(\frac{t}{2})) \sqcup (t \mapsto \omega(\frac{t}{2})).$$

This operation gives a group structure on the set of loops of X based at a given point x_0 . If N is a smooth manifold and $y \in N$, we use $\Omega_y(N)$ to denote the set of all piecewise C^1 -loops $[0, 1] \rightarrow N$ of N based at y . In particular, $(\Omega_y(N), \cdot)$ is a group.

A continuous map $c : I \rightarrow M$ from a real compact interval I into a smooth manifold M is called *absolutely continuous*, or *a.c.* for short if, for every $t_0 \in I$, there is a smooth coordinate chart (ϕ, U) of M such that $c(t_0) \in U$ and $\phi \circ c|_{c^{-1}(U)}$ is absolutely continuous.

Given a smooth distribution \mathcal{D} on a smooth manifold M , we call an absolutely continuous curve $c : I \rightarrow M$, $I \subset \mathbb{R}$, \mathcal{D} -*admissible* if c is tangent to \mathcal{D} almost everywhere (a.e.) i.e., if for almost all $t \in I$ it holds that $\dot{c}(t) \in \mathcal{D}|_{c(t)}$. For $x_0 \in M$, the endpoints of all the \mathcal{D} -admissible curves of M starting at x_0 form the set called \mathcal{D} -*orbit through* x_0 and denoted $\mathcal{O}_{\mathcal{D}}(x_0)$. More precisely,

$$\mathcal{O}_{\mathcal{D}}(x_0) = \{c(1) \mid c : [0, 1] \rightarrow M, \mathcal{D}\text{-admissible}, c(0) = x_0\}. \quad (2)$$

By the Orbit Theorem (see [3]), it follows that $\mathcal{O}_{\mathcal{D}}(x_0)$ is an immersed smooth submanifold of M containing x_0 . It is also known that one may restrict to piecewise smooth curves in the description of the orbit i.e.,

$$\mathcal{O}_{\mathcal{D}}(x_0) = \{c(1) \mid c : [0, 1] \rightarrow M \text{ piecewise smooth and } \mathcal{D}\text{-admissible, } c(0) = x_0\}.$$

We call a smooth distribution \mathcal{D}' on M a subdistribution of \mathcal{D} if $\mathcal{D}' \subset \mathcal{D}$. An immediate consequence of the definition of the orbit shows that in this case

$$\mathcal{O}_{\mathcal{D}'}(x_0) \subset \mathcal{O}_{\mathcal{D}}(x_0), \quad \forall x_0 \in M.$$

If $\pi : E \rightarrow M$, $\eta : F \rightarrow M$ are two smooth maps (e.g. bundles), let $C^\infty(\pi, \eta)$ be the set of all bundle maps $\pi \rightarrow \eta$ i.e., smooth maps $g : E \rightarrow F$ such that $\eta \circ g = \pi$. For a manifold M , let $\pi_{M_{\mathbb{R}}} : M \times \mathbb{R} \rightarrow M$ be the projection onto the first factor i.e., $(x, t) \mapsto x$ (i.e., $\pi_{M_{\mathbb{R}}} = \text{pr}_1$). Recall that there is a canonical bijection between the set $C^\infty(M)$ of smooth functions on M and the set $C^\infty(\text{id}_M, \pi_{M_{\mathbb{R}}})$ given by $f \mapsto f_{\mathbb{R}} := (x \mapsto (x, f(x)))$.

If $\pi : E \rightarrow M$, $\eta : F \rightarrow M$ are any smooth vector bundles over a smooth manifold M , $f \in C^\infty(\pi, \eta)$ and $u, w \in \pi^{-1}(x)$, one defines the vertical derivative f at u in the direction w by

$$\nu(w)|_u(f) := (D_\nu f)(u)(w) := \left. \frac{d}{dt} \right|_0 f(u + tw). \quad (3)$$

Here $w \mapsto (D_\nu f)(u)(w) = \nu(w)|_u(f)$ is an \mathbb{R} -linear map between fibers $\pi^{-1}(x) \rightarrow \eta^{-1}(x)$.

In a similar way, in the case of $f \in C^\infty(E)$ and $u, w \in \pi^{-1}(x)$, one defines the π -vertical derivative $\nu(w)|_u(f) := D_\nu f(u)(w) := \left. \frac{d}{dt} \right|_0 f(u + tw)$ at u in the direction w . This definition agrees with the above one modulo the canonical bijection $C^\infty(E) \cong C^\infty(\text{id}_E, \pi_{E_{\mathbb{R}}})$. This latter definition means that $\nu(w)|_u$ can be viewed as an element of $V|_u(\pi)$ and the mapping $w \mapsto \nu(w)|_u$ gives a (natural) \mathbb{R} -linear isomorphism between $\pi^{-1}(x)$ and $V|_u(\pi)$ where $\pi(u) = x$. If $\tilde{u} \in \Gamma(\pi)$ is a smooth π -section, let $\nu(\tilde{w})$ be the π -vertical vector field on E defined by $\nu(\tilde{w})|_u(f) = \nu(\tilde{w}|_x)|_u(f)$, where $\pi(u) = x$ and $f \in C^\infty(E)$. The same remark holds also locally.

In the case of smooth manifolds M and \hat{M} , $x \in M$, $\hat{x} \in \hat{M}$, we will use freely and without mention the natural inclusions (\subset) and isomorphisms (\cong): $T|_x M, T|_{\hat{x}} \hat{M} \subset T|_{(x, \hat{x})}(M \times \hat{M}) \cong T|_x M \oplus T|_{\hat{x}} \hat{M}$, $T^*|_x M, T^*|_{\hat{x}} \hat{M} \subset T^*|_{(x, \hat{x})}(M \times \hat{M}) \cong T^*|_x M \oplus T^*|_{\hat{x}} \hat{M}$. An element of $T|_{(x, \hat{x})}(M \times \hat{M}) \cong T|_x(M) \oplus T|_{\hat{x}}(\hat{M})$ with respect to the direct sum splitting is denoted usually by (X, \hat{X}) , where $X \in T|_x M$, $\hat{X} \in T|_{\hat{x}} \hat{M}$. Sometimes it is even more convenient to write $X + \hat{X} := (X, \hat{X})$ when we make the identifications $(X, 0) = X$, $(0, \hat{X}) = \hat{X}$.

If (M, g) , (\hat{M}, \hat{g}) are smooth Riemannian manifolds, the space $\overline{M} = M \times \hat{M}$ is a Riemannian manifold, called the Riemannian product manifold of (M, g) , (\hat{M}, \hat{g}) , when endowed with the product metric $\overline{g} := g \oplus \hat{g}$

Let $\nabla, \hat{\nabla}, \overline{\nabla}$ (resp. R, \hat{R}, \overline{R}) denote the Levi-Civita connections (resp. the Riemannian curvature tensors) of (M, g) , (\hat{M}, \hat{g}) , $(\overline{M} = M \times \hat{M}, \overline{g} = g \oplus \hat{g})$ respectively. From Koszul's formula (cf. [15]), one has

$$\overline{\nabla}_{(X, \hat{X})}(Y, \hat{Y}) = (\nabla_X Y, \hat{\nabla}_{\hat{X}} \hat{Y}), \quad (4)$$

when $X, Y \in \text{VF}(M)$, $\hat{X}, \hat{Y} \in \text{VF}(\hat{M})$ and hence from the definition of the Riemannian curvature tensor

$$\overline{R}((X, \hat{X}), (Y, \hat{Y}))(Z, \hat{Z}) = (R(X, Y)Z, \hat{R}(\hat{X}, \hat{Y})\hat{Z}), \quad (5)$$

where $X, Y, Z \in T|_x M$, $\hat{X}, \hat{Y}, \hat{Z} \in T|_{\hat{x}} \hat{M}$.

For any (k, m) -tensor field T on M we define ∇T to be the $(k, m + 1)$ -tensor field such that (see [25], p. 30)

$$(\nabla T)(X_1, \dots, X_m, X) = (\nabla_X T)(X_1, \dots, X_m), \quad (6)$$

$X_1, \dots, X_m, X \in T|_x M$.

Let $x : I \rightarrow M$ and $X : I \rightarrow TM$ be a smooth curve and a smooth vector field along x respectively i.e., a smooth map such that $X(t) \in T|_{x(t)} M$ for all $t \in I$. A *local extension* of X around t_0 is a vector field $\tilde{X} \in \text{VF}(M)$ such that there is an open interval J with $t_0 \in J \subset I$ and $\tilde{X}|_{x(t)} = X(t)$ for all $t \in J$. Then one defines $\nabla_{\dot{x}(t_0)} X$ as $\nabla_{\dot{x}(t_0)} \tilde{X}$ and it is easily seen that this vector does not depend on the choice of a local extension of X around t_0 . The same construction holds true for tensor fields along the path $x(\cdot)$.

The parallel transport of a tensor $T_0 \in T_m^k|_{x(0)}(M)$ from $x(0)$ to $x(t)$ along an absolutely continuous curve $x : I \rightarrow M$ (with $0 \in I$) and with respect to the Levi-Civita connection of (M, g) is denoted by $(P^{\nabla^g})_0^t(x)T_0$. In the notation of the Levi-Civita connection ∇^g (resp. parallel transport P^{∇^g}), the upper index g (resp. ∇^g) referring to the Riemannian metric g (resp. the connection ∇^g) is omitted if it is clear from the context. We also recall the following basic observation.

Proposition 2.1 Let (M, g) be a smooth Riemannian manifold and $t \mapsto x(t)$ an absolutely continuous (a.c. for short) curve on M defined on an open interval $I \ni 0$. Then the parallel transport $T(t) = (P^{\nabla^g})_0^t(x)T_0$ along $t \mapsto x(t)$ w.r.t g of any (k, m) -tensor $T_0 \in T_m^k|_{x(0)}(M)$ uniquely exists and is absolutely continuous.

Let $(x, \hat{x}) : I \rightarrow M \times \hat{M}$ be a smooth curve on $M \times \hat{M}$ defined on an open real interval I containing 0. If $(X(t), \hat{X}(t)) : I \rightarrow T(M \times \hat{M})$ is a smooth vector field on $M \times \hat{M}$ along (x, \hat{x}) i.e., $(X(t), \hat{X}(t)) \in T|_{(x(t), \hat{x}(t))}(M \times \hat{M})$ then one has

$$\overline{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))}(X, \hat{X}) = (\nabla_{\dot{x}(t)} X, \hat{\nabla}_{\dot{\hat{x}}(t)} \hat{X}) \quad (7)$$

only if the covariant derivatives on the right-hand side are well defined (see the next remark).

Remark 2.2 Let $M = \mathbb{R}$, $\hat{M} = \mathbb{R}$ and $(c(t), \hat{c}(t)) = (t, 0)$, $(X(t), \hat{X}(t)) = (1, t)$ and equip M and \hat{M} with the Euclidean metrics: $g(Y, Z) = YZ$, $\hat{g}(\hat{Y}, \hat{Z}) = \hat{Y}\hat{Z}$. Then the left hand side of (7) is defined and equals $(0, 1)$ but on the right hand side the covariant derivative $\hat{\nabla}_{\dot{\hat{c}}(t)} \hat{X} = \hat{\nabla}_0 t$ is not defined: if $\hat{Y} \in \text{VF}(\hat{M})$ were a local extension of \hat{X} around $t = 0$ then $t = \hat{X}(t) = \hat{Y}|_{\hat{c}(t)} = \hat{Y}(0)$ for all t in some open interval containing 0. This is a contradiction. Note that an extension of $(X(t), \hat{X}(t)) = (1, t)$ around $t = 0$ is provided for example by $(x, \hat{x}) \mapsto (1, x)$.

If (N, h) is a Riemannian manifold we define $\text{Iso}(N, h)$ to be the (smooth Lie) group of isometries of (N, h) i.e., the set of diffeomorphisms $F : N \rightarrow N$ such that $F_*|_y : T|_y N \rightarrow T|_{F(y)} N$ is an isometry for all $y \in N$, cf. [25], Lemma III.6.4, p. 118.

It is clear that the isometries respect parallel transport in the sense that for any absolutely continuous $\gamma : [a, b] \rightarrow N$ and $F \in \text{Iso}(N, g)$ one has (cf. [25], p. 41, Eq. (3.5))

$$F_*|_{\gamma(t)} \circ (P^{\nabla^h})_a^t(\gamma) = (P^{\nabla^h})_a^t(F \circ \gamma) \circ F_*|_{\gamma(a)}. \quad (8)$$

The following result is standard.

Theorem 2.3 Let (N, h) be a Riemannian manifold and for any absolutely continuous $\gamma : [0, 1] \rightarrow N$, $\gamma(0) = y_0$, define

$$\Lambda_{y_0}^{\nabla^h}(\gamma)(t) = \int_0^t (P^{\nabla^h})_s^0(\gamma) \dot{\gamma}(s) ds \in T|_{y_0}N, \quad t \in [0, 1].$$

Then the map $\Lambda_{y_0}^{\nabla^h} : \gamma \mapsto \Lambda_{y_0}^{\nabla^h}(\gamma)(\cdot)$ is an injection from the set of absolutely continuous curves $[0, 1] \rightarrow N$ starting at y_0 onto an open subset of the Banach space of absolutely continuous curves $[0, 1] \rightarrow T|_{y_0}N$ starting at 0.

Moreover, the map $\Lambda_{y_0}^{\nabla^h}$ is a bijection onto the latter Banach space if (and only if) (N, h) is a complete Riemannian manifold.

Remark 2.4 (i) For example, in the case where γ is the geodesic $t \mapsto \exp_{y_0}(tY)$ for $Y \in T|_{y_0}N$, one has

$$\Lambda_{y_0}^{\nabla^h}(\gamma)(t) = tY.$$

- (ii) It is directly seen from the definition of $\Lambda_{y_0}^{\nabla^h}$ that it maps injectively (piecewise) C^k -curves, $k = 1, \dots, \infty$, starting at y_0 to (piecewise) C^k -curves starting at 0. Moreover, these correspondences are bijective if (N, h) is complete.
- (iii) The map $\Lambda_{y_0}^{\nabla^h}$ could be used to give the space of absolutely continuous curves $[0, 1] \rightarrow N$ starting at y_0 a structure of a Banach space if (N, h) is complete or an open subset of a Banach space in the case (N, h) is not complete.

3 Characterization of the Rolling Problems as Control Problems

3.1 Definition of the State Space

Following [2], [3] we make the following definition.

Definition 3.1 The *state space* $Q = Q(M, \hat{M})$ for the rolling of two n -dimensional *connected, oriented* smooth Riemannian manifolds $(M, g), (\hat{M}, \hat{g})$ is defined as

$$Q = \{A : T|_xM \rightarrow T|_{\hat{x}}\hat{M} \mid A \text{ o-isometry, } x \in M, \hat{x} \in \hat{M}\},$$

where ‘‘o-isometry’’ stands for ‘‘orientation preserving isometry’’ i.e., if $(X_i)_{i=1}^n$ is a positively oriented g -orthonormal frame of M at x then $(AX_i)_{i=1}^n$ is a positively oriented \hat{g} -orthonormal frame of \hat{M} at \hat{x} .

The linear space of \mathbb{R} -linear map $A : T|_x M \rightarrow T|_{\hat{x}} \hat{M}$ is canonically isomorphic to the tensor product $T^*|_x M \otimes T|_{\hat{x}} \hat{M}$. On the other hand, by using the canonical inclusions $T^*|_x M \subset T^*|_{(x,\hat{x})}(M \times \hat{M})$, $T|_{\hat{x}} \hat{M} \subset T|_{(x,\hat{x})}(M \times \hat{M})$, the space $T^*|_x M \otimes T|_{\hat{x}} \hat{M}$ is canonically included in the space $T_1^1(M \times \hat{M})|_{(x,\hat{x})}$ of $(1, 1)$ -tensors of $M \times \hat{M}$ at (x, \hat{x}) . These inclusions make $T^*M \otimes T\hat{M} := \bigcup_{(x,\hat{x}) \in M \times \hat{M}} T^*|_x M \otimes T|_{\hat{x}} \hat{M}$ a subset of $T_1^1(M \times \hat{M})$ such that $\pi_{T^*M \otimes T\hat{M}} := \pi_{T_1^1(M \times \hat{M})}|_{T^*M \otimes T\hat{M}} : T^*M \otimes T\hat{M} \rightarrow M \times \hat{M}$ is a smooth vector subbundle of the bundle of $(1, 1)$ -tensors $\pi_{T_1^1(M \times \hat{M})}$ on $M \times \hat{M}$.

The state space $Q = Q(M, \hat{M})$ can now be described as a subset of $T^*M \otimes T\hat{M}$ as

$$Q = \{A \in T^*(M) \otimes T(\hat{M})|_{(x,\hat{x})} \mid (x, \hat{x}) \in M \times \hat{M}, \\ \|AX\|_{\hat{g}} = \|X\|_g, \forall X \in T|_x M, \det(A) = 1\}.$$

In the next subsection, we will show that $\pi_Q := \pi_{T^*M \otimes T\hat{M}}|_Q$ is moreover a smooth subbundle of $\pi_{T^*M \otimes T\hat{M}}$ though it is not a vector subbundle.

It is also convenient to consider the manifold $T^*M \otimes T\hat{M}$ and we will refer to it as the *extended state space* for the rolling. This concept of extended state space naturally makes sense also in the case where M and \hat{M} are not assumed to be oriented (or connected).

A point $A \in T^*M \otimes T\hat{M}$ with $\pi_{T^*M \otimes T\hat{M}}(A) = (x, \hat{x})$ (or $A \in Q$ with $\pi_Q(A) = (x, \hat{x})$) will be sometimes denoted by $(x, \hat{x}; A)$ to emphasize the fact that $A : T|_x M \rightarrow T|_{\hat{x}} \hat{M}$. Thus the notation $q = (x, \hat{x}; A)$ simply means that $q = A$.

3.2 The Bundle Structure of Q

In this subsection, it is shown that π_Q is a bundle with typical fiber $\text{SO}(n)$. We will also argue that, even though $\text{SO}(n)$ is a Lie-group, the bundle π_Q cannot in general be given a natural (or useful) $\text{SO}(n)$ -principal bundle if $n > 2$ (see also Theorem 4.14). We will now present the local trivializations of π_Q .

Definition 3.2 Suppose the vector fields $X_i \in \text{VF}(M)$ (resp. $\hat{X}_i \in \text{VF}(\hat{M})$), $i = 1, \dots, n$ form a g -orthonormal (resp. \hat{g} -orthonormal) frame of vector fields on an open subset U of M (resp. \hat{U} of \hat{M}). We denote $F = (X_i)_{i=1}^n$, $\hat{F} = (\hat{X}_i)_{i=1}^n$ and for $x \in U$, $\hat{x} \in \hat{U}$ we let $F|_x = (X_i|_x)_{i=1}^n$, $\hat{F}|_{\hat{x}} = (\hat{X}_i|_{\hat{x}})_{i=1}^n$. Then a local trivialization $\tau = \tau_{F, \hat{F}}$ of Q over $U \times \hat{U}$ induced by F, \hat{F} is given by

$$\tau : \pi_Q^{-1}(U \times \hat{U}) \rightarrow (U \times \hat{U}) \times \text{SO}(n) \\ (x, \hat{x}; A) \mapsto ((x, \hat{x}), \mathcal{M}_{F|_x, \hat{F}|_{\hat{x}}}(A)),$$

where $\mathcal{M}_{F|_x, \hat{F}|_{\hat{x}}}(A)_i^j = \hat{g}(AX_i, \hat{X}_j)$ since $AX_i|_x = \sum_j \hat{g}(AX_i|_x, \hat{X}_j|_{\hat{x}}) \hat{X}_j|_{\hat{x}}$.

For the sake of clarity, we shall write $\mathcal{M}_{F|_x, \hat{F}|_{\hat{x}}}(A)$ as $\mathcal{M}_{F, \hat{F}}(A)$. Obviously $\|AX\|_{\hat{g}} = \|X\|_g$ for all $X \in T|_x M$ is equivalent to $A^{T_{g, \hat{g}}} A = \text{id}_{T|_x M}$ and thus we get

$$\mathcal{M}_{F, \hat{F}}(A)^T \mathcal{M}_{F, \hat{F}}(A) = \mathcal{M}_{\hat{F}, F}(A^{T_{g, \hat{g}}}) \mathcal{M}_{F, \hat{F}}(A) = \mathcal{M}_{F, F}(\text{id}_{T|_x M}) = \text{id}_{\mathbb{R}^n},$$

where T denotes the usual transpose in $\mathfrak{gl}(n)$, the set of Lie algebra of $n \times n$ -real matrices. Since $\det \mathcal{M}_{F, \hat{F}}(A) = \det(A) = +1$, one finally has $\mathcal{M}_{F, \hat{F}}(A) \in \text{SO}(n)$.

Remark 3.3 Notice that the above local trivializations $\tau_{F, \hat{F}}$ of π_Q are just the restrictions of the vector bundle local trivializations

$$(\pi_{T^*(M) \otimes T(\hat{M})})^{-1}(U \times \hat{U}) \rightarrow (U \times \hat{U}) \times \mathfrak{gl}(n)$$

of the bundle $\pi_{T^*(M) \otimes T(\hat{M})}$ induced by F, \hat{F} and defined by the same formula as $\tau_{F, \hat{F}}$. In this setting, one does not even have to assume that the local frames F, \hat{F} are g - or \hat{g} -orthonormal. Hence π_Q is a smooth subbundle of $\pi_{T^*M \otimes T\hat{M}}$ with Q a smooth submanifold of $T^*M \otimes T\hat{M}$.

We next spell out the transition functions of the above defined local trivializations of π_Q (and also of $\pi_{T^*M \otimes T\hat{M}}$ by the above remark). If $F' = ((X'_i), U'), \hat{F}' = ((\hat{X}'_i), \hat{U}')$ are other g -, \hat{g} -orthonormal frames (with $U \cap U' \neq \emptyset, \hat{U} \cap \hat{U}' \neq \emptyset$) and $A = [A_i^j] \in \text{SO}(n)$, then

$$\begin{aligned} & (\tau_{F', \hat{F}'} \circ \tau_{F, \hat{F}}^{-1})((x, \hat{x}), A) = \tau_{F', \hat{F}'}(x, \hat{x}; \sum_{i,j} A_i^j g(X_i, \cdot) \hat{X}_j) \\ = & ((x, \hat{x}), \left[\left(\sum_{i,j} A_i^j g(X_i, X'_k) \hat{g}(\hat{X}_j, \hat{X}'_l) \right)_k^l \right]) \\ = & ((x, \hat{x}), [(\hat{g}(\hat{X}_j, \hat{X}'_l))_j^l] A [(g(X_i, X'_k))_i^k]^T) \\ = & ((x, \hat{x}), \mathcal{M}_{\hat{F}, \hat{F}'}(\text{id}_{T|_{\hat{x}\hat{M}}}) A \mathcal{M}_{F, F'}(\text{id}_{T|xM})^T) \end{aligned}$$

for $x \in U \cap U', \hat{x} \in \hat{U} \cap \hat{U}'$.

Any local trivialization $\tau : \pi_Q^{-1}(\bar{U}) \rightarrow \bar{U} \times \text{SO}(n)$ of π_Q defined on an open set $\bar{U} \subset M \times \hat{M}$ would define a principal $\text{SO}(n)$ -bundle structure on $\pi_Q^{-1}(\bar{U})$ (or rather for $\pi_Q|_{\pi_Q^{-1}(\bar{U})}$) by the formula (see [28], p. 307)

$$\mu((x, \hat{x}; A), B) = \tau^{-1}((x, \hat{x}), (\text{pr}_2 \circ \tau)(x, \hat{x}; A)B), \quad (9)$$

with $\mu : \pi_Q^{-1}(\bar{U}) \times \text{SO}(n) \rightarrow \pi_Q^{-1}(\bar{U})$ the right $\text{SO}(n)$ -action of this principal bundle structure. However, we will show that if we take for the local trivializations τ the ones induced by local orthonormal frames $\tau = \tau_{F, \hat{F}}$ as above, then the (local) actions $\mu_{F, \hat{F}}$ defined by the above formula by these different local trivializations $\tau_{F, \hat{F}}$ do not glue up to form a global principal bundle structure for π_Q if the dimension n of M and \hat{M} is greater than 2. We state this in the following proposition.

Proposition 3.4 The local actions (9) do not render the bundle π_Q to a principal $\text{SO}(n)$ -bundle except when $n \leq 2$.

Proof. If π_Q were a principal $\text{SO}(n)$ -bundle w.r.t local trivializations induced by the orthonormal frames of M and \hat{M} , then the right action $\mu : Q \times \text{SO}(n) \rightarrow Q$ of $\text{SO}(n)$ on Q of this principal bundle structure would be given (locally) by (see above)

$$\mu((x, \hat{x}; A), B) = \tau^{-1}((x, \hat{x}), (\text{pr}_2 \circ \tau)(x, \hat{x}; A)B),$$

for any of the local trivializations $\tau = \tau_{F, \hat{F}}$ induced by orthonormal local frames F, \hat{F} of M, \hat{M} and any $(x, \hat{x}; A)$ with x, \hat{x} in these domains and any $B \in \text{SO}(n)$. Equivalently, the above condition could be written as

$$(\text{pr}_2 \circ \tau)(\mu(q, B)) = (\text{pr}_2 \circ \tau)(q)B,$$

for any $q \in Q$ in the domain of definition of τ and $B \in \text{SO}(n)$.

The formula for the transition maps of these local trivializations as expressed before this proposition shows that the action μ is not well defined if $n \geq 3$. In fact we would be led to an equation of the type

$$(\text{pr}_2 \circ \tau_{F', \hat{F}'})(x, \hat{x}; A)B = (\text{pr}_2 \circ \tau_{F', \hat{F}'} \circ \tau_{F, \hat{F}}^{-1})((x, \hat{x}), (\text{pr}_2 \circ \tau_{F, \hat{F}})(x, \hat{x}; A)B),$$

i.e.,

$$\begin{aligned} \mathcal{M}_{F', \hat{F}'}(A)B &= \mathcal{M}_{\hat{F}, \hat{F}'}(\text{id}_{T|_{\hat{x}}\hat{M}})(\mathcal{M}_{F, \hat{F}}(A)B)\mathcal{M}_{F, F'}(\text{id}_{T|x}M)^{-1} \\ &= \mathcal{M}_{F, \hat{F}'}(A)B\mathcal{M}_{F', F}(\text{id}_{T|x}M) \end{aligned}$$

which, by multiplying by $\mathcal{M}_{F, \hat{F}'}(A)^{-1}$ from the left, is equivalent to

$$\mathcal{M}_{F', F}(\text{id}_{T|x}M)B = B\mathcal{M}_{F', F}(\text{id}_{T|x}M) \quad (10)$$

Since $\text{SO}(n)$ is not commutative for $n \geq 3$, the left and right hand sides are not equal in general: they are equal for all B, F, F' if and only if $\text{SO}(n)$ is commutative i.e. if and only if $n \in \{1, 2\}$. Hence π_Q is not a principal $\text{SO}(n)$ -bundle, at least w.r.t the trivializations that we used, if $n \geq 3$. \square

Remark 3.5 If M and \hat{M} are parallelizable (e.g. if M and \hat{M} are Lie groups) i.e., if there are global frames and hence global orthonormal frames F, \hat{F} , then one can introduce a principal $\text{SO}(n)$ -bundle structure for π_Q by Eq. (9) even for $n > 2$. However, this principal bundle structure then depends on the choice of the global frames F, \hat{F} i.e., we might (and could if $n > 2$) get a different principal bundle structure by the choosing the orthonormal frames differently. We will define on Q a distribution \mathcal{D}_R (see Definition 5.2) that models the natural constraints for the rolling problem and by simple computations one can check that in general for $n \geq 3$ the distribution \mathcal{D}_R is not invariant with respect to this principal bundle action for π_Q .

Hence the principal bundle structure on parallelizable manifolds (or, in the general case, the local principal bundle structures defined by (9)) is (in general) not useful for the study of the rolling model.

We will also study briefly a less restrictive model of rolling (rolling with spinning allowed) where one considers a distribution \mathcal{D}_{NS} on Q . In this case, it will be shown in Theorem 4.14 below that in general there cannot be a principal bundle structure for π_Q which leaves \mathcal{D}_{NS} invariant.

Remark 3.6 Clearly the fact that we chose $\text{SO}(n)$ to act on the right in (9) does not affect the conclusion of the previous Proposition: Left local actions (in an obvious manner) lead to the same conclusion i.e., they don't glue up correctly to give a "natural" global $\text{SO}(n)$ -action.

Indeed, if instead of (9) we tried to define the *left* $\text{SO}(n)$ -action on Q by demanding that locally

$$\lambda(B, (x, \hat{x}; A)) = \tau^{-1}((x, \hat{x}), B(\text{pr}_2 \circ \tau)(x, \hat{x}; A)),$$

we still could not define the action globally. Indeed, it is enough to notice that instead of (10) we would get

$$\begin{aligned} B\mathcal{M}_{F', \hat{F}'}(A) &= \mathcal{M}_{\hat{F}, \hat{F}'}(\text{id}_{T|_{\hat{x}}\hat{M}})(B\mathcal{M}_{F, \hat{F}}(A))\mathcal{M}_{F, F'}(\text{id}_{T|x}M)^{-1} \\ &= \mathcal{M}_{\hat{F}, \hat{F}'}(\text{id}_{T|_{\hat{x}}\hat{M}})B\mathcal{M}_{F', \hat{F}}(A) \end{aligned}$$

i.e.

$$B\mathcal{M}_{\hat{F},\hat{F}'}(\text{id}_{T|_{\hat{x}}\hat{M}}) = \mathcal{M}_{\hat{F},\hat{F}'}(\text{id}_{T|_{\hat{x}}\hat{M}})B$$

which, again, is only true for all B, \hat{F}, \hat{F}' if and only if $n \in \{1, 2\}$.

Despite the lack of a "natural" principal bundle structure for π_Q when $n \geq 3$, we may still make use of the vector bundle structure of the ambient bundle $\pi_{T^*(M) \otimes T(\hat{M})}$ (the extended state space).

Notice that any π_Q -vertical tangent vector (i.e., an element of $V|_q(\pi_Q)$) is of the form $\nu(B)|_q$ for a unique $B \in T^*M \otimes T\hat{M}|_{(x,\hat{x})}$ where $q = (x, \hat{x}; A) \in Q$. The following simple proposition gives the condition when, for a $B \in T^*M \otimes T\hat{M}|_{(x,\hat{x})}$, the vector $\nu(B)|_q \in V|_q(\pi_{T^*M \otimes T\hat{M}})$ is actually tangent to Q i.e., an element of $V|_q(\pi_Q)$.

Proposition 3.7 Let $q = (x, \hat{x}; A) \in Q$ and $B \in T^*(M) \otimes T(\hat{M})|_{(x,\hat{x})}$. Then $\nu(B)|_q$ is tangent to Q (i.e., is an element of $V|_q(\pi_Q)$) if and only if

$$\hat{g}(AX, BY) + \hat{g}(BX, AY) = 0$$

for all $X, Y \in T|_x M$. Denoting $\overline{T} = T_{g,\hat{g}}$, this latter condition can be stated equivalently as $A\overline{T}B + B\overline{T}A = 0$ or more compactly as $B \in A(\mathfrak{so}(T|_x M))$

We will be denoting the (g, \hat{g}) -transpose operation $T_{g,\hat{g}}$ by \overline{T} also in the sequel. The proposition says that $V|_{(x,\hat{x};A)}(\pi_Q)$ is naturally \mathbb{R} -linearly isomorphic to $A(\mathfrak{so}(T|_x M))$.

Remark 3.8 We may reformulate the fact given by the previous proposition as follows. Define $\mathfrak{so}(M) = \bigcup_{x \in M} \mathfrak{so}(T|_x M)$ (with M a Riemannian manifold) i.e.,

$$\mathfrak{so}(M) = \{B \in T_1^1(M) \mid B^{T_g} + B = 0\}.$$

One sees that $\mathfrak{so}(M)$ is a closed embedded submanifold of $T_1^1(M) = T^*M \otimes TM$. Moreover, the map $\pi_{\mathfrak{so}(M)} := \pi_{T_1^1(M)}|_{\mathfrak{so}(M)}$ clearly defines a smooth vector bundle with typical fiber $\mathfrak{so}(n)$, where $n = \dim(M)$.

We may pull back $\pi_{\mathfrak{so}(M)}$ with a map $\pi_{Q,M} := \text{pr}_1 \circ \pi_Q : Q \rightarrow M$ to a smooth bundle $(\pi_{Q,M})^*(\pi_{\mathfrak{so}(M)}) : (\pi_{Q,M})^*(\mathfrak{so}(M)) \rightarrow Q$ over Q . Its elements are all pairs $((x, \hat{x}; A), B) \in Q \times \mathfrak{so}(M)$ where $x = \pi_{\mathfrak{so}(M)}(B)$ and the bundle map is defined by $(\pi_{Q,M})^*(\pi_{\mathfrak{so}(M)})((x, \hat{x}; A), B) = (x, \hat{x}; A)$.

Proposition 3.7 shows that the bundle map $L : (\pi_{Q,M})^*(\pi_{\mathfrak{so}(M)}) \rightarrow V(\pi_Q)$ defined by $L((x, \hat{x}; A), B) = \nu(AB)|_{(x,\hat{x};A)}$ is a diffeomorphism.

3.3 The State Space as a Quotient

In this subsection, we will show that (the n -dimensional version of) the construction of the state space for rolling that has been used e.g. in [5] in dimension two is actually isomorphic to the state space Q .

Proposition 3.9 Let $F_{\text{OON}}(M), F_{\text{OON}}(\hat{M})$ be the oriented orthonormal frame bundles of $(M, g), (\hat{M}, \hat{g})$ (resp. let $F(M), F(\hat{M})$ be the frame bundles of M and \hat{M}). Denote by $\mu, \hat{\mu}$ the right $\text{SO}(n)$ -actions (resp. right $\text{GL}(n)$ -actions) defining the usual

principal bundle structures on these spaces i.e., $\mu((X_k)_{k=1}^n, [A_j^i]) = (\sum_k A_i^k X_k)_{i=1}^n$ and similarly for $\hat{\mu}$. Define a diagonal right $\text{SO}(n)$ -action

$$\Delta : (F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})) \times \text{SO}(n) \rightarrow F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M}),$$

by (resp. right $\text{GL}(n)$ -action $\Delta : (F(M) \times F(\hat{M})) \times \text{GL}(n) \rightarrow F(M) \times F(\hat{M}))$)

$$\Delta(((X_i), (\hat{X}_j)), A) = (\mu((X_i), A), \hat{\mu}((\hat{X}_i), A)).$$

The map $\xi : F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M}) \rightarrow Q(M, \hat{M})$ (resp. $\xi : F(M) \times F(\hat{M}) \rightarrow T^*M \otimes T\hat{M}$) such that

$$\xi((X_i), (\hat{X}_j)) := \left(\sum_i a_i X_i \mapsto \sum_i a_i \hat{X}_i \right)$$

is a smooth surjective submersion. Moreover, for each $q \in Q(M, \hat{M})$ (resp. $q \in T^*M \otimes T\hat{M}$) the inverse image $\xi^{-1}(q)$ coincides with an orbit of Δ . Thus ξ induces a diffeomorphism $\bar{\xi} : (F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M}))/\Delta \rightarrow Q(M, \hat{M})$. (resp. $\bar{\xi} : (F(M) \times F(\hat{M}))/\Delta \rightarrow \{A \in T^*M \otimes T\hat{M} \mid A \text{ is invertible}\}$).

Proof. The smoothness and surjectivity of ξ are obvious and it is also easy to see that ξ is a submersion. Thus it is enough to show that $\xi^{-1}(q)$ coincides with an orbit of Δ . First suppose that

$$\mu((X_i), A)_i = \sum_j A_i^j X_j, \quad \hat{\mu}((\hat{X}_i), A)_i = \sum_j A_i^j \hat{X}_j.$$

Then, for any real numbers a_1, \dots, a_n , one has

$$\begin{aligned} & \xi(\Delta(((X_i), (\hat{X}_i)), A)) \left(\sum_k a_k X_k \right) = \xi(\mu((X_i), A), \hat{\mu}((\hat{X}_i), A)) \left(\sum_k a_k X_k \right) \\ &= \xi(\mu((X_i), A), \hat{\mu}((\hat{X}_i), A)) \left(\sum_{k,i} a_k (A^{-1})_k^i \mu((X_j), A)_i \right) \\ &= \sum_{k,i} a_k (A^{-1})_k^i \hat{\mu}((\hat{X}_j), A)_i = \sum_{k,i,j} a_k (A^{-1})_k^i A_i^j \hat{X}_j = \sum_k a_k \hat{X}_k = \xi((X_i), (\hat{X}_i)) \left(\sum_k a_k X_k \right). \end{aligned}$$

This shows that

$$\Delta(\{((X_i), (\hat{X}_i))\} \times G) \subset \xi^{-1}(\xi((X_i), (\hat{X}_i))),$$

with $G = \text{SO}(n)$ (resp. $G = \text{GL}(n)$). The orbits of Δ all have the same dimension as $\text{SO}(n)$, i.e., $\frac{n(n-1)}{2}$ (resp. $\dim \text{GL}(n) = n^2$) and since

$$\dim \xi^{-1}(q) = \dim(F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})) - \dim Q(M, \hat{M}) = \dim \text{SO}(n),$$

for any $q \in Q(M, \hat{M})$ (resp.

$$\dim \xi^{-1}(q) = \dim(F(M) \times F(\hat{M})) - \dim T^*M \otimes T\hat{M} = \dim \text{GL}(n)),$$

we have that this inclusion is actually an equality. This proves the proposition. \square

Remark 3.10 In the above proposition we implicitly assumed that $(F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M}))/\Delta$ (resp. $(F(M) \times F(\hat{M}))/\Delta$) already has a natural structure of a smooth manifold namely that of a quotient manifold. But it is easily seen that the action Δ is free and proper and hence by a well known result (see [15] Theorem 9.16) it follows that unique smooth quotient manifold structures for the above quotient sets exist. Hence the facts established in the above proof guarantee that $\bar{\xi}$ is a diffeomorphism.

Remark 3.11 Here is the product right action

$$\mu \times \hat{\mu} : (F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})) \times (\text{SO}(n) \times \text{SO}(n)) \rightarrow F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})$$

of $\text{SO}(n) \times \text{SO}(n)$ on $F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M})$ given by

$$\mu \times \hat{\mu}(((X_i), (\hat{X}_i)), (A, \hat{A})) = (\mu((X_i), A), \hat{\mu}((\hat{X}_i), \hat{A})).$$

As it is easily seen, it is unfortunately *not* true that the action $\mu \times \hat{\mu}$ maps a Δ -orbit into a Δ -orbit, unless the dimension n is equal to two (in which case $\text{SO}(n) = \text{SO}(2)$ is commutative) and hence, in the case $n > 2$, the map $\mu \times \hat{\mu}$ does not induce a map $Q \times \text{SO}(n) \rightarrow Q$ (where $Q \cong (F_{\text{OON}}(M) \times F_{\text{OON}}(\hat{M}))/\Delta$ by the above proposition). This is yet another way of seeing that $Q = Q(M, \hat{M})$ cannot be given a "natural" $\text{SO}(n)$ -principal bundle structure for $n \geq 3$ i.e., we cannot induce on Q the principal bundle structures of the frame bundles $F_{\text{OON}}(M)$ and $F_{\text{OON}}(\hat{M})$ if $n > 2$.

Remark 3.12 Notice that on $F(M)$ (resp. on $F_{\text{OON}}(M)$) one may also consider the left $\text{GL}(n)$ (resp. $\text{SO}(n)$) action λ given by $\lambda(A, (X_i))_i = \sum_j A_j^i X_j$. Since $A_j^i = (A^T)_i^j$ it is trivial that this is related to the above right action by $\lambda(A, (X_i)) = \mu((X_i), A^T)$. Notice that $\mu(\lambda(A, (X_i)), B) = \mu(\mu((X_i), A^T), B) = \mu((X_i), A^T B)$ which, if $n \geq 3$ and $A^T B \neq B A^T$, is different from $\lambda(A, \mu((X_i), B)) = \mu(\mu((X_i), B), A^T) = \mu((X_i), B A^T)$. This means that the left and right actions λ and μ do not "commute".

Another way to define naturally a left actions is to use instead of above λ the inverse right-action $\lambda_I(A, (X_i)) := \mu((X_i), A^{-1})$. Also in this case, $\mu(\lambda_I(A, (X_i)), B) = \mu(\mu((X_i), A^{-1}), B) = \mu((X_i), A^{-1} B)$ is not equal, if $n \geq 3$ and $AB \neq BA$, to $\lambda_I(A, \mu((X_i), B)) = \mu(\mu((X_i), B), A^{-1}) = \mu((X_i), B A^{-1})$. On $F_{\text{OON}}(M)$ it is clear that the actions λ and λ_I coincide.

It was proposed in [8] that one could use the inverse left action on $F_{\text{OON}}(M)$ and the left action on $F_{\text{OON}}(\hat{M})$ to induce, respectively, left and right actions on Q . However this is not possible for the following reason (which basically is a repetition of what has been said above). Suppose $q = (x, \hat{x}; A) \in Q$ and let $F, F' \in F_{\text{OON}}(M)$, $\hat{F}, \hat{F}' \in F_{\text{OON}}(\hat{M})$ are such that $\xi(F, \hat{F}) = q$ and $\xi(F', \hat{F}') = q$. Then there is a $B \in \text{SO}(n)$ such that $\mu(F, B) = F'$, $\hat{\mu}(\hat{F}, B) = \hat{F}'$. By using, for example, the left $\text{SO}(n)$ -action λ on $F_{\text{OON}}(M)$ we get $\lambda(C, F') = \lambda(C, \mu(F, B)) = \mu(F, B C^T)$ and also $\mu(\lambda(C, F), B) = \mu(F, C^T B)$. But $\xi(\lambda(C, F'), \hat{F}') = \xi(\lambda(C, F), \hat{F})$ if and only if $\mu(\lambda(C, F), B) = \lambda(C, F')$ which thus is not true unless $C^T B = B C^T$. The case of the inverse left action (which is just the right action $\hat{\mu}$) on $F_{\text{OON}}(\hat{M})$ leads to the same conclusion.

3.4 The Control Problem

Each point $(x, \hat{x}; A)$ of the state space $Q = Q(M, \hat{M})$ can be viewed as describing a contact point of the two manifolds which is given by the points x and \hat{x} of M

and \hat{M} , respectively, and an isometry A of the tangent spaces $T|_x M$, $T|_{\hat{x}} \hat{M}$ at this contact point. The isometry A can be viewed as measuring the relative orientation of these tangent spaces relative to each other in the sense that rotation of, say, $T|_{\hat{x}} \hat{M}$ corresponds to a unique change of the isometry A from $T|_x M$ to $T|_{\hat{x}} \hat{M}$. A curve $t \mapsto (x(t), \hat{x}(t); A(t))$ in Q can then be seen as a motion of M against \hat{M} such that at an instant t , $x(t)$ and $\hat{x}(t)$ represent the common point of contact in M and \hat{M} , respectively, and $A(t)$ measures the relative orientation of coinciding tangent spaces $T|_{x(t)} M$, $T|_{\hat{x}(t)} \hat{M}$ at this point of contact.

In order to call this motion *rolling*, there are two kinematic constraints that will be demanded (see e.g. [2], [3] Chapter 24, [7]) namely

- (i) the *no-spinning* condition;
- (ii) the *no-slipping* condition.

In this section, these conditions will be defined explicitly and it will turn out that they are modeled by certain smooth distributions on the state space Q . The subsequent sections are then devoted to the detailed definitions and analysis of the distribution \mathcal{D}_{NS} and \mathcal{D}_{R} on the state space Q , the former capturing the no-spinning condition (i) while the latter capturing both of the conditions (i) and (ii).

The first restriction (i) for the motion is that the relative orientation of the two manifolds should not change along motion. This *no-spinning condition* (also known as the no-twisting condition) can be formulated as follows.

Definition 3.13 An absolutely continuous (a.c.) curve

$$\begin{aligned} q : I &\rightarrow Q, \\ t &\mapsto (x(t), \hat{x}(t); A(t)), \end{aligned}$$

defined on some real interval $I = [a, b]$, is said to describe a *motion without spinning* of M against \hat{M} if, for every a.c. curve $[a, b] \rightarrow TM; t \mapsto X(t)$ of vectors along $t \mapsto x(t)$, we have

$$\nabla_{\dot{x}(t)} X(t) = 0 \quad \implies \quad \hat{\nabla}_{\dot{\hat{x}}(t)} (A(t)X(t)) = 0 \quad \text{for a.e. } t \in [a, b]. \quad (11)$$

(See also [8] for a similar definition.) Notice that Condition (11) is equivalent to the following: for almost every t and all parallel vector fields $X(\cdot)$ along $x(\cdot)$, one has

$$(\overline{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))} A(t))X(t) = 0.$$

(This is well defined as mentioned in the paragraph immediately below Eq. (6).)

Since the parallel translation $P_0^t(x) : T|_{x(0)} M \rightarrow T|_{x(t)} M$ along $x(\cdot)$ is an (isometric) isomorphism (here $X(t) = P_0^t(x)X(0)$), this shows that (11) is equivalent to

$$\overline{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))} A(t) = 0 \quad \text{for a.e. } t \in [a, b]. \quad (12)$$

The second restriction (ii) is that the manifolds should not slip along each other as they move i.e., the velocity of the contact point should be the same w.r.t both manifolds. This *no-slipping condition* can be formulated as follows.

Definition 3.14 An a.c. curve $I \rightarrow Q; t \mapsto (x(t), \hat{x}(t); A(t))$, defined on some real interval $I = [a, b]$, is said to describe a *motion without slipping* of M against \hat{M} if

$$A(t)\dot{x}(t) = \dot{\hat{x}}(t) \quad \text{for a.e. } t \in [a, b]. \quad (13)$$

Definition 3.15 An a.c. curve $I \rightarrow Q; t \mapsto (x(t), \hat{x}(t); A(t))$, defined on some real interval $I = [a, b]$, is said to describe a *rolling motion* i.e., a *motion without slipping or spinning* of M against \hat{M} if it satisfied both of the conditions (11),(13) (or equivalently (12),(13)). The corresponding curve $t \mapsto (x(t), \hat{x}(t); A(t))$ that satisfies these conditions is called a *rolling curve*.

It is easily seen that $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$, $t \in [a, b]$, is a rolling curve if and only if it satisfies the following driftless control affine system

$$(\Sigma)_R \quad \begin{cases} \dot{x}(t) = u(t), \\ \dot{\hat{x}}(t) = A(t)u(t), \\ \bar{\nabla}_{(u(t), A(t)u(t))} A(t) = 0, \end{cases} \quad \text{for a.e. } t \in [a, b]. \quad (14)$$

where the control u belongs to $\mathcal{U}(M)$, the set of measurable TM -valued functions u defined on some interval $I = [a, b]$ such that there exists a.c. $y : [a, b] \rightarrow M$ verifying $u = \dot{y}$ a.e. on $[a, b]$. Conversely, given any control $u \in \mathcal{U}(M)$ and $q_0 = (x_0, \hat{x}_0; A_0) \in Q$, a solution $q(\cdot)$ to this control system exists on a subinterval $[a, b']$, $a < b' \leq b$ satisfying the initial condition $q(a) = q_0$. The fact that System (14) is driftless and control affine can be seen from its representation in local coordinates (see (90) in Appendix A).

We end up this subsection by the following simple remark.

Remark 3.16 In many cases, it is more convenient to work in the extended state space $T^*(M) \otimes T(\hat{M})$ rather than in (its submanifold) Q because $\pi_{T^*(M) \otimes T(\hat{M})}$ is a vector bundle. Since the above constraints of motion (11) and (13) can also be formulated in this space in verbatim, we will sometimes take this more general approach and then restrict to Q .

4 Study of the Rolling problem (NS)

4.1 The No-Spinning Distribution \mathcal{D}_{NS}

In this section, we build a smooth distribution \mathcal{D}_{NS} on the spaces Q and $T^*M \otimes T\hat{M}$ which plays the role of modelling the no-spinning condition for the rolling, see (11). We will also study the geometry related to this distribution. For more general constructions and some more general results than the ones in this section, see [11], [13].

We begin by recalling some basic observations on parallel transport. As noted in Proposition 2.1, if one starts with a $(1, 1)$ -tensor $A_0 \in T_1^1|_{(x_0, \hat{x}_0)}(M \times \hat{M})$ and has an a.c. curve $t \mapsto (x(t), \hat{x}(t))$ on $M \times \hat{M}$ with $x(0) = x_0$, $\hat{x}(0) = \hat{x}_0$, defined on an open interval $I \ni 0$, then the parallel transport $A(t) = P_0^t(x, \hat{x})A_0$ exists on I and determines an a.c. curve. But now, if A_0 rather belongs to the subspace $T^*M \otimes T\hat{M}$ or Q of $T_1^1(M \times \hat{M})$, it will actually happen that the parallel translate $A(t)$ belongs to this subspace as well for all $t \in I$. This is the content of the next proposition.

Proposition 4.1 Let $t \mapsto (x(t), \hat{x}(t))$ be an absolutely continuous curve in $M \times \hat{M}$ defined on some real interval $I \ni 0$. Then we have

$$\begin{aligned} A_0 \in T^*M \otimes TM &\implies A(t) = P_0^t(x, \hat{x})A_0 \in T^*M \otimes T\hat{M} \quad \forall t \in I, \\ A_0 \in Q &\implies A(t) = P_0^t(x, \hat{x})A_0 \in Q \quad \forall t \in I, \end{aligned}$$

and

$$P_0^t(x, \hat{x})A_0 = P_0^t(\hat{x}) \circ A_0 \circ P_t^0(x) \quad \forall t \in I. \quad (15)$$

Proof. Let $Y \in T|_{x(0)}M$, $\hat{Y} \in T|_{\hat{x}(0)}\hat{M}$ and let $Y(t) = P_0^t(x)Y$, $\hat{Y}(t) = P_0^t(\hat{x})\hat{Y}$ be their parallel translates along $t \mapsto x(t)$ and $t \mapsto \hat{x}(t)$ respectively. Similarly, choose $\omega \in T^*|_{x(0)}M$ and denote $\omega(t) = P_0^t(x)\omega$ its parallel translate. Then $Y(t)$, $\hat{Y}(t)$ and $\omega(t)$ can be viewed as a curves in $T(M \times \hat{M})$ and $T^*(M \times \hat{M})$ using the canonical inclusions $T|_{x(t)}M, T|_{\hat{x}(t)}\hat{M} \subset T|_{(x(t), \hat{x}(t))}(M \times \hat{M})$, $T^*|_{x(t)}M \subset T^*|_{(x(t), \hat{x}(t))}(M \times \hat{M})$.

With $A_0 \in T^*M \otimes T\hat{M}|_{(x(0), \hat{x}(0))} \subset T_1^1(M \times \hat{M})|_{(x(0), \hat{x}(0))}$ and $A(t) = P_0^t(x, \hat{x})A_0 \in T_1^1(M \times \hat{M})|_{(x(t), \hat{x}(t))}$, we have, for a.e. t (the contractions that use are obvious),

$$\bar{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))}(A(\cdot)\omega(\cdot)) = (\bar{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))}A(\cdot))\omega(t) + A(t)(\nabla_{\dot{x}(t)}\omega(\cdot)) = 0,$$

and similarly $\bar{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))}(A(\cdot)\hat{Y}(\cdot)) = 0$. It implies that $A(t)\omega(t)$, $A(t)\hat{Y}(t)$ (as elements of $T_1^1(M \times \hat{M})$) are parallel to $t \mapsto (\dot{x}(t), \dot{\hat{x}}(t))$ with initial conditions $A_0\omega = 0$ and $A_0\hat{Y} = 0$ since $A_0 \in T^*M \otimes T\hat{M}$. By the uniqueness of solutions of ODEs, this shows that $A(t)\omega(t) = 0$ and $A(t)\hat{Y}(t) = 0$ for all $t \in I$ i.e., since \hat{Y}, ω were arbitrary, $A(t) \in T^*M \otimes T\hat{M}$ for all $t \in I$.

Suppose next that $A_0 \in Q|_{(x(0), \hat{x}(0))}$ and denote $A(t) = P_0^t(x, \hat{x})A_0$. Then $A_0 \in T^*M \otimes T\hat{M}$ and, by what we just proved, $A(t) \in T^*M \otimes T\hat{M}$ for all $t \in I$. It follows that $A(t)Y(t) \in T|_{\hat{x}(t)}\hat{M}$ and thus taking its norm w.r.t \hat{g} allows us to compute a.e.

$$\frac{d}{dt} \|A(t)Y(t)\|_{\hat{g}}^2 = 2\hat{g}((\bar{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))}A(\cdot))Y(t) + A(t)\nabla_{\dot{x}(t)}Y(\cdot), A(t)Y(t)) = 0.$$

The initial condition for $\|A(t)Y(t)\|_{\hat{g}}^2$ at $t = 0$ is $\|(A(0)Y(0))\|_{\hat{g}}^2 = \|Y\|_g^2$, since $A_0 = A(0)$ is an isometry (and $Y(0) = Y$). Since $\|Y(t)\|_{\hat{g}}^2$ also satisfies $\frac{d}{dt} \|Y(t)\|_{\hat{g}}^2 = 0$ and the initial condition $\|Y(0)\|_{\hat{g}}^2 = \|Y\|_g^2$, we see that $\|A(t)Y(t)\|_{\hat{g}}^2 = \|Y(t)\|_{\hat{g}}^2$ for all $t \in I$ (since the maps $t \mapsto \|A(t)Y(t)\|_{\hat{g}}^2$, $t \mapsto \|Y(t)\|_{\hat{g}}^2$ were a.c.). Since the parallel translation $P_0^t(x) : T|_{x(0)}M \rightarrow T|_{x(t)}M$ is a linear (isometric) isomorphism for every t , this proves that $A(t) : T|_{x(t)}M \rightarrow T|_{\hat{x}(t)}\hat{M}$ is an isometry for every t . Because $t \mapsto \det(A(t))$ is a continuous map $I \rightarrow \{-1, +1\}$ and $\det(A(0)) = \det(A_0) = +1$, it follows that $\det(A(t)) = +1$ for all t . Hence $A(t) \in Q$ for all t .

Finally Eq. (15) is proved as follows. Consider $B(t) := P_0^t(\hat{x}) \circ A_0 \circ P_t^0(x)$, which is an a.c. curve in $T^*M \otimes T\hat{M}$ (or even in Q if $A_0 \in Q$) along $t \mapsto (x(t), \hat{x}(t))$. Now $B(0) = A_0$ and, for $X_0 \in T|_{x(0)}M$, $X(t) := P_0^t(x)X_0$, we have

$$\begin{aligned} 0 &= \hat{\nabla}_{\dot{\hat{x}}(t)}(P_0^t(\hat{x})(A_0X_0)) = \hat{\nabla}_{\dot{\hat{x}}(t)}(B(t)X(t)) \\ &= (\bar{\nabla}_{(\dot{x}, \dot{\hat{x}})(t)}B(t))X(t) + B(t)\nabla_{\dot{x}(t)}X(t) = (\bar{\nabla}_{(\dot{x}, \dot{\hat{x}})(t)}B(t))X(t), \end{aligned}$$

from which it follows, since X_0 was arbitrary, that $\overline{\nabla}_{(\hat{x}, \hat{x})(t)} B(t) = 0$ for a.e. $t \in I$. Thus $t \mapsto A(t)$ and $t \mapsto B(t)$ solve the same initial value problem and hence (being a.c.) are equal $A(t) = B(t)$ i.e.,

$$P_0^t(x, \hat{x})A_0 = P_0^t(\hat{x}) \circ A_0 \circ P_t^0(x), \quad \forall t \in I,$$

which is what we wished to prove. \square

Let $T(M \times \hat{M}) \times_{M \times \hat{M}} (T^*(M) \otimes T(\hat{M}))$ be the total space of the product vector bundle $\pi_{T(M \times \hat{M})} \times_{M \times \hat{M}} \pi_{T^*(M) \otimes T(\hat{M})}$ over $M \times \hat{M}$. We will define certain *lift* operations corresponding to parallel translation of elements of $T^*M \otimes T\hat{M}$.

Definition 4.2 The *No-Spinning lift* is defined to be the map

$$\mathcal{L}_{\text{NS}} : T(M \times \hat{M}) \times_{M \times \hat{M}} (T^*(M) \otimes T(\hat{M})) \rightarrow T(T^*(M) \otimes T(\hat{M})),$$

such that, if $q = (x, \hat{x}; A) \in T^*(M) \otimes T(\hat{M})$, $X \in T|_x M$, $\hat{X} \in T|_{\hat{x}} \hat{M}$ and $t \mapsto (x(t), \hat{x}(t))$ is a smooth curve on in $M \times \hat{M}$ defined on an open interval $I \ni 0$ s.t. $\dot{x}(0) = X$, $\dot{\hat{x}}(0) = \hat{X}$, then one has

$$\mathcal{L}_{\text{NS}}((X, \hat{X}), q) = \left. \frac{d}{dt} \right|_0 P_0^t(x, \hat{x})A \in T|_q (T^*(M) \otimes T(\hat{M})). \quad (16)$$

The smoothness of the map \mathcal{L}_{NS} can be easily seen by using fiber or local coordinates (see Appendix A). We will usually use a notation $\mathcal{L}_{\text{NS}}(\overline{X})|_q$ for $\mathcal{L}_{\text{NS}}(\overline{X}, q)$ when $\overline{X} \in T|_{(x, \hat{x})}(M \times \hat{M})$ and $q = (x, \hat{x}; A) \in T^*(M) \otimes T(\hat{M})$. In particular, when $\overline{X} \in \text{VF}(M \times \hat{M})$, we get a *lifted vector field* on $T^*(M) \otimes T(\hat{M})$ given by $q \mapsto \mathcal{L}_{\text{NS}}(\overline{X})|_q$. The smoothness of $\mathcal{L}_{\text{NS}}(\overline{X})$ for $\overline{X} \in \text{VF}(M \times \hat{M})$ follows immediately from the smoothness of the map \mathcal{L}_{NS} . Notice that, by Proposition 4.1, the No-Spinning lift map \mathcal{L}_{NS} restricts to

$$\mathcal{L}_{\text{NS}} : T(M \times \hat{M}) \times_{M \times \hat{M}} Q \rightarrow TQ,$$

where $T(M \times \hat{M}) \times_{M \times \hat{M}} Q$ is the total space of the fiber product $\pi_{T(M \times \hat{M})} \times_{M \times \hat{M}} \pi_Q$.

We now define the distribution \mathcal{D}_{NS} on $T^*(M) \otimes T(\hat{M})$ and Q capturing the no-spinning condition (see Eq. (11)).

Definition 4.3 The *No-Spinning (NS) distribution* \mathcal{D}_{NS} on $T^*(M) \otimes T(\hat{M})$ is a $2n$ -dimensional smooth distribution defined pointwise by

$$\mathcal{D}_{\text{NS}}|_{(x, \hat{x}; A)} = \mathcal{L}_{\text{NS}}(T|_{(x, \hat{x})}(M \times \hat{M}))|_{(x, \hat{x}; A)}, \quad (17)$$

with $(x, \hat{x}; A) \in T^*(M) \otimes T(\hat{M})$. Since $\mathcal{D}_{\text{NS}}|_Q \subset T(Q)$ (by Proposition 4.1) this distribution restricts to a $2n$ -dimensional smooth distribution on Q which we also denote by \mathcal{D}_{NS} (instead of $\mathcal{D}_{\text{NS}}|_Q$).

The No-Spinning lift \mathcal{L}_{NS} will also be called \mathcal{D}_{NS} -lift since it maps vectors of $M \times \hat{M}$ to vectors in \mathcal{D}_{NS} .

The distribution \mathcal{D}_{NS} is smooth since $\mathcal{L}_{\text{NS}}(\overline{X})$ is smooth for any smooth vector field $\overline{X} \in \text{VF}(M \times \hat{M})$. Also, the fact that the rank of \mathcal{D}_{NS} exactly is $2n$ follows from the next proposition, which itself follows immediately from Eq. (16).

Proposition 4.4 For every $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$ and $\bar{X} \in T|_{(x, \hat{x})}M \times \hat{M}$, one has

$$(\pi_{T^*M \otimes T\hat{M}})_*(\mathcal{L}_{\text{NS}}(\bar{X})|_q) = \bar{X},$$

and in particular $(\pi_Q)_*(\mathcal{L}_{\text{NS}}(\bar{X})|_q) = \bar{X}$ if $q \in Q$.

Thus $(\pi_{T^*M \otimes T\hat{M}})_*$ (resp. π_Q) maps $\mathcal{D}_{\text{NS}}|_{(x, \hat{x}; A)}$ isomorphically onto $T|_{(x, \hat{x})}(M \times \hat{M})$ for every $(x, \hat{x}; A) \in T^*M \otimes T\hat{M}$ (resp. $(x, \hat{x}; A) \in Q$) and the inverse map of $(\pi_{T^*M \otimes T\hat{M}})_*|_{\mathcal{D}_{\text{NS}}|_q}$ (resp. $(\pi_Q)_*|_{\mathcal{D}_{\text{NS}}|_q}$) is $\bar{X} \mapsto \mathcal{L}_{\text{NS}}(\bar{X})|_q$.

Remark 4.5 It should now be clear that an a.c. map $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ in $T^*M \otimes T\hat{M}$ or Q satisfies (11) if and only if q is tangent a.e. to \mathcal{D}_{NS} i.e., for a.e. t it holds that $\dot{q}(t) \in \mathcal{D}_{\text{NS}}|_{q(t)}$.

The following basic formula for the lift \mathcal{L}_{NS} will be useful.

Theorem 4.6 For $\bar{X} \in T|_{(x, \hat{x})}(M \times \hat{M})$ and $A \in \Gamma(\pi_{T^*M \otimes T\hat{M}})$, we have

$$\mathcal{L}_{\text{NS}}(\bar{X})|_{A|_{(x, \hat{x})}} = A_*(\bar{X}) - \nu(\bar{\nabla}_{\bar{X}}A)|_{A|_{(x, \hat{x})}}, \quad (18)$$

where ν denotes the vertical derivative in the vector bundle $\pi_{T^*M \otimes T\hat{M}}$ and A_* is the map $T(M \times \hat{M}) \rightarrow T(T^*M \otimes T\hat{M})$.

Proof. Choose smooth paths $c : [-1, 1] \rightarrow M$, $\hat{c} : [-1, 1] \rightarrow \hat{M}$ such that $(\dot{c}(0), \dot{\hat{c}}(0)) = \bar{X}$ and take an arbitrary $f \in C^\infty(T^*M \otimes T\hat{M})$. Define $\tilde{A}(t) = P_0^t(c, \hat{c})A|_{(x, \hat{x})}$. Then

$$\mathcal{L}_{\text{NS}}(\bar{X})|_{A|_{(x, \hat{x})}} = \dot{\tilde{A}}(0) = \tilde{A}_*\left(\frac{\partial}{\partial t}\right).$$

Also, it is known that (see e.g. [25], p.29)

$$P_t^0(c, \hat{c})(A|_{(c(t), \hat{c}(t))}) = A|_{(x, \hat{x})} + t\bar{\nabla}_{\bar{X}}A + t^2F(t), \quad (19)$$

with $t \mapsto F(t)$ a C^∞ -function $] - 1, 1[\rightarrow T^*|_xM \otimes T|_{\hat{x}}\hat{M}$. On the other hand, one has

$$\begin{aligned} (A_*(\bar{X}) - \tilde{A}_*\left(\frac{\partial}{\partial t}\right))f &= \lim_{t \rightarrow 0} \frac{f(A|_{(c(t), \hat{c}(t))}) - f(P_0^t(c, \hat{c})A|_{(x, \hat{x})})}{t} \\ &= \lim_{t \rightarrow 0} \frac{f(P_0^t(c, \hat{c})A|_{(x, \hat{x})} + tP_0^t(c, \hat{c})\bar{\nabla}_{\bar{X}}A + t^2P_0^t(c, \hat{c})F(t)) - f(P_0^t(c, \hat{c})A|_{(x, \hat{x})})}{t} \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \int_0^t \frac{d}{ds} f(P_0^t(c, \hat{c})A|_{(x, \hat{x})} + sP_0^t(c, \hat{c})\bar{\nabla}_{\bar{X}}A + s^2P_0^t(c, \hat{c})F(t)) ds \\ &= \left. \frac{d}{ds} \right|_{s=0} f(A|_{(x, \hat{x})} + s\bar{\nabla}_{\bar{X}}A + s^2F(0)) = \nu(\bar{\nabla}_{\bar{X}}A)|_A f. \end{aligned}$$

□

We shall write Eq. (18) from now on with a compressed notation

$$\mathcal{L}_{\text{NS}}(\bar{X})|_A = A_*(\bar{X}) - \nu(\bar{\nabla}_{\bar{X}}A)|_A.$$

Remark 4.7 If $A \in \Gamma(\pi_{T^*(M) \otimes T(\hat{M})})$ and $q := A|_{(x, \hat{x})} \in Q$ (e.g. if $A \in \Gamma(\pi_Q)$), then on the right hand side of (18), both terms are elements of $T|_q(T^*M \otimes T\hat{M})$ but their difference is actually an element of $T|_q Q$.

Also, it is clear that Eq. (18) only indicates the decomposition of the map A_* w.r.t to the direct sum decomposition

$$T(T^*M \otimes T\hat{M}) = \mathcal{D}_{\text{NS}} \oplus_{T^*M \otimes T\hat{M}} V(\pi_{T^*M \otimes T\hat{M}}), \quad (20)$$

when $A \in \Gamma(\pi_{T^*(M) \otimes T(\hat{M})})$ and

$$TQ = \mathcal{D}_{\text{NS}} \oplus_Q V(\pi_Q), \quad (21)$$

when $A \in \Gamma(\pi_Q)$ respectively.

As a trivial corollary of the theorem, one gets the following.

Corollary 4.8 Suppose $t \mapsto (x(t), \hat{x}(t); A(t))$ is an a.c. curve on $T^*M \otimes T\hat{M}$ or Q defined on an open real interval I . Then, for a.e. $t \in I$,

$$\mathcal{L}_{\text{NS}}(\dot{x}(t), \dot{\hat{x}}(t))|_{(x(t), \hat{x}(t); A(t))} = \dot{A}(t) - \nu(\overline{\nabla}_{(\dot{x}(t), \dot{\hat{x}}(t))} A)|_{(x(t), \hat{x}(t); A(t))}.$$

Hence $t \mapsto (x(t), \hat{x}(t); A(t))$ is tangent to \mathcal{D}_{NS} at $t_0 \in I$ if and only if $\overline{\nabla}_{(\dot{x}(t_0), \dot{\hat{x}}(t_0))} A = 0$.

4.2 The Control System Associated to the No-Spinning Problem

We next parameterize the set of all absolutely continuous curves which are tangent to the distribution \mathcal{D}_{NS} as a driftless control affine system.

An a.c. curve $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ in Q describes a rolling motion of M against \hat{M} without spinning if and only if $\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{x}(t), \dot{\hat{x}}(t))|_{q(t)}$ for a.e. t . This can be expressed equivalently by saying that $q(\cdot)$ is a solution of a control affine driftless system

$$(\Sigma)_{\text{NS}} \quad \begin{cases} \dot{x}(t) = u(t), \\ \dot{\hat{x}}(t) = \hat{u}(t), \\ \overline{\nabla}_{(u(t), \hat{u}(t))} A(\cdot) = 0 \end{cases}, \quad \text{for a.e. } t \in [a, b], \quad (22)$$

where the control (u, \hat{u}) belongs to the set $\mathcal{U}([a, b], M) \times \mathcal{U}([a, b], \hat{M})$. The fact that System (22) is driftless and control affine can also be seen from its representation in local coordinates; see (90) in Appendix A.

4.3 The Analysis of the Control System $(\Sigma)_{\text{NS}}$

In this subsection, we investigate the structure of the reachable sets associated to $(\Sigma)_{\text{NS}}$ and relate them to the holonomy groups of the Riemannian manifolds (M, g) and (\hat{M}, \hat{g}) .

4.3.1 Description of the Orbits of $(\Sigma)_{NS}$

We begin this section by recalling some standard definitions and introducing some notation concerning the subsequent subsections. If (N, h) is a Riemannian manifold, then the holonomy group $H^{\nabla^h}|_y$ of it at y is defined by

$$H^{\nabla^h}|_y = \{(P^{\nabla^h})_0^1(\gamma) \mid \gamma \in \Omega_y(N)\},$$

and it is a subgroup $O(T|_y N)$ of all h -orthogonal transformations of $T|_y N$. If N is oriented, then one can easily prove that $H^{\nabla^h}|_y$ is actually a subgroup of $SO(T|_y N)$. If $F = (Y_i)_{i=1}^n$, $n = \dim N$, is an orthonormal frame of N at y we write

$$H^{\nabla^h}|_F = \{\mathcal{M}_{F,F}(A) \mid A \in H^{\nabla^h}|_y\}.$$

This is a subgroup of $SO(n)$, isomorphic (as Lie group) to $H^{\nabla^h}|_y$. Lie algebra of the holonomy group $H^{\nabla^h}|_y$ (resp. $H^{\nabla^h}|_F$) will be denoted by $\mathfrak{h}^{\nabla^h}|_y$ (resp. $\mathfrak{h}^{\nabla^h}|_F$). The Lie algebra $\mathfrak{h}^{\nabla^h}|_y$ is a Lie subalgebra of the Lie algebra $\mathfrak{so}(T|_y N)$ of h -antisymmetric linear maps $T|_y N \rightarrow T|_y N$ while $\mathfrak{h}^{\nabla^h}|_F$ is a Lie subalgebra of $\mathfrak{so}(n)$.

In this setting, we will be using the notations $H|_x = H^{\nabla}|_x$ and $\hat{H}|_{\hat{x}} = H^{\hat{\nabla}}|_{\hat{x}}$ respectively for the holonomy groups of (M, g) and (\hat{M}, \hat{g}) at $x \in M$ and $\hat{x} \in \hat{M}$. If F and \hat{F} are respectively orthonormal frames of M and \hat{M} we use $H|_F$ and $\hat{H}|_{\hat{F}}$ respectively to denote $H^{\nabla}|_F$ and $H^{\hat{\nabla}}|_{\hat{F}}$. The corresponding Lie algebras will be written as $\mathfrak{h}|_x$, $\hat{\mathfrak{h}}|_{\hat{x}}$, $\mathfrak{h}|_F$, $\hat{\mathfrak{h}}|_{\hat{F}}$.

We now describe the structure of the orbit $\mathcal{O}_{\mathcal{D}_{NS}}(A_0)$ of \mathcal{D}_{NS} through a point $(x_0, \hat{x}_0; A_0) \in Q$ as follows.

Theorem 4.9 Let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. Then the part of the orbit $\mathcal{O}_{\mathcal{D}_{NS}}(q_0)$ of \mathcal{D}_{NS} through q_0 that lies in the π_Q -fiber over (x_0, \hat{x}_0) is given by

$$\begin{aligned} \mathcal{O}_{\mathcal{D}_{NS}}(q_0) \cap Q|_{(x_0, \hat{x}_0)} &= \{\hat{h} \circ A_0 \circ h \mid \hat{h} \in \hat{H}|_{\hat{x}_0}, h \in H|_{x_0}^{-1}\} \\ &=: \hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}, \end{aligned} \quad (23)$$

and is an immersed submanifold of the fiber $Q|_{(x_0, \hat{x}_0)} = \pi_Q^{-1}(x_0, \hat{x}_0)$.

Moreover, if F, \hat{F} are orthonormal frames at x_0, \hat{x}_0 , as above, then there is a diffeomorphism (depending on F and \hat{F})

$$\mathcal{O}_{\mathcal{D}_{NS}}(q_0) \cap Q|_{(x_0, \hat{x}_0)} \cong \hat{H}|_{\hat{F}} \mathcal{M}_{F, \hat{F}}(A_0) H|_F^{-1}, \quad (24)$$

where the groups on the right hand side are Lie subgroups of $SO(n)$.

In the previous statement, we have used the following notation. If G is a group and S is a subset of G , then $S^{-1} := \{g^{-1} \mid g \in S\}$. Of course $G^{-1} = G$ but, in Eq. (23), it is somewhat more convenient to leave $H|_{x_0}^{-1}$ and not to replace it by $H|_{x_0}$.

Proof. Notice that $q_1 = (x_0, \hat{x}_0; A_1) \in \mathcal{O}_{\mathcal{D}_{NS}}(q_0) \cap \pi_Q^{-1}(x_0, \hat{x}_0)$ if and only if there is a piecewise C^1 path $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$, $t \in [0, 1]$, with $q(0) = q_0$, $q(1) = q_1$ and tangent to \mathcal{D}_{NS} . This is, on the other hand, equivalent, by the definition of \mathcal{D}_{NS} , to the fact that $A(t) = P_0^t(x, \hat{x})A_0$. It is also clear that $t \mapsto (x(t), \hat{x}(t))$ is a piecewise C^1 loop of $M \times \hat{M}$ based at (x_0, \hat{x}_0) i.e., it belongs to $\Omega_{(x_0, \hat{x}_0)}(M \times \hat{M})$ which can

be identified, in a natural way, with $\Omega_{x_0}(M) \times \Omega_{\hat{x}_0}(\hat{M})$. By these remarks, Eq. (15) and the above definition of the holonomy groups, we get

$$\begin{aligned} \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap \pi_Q^{-1}(x_0, \hat{x}_0) &= \{P_0^1(\bar{x})A_0 \mid \bar{x} \in \Omega_{(x_0, \hat{x}_0)}(M \times \hat{M})\} \\ &= \{P_0^1(x, \hat{x})A_0 \mid x \in \Omega_{x_0}(M), \hat{x} \in \Omega_{\hat{x}_0}(\hat{M})\} \\ &= \{P_0^1(\hat{x}) \circ A_0 \circ P_1^0(x) \mid x \in \Omega_{x_0}(M), \hat{x} \in \Omega_{\hat{x}_0}(\hat{M})\} = \hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}. \end{aligned}$$

We next prove that $\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}$ is an immersed submanifold of $Q|_{(x_0, \hat{x}_0)}$. Let $f : \hat{H}|_{\hat{x}_0} \times H|_{x_0} \rightarrow Q|_{(x_0, \hat{x}_0)}$ be a map given by $f(\hat{h}, h) := \hat{h} \circ A_0 \circ h^{-1}$. The map f is clearly smooth, when we consider $H|_{x_0}$ (resp. $\hat{H}|_{\hat{x}_0}$) as a Lie subgroup of $\text{SO}(T|_{x_0}M)$ (resp. $\text{SO}(T|_{\hat{x}_0}\hat{M})$). Moreover, denote $G = \hat{H}|_{\hat{x}_0} \times H|_{x_0}$ and consider the smooth (left) group actions $\mu : G \times Q|_{(x_0, \hat{x}_0)} \rightarrow Q|_{(x_0, \hat{x}_0)}$ and $m : G \times G \rightarrow G$ of G on $Q|_{(x_0, \hat{x}_0)}$ and itself given by $\mu((\hat{h}, h), A) = \hat{h} \circ A \circ h^{-1}$, $m((\hat{h}, h), (\hat{k}, k)) = (\hat{h}\hat{k}, hk)$. Then we see that

$$\begin{aligned} \mu((\hat{h}, h), f(\hat{k}, k)) &= \hat{h} \circ (\hat{k} \circ A_0 \circ k^{-1}) \circ h^{-1} \\ &= (\hat{h}\hat{k}) \circ A_0 \circ (hk)^{-1} = f(\hat{h}\hat{k}, hk) = f(m((\hat{h}, h), (\hat{k}, k))), \end{aligned}$$

which shows that f is G -equivariant map. Since G acts transitively (by the action m) on itself, it follows that f has constant rank (see [15] Theorem 9.7).

Unfortunately f is not injective but there is an easy solution to this obstacle. Notice that $K := f^{-1}(A_0)$ is a closed subgroup of G , hence G/K (the right coset space) is a smooth manifold and f induces a smooth map $\bar{f} : G/K \rightarrow Q|_{(x_0, \hat{x}_0)}$, which is still G -equivariant, when one uses the (left) G -action \bar{m} on G/K induced by m . Now \bar{f} is injective and constant rank, hence an injective immersion (see [15] Theorem 7.14) into $Q|_{(x_0, \hat{x}_0)}$. But the image of \bar{f} is exactly $\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}$.

Moreover, given orthonormal frames F and \hat{F} , we clearly see that

$$\hat{h} \circ A_0 \circ h \mapsto \mathcal{M}_{\hat{F}, \hat{F}}(\hat{h}) \mathcal{M}_{F, \hat{F}}(A_0) \mathcal{M}_{F, F}(h)$$

gives the desired diffeomorphism

$$\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1} \rightarrow \hat{H}|_{\hat{F}} \mathcal{M}_{F, \hat{F}}(A_0) H|_F^{-1}.$$

□

Corollary 4.10 If M and \hat{M} are simply-connected, then each π_Q -fiber $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap Q|_{(x, \hat{x})}$, with $(x, \hat{x}) \in M \times \hat{M}$, of any orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$, $q_0 = (x_0, \hat{x}_0; A_0)$, is a compact connected embedded smooth submanifold Q . In particular, if a \mathcal{D}_{NS} -orbit is open in Q then it is equal to Q .

Proof. Without loss of generality we may assume that $(x, \hat{x}) = (x_0, \hat{x}_0)$. By Theorem 3.2.8 in [11] (in this relation, see also Appendix 5 in [13]), the simply connectedness assumption implies that $H|_{x_0}$ and $\hat{H}|_{\hat{x}_0}$ are respectively (closed and hence) compact connected Lie-subgroups of $\text{SO}(T|_{x_0}M)$ and $\text{SO}(T|_{\hat{x}_0}\hat{M})$.

Now $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap Q|_{(x_0, \hat{x}_0)}$ is compact (as a subset of Q) and connected since it is a continuous image (by the map f in the proof of Theorem 4.9) of the compact connected set $\hat{H}|_{\hat{x}_0} \times H|_{x_0}$. Finally notice that a compact immersed submanifold is embedded.

The last claim follows from the fact that an open orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ has a open fiber $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap Q|_{(x_0, \hat{x}_0)}$ in $Q|_{(x_0, \hat{x}_0)}$. This fiber is also compact by what we just proved and hence $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap Q|_{(x_0, \hat{x}_0)} = Q|_{(x_0, \hat{x}_0)}$ by connectedness of $Q|_{(x_0, \hat{x}_0)}$. This clearly implies that $Q = \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$. \square

The next corollary gives the infinitesimal version of Theorem 4.9.

Corollary 4.11 Let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. Then

$$\begin{aligned} T|_{q_0} \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap V|_{q_0}(\pi_Q) &= \nu(\{\hat{k} \circ A_0 - A_0 \circ k \mid k \in \mathfrak{h}|_{x_0}, \hat{k} \in \hat{\mathfrak{h}}|_{\hat{x}_0}\})|_{q_0} \\ &=: \nu(\hat{\mathfrak{h}}|_{\hat{x}_0} \circ A_0 - A_0 \circ \mathfrak{h}|_{x_0})|_{q_0}, \end{aligned} \quad (25)$$

where $\mathfrak{h}|_{x_0}, \hat{\mathfrak{h}}|_{\hat{x}_0}$ are the Lie algebras of the holonomy groups $H|_{x_0}, \hat{H}|_{\hat{x}_0}$ of M, \hat{M} .

Proof. As in the previous proof, consider the map

$$\begin{aligned} f : \hat{H}|_{\hat{x}_0} \times H|_{x_0} &\rightarrow \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap \pi_Q^{-1}(x_0, \hat{x}_0), \\ (\hat{h}, h) &\mapsto \hat{h} \circ A_0 \circ h^{-1}, \end{aligned}$$

which is known to be a submersion by the previous considerations. We deduce that

$$f_*(\hat{\mathfrak{h}} \times \mathfrak{h}) = T|_{q_0}(\mathcal{O}_{\mathcal{D}_{\text{NS}}}(A_0) \cap \pi_Q^{-1}(x_0, \hat{x}_0)) = T|_{q_0} \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap V|_{q_0}(\pi_Q).$$

But it is obvious that $f_*|_{\hat{\mathfrak{h}} \times \mathfrak{h}}(\hat{k}, k) = \nu(\hat{k} \circ A_0 - A_0 \circ k)|_{q_0}$, which then proves the claim. \square

Remark 4.12 By the previous corollary and the Ambrose-Singer holonomy theorem (see [11], [13]), we have for $q_0 = (x_0, \hat{x}_0; A_0) \in Q$,

$$\begin{aligned} T|_{q_0} \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap V|_{q_0}(\pi_Q) &= \{P_1^0(\hat{c})\hat{R}|_{\hat{x}}(\hat{X}, \hat{Y})P_0^1(\hat{c})A_0 - A_0P_1^0(c)R|_x(X, Y)P_0^1(c) \mid \\ &\quad x \in M, \hat{x} \in \hat{M}, X, Y \in T|_x M, \hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}, \\ &\quad c \in C_{\text{pw}}^1([0, 1], M), c(0) = x_0, c(1) = x, \\ &\quad \hat{c} \in C_{\text{pw}}^1([0, 1], \hat{M}), \hat{c}(0) = \hat{x}_0, \hat{c}(1) = \hat{x}\}, \end{aligned}$$

where $C_{\text{pw}}^1([0, 1], M)$ (resp. $C_{\text{pw}}^1([0, 1], \hat{M})$) is the set of piecewise continuously differentiable maps $[0, 1] \rightarrow M$ (resp. $[0, 1] \rightarrow \hat{M}$).

Theorem 4.9 shows that, since M, \hat{M} are connected, all the π_Q -fibers of the reachable set $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ are diffeomorphic i.e.,

$$\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap \pi_Q^{-1}(x_0, \hat{x}_0) \cong \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \cap \pi_Q^{-1}(x_1, \hat{x}_1),$$

for every $(x_1, \hat{x}_1) \in M \times \hat{M}$. This follows from the fact that if points $x, y \in M$, then (since M is connected) $H|_x$ and $H|_y$ are isomorphic, the same observation holding in \hat{M} . We will now prove that the reachable set $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ has actually a bundle structure over $M \times \hat{M}$.

Proposition 4.13 For $q_0 = (x_0, \hat{x}_0; A_0) \in Q$, denote $\pi_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)} := \pi_Q|_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)}$. Then $\pi_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)} : \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) \rightarrow M \times \hat{M}$ is a smooth subbundle of π_Q with typical fiber $\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}$ and $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ is a smooth immersed submanifold of Q .

Proof. The surjectivity of $\pi_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)}$ onto $M \times \hat{M}$ follows immediately from the connectivity of M, \hat{M} .

Choose local charts (ϕ, U) and $(\hat{\phi}, \hat{U})$ of M and \hat{M} around x_0, \hat{x}_0 centered at x_0, \hat{x}_0 (i.e., $\phi(x_0) = 0, \hat{\phi}(\hat{x}_0) = 0$) and so that $\phi(U)$ and $\hat{\phi}(\hat{U})$ are convex. Then, define

$$\begin{aligned} \tau_{(\phi, \hat{\phi})} : \pi_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)}^{-1}(U \times \hat{U}) &\rightarrow (U \times \hat{U}) \times (\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}) \\ (x, \hat{x}; A) &\mapsto ((x, \hat{x}), P_1^0(t \mapsto (\phi^{-1}(t\phi(x)), \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))))A), \end{aligned}$$

where we notice that, since $A = P_0^1(c, \hat{c})A_0 = P_0^1(\hat{c}) \circ A_0 \circ P_1^0(c)$ for some piecewise C^1 paths $c : [0, 1] \rightarrow M$ and $\hat{c} : [0, 1] \rightarrow \hat{M}$ with $c(0) = x_0, \hat{c}(0) = \hat{x}_0$, one has

$$\begin{aligned} &P_1^0(t \mapsto (\phi^{-1}(t\phi(x)), \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))))A \\ &= P_0^1(t \mapsto \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))) \circ A \circ P_1^0(t \mapsto \phi^{-1}(t\phi(x))) \\ &= P_0^1(t \mapsto \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))) \circ P_0^1(\hat{c}) \circ A_0 \circ P_1^0(c) \circ P_1^0(t \mapsto \phi^{-1}(t\phi(x))). \end{aligned}$$

The concatenation of the path c and $t \mapsto \phi^{-1}(t\phi(x))$ is a piecewise C^1 loop of M based at x_0 and the concatenation of \hat{c} and $t \mapsto \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))$ is a piecewise C^1 loop based of \hat{M} at \hat{x}_0 . Thus $P_1^0(t \mapsto (\phi^{-1}(t\phi(x)), \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))))A$ is an element of $\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1}$.

It is clear that $\tau_{(\phi, \hat{\phi})}$ is a smooth bijection onto $(U \times \hat{U}) \times (\hat{H}|_{\hat{x}_0} \circ A_0 \circ H|_{x_0}^{-1})$. Its inverse map is given by $\psi_{(\phi, \hat{\phi})}$,

$$\psi_{(\phi, \hat{\phi})}((x, \hat{x}), B) = (x, \hat{x}; P_1^0(t \mapsto (\phi^{-1}(t\phi(x)), \hat{\phi}^{-1}(t\hat{\phi}(\hat{x}))))B),$$

which is clearly smooth into Q with image contained in $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ and hence it is smooth into $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ by the basic properties of an orbit. This shows that $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ is a smooth bundle.

Since the maps $\bar{\tau}_{(\phi, \hat{\phi})}$ defined on $\pi_Q^{-1}(U \times \hat{U})$ by the same formula as $\tau_{(\phi, \hat{\phi})}$ are diffeomorphisms (by an identical argument as above) onto $(U \times \hat{U}) \times \pi_Q^{-1}(x_0, \hat{x}_0)$, we see that $\pi_{\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)}$ is a smooth (immersed) subbundle of π_Q . \square

We may now also prove that $\pi_Q : Q \rightarrow M \times \hat{M}$ cannot be equipped with a principal bundle structure leaving the distribution \mathcal{D}_{NS} invariant except in special cases.

Theorem 4.14 Generically, in dimension $n \geq 3$, π_Q cannot be equipped with a principal bundle structure which leaves \mathcal{D}_{NS} invariant.

More precisely, if $n \geq 3$ and F, \hat{F} are oriented orthonormal frames of M and \hat{M} at x_0 and \hat{x}_0 , respectively, and if $H|_F \subset \text{SO}(n), \hat{H}|_{\hat{F}} \subset \text{SO}(n)$ are the holonomy groups with respect to these frames, then $H|_F \cap \hat{H}|_{\hat{F}} \neq \{\text{id}_{\mathbb{R}^n}\}$ implies that there is no principal bundle structure on π_Q which leaves \mathcal{D}_{NS} invariant.

Especially this holds if M (resp. \hat{M}) has full holonomy $\text{SO}(n)$ and \hat{M} (resp. M) is not flat.

Proof. Suppose $\mu : G \times Q \rightarrow Q$ is a left principal bundle structure for π_Q leaving \mathcal{D}_{NS} invariant. Notice that G is diffeomorphic to the π_Q -fibers i.e., to $\text{SO}(n)$ (but, of course, does not need to be isomorphic to it as a Lie group). The fact that for

all $g \in G$ we have $(\mu_g)_* \mathcal{D}_{\text{NS}} \subset \mathcal{D}_{\text{NS}}$ is clearly equivalent to $(\mu_g)_* \mathcal{L}_{\text{NS}}(X, \hat{X})|_q = \mathcal{L}_{\text{NS}}(X, \hat{X})|_{\mu(g,q)}$ for all $q = (x, \hat{x}; A) \in Q$ and $X \in T|_x M$, $\hat{X} \in T|_{\hat{x}} \hat{M}$. But this means that for all $g \in G$, $(x, \hat{x}; A) \in Q$ and a.c. paths $\gamma, \hat{\gamma}$ starting at x, \hat{x} respectively, we have $\mu(g, q_{\mathcal{D}_{\text{NS}}}(q)(t)) = q_{\mathcal{D}_{\text{NS}}}(\mu(g, q))(t)$ where $q_{\mathcal{D}_{\text{NS}}}(q)$ is the unique solution to $\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))|_{q(t)}$, $q(0) = q$. Since we know that if $q = (x, \hat{x}; A) \in Q$, then $q_{\mathcal{D}_{\text{NS}}}(q)(t) = (\gamma(t), \hat{\gamma}(t); P_0^t(\hat{\gamma}) \circ A \circ P_t^0(\gamma))$ for all t , we get that

$$\mu(g, P_0^t(\hat{\gamma}) \circ A \circ P_t^0(\gamma)) = P_0^t(\hat{\gamma}) \circ \mu(g, A) \circ P_t^0(\gamma).$$

Let F, \hat{F} be chosen as in the statement above. Define $(x_0, \hat{x}_0; A_0) \in Q$ by $A_0 = \sum_i g(X_i, \cdot) \hat{X}_i$ and choose $B \in H|_F \cap \hat{H}|_{\hat{F}}$, $B \neq \text{id}_{\mathbb{R}^n}$. Choose loops $\gamma, \hat{\gamma}$ based at x_0, \hat{x}_0 such that $\mathcal{M}_{F,F}(P_0^1(\gamma)) = B$, $\mathcal{M}_{\hat{F},\hat{F}}(P_0^1(\hat{\gamma})) = B$. Since $\mathcal{M}_{F,\hat{F}}(A_0) = \text{id}_{\mathbb{R}^n}$ by the definition of A_0 , we have

$$\begin{aligned} \mathcal{M}_{F,\hat{F}}(A_0) &= \text{id}_{\mathbb{R}^n} = \text{Bid}_{\mathbb{R}^n} B^{-1} = \mathcal{M}_{\hat{F},\hat{F}}(P_0^1(\hat{\gamma})) \mathcal{M}_{F,\hat{F}}(A_0) \underbrace{\mathcal{M}_{F,F}(P_0^1(\gamma))^{-1}}_{=\mathcal{M}_{F,F}(P_1^0(\gamma))} \\ &= \mathcal{M}_{F,\hat{F}}(P_0^1(\hat{\gamma}) \circ A_0 \circ P_1^0(\gamma)) \end{aligned}$$

i.e.,

$$A_0 = P_0^1(\hat{\gamma}) \circ A_0 \circ P_1^0(\gamma).$$

Applying to this what was done above, we get

$$\mu(g, A_0) = \mu(g, P_0^1(\hat{\gamma}) \circ A_0 \circ P_1^0(\gamma)) = P_0^1(\hat{\gamma}) \circ \mu(g, A_0) \circ P_1^0(\gamma), \quad \forall g \in G$$

i.e.,

$$\mathcal{M}_{F,\hat{F}}(\mu(g, A_0)) = B \mathcal{M}_{F,\hat{F}}(\mu(g, A_0)) B^{-1}, \quad g \in G.$$

But $\mu(G, A_0) = \pi_Q^{-1}(x_0, \hat{x}_0)$ whence $\mathcal{M}_{F,\hat{F}}(\mu(G, A_0)) = \text{SO}(n)$ and thus we have found a $B \in \text{SO}(n)$ which is not the identity $\text{id}_{\mathbb{R}^n}$ such that $C = BCB^{-1}$ for all $C \in \text{SO}(n)$ i.e., B belongs to the center of $\text{SO}(n)$. But in dimension $n \geq 3$ the center of $\text{SO}(n)$ is $\{\text{id}_{\mathbb{R}^n}\}$, contradicting the fact that $B \neq \text{id}_{\mathbb{R}^n}$. This contradiction shows that the existence of a principal bundle structure μ on π_Q that preserves \mathcal{D}_{NS} is impossible in this case. \square

4.3.2 Consequences for Controllability

From the previous characterizations of the reachable set of $(\Sigma)_{\text{NS}}$, we now derive consequences for the controllability of the control system $(\Sigma)_{\text{NS}}$.

We start with the following remark.

Remark 4.15 All the results, except Theorem 4.14, of the previous section can obviously be formulated in verbatim in the space $T^*M \otimes T\hat{M}$ instead of Q (i.e., we may replace Q by $T^*M \otimes T\hat{M}$ everywhere) and the statements hold true in this setting. However, Theorem 4.9 (formulated in $T^*M \otimes T\hat{M}$) then implies each orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ of \mathcal{D}_{NS} in $T^*M \otimes T\hat{M}$, $q_0 = (x_0, \hat{x}_0; A_0) \in T^*M \otimes T\hat{M}$, can have dimension of at most $2n + \dim H|_{x_0} + \dim \hat{H}|_{\hat{x}_0} \leq n^2 + n$. Since the dimension of $T^*M \otimes T\hat{M}$ is $n^2 + 2n$, the orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ has a codimension of at least n . This shows that \mathcal{D}_{NS} (or the related control problem) is never completely controllable in $T^*M \otimes T\hat{M}$.

Theorem 4.9 states that the controllability of \mathcal{D}_{NS} is completely determined by the holonomy groups of M and \hat{M} . The next theorem highlights that fact at the Lie algebraic level.

Theorem 4.16 The control system $(\Sigma)_{\text{NS}}$ is completely controllable if and only if, for every $A \in \text{SO}(n)$, the following holds:

$$\mathfrak{h} + A^{-1}\hat{\mathfrak{h}}A = \mathfrak{so}(n), \quad (26)$$

where \mathfrak{h} and $\hat{\mathfrak{h}}$ are respectively the Lie subalgebras of $\mathfrak{so}(n)$ isomorphic (as Lie algebras) to the holonomy Lie algebras of ∇ and $\hat{\nabla}$.

Proof. Clearly, an orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) = Q$, where $(x_0, \hat{x}_0; A_0) = q_0 \in Q$, is an open subset of Q if and only if $T|_q \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) = T|_q Q$ for some (and hence every) $q \in \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$. Thus the decomposition given by Eq. (21) implies that an orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ is open in Q if and only if $V|_q(\pi_Q) \subset T|_q \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$ for some $q \in \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0)$.

By connectedness of Q , we get that \mathcal{D}_{NS} is controllable i.e., $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q_0) = Q$ for some (and hence every) $(x_0, \hat{x}_0; A_0) = q_0 \in Q$ if and only if every orbit $\mathcal{O}_{\mathcal{D}_{\text{NS}}}(q)$, $(x, \hat{x}; A) = q \in Q$ is open in Q .

From now on, fix $(x_0, \hat{x}_0) \in M \times \hat{M}$. Proposition 4.13 implies that every \mathcal{D}_{NS} orbit intersects every π_Q -fiber. Hence \mathcal{D}_{NS} is controllable if and only if $V|_q(\pi_Q) \subset T|_q \mathcal{O}_{\mathcal{D}_{\text{NS}}}(q)$ for every $q = (x_0, \hat{x}_0; A) \in Q|_{(x_0, \hat{x}_0)}$. By Corollary 4.11, this condition is equivalent to the condition that, for every $q = (x_0, \hat{x}_0; A) \in Q|_{(x_0, \hat{x}_0)}$,

$$\nu(\hat{\mathfrak{h}}|_{\hat{x}_0} \circ A - A \circ \mathfrak{h}|_{x_0}) = V|_q(\pi_Q).$$

Next, by Proposition 3.7, one deduces that, for every $q \in Q$,

$$V|_q(\pi_Q) = \nu(A(\mathfrak{so}(T|_x M)))|_q$$

and thus we conclude that \mathcal{D}_{NS} is controllable if and only if, for all $q = (x_0, \hat{x}_0; A) \in Q|_{(x_0, \hat{x}_0)}$,

$$A^{-1} \circ \hat{\mathfrak{h}}|_{\hat{x}_0} \circ A - \mathfrak{h}|_{x_0} = \mathfrak{so}(T|_x M).$$

Choosing arbitrary orthonormal local frames F and \hat{F} of M and \hat{M} at x_0 and \hat{x}_0 , respectively, we see that the above condition is equivalent to

$$\mathcal{M}_{F, \hat{F}}(A)^{-1} \hat{\mathfrak{h}}|_{\hat{F}} \mathcal{M}_{F, \hat{F}}(A) - \mathfrak{h}|_F = \mathfrak{so}(n), \quad \forall A \in Q|_{(x_0, \hat{x}_0)},$$

where

$$\mathfrak{h}|_F = \{\mathcal{M}_F(k) \mid k \in \mathfrak{h}\}, \quad \hat{\mathfrak{h}}|_{\hat{F}} = \{\mathcal{M}_{\hat{F}}(\hat{k}) \mid \hat{k} \in \hat{\mathfrak{h}}\},$$

are the holonomy Lie algebras as subalgebras of $\mathfrak{so}(n)$ w.r.t. the frames F and \hat{F} respectively.

The proof is finished by noticing that $\{\mathcal{M}_{F, \hat{F}}(A) \mid A \in Q|_{(x_0, \hat{x}_0)}\} = \text{SO}(n)$ and that the orthonormal frames F, \hat{F} were arbitrary chosen. \square

Theorem 4.17 Suppose M, \hat{M} are simply connected. Then $(\Sigma)_{NS}$ is completely controllable if and only if

$$\mathfrak{h} + \hat{\mathfrak{h}} = \mathfrak{so}(n) \quad (27)$$

where $\mathfrak{h}, \hat{\mathfrak{h}}$ are the Lie subalgebra of $\mathfrak{so}(n)$ isomorphic (as Lie algebras) to the holonomy Lie algebras of ∇ and $\hat{\nabla}$ respectively.

Proof. By Theorem 4.16, necessity of the condition is obvious.

Conversely suppose that the condition in Eq. (27) holds. This condition implies that for $(x_0, \hat{x}_0) \in M \times \hat{M}$ there is an $q_0 = (x_0, \hat{x}_0; A_0) \in Q|_{(x_0, \hat{x}_0)}$ such that

$$A_0^{-1} \circ \hat{\mathfrak{h}}|_{\hat{x}_0} \circ A_0 - \mathfrak{h}|_{x_0} = \mathfrak{so}(T|_{x_0}M).$$

By Proposition 3.7 and Corollary 4.11 this means that $T|_{q_0}\mathcal{O}_{\mathcal{D}_{NS}}(q_0) \cap V|_{q_0}(\pi_Q) = V|_{q_0}(\pi_Q)$ and hence $T|_{q_0}\mathcal{O}_{\mathcal{D}_{NS}}(q_0) = T|_{q_0}Q$ by Eq. (21) which implies that $\mathcal{O}_{\mathcal{D}_{NS}}(q_0)$ is open in Q . Corollary 4.10 then implies that $\mathcal{O}_{\mathcal{D}_{NS}}(q_0) = Q$ i.e., $(\Sigma)_{NS}$ is completely controllable. \square

There is a complete classification of holonomy groups of Riemannian manifolds by Cartan (for symmetric spaces, see [9]) and Berger (for non-symmetric spaces, see [11]). Hence the above theorems reduce the question of complete controllability of $(\Sigma)_{NS}$ to an essentially linear algebraic problem.

For instance, in the case where both manifolds are non-symmetric, simply connected and irreducible, we get the following proposition.

Theorem 4.18 Assume that the manifolds M and \hat{M} are complete non-symmetric, simply connected, irreducible and $n \neq 8$. Then, the control system $(\Sigma)_{NS}$ is completely controllable if and only if either H or \hat{H} is equal to $\text{SO}(n)$ (w.r.t some orthonormal frames).

Proof. Suppose first that $H|_F = \text{SO}(n)$. Choose any $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and define $\hat{F} = A_0F$ (which is an orthonormal frame of \hat{M} at \hat{x}_0 since $A_0 \in Q$) and compute, noticing that $\mathcal{M}_{F, \hat{F}}(A_0) = \text{id}_{\mathbb{R}^n}$,

$$\pi_Q^{-1}(x_0, \hat{x}_0) \cap \mathcal{O}_{\mathcal{D}_{NS}}(q_0) \cong \hat{H}|_{\hat{F}}H|_F = H|_{\hat{F}}\text{SO}(n) = \text{SO}(n),$$

where the first diffeomorphism comes from Theorem 4.9. But the π_Q -fibers of Q are diffeomorphic to $\text{SO}(n)$ and hence $\pi_Q^{-1}(x_0, \hat{x}_0) \cap \mathcal{O}_{\mathcal{D}_{NS}}(q_0) = \pi_Q^{-1}(x_0, \hat{x}_0)$. By connectedness of M, \hat{M} it follows that $Q = \mathcal{O}_{\mathcal{D}_{NS}}(q_0)$.

Assume now that both holonomy groups are different from $\text{SO}(n)$. We also remark that if one holonomy group is included in the other one, then complete controllability cannot hold according to Eq. (26). Using Berger's list, see [11], and taking into account that

$$\text{Sp}(m) \subset \text{SU}(2m) \subset \text{U}(2m) \subset \text{SO}(4m)$$

where $n = 4m$, it only remains to study the following case: $n = 4m$ with $m \geq 2$, one group is equal to $\text{U}(2m)$ and the other one to $\text{Sp}(m) \cdot \text{Sp}(1)$. Recall that

$$\dim \underbrace{\left(\text{U}(2m) \left(\text{Sp}(m) \cdot \text{Sp}(1) \right) \right)}_{\text{U}(2m) \cdot \text{Sp}(1)} \leq \dim \text{U}(2m) + \dim \text{Sp}(1) = 4m^2 + 3.$$

On the other hand $\dim \mathrm{SO}(4m) = 8m^2 - 2m$ which is always strictly larger than $4m^2 + 3$ for all $m \geq 2$. □

Remark 4.19 If $n = 8$, one is left with the study of the case where one of the holonomy groups is equal to $\mathrm{Spin}(7)$ and the other one is either equal to $\mathrm{U}(4)$ or to $\mathrm{Sp}(2) \cdot \mathrm{Sp}(1)$.

As a corollary to Theorem 4.9 and Theorem 4.16, we get the following result of non controllability in the case where both manifolds are reducible.

Proposition 4.20 Suppose that both (M, g) and (\hat{M}, \hat{g}) are reducible Riemannian manifolds. Then $(\Sigma)_{NS}$ is not completely controllable.

Proof. We need to show that, under the assumptions, there exists $q_0 = (x_0; \hat{x}_0; A_0) \in Q$ so that the orbit $\mathcal{O}_{\mathcal{D}_{NS}}(q_0)$ is a proper subset of Q .

Fix $x_0 \in M$ and $\hat{x}_0 \in \hat{M}$. Since (M, g) and (\hat{M}, \hat{g}) are reducible, there exist subspaces $V_1, V_2 \subset T|_{x_0}M$ and $\hat{V}_1, \hat{V}_2 \subset T|_{\hat{x}_0}\hat{M}$, with $n_i = \dim(V_i) \geq 1$, $\hat{n}_i = \dim(\hat{V}_i) \geq 1$ and such that $H|_{x_0}(V_i) \subset V_i$, $\hat{H}|_{\hat{x}_0}(\hat{V}_i) \subset \hat{V}_i$, for $i = 1, 2$.

Let $X_1^1, \dots, X_{n_1}^1$ and $X_1^2, \dots, X_{n_2}^2$ be an orthonormal basis of V_1 and an orthonormal basis of V_2 respectively and, similarly, let $\hat{X}_1^1, \dots, \hat{X}_{\hat{n}_1}^1$ and $\hat{X}_1^2, \dots, \hat{X}_{\hat{n}_2}^2$ be an orthonormal basis of \hat{V}_1 and an orthonormal basis of \hat{V}_2 respectively. Here, V_i and \hat{V}_i , $i = 1, 2$, are equipped with the metrics induced by $g|_{x_0}$ and $\hat{g}|_{\hat{x}_0}$ respectively. These vectors form orthonormal frames F and \hat{F} of M and \hat{M} at x_0 and \hat{x}_0 respectively.

It follows from the Ambrose-Singer Holonomy Theorem (cf. [11] Theorem 2.4.3, [13] Theorem 8.1) that the Lie algebras $\mathfrak{h}|_F$ and $\hat{\mathfrak{h}}|_{\hat{F}}$ of $H|_F$ and $\hat{H}|_{\hat{F}}$ respectively split into direct sums of Lie-subalgebras,

$$\begin{aligned}\mathfrak{h}|_F &= \mathfrak{h}_1 \oplus \mathfrak{h}_2 \subset \mathfrak{so}(n_1) \oplus \mathfrak{so}(n_2), \\ \hat{\mathfrak{h}}|_{\hat{F}} &= \hat{\mathfrak{h}}_1 \oplus \hat{\mathfrak{h}}_2 \subset \mathfrak{so}(\hat{n}_1) \oplus \mathfrak{so}(\hat{n}_2).\end{aligned}$$

Without loss of generality, we assume that $\hat{n}_1 \geq n_1$.

Finally, we define the linear map $A_0 : T|_{x_0}M \rightarrow T|_{\hat{x}_0}\hat{M}$ by

$$A_0(X_j^1) = \hat{X}_j^1, \quad j = 1, \dots, n_1, \quad A_0(X_j^2) = \hat{X}_{n_1+j}^1, \quad j = 1, \dots, \hat{n}_1 - n_1,$$

and

$$A_0(X_j^2) = \hat{X}_{j-(\hat{n}_1-n_1)}^2, \quad j = \hat{n}_1 - n_1 + 1, \dots, n_2.$$

Thus, we have $\mathcal{M}_{F, \hat{F}}(A_0) = \mathrm{id}_{\mathbb{R}^n}$ and hence

$$\mathfrak{h}|_F + \mathcal{M}_{F, \hat{F}}(A_0)^{-1} \hat{\mathfrak{h}}|_{\hat{F}} \mathcal{M}_{F, \hat{F}}(A_0) = \mathfrak{h}_1 \oplus \mathfrak{h}_2 + \hat{\mathfrak{h}}_1 \oplus \hat{\mathfrak{h}}_2.$$

The latter linear vector space is necessarily a proper subset of $\mathfrak{so}(n)$. In fact, if E_{ij} is the $n \times n$ -matrix with 1 at the i -th row, j -th column and zero otherwise, then the above linear space does not contain $E_{n_1} - E_{1n} \in \mathfrak{so}(n)$. Therefore, the claim follows from Theorem 4.16. □

Corollary 4.21 Suppose that (M, g) and (\hat{M}, \hat{g}) are equal to the Riemannian products $(M_1 \times M_2, g_1 \oplus g_2)$ and $(\hat{M}_1 \times \hat{M}_2, \hat{g}_1 \oplus \hat{g}_2)$, with $\dim M_i \geq 1$, $\dim \hat{M}_i \geq 1$, $i = 1, 2$ respectively. Then, $(\Sigma)_{NS}$ is not controllable on Q .

Proof. From the basic result on holonomy groups, we get the following decomposition $H|_x = H^{\nabla^{g_1}}|_{x_1} \times H^{\nabla^{g_2}}|_{x_2}$, where $x = (x_1, x_2) \in M$, and $\hat{H}|_x = H^{\nabla^{g_1}}|_{\hat{x}_1} \times H^{\nabla^{g_2}}|_{\hat{x}_2}$, where $\hat{x} = (\hat{x}_1, \hat{x}_2) \in \hat{M}$. This shows that the actions of H and \hat{H} on $T|_x M$, $T|_{\hat{x}} \hat{M}$, respectively, are both reducible. Thus, the claim follows from the previous proposition. \square

4.4 Computations of Lie Brackets

In this section, we compute commutators of the vectors fields of $T^*M \otimes T\hat{M}$ and Q with respect to the splitting of $T(T^*M \otimes T\hat{M})$ (resp. TQ) as a direct sum $\mathcal{D}_{\text{NS}} \oplus V(\pi_{T^*M \otimes T\hat{M}})$ (resp. $\mathcal{D}_{\text{NS}} \oplus V(\pi_Q)$) as given in Remark 4.7 above. The main results are Propositions 4.25, 4.25 and 4.28. These computations will serve as preliminaries for the Lie bracket computations relative to the rolling distribution \mathcal{D}_{R} studied in the next section. It is convenient to make the computations in $T^*M \otimes T\hat{M}$ and then to simply restrict the results to Q .

We will now introduce some notations that will be used in the sequel. If $\bar{T} \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T(M \times \hat{M})})$ (resp. $\bar{T} \in C^\infty(\pi_Q, \pi_{T(M \times \hat{M})})$), we will denote by $\mathcal{L}_{\text{NS}}(\bar{T}(\cdot))$ the vector field on $T^*M \otimes T\hat{M}$ (resp. Q) defined by

$$(x, \hat{x}; A) \mapsto \mathcal{L}_{\text{NS}}(\bar{T}(A))|_{(x, \hat{x}; A)}.$$

If $U \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}})$ (resp. $U \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}})$) such that $U(A) \in A(\mathfrak{so}(T|_x M))$ for all $(x, \hat{x}; A) \in Q$, see Proposition 3.7) we will use $\nu(U(\cdot))$ to denote the (vertical) vector field on $T^*M \otimes T\hat{M}$ (resp. Q) defined by

$$(x, \hat{x}; A) \mapsto \nu(U(A))|_{(x, \hat{x}; A)}.$$

Next, for $(x_0, \hat{x}_0; A_0) \in T^*M \otimes T\hat{M}$ and

$$\bar{X} \in T|_{(x_0, \hat{x}_0)}(M \times \hat{M}), \quad U \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}}),$$

(resp. $(x_0, \hat{x}_0; A_0) \in Q$ and $U \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}})$ such that $U(A) \in A(\mathfrak{so}(T|_x M))$ for all $(x, \hat{x}; A) \in Q$), we define $\mathcal{L}_{\text{NS}}(\bar{X})|_{(x_0, \hat{x}_0; A_0)} U \in T^*M \otimes T\hat{M}$ by

$$\mathcal{L}_{\text{NS}}(\bar{X})|_{(x_0, \hat{x}_0; A_0)} U = \bar{\nabla}_{\bar{X}}(U(\tilde{A})) - \nu(\bar{\nabla}_{\bar{X}} \tilde{A})|_{(x_0, \hat{x}_0; A_0)} U, \quad (28)$$

where \tilde{A} is any local smooth $\pi_{T^*M \otimes T\hat{M}}$ -section (resp. π_Q -section) such that $\tilde{A}|_{(x_0, \hat{x}_0)} = A_0$ and we have written $U(\tilde{A})$ for the composition $U \circ \tilde{A}$. We will use this latter compactified notation also in the sequel. It is easy to check that Formula (28) is well defined (i.e., does not depend on the choice of the extension \tilde{A}).

We next have the following simple proposition.

Proposition 4.22 Let $\mathcal{X} \in \text{VF}(T^*M \otimes T\hat{M})$. Then, there are unique smooth bundle maps $\bar{T} \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T(M \times \hat{M})})$, $U \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}})$ such that

$$\mathcal{X}|_{(x, \hat{x}; A)} = \mathcal{L}_{\text{NS}}(\bar{T}(A))|_{(x, \hat{x}; A)} + \nu(U(A))|_{(x, \hat{x}; A)}, \quad (x, \hat{x}; A) \in T^*M \otimes T\hat{M}. \quad (29)$$

Conversely, the smooth bundle maps \bar{T} and U defined previously determine a unique smooth vector field on \mathcal{X} on $T^*M \otimes T\hat{M}$ by the above formula.

The same result also holds on Q by assuming that

$$\mathcal{X} \in \text{VF}(Q), \quad \bar{T} \in C^\infty(\pi_Q, \pi_{T(M \times \hat{M})}), \quad U \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}}),$$

such that $U(A) \in A(\mathfrak{so}(T|_x M))$ for all $(x, \hat{x}; A) \in Q$ (see Proposition 3.7).

Proof. First of all, there are unique smooth vector fields \mathcal{X}^h and \mathcal{X}^v such that

$$\mathcal{X}^h|_{(x,\hat{x};A)} \in \mathcal{D}_{\text{NS}}|_{(x,\hat{x};A)}, \quad \mathcal{X}^v|_{(x,\hat{x};A)} \in V|_{(x,\hat{x};A)}(\pi_{T^*M \otimes T\hat{M}}),$$

for all $(x, \hat{x}; A) \in T^*M \otimes T\hat{M}$ and $\mathcal{X} = \mathcal{X}^h + \mathcal{X}^v$. Then, we define

$$\bar{T}(A) = (\pi_{T^*M \otimes T\hat{M}})_* \mathcal{X}^h|_{(x,\hat{x};A)}, \quad U(A) = \nu|_{(x,\hat{x};A)}^{-1}(\mathcal{X}^v|_{(x,\hat{x};A)}),$$

where $\nu|_{(x,\hat{x};A)}$ is the isomorphism

$$T^*M \otimes T\hat{M}|_{(x,\hat{x};A)} \rightarrow V|_{(x,\hat{x};A)}(\pi_{T^*M \otimes T\hat{M}}); \quad B \mapsto \nu(B)|_{(x,\hat{x};A)}.$$

In the case of Q this last isomorphism is replaced by the isomorphism

$$A(\mathfrak{so}(T|_x M)) \rightarrow V|_{(x,\hat{x};A)}(\pi_Q); \quad B \mapsto \nu(B)|_{(x,\hat{x};A)}.$$

This clearly proves the claims. \square

The next lemma will be useful in the subsequent calculations.

Lemma 4.23 Let $(x, \hat{x}; A) \in T^*M \otimes T\hat{M}$ (resp. $(x, \hat{x}; A) \in Q$). Then there exists a local $\pi_{T^*M \otimes T\hat{M}}$ -section (resp. π_Q -section) \tilde{A} around (x, \hat{x}) such that $\tilde{A}|_{(x,\hat{x})} = A$ and $\bar{\nabla}_{\bar{X}} \tilde{A} = 0$ for all $\bar{X} \in T|_{(x,\hat{x})}(M \times \hat{M})$.

Proof. Let U be an open neighborhood of the origin of $T|_{(x,\hat{x})}(M \times \hat{M})$, where the \bar{g} -exponential map $\bar{\text{exp}} : U \rightarrow M \times \hat{M}$ is a diffeomorphism onto its image. Parallel translate A along geodesics $t \mapsto \bar{\text{exp}}(t\bar{X})$, $\bar{X} \in U$, to get a local section \tilde{A} of $T^*(M) \otimes T(\hat{M})$ in a neighborhood of $\bar{x} = (x, \hat{x})$. More explicitly, one has

$$\tilde{A}|_{\bar{y}} = P_0^1(t \mapsto \bar{\text{exp}}(t(\bar{\text{exp}}_{\bar{x}})^{-1}(\bar{y})))A,$$

for $\bar{y} \in U$. If $(x, \hat{x}; A) \in Q$, this actually provides a local π_Q -section. Moreover, we clearly have $\bar{\nabla}_{\bar{X}} \tilde{A} = 0$ for all $\bar{X} \in T|_{(x,\hat{x})}(M \times \hat{M})$. \square

Notice that the choice of \tilde{A} corresponding to $(x, \hat{x}; A)$ is, of course, not unique.

Lemma 4.24 Let \tilde{A} be a smooth local $\pi_{T^*M \otimes T\hat{M}}$ -section and $\tilde{A}|_{(x,\hat{x})} = A$. Then, for any vector fields $\bar{X}, \bar{Y} \in \text{VF}(M \times \hat{M})$ such that $\bar{X}|_{(x,\hat{x})} = (X, \hat{X})$, $\bar{Y}|_{(x,\hat{x})} = (Y, \hat{Y})$, one has

$$([\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}] \tilde{A})|_{(x,\hat{x})} = -AR(X, Y) + \hat{R}(\hat{X}, \hat{Y})A + (\bar{\nabla}_{[\bar{X}, \bar{Y}]} \tilde{A})|_{(x,\hat{x})}. \quad (30)$$

Here $[\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}]$ is given by $\bar{\nabla}_{\bar{X}} \circ \bar{\nabla}_{\bar{Y}} - \bar{\nabla}_{\bar{Y}} \circ \bar{\nabla}_{\bar{X}}$ and is an \mathbb{R} -linear map on the set of local sections of $\pi_{T^*M \otimes T\hat{M}}$ around (x, \hat{x}) .

Proof. For an arbitrary $Z \in \text{VF}(M)$, which we may interpret as a vector field on $M \times \hat{M}$ as usual, we calculate

$$\begin{aligned}
& ([\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}]\tilde{A})Z = \bar{\nabla}_{\bar{X}}((\bar{\nabla}_{\bar{Y}}\tilde{A})Z) - (\bar{\nabla}_{\bar{Y}}\tilde{A})(\bar{\nabla}_{\bar{X}}Z) - \bar{\nabla}_{\bar{Y}}((\bar{\nabla}_{\bar{X}}\tilde{A})Z) + (\bar{\nabla}_{\bar{X}}\tilde{A})(\bar{\nabla}_{\bar{Y}}Z) \\
&= \bar{\nabla}_{\bar{X}}(\bar{\nabla}_{\bar{Y}}(\tilde{A}Z)) - \tilde{A}\bar{\nabla}_{\bar{Y}}Z - (\bar{\nabla}_{\bar{Y}}\tilde{A})(\bar{\nabla}_{\bar{X}}Z) \\
&\quad - \bar{\nabla}_{\bar{Y}}(\bar{\nabla}_{\bar{X}}(\tilde{A}Z)) - \tilde{A}\bar{\nabla}_{\bar{X}}Z + (\bar{\nabla}_{\bar{X}}\tilde{A})(\bar{\nabla}_{\bar{Y}}Z) \\
&= [\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}](\tilde{A}Z) - (\bar{\nabla}_{\bar{X}}\tilde{A})(\bar{\nabla}_{\bar{Y}}Z) - \tilde{A}\bar{\nabla}_{\bar{X}}(\bar{\nabla}_{\bar{Y}}Z) - (\bar{\nabla}_{\bar{Y}}\tilde{A})(\bar{\nabla}_{\bar{X}}Z) \\
&\quad + (\bar{\nabla}_{\bar{Y}}\tilde{A})(\bar{\nabla}_{\bar{X}}Z) + \tilde{A}\bar{\nabla}_{\bar{Y}}(\bar{\nabla}_{\bar{X}}Z) + (\bar{\nabla}_{\bar{X}}\tilde{A})(\bar{\nabla}_{\bar{Y}}Z) \\
&= [\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}](\tilde{A}Z) + \tilde{A}[\bar{\nabla}_{\bar{Y}}, \bar{\nabla}_{\bar{X}}]Z \\
&= \bar{R}(\bar{X}, \bar{Y})(\tilde{A}Z) + \bar{\nabla}_{[\bar{X}, \bar{Y}]}(\tilde{A}Z) + \tilde{A}\bar{R}(\bar{Y}, \bar{X})Z + \tilde{A}\bar{\nabla}_{[\bar{Y}, \bar{X}]}Z \\
&= -\tilde{A}\bar{R}(\bar{X}, \bar{Y})Z + \bar{R}(\bar{X}, \bar{Y})(\tilde{A}Z) + (\bar{\nabla}_{[\bar{X}, \bar{Y}]} \tilde{A})Z,
\end{aligned}$$

and evaluating the above quantity at (x, \hat{x}) , we get

$$([\bar{\nabla}_{\bar{X}}, \bar{\nabla}_{\bar{Y}}]\tilde{A})Z|_{(x, \hat{x})} = -A(R(X, Y)Z) + \hat{R}(\hat{X}, \hat{Y})(AZ) + (\bar{\nabla}_{[\bar{X}, \bar{Y}]} \tilde{A})Z|_{(x, \hat{x})}.$$

Since the value $Z|_x$ can be chosen arbitrarily in $T|_x M$, the claim follows. \square

Proposition 4.25 Let $\bar{T} = (T, \hat{T}), \bar{S} = (S, \hat{S}) \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T(M \times \hat{M})})$ and $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$. Then

$$\begin{aligned}
[\mathcal{L}_{\text{NS}}(\bar{T}(\cdot)), \mathcal{L}_{\text{NS}}(\bar{S}(\cdot))]|_q &= \mathcal{L}_{\text{NS}}([\bar{T}(\tilde{A}), \bar{S}(\tilde{A})])|_q \\
&\quad + \nu(AR(T(A), S(A)) - \hat{R}(\hat{T}(A), \hat{S}(A))A)|_q, \quad (31)
\end{aligned}$$

where \tilde{A} is chosen to be any local section as in Lemma 4.23 corresponding to $q = (x, \hat{x}; A)$. The same result also holds on Q by assuming that $\bar{T}, \bar{S} \in C^\infty(\pi_Q, \pi_{T(M \times \hat{M})})$.

Proof. Let $f \in C^\infty(T^*M \otimes T\hat{M})$. By using the definition of \mathcal{L}_{NS} and ν , one obtains that

$$\begin{aligned}
& \mathcal{L}_{\text{NS}}(\bar{T}(A))|_q(\mathcal{L}_{\text{NS}}(\bar{S}(\cdot))(f)) \\
&= \bar{T}(A)(\mathcal{L}_{\text{NS}}(\bar{S}(\tilde{A}))|_{\tilde{A}}(f)) - \frac{d}{dt}\Big|_0 \mathcal{L}_{\text{NS}}(\bar{S}(A + t\bar{\nabla}_{\bar{T}(A)}\tilde{A}))|_{A+t\bar{\nabla}_{\bar{T}(A)}\tilde{A}}\tilde{A}(f) \\
&= \bar{T}(A)(\bar{S}(\tilde{A})(f(\tilde{A})) - \frac{d}{dt}\Big|_0 f(\tilde{A} + t\bar{\nabla}_{\bar{S}(\tilde{A})}\tilde{A})) \\
&\quad - \frac{d}{dt}\Big|_0 \bar{S}(A + t\bar{\nabla}_{\bar{T}(A)}\tilde{A})(f(\tilde{A} + t\bar{\nabla}_{\bar{T}(\tilde{A})}\tilde{A})) \\
&\quad + \frac{\partial^2}{\partial t \partial s}\Big|_0 f(A + t\bar{\nabla}_{\bar{T}(A)}\tilde{A} + s\bar{\nabla}_{\bar{S}(A+t\bar{\nabla}_{\bar{T}(A)}\tilde{A})}\tilde{A})(\tilde{A} + t\bar{\nabla}_{\bar{T}(\tilde{A})}\tilde{A}).
\end{aligned}$$

Here, we used the fact that $\bar{\nabla}_{\bar{X}}\tilde{A} = 0$ for all $\bar{X} \in T|_{(x, \hat{x})}(M \times \hat{M})$ and the fact that $\frac{\partial}{\partial t}$ and $\bar{T}(\tilde{A})$ commute (as the obvious vector fields on $M \times \hat{M} \times \mathbb{R}$ with points (x, \hat{x}, t)) to write the last expression in the form

$$\begin{aligned}
& \bar{T}(A)(\bar{S}(\tilde{A})(f(\tilde{A}))) - \frac{d}{dt}\Big|_0 \bar{T}(A)(f(\tilde{A} + t\bar{\nabla}_{\bar{S}(\tilde{A})}\tilde{A})) - \frac{d}{dt}\Big|_0 \bar{S}(A)(f(\tilde{A} + t\bar{\nabla}_{\bar{T}(\tilde{A})}\tilde{A})) \\
&\quad + \frac{\partial^2}{\partial t \partial s}\Big|_0 f(A + st\bar{\nabla}_{\bar{S}(A)}(\bar{\nabla}_{\bar{T}(\tilde{A})}\tilde{A})).
\end{aligned}$$

By interchanging the roles of \overline{T} and \overline{S} and using the definition of commutator of vector fields, we get from this

$$\begin{aligned}
& [\mathcal{L}_{\text{NS}}(\overline{T}(\cdot)), \mathcal{L}_{\text{NS}}(\overline{S}(\cdot))]|_q(f) \\
&= [\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_q(f(\tilde{A})) + \frac{\partial^2}{\partial t \partial s} \Big|_0 f(A + st \overline{\nabla}_{\overline{S}(A)}(\overline{\nabla}_{\overline{T}(\tilde{A})}\tilde{A})) \\
&\quad - \frac{\partial^2}{\partial t \partial s} \Big|_0 f(A + st \overline{\nabla}_{\overline{T}(A)}(\overline{\nabla}_{\overline{S}(\tilde{A})}\tilde{A})) \\
&= [\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_q(f(\tilde{A})) + \frac{d}{dt} \Big|_0 \nu(t \overline{\nabla}_{\overline{S}(A)}(\overline{\nabla}_{\overline{T}(\tilde{A})}\tilde{A}))|_q(f) \\
&\quad - \frac{d}{dt} \Big|_0 \nu(t \overline{\nabla}_{\overline{T}(A)}(\overline{\nabla}_{\overline{S}(\tilde{A})}\tilde{A}))|_q(f) \\
&= [\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_q(f(\tilde{A})) + \nu(\overline{\nabla}_{\overline{S}(A)}(\overline{\nabla}_{\overline{T}(\tilde{A})}\tilde{A}))|_q(f) - \nu(\overline{\nabla}_{\overline{T}(A)}(\overline{\nabla}_{\overline{S}(\tilde{A})}\tilde{A}))|_q(f) \\
&= [\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_q(f(\tilde{A})) - \nu([\overline{\nabla}_{\overline{T}(\tilde{A})}, \overline{\nabla}_{\overline{S}(\tilde{A})}]\tilde{A})|_q(f).
\end{aligned}$$

Finally, using Lemma 4.24, we get that the last line is equal to

$$\begin{aligned}
& [\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_{(x, \hat{x})}(f(\tilde{A})) \\
&\quad - \nu(\overline{\nabla}_{[\overline{T}(\tilde{A}), \overline{S}(\tilde{A})]|_{(x, \hat{x})}}\tilde{A} - AR(T(A), S(A)) + \hat{R}(\hat{T}(A), \hat{S}(A))A)|_q(f),
\end{aligned}$$

from which the claim follows by using the definition of \mathcal{L}_{NS} and the linearity of $\nu(\cdot)|_q$. \square

Notice that the expression $\mathcal{L}_{\text{NS}}([\overline{T}(\tilde{A}), \overline{S}(\tilde{A})])|_q$ in the statement of the proposition only depends on A but not on the choice of the local smooth section \tilde{A} as long as it has the properties: $\tilde{A}|_{(x, \hat{x})} = A$ and $\overline{\nabla}_{\overline{X}}\tilde{A} = 0 \forall \overline{X} \in T|_{(x, \hat{x})}(M \times \hat{M})$ as in Lemma 4.23.

Proposition 4.26 Consider

$$\overline{T} \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T(M \times \hat{M})}), \quad U \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}}),$$

and $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$. Then

$$[\mathcal{L}_{\text{NS}}(\overline{T}(\cdot)), \nu(U(\cdot))]|_q = -\mathcal{L}_{\text{NS}}(\nu(U(A))|_q \overline{T})|_q + \nu(\overline{\nabla}_{\overline{T}(A)}(U(\tilde{A})))|_q, \quad (32)$$

where \tilde{A} is chosen to be any local section as in Lemma 4.23 corresponding to $q = (x, \hat{x}; A)$. The same result also holds on Q by assuming that

$$\overline{T} \in C^\infty(\pi_Q, \pi_{T(M \times \hat{M})}), \quad U \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}}),$$

such that $U(A) \in A(\mathfrak{so}(T|_x M))$ for all $(x, \hat{x}; A) \in Q$.

Proof. Let $f \in C^\infty(T^*M \otimes T\hat{M})$. Then $\mathcal{L}_{\text{NS}}(\overline{T}(A))|_q(\nu(U(\cdot))(f))$ is equal to

$$\overline{T}(A)(\nu(U(\tilde{A}))|_{\tilde{A}}(f)) - \frac{d}{dt} \Big|_0 \nu(U(A + t \overline{\nabla}_{\overline{T}(A)}\tilde{A}))|_{A+t \overline{\nabla}_{\overline{T}(A)}\tilde{A}}(f),$$

which is equal to $\bar{T}(A)(\nu(U(\tilde{A}))|_{\tilde{A}}(f))$ once we recall that $\bar{\nabla}_{\bar{T}(A)}\tilde{A} = 0$. In addition, one has

$$\begin{aligned} \nu(U(A))|_q(\mathcal{L}_{\text{NS}}(\bar{T}(\cdot))(f)) &= \frac{d}{dt}\Big|_0 \mathcal{L}_{\text{NS}}(\bar{T}(A + tU(A))|_{A+tU(A)}(f)) \\ &= \frac{d}{dt}\Big|_0 \bar{T}(A + tU(A))(f(\tilde{A} + tU(\tilde{A}))) \\ &\quad - \frac{\partial^2}{\partial s \partial t}\Big|_0 f(A + tU(A) + s\bar{\nabla}_{\bar{T}(A+tU(A))}(\tilde{A} + tU(\tilde{A}))) \\ &= \frac{d}{dt}\Big|_0 \bar{T}(A + tU(A))(f(\tilde{A} + tU(\tilde{A}))) - \frac{\partial^2}{\partial s \partial t}\Big|_0 f(A + tU(A) + st\bar{\nabla}_{\bar{T}(A+tU(A))}(U(\tilde{A}))), \end{aligned}$$

since $\bar{\nabla}_{\bar{T}(A+tU(A))}\tilde{A} = 0$.

We next simplify the first term on the last line to get

$$\begin{aligned} \frac{d}{dt}\Big|_0 \bar{T}(A + tU(A))(f(\tilde{A} + tU(\tilde{A}))) \\ = D_\nu \bar{T}(A)(U(A))(f(\tilde{A})) + \bar{T}(A)(\nu(U(\tilde{A}))|_{\tilde{A}}(f)), \end{aligned}$$

and then, for the second term, one obtains

$$\begin{aligned} \frac{\partial^2}{\partial s \partial t}\Big|_0 f(A + tU(A) + st\bar{\nabla}_{\bar{T}(A+tU(A))}(U(\tilde{A}))) \\ = \frac{d}{ds}\Big|_0 D_\nu f(A)(U(A) + s\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A}))) = D_\nu f(A)(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A}))). \end{aligned}$$

Therefore one deduces

$$\begin{aligned} [\mathcal{L}_{\text{NS}}(\bar{T}(\cdot)), \nu(U(\cdot))]|_q(f) &= -D_\nu \bar{T}(A)(U(A))(f(\tilde{A})) + D_\nu f(A)(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A}))) \\ &= -\tilde{A}_*(\nu(U(A))|_q \bar{T})(f) + \nu(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A})))|_q(f) \\ &= -\mathcal{L}_{\text{NS}}(\nu(U(A))|_q \bar{T})|_q(f) + \nu(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A})))|_q(f), \end{aligned}$$

where the last line follows from the definition of \mathcal{L}_{NS} and the fact that $\bar{\nabla}_{\nu(U(A))|_q \bar{T}}\tilde{A} = 0$. This completes the proof. \square

Remark 4.27 Notice that the term $\nu(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A})))|_q$ in the statement of the proposition only depends on A but not on the choice of the local smooth section \tilde{A} as long as it has the properties: $\tilde{A}|_{(x,\hat{x})} = A$ and $\bar{\nabla}_{\bar{X}}\tilde{A} = 0 \forall \bar{X} \in T|_{(x,\hat{x})}(M \times \hat{M})$ as in Lemma 4.23. There is an alternative way of seeing this: If $f \in C^\infty(T^*M \otimes T\hat{M})$, then from the definition of \mathcal{L}_{NS} , one has

$$\begin{aligned} \nu(\bar{\nabla}_{\bar{T}(A)}(U(\tilde{A})))|_{U(A)}(f) &= -\mathcal{L}_{\text{NS}}(\bar{T}(A))|_{U(A)}(f) + \bar{T}(A)(f(U(\tilde{A}))) \\ &= -\mathcal{L}_{\text{NS}}(\bar{T}(A))|_{U(A)}(f) + \tilde{A}_*(\bar{T}(A))(f \circ U) \\ &= -\mathcal{L}_{\text{NS}}(\bar{T}(A))|_{U(A)}(f) + \mathcal{L}_{\text{NS}}(\bar{T}(A))|_q(f \circ U), \end{aligned}$$

where the right hand side only depends on A and not on the choice of the local section \tilde{A} as explained above. Notice also that the map $v \mapsto \nu(v)|_u$ is always injective on any vector bundle.

From the above calculation, we actually get the following formula

$$\nu(\overline{\nabla}_{\overline{T}(A)}(U(\tilde{A})))|_{U(A)} = -\mathcal{L}_{\text{NS}}(\overline{T}(A))|_{U(A)} + U_*(\mathcal{L}_{\text{NS}}(\overline{T}(A))|_q). \quad (33)$$

Unfortunately, this is not much use to simplify Formula (32) since there one has to calculate $\nu(\overline{\nabla}_{\overline{T}(A)}(U(\tilde{A})))|_{(x,\hat{x};A)}$ but not $\nu(\overline{\nabla}_{\overline{T}(A)}(U(\tilde{A})))|_{U(A)}$.

By using the notation introduced at the beginning of this section, the result of the previous proposition can be written into the form ($q = (x, \hat{x}; A)$)

$$[\mathcal{L}_{\text{NS}}(\overline{T}(\cdot)), \nu(U(\cdot))]_q = \nu(\mathcal{L}_{\text{NS}}(\overline{T}(A))|_q U)|_q - \mathcal{L}_{\text{NS}}(\nu(U(A))|_q \overline{T})|_q, \quad (34)$$

and this provides us with another argument for seeing the fact explained in the previous remark.

Finally, we derive a formula for the commutators of two vertical vector fields.

Proposition 4.28 Let $U, V \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}})$ and $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$. Then

$$[\nu(U(\cdot)), \nu(V(\cdot))]_q = \nu(\nu(U(A))|_q V - \nu(V(A))|_q U)|_q. \quad (35)$$

The same result also holds on Q by assuming that $U, V \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}})$ such that $U(A), V(A) \in A(\mathfrak{so}(T|_x M))$ for all $(x, \hat{x}; A) \in Q$.

Proof. Let $f \in C^\infty(T^*M \otimes T\hat{M})$. Then,

$$\begin{aligned} \nu(U(A))|_q(\nu(V(\cdot))(f)) &= \frac{d}{dt}\Big|_0 \nu(V(A + tU(A))|_{A+tU(A)}(f)) \\ &= \frac{\partial^2}{\partial t \partial s}\Big|_0 f(A + tU(A) + sV(A + tU(A))) = \frac{d}{ds}\Big|_0 D_\nu(f)(A)(V(A + tU(A))) \\ &= D_\nu(f)(A)(\nu(U(A))|_q V) = \frac{d}{dt}\Big|_0 f(A + t\nu(U(A))|_q V) = \nu(\nu(U(A))|_q V)|_q(f), \end{aligned}$$

from which the result follows. □

5 Study of the Rolling problem (R)

5.1 The Rolling Distribution \mathcal{D}_R

In this section, we investigate the rolling problem as a control system $(\Sigma)_R$ associated to a subdistribution \mathcal{D}_R of \mathcal{D}_{NS} defined as follows.

Recall that the no-spinning distribution \mathcal{D}_{NS} defined on Q models the fact that the admissible curves $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ inscribed on Q , i.e., the curves describing the motion of M against \hat{M} , must verify the no-spinning condition (11). The latter is equivalent to the condition that $t \mapsto q(t)$ is tangent (a.e.) to \mathcal{D}_{NS} , $\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{x}(t), \dot{\hat{x}}(t))|_{q(t)}$ for a.e. t . As regards the rolling of one manifold onto another one, the admissible curve $q(\cdot)$ must also verify the no-slipping condition (13) that we recall next. Since $q(\cdot)$ is tangent to \mathcal{D}_{NS} , we have $A(t) = P_0^t(x, \hat{x})A(0)$, and

hence the no-slipping condition (13) writes $A(t)\dot{x}(t) = \hat{\dot{x}}(t)$. It forces one to have, for a.e. t ,

$$\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{x}(t), A(t)\dot{x}(t))|_{q(t)}.$$

Evaluating at $t = 0$ and noticing that if $q_0 := q(0)$, with $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and $\dot{x}(0) =: X \in T|_{x_0}M$ are arbitrary, we get

$$\dot{q}(0) = \mathcal{L}_{\text{NS}}(X, A_0X)|_{q_0}.$$

This motivates the following definition.

Definition 5.1 For $q = (x, \hat{x}; A) \in Q$, we define the *Rolling lift* or \mathcal{D}_{R} -lift as a bijective linear map

$$\mathcal{L}_{\text{R}} : T|_xM \times Q|_{(x, \hat{x})} \rightarrow T|_qQ,$$

given by

$$\mathcal{L}_{\text{R}}(X, q) = \mathcal{L}_{\text{NS}}(X, AX)|_q. \quad (36)$$

This map naturally induces $\mathcal{L}_{\text{R}} : \text{VF}(M) \rightarrow \text{VF}(Q)$ as follows. For $X \in \text{VF}(M)$ we define $\mathcal{L}_{\text{R}}(X)$, the *Rolling lifted* vector field associated to X , by

$$\begin{aligned} \mathcal{L}_{\text{R}}(X) : Q &\rightarrow T(Q); \\ q &\mapsto \mathcal{L}_{\text{R}}(X)|_q, \end{aligned}$$

where $\mathcal{L}_{\text{R}}(X)|_q := \mathcal{L}_{\text{R}}(X, q)$.

The Rolling lift map \mathcal{L}_{R} allows one to construct a distribution on Q (see [6]) reflecting both of the rolling restrictions of motion defined by the no-spinning condition, Eq. (11), and the no-slipping condition, Eq. (13).

Definition 5.2 The *rolling distribution* \mathcal{D}_{R} on Q is the n -dimensional smooth distribution defined pointwise by

$$\mathcal{D}_{\text{R}}|_{(x, \hat{x}; A)} = \mathcal{L}_{\text{R}}(T|_xM)|_{(x, \hat{x}; A)}, \quad (37)$$

for $(x, \hat{x}; A) \in Q$.

The Rolling lift \mathcal{L}_{R} will also be called \mathcal{D}_{R} -lift since it maps vectors of M to vectors in \mathcal{D}_{R} . Thus an absolutely continuous curve $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ in Q is a rolling curve if and only if it is a.e. tangent to \mathcal{D}_{R} i.e., $\dot{q}(t) \in \mathcal{D}_{\text{R}}|_{q(t)}$ for a.e. t or, equivalently, if $\dot{q}(t) = \mathcal{L}_{\text{R}}(\dot{x}(t))|_{q(t)}$ for a.e. t .

Define $\pi_{Q, M} = \text{pr}_1 \circ \pi_Q : Q \rightarrow M$ and notice that its differential $(\pi_{Q, M})_*$ maps each $\mathcal{D}_{\text{R}}|_{(x, \hat{x}; A)}$, $(x, \hat{x}; A) \in Q$, isomorphically onto $T|_xM$. This implies the following standard result.

Proposition 5.3 For any $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and absolutely continuous $\gamma : [0, a] \rightarrow M$, $a > 0$, such that $c(0) = x_0$, there exists a unique absolutely continuous $q : [0, a'] \rightarrow Q$, $q(t) = (\gamma(t), \hat{\gamma}(t); A(t))$, with $0 < a' \leq a$ (and a' maximal with

the latter property), which is tangent to \mathcal{D}_R a.e. and $q(0) = q_0$. We denote this unique curve q by

$$t \mapsto q_{\mathcal{D}_R}(\gamma, q_0)(t) = (\gamma(t), \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t); A_{\mathcal{D}_R}(\gamma, q_0)(t)),$$

and refer to it as the *rolling curve* with initial conditions (γ, q_0) or *along γ with initial position q_0* . In the case that \hat{M} is a complete manifold one has $a' = a$.

Conversely, any absolutely continuous curve $q : [0, a] \rightarrow Q$, which is a.e. tangent to \mathcal{D}_R , is a rolling curve along $\gamma = \pi_{Q, M} \circ q$ i.e., has the form $q_{\mathcal{D}_R}(\gamma, q(0))$.

Proof. We need to show only that completeness of (\hat{M}, \hat{g}) implies that $a' = a$. In fact, $\hat{X}(t) := A_0 \int_0^t P_s^0(\gamma) \dot{\gamma}(s) ds$ defines an a.c. curve $t \mapsto \hat{X}(t)$ in $T|_{\hat{x}_0} \hat{M}$ defined on $[0, a]$ and the completeness of \hat{M} implies that there is a unique a.c. curve $\hat{\gamma}$ on \hat{M} defined on $[0, a]$ such that $\hat{X}(t) = \int_0^t P_s^0(\hat{\gamma}) \dot{\hat{\gamma}}(s) ds$ for all $t \in [0, a]$ (see also Remark 5.5 below). Defining $A(t) = P_0^t(\hat{\gamma}) \circ A_0 \circ P_t^0(\gamma)$, $t \in [0, a]$ (parallel transports are always defined on the same interval as the a.c. curve along which the parallel transport takes place) we notice that $t \mapsto (\gamma(t), \hat{\gamma}(t); A(t))$ is the rolling curve along γ starting at q_0 that is defined on the interval $[0, a]$. Hence $a' = a$. \square

Of course, it is not important in the previous result that we start the parametrization of the curve γ at $t = 0$.

Remark 5.4 It follows immediately from the uniqueness statement of the previous theorem that, if $\gamma : [a, b] \rightarrow M$ and $\omega : [c, d] \rightarrow M$ are two a.c. curves with $\gamma(b) = \omega(c)$ and $q_0 \in Q$, then

$$q_{\mathcal{D}_R}(\omega \sqcup \gamma, q_0) = q_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q_0)(b)) \sqcup q_{\mathcal{D}_R}(\gamma, q_0). \quad (38)$$

On the group $\Omega_{x_0}(M)$ of piecewise differentiable loops of M based at x_0 one has

$$q_{\mathcal{D}_R}(\omega \cdot \gamma, q_0) = q_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q_0)(1)) \cdot q_{\mathcal{D}_R}(\gamma, q_0),$$

where $\gamma, \omega \in \Omega_{x_0}(M)$.

Specializing to (M, g) and (\hat{M}, \hat{g}) , we will write in the sequel Λ_{x_0} and $\hat{\Lambda}_{\hat{x}_0}$ for $\Lambda_{x_0}^\nabla$ and $\hat{\Lambda}_{\hat{x}_0}^{\hat{\nabla}}$ respectively, where $x_0 \in M$, $\hat{x}_0 \in \hat{M}$.

Remark 5.5 It follows from Proposition 4.1 that, for $q_0 = (x_0, \hat{x}_0; A_0)$ and an a.c. curve γ starting from x_0 , the corresponding rolling curve is given by

$$q_{\mathcal{D}_R}(\gamma, q_0)(t) = (\gamma(t), \hat{\Lambda}_{\hat{x}_0}^{-1}(A_0 \circ \Lambda_{x_0}(\gamma))(t); P_0^t(\hat{\Lambda}_{\hat{x}_0}^{-1}(A_0 \circ \Lambda_{x_0}(\gamma))) \circ A_0 \circ P_t^0(\gamma)). \quad (39)$$

In the case where the curve γ on M is a geodesic, we can give a more precise form of the rolling curve along γ with a given initial position.

Proposition 5.6 Consider $q_0 = (x_0, \hat{x}_0; A_0) \in Q$, $X \in T|_{x_0} M$ and $\gamma : [0, a] \rightarrow M$; $\gamma(t) = \exp_{x_0}(tX)$, a geodesic of (M, g) with $\gamma(0) = x_0$, $\dot{\gamma}(0) = X$. Then the rolling curve $q_{\mathcal{D}_R}(\gamma, q_0) = (\gamma, \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0); A_{\mathcal{D}_R}(\gamma, q_0)) : [0, a'] \rightarrow Q$, $0 < a' \leq a$, along γ with initial position q_0 is given by

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t) = \widehat{\exp}_{\hat{x}_0}(tA_0X), \quad A_{\mathcal{D}_R}(\gamma, q_0)(t) = P_0^t(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \circ A_0 \circ P_t^0(\gamma).$$

Of course, $a' = a$ if \hat{M} is complete.

Proof. Let $0 < a' \leq a$ such that $\hat{\gamma}(t) := \widehat{\exp}_{\hat{x}_0}(tA_0X)$ is defined on $[0, a']$. Then, by proposition 4.1, $q(t) := (\gamma(t), \hat{\gamma}(t); A(t))$ with $A(t) := P_0^t(\hat{\gamma}) \circ A_0 \circ P_t^0(\gamma)$, $t \in [0, a']$, is a curve on Q and $A(t)$ is parallel to $(\gamma, \hat{\gamma})$ in $M \times \hat{M}$. Therefore $t \mapsto q(t)$ is tangent to \mathcal{D}_{NS} on $[0, a']$ and thus $\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))|_{q(t)}$. Moreover, since γ and $\hat{\gamma}$ are geodesics,

$$A(t)\dot{\gamma}(t) = (P_0^t(\hat{\gamma}) \circ A_0)(P_t^0(\gamma)\dot{\gamma}(t)) = P_0^t(\hat{\gamma})(A_0X) = \dot{\hat{\gamma}}(t),$$

which shows that for $t \in [0, a']$,

$$\begin{aligned} \dot{q}(t) &= \mathcal{L}_{\text{NS}}(\dot{\gamma}(t), A(t)\dot{\gamma}(t))|_{q(t)} \\ &= \mathcal{L}_{\text{R}}(\dot{\gamma}(t))|_{q(t)}. \end{aligned}$$

Hence $t \mapsto q(t)$ is tangent to \mathcal{D}_{R} i.e., it is a rolling curve along γ with initial position $q(0) = (\gamma(0), \hat{\gamma}(0); A(0)) = (x_0, \hat{x}_0; A_0) = q_0$. □

Remark 5.7 If $\gamma(t) = \exp_{x_0}(tA_0X)$ and $q_0 = (x_0, \hat{x}_0; A_0)$, the statement of the proposition can be written in a compact form as

$$A_{\mathcal{D}_{\text{R}}}(\gamma, q_0)(t) = P_0^t(s \mapsto \overline{\exp}_{(x_0, \hat{x}_0)}(s(X, A_0X)))A_0,$$

for all t where defined.

The next proposition describes the symmetry of the study of the rolling problem of (M, g) rolling against (\hat{M}, \hat{g}) to the problem of (\hat{M}, \hat{g}) rolling against (M, g) .

Proposition 5.8 Let $\widehat{\mathcal{D}}_{\text{R}}$ be the rolling distribution in $\hat{Q} := Q(\hat{M}, M)$. Then the map

$$\iota : Q \rightarrow \hat{Q}; \quad \iota(x, \hat{x}; A) = (\hat{x}, x; A^{-1})$$

is a diffeomorphism of Q onto \hat{Q} and

$$\iota_*\mathcal{D}_{\text{R}} = \widehat{\mathcal{D}}_{\text{R}}.$$

In particular, $\iota(\mathcal{O}_{\mathcal{D}_{\text{R}}}(q)) = \mathcal{O}_{\widehat{\mathcal{D}}_{\text{R}}}(\iota(q))$.

Proof. It is obvious that ι is a diffeomorphism (with the obvious inverse map) and for an a.c. path $q(t) = (\gamma(t), \hat{\gamma}(t); A(t))$ in Q , $(\iota \circ q)(t) = (\hat{\gamma}(t), \gamma(t); A(t)^{-1})$ is a.c. in \hat{Q} and for a.e. t ,

$$\begin{cases} \dot{\hat{\gamma}}(t) = A(t)\dot{\gamma}(t) \\ A(t) = P_0^t(\hat{\gamma}) \circ A(0) \circ P_t^0(\gamma) \end{cases} \iff \begin{cases} \dot{\gamma}(t) = A(t)^{-1}\dot{\hat{\gamma}}(t) \\ A(t)^{-1} = P_0^t(\gamma) \circ A(0)^{-1} \circ P_t^0(\hat{\gamma}) \end{cases} .$$

These simple remarks prove the claims. □

Remark 5.9 Notice that Definitions 5.1 and 5.2 make sense not only in Q but also in the space $T^*M \otimes T\hat{M}$. It is easily seen that \mathcal{D}_{R} defined on $T^*M \otimes T\hat{M}$ by Eq. (37) is actually tangent to Q so its restriction to Q gives exactly \mathcal{D}_{R} on Q as defined above. Similarly, Propositions 5.3, 5.6 and 5.8 still hold if we replace Q by $T^*M \otimes T\hat{M}$ and \hat{Q} by $T^*\hat{M} \otimes TM$ everywhere in their statements.

5.2 The Bundle Structure of a \mathcal{D}_R -orbit

We begin with the following remark.

Remark 5.10 Notice that the map $\pi_{Q,M} : Q \rightarrow M$ is in fact a bundle. Indeed, let $F = (X_i)_{i=1}^n$ be a local oriented orthonormal frame of M defined on an open set U . Then the local trivialization of $\pi_{Q,M}$ induced by F is

$$\tau_F : \pi_{Q,M}^{-1}(U) \rightarrow U \times F_{\text{OON}}(\hat{M}); \quad \tau_F(x, \hat{x}; A) = (x, (AX_i|_x)_{i=1}^n),$$

is a diffeomorphism.

We also notice that since $\pi_{Q,M}$ -fibers are diffeomorphic to $F_{\text{OON}}(\hat{M})$, in order that there would be a principal G -bundle structure for $\pi_{Q,M}$, it is necessary (but not sufficient) that $F_{\text{OON}}(\hat{M})$ is diffeomorphic to the Lie-group G . In section 6 we consider special cases where there is indeed a principal bundle structure on $\pi_{Q,M}$ which moreover leaves \mathcal{D}_R invariant.

From Proposition 5.6, we deduce that each \mathcal{D}_R -orbit is a smooth bundle over M . This is given in the next proposition (the proof being similar to that of Proposition 4.13).

Proposition 5.11 Let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and suppose that \hat{M} is complete. Then

$$\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M} := \pi_{Q,M}|_{\mathcal{O}_{\mathcal{D}_R}(q_0)} : \mathcal{O}_{\mathcal{D}_R}(q_0) \rightarrow M,$$

is a smooth subbundle of $\pi_{Q,M}$.

Proof. We first show that $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}$ is surjective. If $x \in M$, there is a piecewise smooth path $\gamma : [a, b] \rightarrow M$ from x_0 to x such that each smooth piece is a g -geodesic. By Proposition 5.6 and completeness of \hat{M} it follows that there is a rolling path $q_{\mathcal{D}_R}(\gamma, q_0) : [a, b] \rightarrow Q$ along γ with initial position q_0 defined on the whole interval $[a, b]$. But then $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}(q_{\mathcal{D}_R}(\gamma, q_0)(b)) = x$ which proves the claimed surjectivity.

Since $\mathcal{D}_R|_q \subset T|_q \mathcal{O}_{\mathcal{D}_R}(A_0)$ for every $q \in \mathcal{O}_{\mathcal{D}_R}(q_0)$ and $(\pi_{Q,M})_*$ maps $\mathcal{D}_R|_q$ isomorphically onto $T|_{\pi_{Q,M}(q)} M$, one immediately deduces that $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}$ is also a submersion. This implies that each fiber $(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M})^{-1}(x) = \mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(x)$, $x \in M$, is a smooth closed submanifold of $\mathcal{O}_{\mathcal{D}_R}(q_0)$.

Choose next, for each $x \in M$, an open convex $U_x \subset T|_x M$ such that $\exp_x|_{U_x}$ is a diffeomorphism onto its image and $0 \in U$. Define

$$\begin{aligned} \tau_x : \pi_{Q,M}^{-1}(U_x) &\rightarrow U_x \times \pi_{Q,M}^{-1}(x), \\ q &= (y, \hat{y}; A) \mapsto (y, (x, \hat{\gamma}_{\mathcal{D}_R}(\gamma_{y,x}, q)(1); A_{\mathcal{D}_R}(\gamma_{y,x}, q)(1))), \end{aligned}$$

where $\gamma_{y,x} : [0, 1] \rightarrow M$; $\gamma_{y,x}(t) = \exp_x((1-t)\exp_x^{-1}(y))$ is a geodesic from y to x . It is obvious that τ_x is a smooth bijection. Moreover, restricting τ_x to $\mathcal{O}_{\mathcal{D}_R}(q_0)$ clearly gives a smooth bijection

$$\mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(U_x) \rightarrow U_x \times (\mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(x)).$$

The inverse of τ_x , $\tau_x^{-1} : U_x \times \pi_{Q,M}^{-1}(x) \rightarrow \pi_{Q,M}^{-1}(U_x)$ is constructed with a formula similar to that of τ_x and is seen, in the same way, to be smooth. This inverse restricted to $U_x \times (\mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(x))$ maps bijectively onto $\mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(U_x)$ and thus τ_x is a smooth local trivialization of $\mathcal{O}_{\mathcal{D}_R}(q_0)$. This completes the proof. \square

In the case where \hat{M} is not complete, the result of Proposition 5.11 remains valid if we just claim that $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}$ is a bundle over its image, which is an open connected subset of M . The next remark illustrates this point.

Remark 5.12 In the previous proposition, the assumption of completeness of \hat{M} cannot be removed. In fact, choose $M = \mathbb{R}^2$, $\hat{M} = \{\hat{x} \in \mathbb{R}^2 \mid \|\hat{x}\| < 1\}$ (with $\|\cdot\|$ the Euclidean norm). Then

$$Q \cong M \times \hat{M} \times \text{SO}(2), \quad T(Q) \cong Q \times \mathbb{R}^2 \times \mathbb{R}^2 \times \mathfrak{so}(2)$$

and \mathcal{D}_R is given by

$$\mathcal{D}_R|_{(x, \hat{x}; A)} = \{(v, Av, 0) \mid v \in \mathbb{R}^2\},$$

as a subspace of $T|_{(x, \hat{x}; A)}Q \cong \mathbb{R}^2 \times \mathbb{R}^2 \times \mathfrak{so}(2)$. If $x_0 = 0$, $\hat{x}_0 = 0$ and $A_0 = \text{id}_{\mathbb{R}^2}$ is the identity map $T|_0M \cong \mathbb{R}^2 \rightarrow T|_0\hat{M} \cong \mathbb{R}^2$, we have that the orbit is equal to the 2-dimensional submanifold of Q given by $\{(x, A_0x, A_0) \mid \|x\| < 1\}$ and its image under the projection on the first factor, $\pi_{Q, M}$ is a proper open subset $\{x \in \mathbb{R}^2 \mid \|x\| < 1\}$ of M . Thus $\pi_{Q, M}|_{\mathcal{O}_{\mathcal{D}_R}(x_0, \hat{x}_0; A_0)}$ is not a bundle over M , since this map is not surjective.

Proposition 5.13 For any Riemannian isometries $F \in \text{Iso}(M, g)$ and $\hat{F} \in \text{Iso}(\hat{M}, \hat{g})$ of (M, g) , (\hat{M}, \hat{g}) respectively, one defines smooth free right and left actions of $\text{Iso}(M, g)$, $\text{Iso}(\hat{M}, \hat{g})$ on Q by

$$q_0 \cdot F := (F^{-1}(x_0), \hat{x}_0; A_0 \circ F_*|_{F^{-1}(x_0)}), \quad \hat{F} \cdot q_0 := (x_0, \hat{F}(\hat{x}_0); \hat{F}_*|_{\hat{x}_0} \circ A_0),$$

where $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. We also set

$$\hat{F} \cdot q_0 \cdot F := (\hat{F} \cdot q_0) \cdot F = \hat{F} \cdot (q_0 \cdot F).$$

Then for any $q_0 = (x_0, \hat{x}_0; A_0) \in Q$, a.c. $\gamma : [0, 1] \rightarrow M$, $\gamma(0) = x_0$, and $F \in \text{Iso}(M, g)$, $\hat{F} \in \text{Iso}(\hat{M}, \hat{g})$, one has

$$\hat{F} \cdot q_{\mathcal{D}_R}(\gamma, q_0)(t) \cdot F = q_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F)(t), \quad (40)$$

for all $t \in [0, 1]$. In particular,

$$\hat{F} \cdot \mathcal{O}_{\mathcal{D}_R}(q_0) \cdot F = \mathcal{O}_{\mathcal{D}_R}(\hat{F} \cdot q_0 \cdot F).$$

Proof. The fact that the group actions are well defined is clear and the smoothness of these actions can be proven by writing out the Lie-group structures of the isometry groups (using e.g. Lemma III.6.4 in [25]). If $q_0 \cdot F = q_0 \cdot F'$ for some $F, F' \in \text{Iso}(M, g)$ and $q_0 \in Q$, then $F^{-1}(x_0) = F'^{-1}(x_0)$, $F_*|_{x_0} = F'_*|_{x_0}$ and hence $F = F'$ since M is connected (see [25], p. 43). This proves the freeness of the right $\text{Iso}(M, g)$ -action. The same argument proves the freeness of the left $\text{Iso}(\hat{M}, \hat{g})$ -action.

Finally, Eq. (40) follows from a simple application of Eq. (8). In fact, by Remark 5.5 the rolling curve $q_{\mathcal{D}_R}(\gamma, q_0) = (\gamma, \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0); A_{\mathcal{D}_R}(\gamma, q_0))$ is defined by

$$\begin{aligned} P_t^0(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0))\dot{\hat{\gamma}}_{\mathcal{D}_R}(\gamma, q_0)(t) &= A_0 P_t^0(\gamma)\dot{\gamma}(t), \\ A_{\mathcal{D}_R}(\gamma, q_0)(t) &= P_0^t(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \circ A_0 \circ P_t^0(\gamma). \end{aligned}$$

First, by using (8), we get

$$\begin{aligned}
& P_t^0(\hat{F} \circ \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \frac{d}{dt}(\hat{F} \circ \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0))(t) = \hat{F}_* P_t^0(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \hat{F}_*^{-1}(\hat{F}_* \dot{\hat{\gamma}}_{\mathcal{D}_R}(\gamma, q_0)(t)) \\
& = \hat{F}_* A_0 P_t^0(\gamma) \dot{\gamma}(t) = (\hat{F}_* A_0 F_*)(F_*^{-1} P_t^0(\gamma) F_*) F_*^{-1} \dot{\gamma}(t) \\
& = (\hat{F}_* A_0 F_*) P_t^0(F^{-1} \circ \gamma) \frac{d}{dt}(F^{-1} \circ \gamma)(t),
\end{aligned}$$

and since by definition one has

$$\begin{aligned}
& P_t^0(\hat{\gamma}_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F)) \dot{\hat{\gamma}}_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F) \\
& = (\hat{F}_* A_0 F_*) P_t^0(F^{-1} \circ \gamma) \frac{d}{dt}(F^{-1} \circ \gamma)(t),
\end{aligned}$$

the uniqueness of solutions of a system of ODEs gives that

$$\hat{F} \circ \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) = \hat{\gamma}_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F).$$

Hence

$$\begin{aligned}
& \hat{F}_* A_{\mathcal{D}_R}(\gamma, q_0) F_* = \hat{F}_*(P_0^t(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \circ A_0 \circ P_t^0(\gamma)) F_* \\
& = P_0^t(\hat{F} \circ \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)) \circ (\hat{F}_* A_0 F_*) \circ P_t^0(F^{-1} \circ \gamma) \\
& = P_0^t(\hat{\gamma}_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F)) \circ (\hat{F}_* A_0 F_*) \circ P_t^0(F^{-1} \circ \gamma) = A_{\mathcal{D}_R}(F^{-1} \circ \gamma, \hat{F} \cdot q_0 \cdot F)
\end{aligned}$$

which proves (40). \square

Corollary 5.14 Let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and $\gamma, \omega : [0, 1] \rightarrow M$ be absolutely continuous such that $\gamma(0) = \omega(0) = x_0$, $\gamma(1) = \omega(1)$. Then assuming that $q_{\mathcal{D}_R}(\gamma, q_0)$, $q_{\mathcal{D}_R}(\omega, q_0)$, $q_{\mathcal{D}_R}(\omega^{-1} \cdot \gamma, q_0)$ exist and if there exists $\hat{F} \in \text{Iso}(\hat{M}, \hat{g})$ such that

$$\hat{F} \cdot q_0 = q_{\mathcal{D}_R}(\omega^{-1} \cdot \gamma, q_0)(1),$$

then

$$\hat{F} \cdot q_{\mathcal{D}_R}(\omega, q_0)(1) = q_{\mathcal{D}_R}(\gamma, q_0)(1).$$

Proof.

$$\begin{aligned}
& q_{\mathcal{D}_R}(\gamma, q_0)(1) = q_{\mathcal{D}_R}(\omega \cdot \omega^{-1} \cdot \gamma, q_0)(1) = (q_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\omega^{-1} \cdot \gamma, q_0)(1)) \cdot q_{\mathcal{D}_R}(\omega^{-1} \cdot \gamma, q_0))(1) \\
& = (q_{\mathcal{D}_R}(\omega, \hat{F} \cdot q_0) \cdot q_{\mathcal{D}_R}(\omega^{-1} \cdot \gamma, q_0))(1) = q_{\mathcal{D}_R}(\omega, \hat{F} \cdot q_0)(1) = \hat{F} \cdot q_{\mathcal{D}_R}(\omega, q_0)(1).
\end{aligned}$$

\square

Proposition 5.15 Let $\pi_1 : (M_1, g_1) \rightarrow (M, g)$ and $\hat{\pi} : (\hat{M}_1, \hat{g}_1) \rightarrow (\hat{M}, \hat{g})$ be Riemannian coverings. Write $Q_1 = Q(M_1, \hat{M}_1)$ and $(\mathcal{D}_R)_1$ for the rolling distribution in Q_1 . Then the map

$$\Pi : Q_1 \rightarrow Q; \quad \Pi(x_1, \hat{x}_1; A_1) = (\pi(x_1), \hat{\pi}(\hat{x}_1); \hat{\pi}_*|_{\hat{x}_1} \circ A_1 \circ (\pi_*|_{x_1})^{-1})$$

is a covering map of Q_1 over Q and

$$\Pi_*(\mathcal{D}_R)_1 = \mathcal{D}_R.$$

Moreover, for every $q_1 \in Q_1$ the restriction onto $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)$ of Π is a covering map $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1) \rightarrow \mathcal{O}_{\mathcal{D}_R}(\Pi(q_1))$. Then, for every $q_1 \in Q_1$, $\Pi(\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)) = \mathcal{O}_{\mathcal{D}_R}(\Pi(q_1))$ and one has $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1) = Q_1$ if and only if $\mathcal{O}_{\mathcal{D}_R}(\Pi(q_1)) = Q$.

As an immediate corollary of the above proposition, we obtain the following result regarding the complete controllability of (\mathcal{D}_R) .

Corollary 5.16 Let $\pi_1 : (M_1, g_1) \rightarrow (M, g)$ and $\hat{\pi} : (\hat{M}_1, \hat{g}_1) \rightarrow (\hat{M}, \hat{g})$ be Riemannian coverings. Write $Q = Q(M, \hat{M})$, \mathcal{D}_R and $Q_1 = Q(M_1, \hat{M}_1)$, $(\mathcal{D}_R)_1$ respectively for the state space and for the rolling distribution in the respective state space. Then the control system associated to \mathcal{D}_R is completely controllable if and only if the control system associated to $(\mathcal{D}_R)_1$ is completely controllable. As a consequence, when one addresses the complete controllability issue for the rolling distribution \mathcal{D}_R , one can assume with no loss of generality that both manifolds M and \hat{M} are simply connected.

We now proceed with the proof of Proposition 5.15.

Proof. It is clear that Π is a local diffeomorphism onto Q . To show that it is a covering map, let $q_1 = (x_1, \hat{x}_1; A_1)$ and choose evenly covered w.r.t $\pi, \hat{\pi}$ open sets U and \hat{U} of M, \hat{M} containing $\pi(x_1), \hat{\pi}(\hat{x}_1)$, respectively. Thus $\pi^{-1}(U) = \bigcup_{i \in I} U_i$ and $\hat{\pi}^{-1}(\hat{U}) = \bigcup_{i \in \hat{I}} \hat{U}_i$ where $U_i, i \in I$ (resp. $\hat{U}_i, i \in \hat{I}$) are mutually disjoint connected open subsets of M_1 (resp. \hat{M}_1) such that π (resp. $\hat{\pi}$) maps each U_i (resp. \hat{U}_i) diffeomorphically onto U (resp. \hat{U}). Then

$$\Pi^{-1}(\pi_Q^{-1}(U \times \hat{U})) = \pi_{Q_1}^{-1}((\pi \times \hat{\pi})^{-1}(U \times \hat{U})) = \bigcup_{i \in I, j \in \hat{I}} \pi_{Q_1}^{-1}(U_i \times \hat{U}_j),$$

where $\pi_{Q_1}^{-1}(U_i \times \hat{U}_j)$ for $(i, j) \in I \times \hat{I}$ are clearly mutually disjoint and connected. Now if for a given $(i, j) \in I \times \hat{I}$ we have $(y_1, \hat{y}_1, B_1), (z_1, \hat{z}_1, C_1) \in \pi_{Q_1}^{-1}(U_i \times \hat{U}_j)$ such that $\Pi(y_1, \hat{y}_1; B_1) = \Pi(z_1, \hat{z}_1; C_1)$, then $y_1 = z_1, \hat{y}_1 = \hat{z}_1$ and hence $B_1 = C_1$, which shows that Π restricted to $\pi_{Q_1}^{-1}(U_i \times \hat{U}_j)$ is injective. It is also a local diffeomorphism, as mentioned above, and clearly surjective onto $\pi_Q^{-1}(U \times \hat{U})$, which proves that $\pi_Q^{-1}(U \times \hat{U})$ is evenly covered with respect to Π . This finishes the proof that Π is a covering map.

Suppose next that $q_1(t) = (\gamma_1(t), \hat{\gamma}_1(t); A_1(t))$ is a smooth path on Q_1 tangent to $(\mathcal{D}_R)_1$ and defined on an interval containing $0 \in \mathbb{R}$. Define $q(t) = (\gamma(t), \hat{\gamma}(t); A(t)) := (\Pi \circ q_1)(t)$. Then

$$\begin{aligned} \dot{\hat{\gamma}}(t) &= \hat{\pi}_* \dot{\hat{\gamma}}_1(t) = \hat{\pi}_* A_1(t) \dot{\gamma}_1(t) = A(t) \pi_* \dot{\gamma}_1(t) = A(t) \dot{\gamma}(t) \\ A(t) &= \hat{\pi}_*|_{\hat{\gamma}_1(t)} \circ P_0^t(\hat{\gamma}_1(t)) \circ A_1(0) \circ P_t^0(\gamma_1) \circ (\pi_*|_{\gamma_1(t)})^{-1} \\ &= P_0^t(\hat{\gamma}(t)) \circ \hat{\pi}_*|_{\hat{\gamma}_1(t)} \circ A_1(0) \circ (\pi_*|_{\gamma_1(t)})^{-1} \circ P_t^0(\gamma) \\ &= P_0^t(\hat{\gamma}(t)) \circ A(0) \circ P_t^0(\gamma), \end{aligned}$$

which shows that $q(t)$ is tangent to \mathcal{D}_R . This shows that $\Pi_*(\mathcal{D}_R)_1 \subset \mathcal{D}_R$ and the equality follows from the fact that Π is a local diffeomorphism and the ranks of $(\mathcal{D}_R)_1$ and \mathcal{D}_R are the same i.e., $= n$.

Let $q_1 = (x_1, \hat{x}_1; A_1)$. We proceed to show that the restriction of Π gives a covering $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1) \rightarrow \mathcal{O}_{\mathcal{D}_R}(\Pi(q_1))$. First, since $\Pi_*(\mathcal{D}_R)_1 = \mathcal{D}_R$ and $\Pi : Q_1 \rightarrow Q$ is a covering map, it follows that $\Pi(\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)) = \mathcal{O}_{\mathcal{D}_R}(\Pi(q_1))$.

Let $q := \Pi(q_1)$ and let $U \subset Q$ be an evenly covered neighbourhood of q w.r.t. Π . By the Orbit Theorem, there exists vector fields $Y_1, \dots, Y_d \in \text{VF}(Q)$ tangent to \mathcal{D}_R and $(u_1, \dots, u_d) \in (L^1([0, 1]))^d$ and a connected open neighbourhood W of (u_1, \dots, u_d) in $(L^1([0, 1]))^d$ such that the image of the end point map

$\text{end}_{(Y_1, \dots, Y_d)}(q, W)$ is an open subset of the orbit $\mathcal{O}_{\mathcal{D}_R}(q)$ containing q and included in the Π -evenly covered set U . Let $(Y_i)_1, i = 1, \dots, d$, be the unique vector fields on Q_1 defined by $\Pi_*(Y_i)_1 = Y_i, i = 1, \dots, d$. Since $\Pi_*(\mathcal{D}_R)_1 = \mathcal{D}_R$, it follows that $(Y_i)_1$ are tangent to $(\mathcal{D}_R)_1$ and also, $\Pi \circ \text{end}_{(Y_1)_1, \dots, (Y_d)_1} = \text{end}_{(Y_1, \dots, Y_d)} \circ (\Pi \times \text{id})$. It follows that $\text{end}_{(Y_1)_1, \dots, (Y_d)_1}(q'_1, W)$ is an open subset of $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)$ contained in $\Pi^{-1}(U)$ for every $q'_1 \in (\Pi|_{\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)})^{-1}(q)$.

Since $\text{end}_{(Y_1)_1, \dots, (Y_d)_1}$ is continuous and W is connected, it thus follows that for each $q'_1 \in (\Pi|_{\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)})^{-1}(q)$, the connected set $\text{end}_{(Y_1)_1, \dots, (Y_d)_1}(q'_1, W)$ is contained in a single component of $\Pi^{-1}(U)$ which, since U was evenly covered, is mapped diffeomorphically by Π onto U . But then Π maps $\text{end}_{(Y_1)_1, \dots, (Y_d)_1}(q'_1, W)$ diffeomorphically onto $\text{end}_{(Y_1, \dots, Y_d)}(q, W)$. Since it is also obvious that

$$(\Pi|_{\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)})^{-1}(\text{end}_{(Y_1, \dots, Y_d)}(q, W)) = \bigcup_{q'_1 \in (\Pi|_{\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)})^{-1}(q)} \text{end}_{(Y_1)_1, \dots, (Y_d)_1}(q'_1, W),$$

we have proved that $\text{end}_{(Y_1, \dots, Y_d)}(q, W)$ is an evenly covered neighbourhood of q in $\mathcal{O}_{\mathcal{D}_R}(q)$ w.r.t $\Pi|_{\mathcal{O}_{(\mathcal{D}_R)_1}(q_1)}$.

Finally, let us prove that for every $q_1 \in Q_1$, the following implication holds true,

$$\mathcal{O}_{\mathcal{D}_R}(\Pi(q_1)) = Q \implies \mathcal{O}_{(\mathcal{D}_R)_1}(q_1) = Q_1,$$

(the converse statement being trivial). Indeed, if $\mathcal{O}_{\mathcal{D}_R}(\Pi(q_1)) = Q$, then, for every $q \in Q$, $\mathcal{O}_{\mathcal{D}_R}(q) = Q$ and, on the other hand, the fact that Π restricts to a covering map $\mathcal{O}_{(\mathcal{D}_R)_1}(q'_1) \rightarrow \mathcal{O}_{\mathcal{D}_R}(\Pi(q'_1)) = Q$ for any $q'_1 \in Q_1$ implies that all the orbits $\mathcal{O}_{(\mathcal{D}_R)_1}(q'_1), q'_1 \in Q_1$, are open on Q_1 . But Q_1 is connected (and orbits are non-empty) and hence there cannot be but one orbit. In particular, $\mathcal{O}_{(\mathcal{D}_R)_1}(q_1) = Q_1$. \square

5.3 Computations of some Lie Brackets

We compute some commutators of the vector fields of the form $\mathcal{L}_R(X)$ with $X \in \text{VF}(M)$. The formulas obtained hold both in Q and $T^*M \otimes T\hat{M}$ and thus we do them in the latter space.

The first commutators of the \mathcal{D}_R -lifted fields are given in the following theorem.

Proposition 5.17 If $X, Y \in \text{VF}(M)$, $q = (x_0, \hat{x}_0; A) \in T^*(M) \otimes T(\hat{M})$, then the commutator of the lifts $\mathcal{L}_R(X)$ and $\mathcal{L}_R(Y)$ at q is given by

$$[\mathcal{L}_R(X), \mathcal{L}_R(Y)]|_q = \mathcal{L}_R([X, Y])|_q + \nu(AR(X, Y) - \hat{R}(AX, AY)A)|_q. \quad (41)$$

Proof. Choosing $\bar{T}(B) = (X, BX), \bar{S}(B) = (Y, BY)$ for $B \in T^*(M) \otimes T(\hat{M})$ in proposition 4.25 we have

$$[\mathcal{L}_R(X), \mathcal{L}_R(Y)]|_q = \mathcal{L}_{\text{NS}}([X + \tilde{A}X, Y + \tilde{A}Y])|_q + \nu(AR(X, Y) - \hat{R}(AX, AY)A)|_q,$$

where

$$\begin{aligned} & [(X, \tilde{A}X), (Y, \tilde{A}Y)]|_{(x_0, \hat{x}_0)} = \bar{\nabla}_{(X, \tilde{A}X)}(Y, \tilde{A}Y) - \bar{\nabla}_{(Y, \tilde{A}Y)}(X, \tilde{A}X) \\ & = (\nabla_X Y - \nabla_Y X, \hat{\nabla}_{AX}(\tilde{A}Y) - \hat{\nabla}_{AX}(\tilde{A}Y))|_{(x_0, \hat{x}_0)} + \bar{\nabla}_{(0, AX)}Y - \bar{\nabla}_{(0, AY)}X \\ & \quad + \bar{\nabla}_{(X, 0)}(\tilde{A}Y)|_{(x_0, \hat{x}_0)} - \bar{\nabla}_{(Y, 0)}(\tilde{A}X)|_{(x_0, \hat{x}_0)}, \end{aligned}$$

in which e.g.

$$\begin{aligned}\hat{\nabla}_{AX}(\tilde{A}Y)|_{\hat{x}_0} &= (\bar{\nabla}_{(0,AX)}\tilde{A})|_{(x_0,\hat{x}_0)}\tilde{Y}|_{\hat{x}_0} + \tilde{A}|_{(x_0,\hat{x}_0)}(\nabla_0 Y)|_{x_0} = 0, \\ \bar{\nabla}_{(0,AX)}Y &= 0, \\ \bar{\nabla}_{(X,0)}(\tilde{A}Y)|_{(x_0,\hat{x}_0)} &= (\bar{\nabla}_{(X,0)}\tilde{A})|_{(x_0,\hat{x}_0)}Y|_{x_0} + A\nabla_X Y|_{x_0} = A\nabla_X Y|_{x_0}.\end{aligned}$$

Therefore

$$\begin{aligned}[\mathcal{L}_R(X), \mathcal{L}_R(Y)]|_q &= \mathcal{L}_{NS}\left((\nabla_X Y - \nabla_Y X, 0) + 0 + (0, A\nabla_X Y - A\nabla_Y X)\right)|_q \\ &\quad + \nu(AR(X, Y) - \hat{R}(AX, AY)A)|_q,\end{aligned}$$

which proves the claim after noticing that, by torsion freeness of ∇ , one has $\nabla_X Y - \nabla_Y X = [X, Y]$. \square

Proposition 5.17 justifies the next definition.

Definition 5.18 Given vector fields $X, Y, Z_1, \dots, Z_k \in \text{VF}(M)$, we define the *Rolling Curvature* of the rolling of M against \hat{M} as the smooth mapping

$$\text{Rol}(X, Y) : \pi_{T^*M \otimes T\hat{M}} \rightarrow \pi_{T^*M \otimes T\hat{M}},$$

by

$$\text{Rol}(X, Y)(A) := AR(X, Y) - \hat{R}(AX, AY)A, \quad (42)$$

and similarly, the smooth mapping

$$\bar{\nabla}^k \text{Rol}(X, Y, Z_1, \dots, Z_k) : \pi_{T^*M \otimes T\hat{M}} \rightarrow \pi_{T^*M \otimes T\hat{M}},$$

by

$$\begin{aligned}\bar{\nabla}^k \text{Rol}(X, Y, Z_1, \dots, Z_k)(A) &:= A\nabla^k R(X, Y, (\cdot), Z_1, \dots, Z_k) \\ &\quad - \hat{\nabla}^k \hat{R}(AX, AY, A(\cdot), AZ_1, \dots, AZ_k).\end{aligned} \quad (43)$$

Restricting to Q , we have

$$\text{Rol}(X, Y), \bar{\nabla}^k \text{Rol}(X, Y, Z_1, \dots, Z_k)(A) \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}}),$$

such that, for all $(x, \hat{x}; A) \in Q$,

$$\text{Rol}(X, Y)(A), \bar{\nabla}^k \text{Rol}(X, Y, Z_1, \dots, Z_k)(A) \in A(\mathfrak{so}(T|_x M)).$$

Remark 5.19 With this notation, Eq. (41) of Proposition 5.17 can be written as

$$[\mathcal{L}_R(X), \mathcal{L}_R(Y)]|_q = \mathcal{L}_R([X, Y])|_q + \nu(\text{Rol}(X, Y)(A))|_q.$$

Recall that, on a Riemannian manifold (N, h) , a smooth vector field $t \mapsto Y(t)$ along a smooth curve $t \mapsto \gamma(t)$ is a Jacobi field if Y satisfies the following second order ODE:

$$\nabla_{\dot{\gamma}(t)}^h \nabla_{\dot{\gamma}(t)}^h Y(\cdot) = R^h(\dot{\gamma}(t), Y(t))\dot{\gamma}(t).$$

The next lemma relates the rolling curvature Rol to the Jacobi fields of M and \hat{M} .

Lemma 5.20 Suppose that $q_0 = (x_0, \hat{x}_0; A_0) \in T^*M \otimes T\hat{M}$, $\gamma : [a, b] \rightarrow M$ is a smooth curve with $\gamma(a) = x_0$ and that the rolling problem along γ has a solution $q_{\mathcal{D}_R}(\gamma, q_0) = (\gamma, \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0); A_{\mathcal{D}_R}(\gamma, q_0))$ on the interval $[a, b]$. If $t \mapsto Y(t)$ is a Jacobi field of (M, g) along γ , then $\hat{Y}(t) = A_{\mathcal{D}_R}(\gamma, q_0)(t)Y(t)$ is a vector field along $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)$ and, for all $t \in [a, b]$,

$$\begin{aligned} \hat{\nabla}_{\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t)} \hat{\nabla}_{\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(\cdot)} \hat{Y}(\cdot) &= \hat{R}(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t), \hat{Y}(t)) \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t) \\ &\quad + \text{Rol}(\dot{\gamma}(t), Y(t))(A_{\mathcal{D}_R}(\gamma, q_0)(t)) \dot{\gamma}(t). \end{aligned}$$

Proof. Since $\bar{\nabla}_{(\dot{\gamma}(t), \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t))} A_{\mathcal{D}_R}(\gamma, q_0)(\cdot) = 0$ and Y is a Jacobi field, one has

$$\begin{aligned} &\hat{\nabla}_{\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t)} \hat{\nabla}_{\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(\cdot)} \hat{Y}(\cdot) \\ &= A_{\mathcal{D}_R}(\gamma, q_0)(t) \nabla_{\dot{\gamma}(t)} \nabla_{\dot{\gamma}(\cdot)} Y(\cdot) = A_{\mathcal{D}_R}(\gamma, q_0)(t) R(\dot{\gamma}(t), Y(t)) \dot{\gamma}(t) \\ &= \text{Rol}(\dot{\gamma}(t), Y(t))(A_{\mathcal{D}_R}(\gamma, q_0)(t)) \dot{\gamma}(t) \\ &\quad + \hat{R}(A_{\mathcal{D}_R}(\gamma, q_0)(t) \dot{\gamma}(t), A_{\mathcal{D}_R}(\gamma, q_0)(t) Y(t)) A_{\mathcal{D}_R}(\gamma, q_0)(t) \dot{\gamma}(t) \end{aligned}$$

from which the claim follows by using the facts that

$$A_{\mathcal{D}_R}(\gamma, q_0)(t) \dot{\gamma}(t) = \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(t) \text{ and } A_{\mathcal{D}_R}(\gamma, q_0)(t) Y(t) = \hat{Y}(t).$$

□

We will use Lemma 5.20 to prove Theorem 5.44.

Remark 5.21 Notice that if, in Lemma 5.20, it held that

$$\text{Rol}(Y(t), \dot{\gamma}(t))(A_{\mathcal{D}_R}(\gamma, q_0)(t)) \dot{\gamma}(t) = 0,$$

for all $t \in [a, b]$, then \hat{Y} defined there would be a Jacobi field along $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)$. Hence, Rol measures the obstruction for $\hat{Y} = A_{\mathcal{D}_R}(\gamma, q_0)(t)Y(t)$ to be a Jacobi field of \hat{M} , if $Y(t)$ is a Jacobi field on M along γ .

Before proceeding with the computations of higher order brackets of the vector fields $\mathcal{L}_R(X)$, we prove the following lemma.

Lemma 5.22 Let $\tilde{A} \in \Gamma(\pi_{T^*M \otimes T\hat{M}})$ and $(x, \hat{x}; A) \in T^*M \otimes T\hat{M}$ such that $\tilde{A}|_{(x, \hat{x})} = A$ and $\bar{\nabla}_{\tilde{X}} \tilde{A} = 0$ for all $\tilde{X} \in T|_{(x, \hat{x})}(M \times \hat{M})$. Then, for $X_1, \dots, X_{k+2}, Y \in \text{VF}(M)$,

$$\bar{\nabla}_{(Y, AY)} (\bar{\nabla}^k \text{Rol}(X_1, X_2, X_3, \dots, X_{k+2}))(\tilde{A}) \quad (44)$$

$$= \bar{\nabla}^{k+1} \text{Rol}(X_1, \dots, X_{k+2}, Y)(A) + \sum_{i=1}^{k+2} \bar{\nabla}^k \text{Rol}(X_1, \dots, \nabla_Y X_i, \dots, X_{k+2})(A) \quad (45)$$

Proof. If $k = 0$, we have $\bar{\nabla}^0 \text{Rol}(X_1, X_2)(\tilde{A}) = \text{Rol}(X_1, X_2)(\tilde{A})$ and since $\bar{\nabla}_{(Y, AY)} \tilde{A} = 0$, one gets

$$\begin{aligned} \bar{\nabla}_{(Y, AY)} \text{Rol}(X_1, X_2)(\tilde{A}) &= \bar{\nabla}_{(Y, AY)} (\tilde{A} R(X_1, X_2) - \hat{R}(\tilde{A} X_1, \tilde{A} X_2) \tilde{A}) \\ &= A \nabla_Y (R(X_1, X_2)) - (\bar{\nabla}_{(Y, AY)} \hat{R}(\tilde{A} X_1, \tilde{A} X_2)) \tilde{A} \\ &= A \nabla R(X_1, X_2, (\cdot), Y) + AR(\nabla_Y X_1, X_2) + AR(X_1, \nabla_Y X_2) \\ &\quad - \hat{\nabla} \hat{R}(AX_1, AX_2, A(\cdot), AY) - \hat{R}(A \nabla_Y X_1, AX_2) A - \hat{R}(AX_1, A \nabla_Y X_2), \end{aligned}$$

where on the last line we have computed $\bar{\nabla}_{(Y, AY)} (\tilde{A} X_i) = A \nabla_Y X_i$. The case $k > 0$ is proved by induction and similar computations. □

Proposition 5.23 Let $X, Y, Z \in \text{VF}(M)$. Then, for $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$, one has

$$[\mathcal{L}_R(Z), \nu(\text{Rol}(X, Y)(\cdot))] \Big|_q = -\mathcal{L}_{\text{NS}}(\text{Rol}(X, Y)(A)Z) \Big|_q + \nu(\bar{\nabla}^1 \text{Rol}(X, Y, Z)(A)) \Big|_q \\ + \nu(\text{Rol}(\nabla_Z X, Y)(A)) \Big|_q + \nu(\text{Rol}(X, \nabla_Z Y)(A)) \Big|_q.$$

Proof. Taking $\bar{T}(B) = (Z, BZ)$ and $U = \text{Rol}(X, Y)$ for $B \in T^*M \otimes T\hat{M}$ in Proposition 4.26, we get

$$[\mathcal{L}_R(Z), \nu(\text{Rol}(X, Y)(\cdot))] \Big|_q \\ = -\mathcal{L}_{\text{NS}}(\nu(\text{Rol}(X, Y)(A)) \Big|_q (Z + (\cdot)Z)) \Big|_q + \nu(\bar{\nabla}_{Z+AZ}(\text{Rol}(X, Y)(\tilde{A}))) \Big|_q.$$

From here, one easily computes that

$$\nu(\text{Rol}(X, Y)(A)) \Big|_q (Z + (\cdot)Z) = \frac{d}{dt} \Big|_0 (Z + (A + t\text{Rol}(X, Y)(A))Z) = \text{Rol}(X, Y)(A)Z,$$

and by Lemma 5.22, one gets

$$\bar{\nabla}_{Z+AZ}(\text{Rol}(X, Y)(\tilde{A})) = \bar{\nabla}^1 \text{Rol}(X, Y, Z)(A) + \text{Rol}(\nabla_Z X, Y)(A) + \text{Rol}(X, \nabla_Z Y)(A).$$

□

By Proposition 5.17, the last two terms (when considered as vector fields on $T^*M \otimes T\hat{M}$) on the right hand side belong to $\text{VF}_{\mathcal{D}_R}^2$.

Since for $X, Y \in \text{VF}(M)$ and $q = (x, \hat{x}; A) \in \hat{Q}$ we have $\nu(\text{Rol}(X, Y)(A)) \Big|_q \in \mathcal{O}_{\mathcal{D}_R}(q)$ by Proposition 5.17, it is reasonable to compute the Lie-bracket of two elements of this type. This is given in the following proposition.

Proposition 5.24 For any $q = (x, \hat{x}; A) \in \hat{Q}$ and $X, Y, Z, W \in \text{VF}(M)$ we have

$$[\nu(\text{Rol}(X, Y)(\cdot)), \nu(\text{Rol}(Z, W)(\cdot))] \Big|_q \\ = \nu(\text{Rol}(X, Y)(A)R(Z, W) - \hat{R}(\text{Rol}(X, Y)(A)Z, AW)A - \hat{R}(AZ, \text{Rol}(X, Y)(A)W)A \\ - \hat{R}(AZ, AW)\text{Rol}(X, Y)(A) - \text{Rol}(Z, W)(A)R(X, Y) + \hat{R}(\text{Rol}(Z, W)(A)X, AY)A \\ + \hat{R}(AX, \text{Rol}(Z, W)(A)Y)A + \hat{R}(AX, AY)\text{Rol}(Z, W)(A)) \Big|_q.$$

Proof. We use Proposition 4.28 where for U, V we take $U(A) = \text{Rol}(X, Y)(A)$ and $V(A) = \text{Rol}(Z, W)(A)$. First compute for B such that $\nu(B) \Big|_q \in V \Big|_q(\hat{Q})$ that

$$\nu(B) \Big|_q U = \nu(B) \Big|_q (\tilde{A} \mapsto \tilde{A}R(X, Y) - \hat{R}(\tilde{A}X, \tilde{A}Y)\tilde{A}) \\ = \frac{d}{dt} \Big|_0 ((A + tB)R(X, Y) - \hat{R}((A + tB)X, (A + tB)Y)(A + tB)) \\ = BR(X, Y) - \hat{R}(BX, AY)A - \hat{R}(AX, BY)A - \hat{R}(AX, AY)B$$

So by taking $B = V(A)$ we get

$$\nu(V(A)) \Big|_q U = \text{Rol}(Z, W)(A)R(X, Y) - \hat{R}(\text{Rol}(Z, W)(A)X, AY)A \\ - \hat{R}(AX, \text{Rol}(Z, W)(A)Y)A - \hat{R}(AX, AY)\text{Rol}(Z, W)(A)$$

and similarly for $\nu(U(A)) \Big|_q V$.

□

From Proposition 4.26 we get the following proposition.

Proposition 5.25 Let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. Suppose that, for some $X \in \text{VF}(M)$ and a real sequence $(t_n)_{n=1}^\infty$ s.t. $t_n \neq 0$ for all n , $\lim_{n \rightarrow \infty} t_n = 0$, we have

$$V|_{\Phi_{\mathcal{L}_R(X)}(t_n, q_0)}(\pi_Q) \subset T(\mathcal{O}_{\mathcal{D}_R}(q_0)), \quad \forall n. \quad (46)$$

Then $\mathcal{L}_{\text{NS}}(Y, \hat{Y})|_{q_0} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ for every $Y \in T|_{x_0} M$ that is g -orthogonal to $X|_{x_0}$ and every $\hat{Y} \in T|_{\hat{x}_0} \hat{M}$ that is \hat{g} -orthogonal to $A_0 X|_{x_0}$. Hence the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ has codimension at most 1 inside Q .

Proof. Letting n tend to infinity, it follows from (46) that $V|_{q_0}(\pi_Q) \subset T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$. Recall, from Proposition 3.7, that every element of $V|_{q_0}(\pi_Q)$ is of the form $\nu(B)|_{q_0}$, with a unique $B \in Q|_{(x_0, \hat{x}_0)}$ satisfying $A_0^T B \in \mathfrak{so}(T|_{x_0} M)$. Fix such a B and define a smooth local section \tilde{S} of $\mathfrak{so}(TM) \rightarrow M$ defined on an open set $W \ni x_0$ by

$$\tilde{S}|_x = P_0^1(t \mapsto \exp_{x_0}(t \exp_{x_0}^{-1}(x)))(A_0^T B).$$

Then clearly, $\tilde{S}|_{x_0} = A_0^T B$ and $\nabla_Y \tilde{S} = 0$ for all $Y \in T|_{x_0} M$ and it is easy to verify that $\tilde{S}|_x \in \mathfrak{so}(T|_x M)$ for all $x \in W$.

We next define a smooth map $U : \pi_Q^{-1}(W \times \hat{M}) \rightarrow T^*M \otimes T\hat{M}$ by $U(x, \hat{x}; A) = A\tilde{S}|_x$. Obviously $\nu(U(x, \hat{x}; A)) \in V|_{(x, \hat{x}; A)}(\pi_Q)$ for all $(x, \hat{x}; A)$. Then, choosing in Proposition 4.26, $\bar{T} = X + (\cdot)X$ (and the above U) and noticing that

$$\nu(U(A_0))|_{q_0} \bar{T} = U(A_0)X = BX,$$

one gets

$$[\mathcal{L}_R(X), \nu(U(\cdot))]|_{q_0} = -\mathcal{L}_{\text{NS}}(BX)|_{q_0} + \nu(\bar{\nabla}_{(X, A_0 X)}(U(\tilde{A})))|_{q_0} \quad (47)$$

where $\tilde{A}|_{(x_0, \hat{x}_0)} = A_0$. By the choice of \tilde{S} and \tilde{A} , we have, for all $\bar{Y} = (Y, \hat{Y}) \in T|_{(x_0, \hat{x}_0)} M \times \hat{M}$,

$$\nabla_{\bar{Y}}(U(\tilde{A})) = \nabla_{\bar{Y}}(\tilde{A}\tilde{S}) = (\nabla_{\bar{Y}}\tilde{A})\tilde{S}|_{(x_0, \hat{x}_0)} + \tilde{A}|_{(x_0, \hat{x}_0)} \nabla_Y \tilde{S} = 0,$$

and hence the last term on the right hand side of (47) actually vanishes.

By definition, the vector field $q \mapsto \mathcal{L}_R(X)|_q$ is tangent to the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ and, by the assumption of Equation (46), the values of the map $q = (x, \hat{x}; A) \mapsto \nu(U(A))|_q$ are also tangent to $\mathcal{O}_{\mathcal{D}_R}(q_0)$ at the points $\Phi_{\mathcal{L}_R(X)}(t_n, q_0)$, $n \in \mathbb{N}$. Hence $((\Phi_{\mathcal{L}_R(X)})_{-t_n})_* \nu(U(\cdot))|_{\Phi_{\mathcal{L}_R(X)}(t_n, q_0)} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ and therefore,

$$\begin{aligned} & [\mathcal{L}_R(X), \nu(U(\cdot))]|_{q_0} \\ &= \lim_{n \rightarrow \infty} \frac{((\Phi_{\mathcal{L}_R(X)})_{-t_n})_* \nu(U(\cdot))|_{\Phi_{\mathcal{L}_R(X)}(t_n, q_0)} - \nu(B)|_{q_0}}{t_n} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0), \end{aligned}$$

i.e., the left hand side of (47) must belong to $T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$. But this implies that

$$\mathcal{L}_{\text{NS}}(BX)|_{q_0} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0), \quad \forall B \text{ s.t. } \nu(B) \in V|_{q_0}(\pi_Q)$$

i.e.,

$$\mathcal{L}_{\text{NS}}(A_0 \mathfrak{so}(T|_{x_0} M) X)|_{q_0} \subset T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0).$$

Notice next that $\mathfrak{so}(T|_{x_0} M) X$ is exactly the set $X|_{x_0}^\perp$ of vectors of $T|_{x_0} M$ that are g -perpendicular to $X|_{x_0}$. Since $A_0 \in Q$, it follows that the set $A_0 \mathfrak{so}(T|_{x_0} M) X$ is equal to $A_0 X|_{x_0}^\perp$ which is the set of vectors of $T|_{\hat{x}_0} \hat{M}$ that are \hat{g} -perpendicular to $A_0 X|_{x_0}$. We conclude that $\mathcal{L}_{\text{NS}}(Y)|_{q_0} = \mathcal{L}_R(Y)|_{q_0} - \mathcal{L}_{\text{NS}}(A_0 Y)|_{q_0} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ for all $Y \in X|_{x_0}^\perp$.

Finally notice that since the subspaces $X^\perp \times \{0\}$, $\mathbb{R}(X, A_0 X)$ and $\{0\} \times (A_0 X)^\perp$ of $T|_{(x_0, \hat{x}_0)}(M \times \hat{M})$ are linearly independent, their \mathcal{L}_{NS} -lifts at q_0 are that also and hence these lifts span a $(n-1) + 1 + (n-1) = 2n-1$ dimensional subspace of $T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$. This combined with the fact that $V|_{q_0}(\pi_Q) \subset T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ shows $\dim \mathcal{O}_{\mathcal{D}_R}(q_0) \geq 2n-1 + \dim V|_{q_0}(\pi_Q) = \dim(Q) - 1$ i.e., the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ has codimension at most 1 in Q . This finishes the proof. \square

Corollary 5.26 Suppose there is a point $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and $\epsilon > 0$ such that for every $X \in \text{VF}(M)$ with $\|X\|_g < \epsilon$ on M one has

$$V|_{\Phi_{\mathcal{L}_R(X)}(t, q_0)}(\pi_Q) \subset T(\mathcal{O}_{\mathcal{D}_R}(q_0)), \quad |t| < \epsilon.$$

Then the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is open in Q .

As a consequence, we have the following characterization of complete controllability: the control system $(\Sigma)_R$ is completely controllable if and only if

$$\forall q \in Q, \quad V|_q(\pi_Q) \subset T|_q \mathcal{O}_{\mathcal{D}_R}(q). \quad (48)$$

Proof. For the first part of the corollary, the assumptions and the previous proposition imply that for every $X \in T|_{x_0} M$ we have $\mathcal{L}_{\text{NS}}(Y, \hat{Y})|_{q_0} \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ for every $Y \in X^\perp$, $\hat{Y} \in A_0 X^\perp$. But since X is an arbitrary element of $T|_{x_0} M$, this means that $\mathcal{D}_{\text{NS}}|_{q_0} \subset T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$ and because $T|_{q_0} Q = \mathcal{D}_{\text{NS}}|_{q_0} \oplus V|_{q_0}(\pi_Q)$, we get $T|_{q_0} Q = T|_{q_0}(\mathcal{O}_{\mathcal{D}_R}(q_0))$. This implies that $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is open in Q . The last part of the corollary is an immediate consequence of this and the fact that Q is connected. \square

Remark 5.27 The above corollary is intuitively obvious. Assumption given by Eq. (48) simply means that there is complete freedom for infinitesimal spinning, i.e., for reorienting one manifold with respect to the other one without moving in $M \times \hat{M}$. In that case, proving complete controllability is easy, by using a crab-like motion.

5.4 Controllability Properties of \mathcal{D}_R

5.4.1 First Results

Proposition 5.17 has the following simple consequence.

Corollary 5.28 The following cases are equivalent:

- (i) The rolling distribution \mathcal{D}_R on Q is involutive.

(ii) For all $X, Y, Z \in T|_x M$ and $(x, \hat{x}; A) \in T^*(M) \otimes T(\hat{M})$

$$\text{Rol}(X, Y)(A) = 0.$$

(iii) (M, g) and (\hat{M}, \hat{g}) both have constant and equal curvature.

The same result holds when one replaces Q by $T^*M \otimes T\hat{M}$.

Proof. (i) \iff (ii) follows from Proposition 5.17.

For the rest of the proof, we use

$$\sigma_{(X,Y)} = g(R(X, Y)Y, X), \text{ and } \sigma_{(\hat{X}, \hat{Y})} = \hat{g}(\hat{R}(\hat{X}, \hat{Y})\hat{Y}, \hat{X}),$$

to denote the sectional curvature of M w.r.t orthonormal vectors $X, Y \in T|_x M$ and the sectional curvature of \hat{M} w.r.t. orthonormal vectors $\hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}$ respectively. We have seen that the involutivity of \mathcal{D}_R is equivalent to the condition in (ii) which is again equivalent (since sectional curvatures completely determine Riemannian curvatures) to the equation

$$\sigma_{(X,Y)} = \hat{\sigma}_{(AX, AY)}, \quad \forall (x, \hat{x}; A) \in Q, \quad X, Y \in T|_x M. \quad (49)$$

(ii) \implies (iii) If we fix $x \in M$ and g -orthonormal vectors $X, Y \in T|_x M$, then, for any $\hat{x} \in \hat{M}$ and any \hat{g} -orthonormal vectors $\hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}$, we may choose $A \in Q|_{(x, \hat{x})}$ such that $AX = \hat{X}$, $AY = \hat{Y}$ (in the case $n = 2$ we may have to replace, say, \hat{X} by $-\hat{X}$ but this does not change anything in the argument below). Hence the above equation (49) shows that the sectional curvatures at every point $\hat{x} \in \hat{M}$ and w.r.t every orthonormal pair \hat{X}, \hat{Y} are all the same i.e., $\sigma_{(X,Y)}$. Thus (\hat{M}, \hat{g}) has constant sectional curvatures i.e., it has a constant curvature. Changing the roles of M and \hat{M} we see that (M, g) also has constant curvature and the constants of curvatures are the same.

(iii) \implies (ii) Suppose that M, \hat{M} have constant and equal curvatures. By a standard result (see [25] Lemma II.3.3), this is equivalent to the fact that there exists $k \in \mathbb{R}$ such that

$$\begin{aligned} R(X, Y)Z &= k(g(Y, Z)X - g(X, Z)Y), \quad X, Y, Z \in T|_x M, \quad x \in M, \\ \hat{R}(\hat{X}, \hat{Y})\hat{Z} &= k(\hat{g}(\hat{Y}, \hat{Z})\hat{X} - \hat{g}(\hat{X}, \hat{Z})\hat{Y}), \quad \hat{X}, \hat{Y}, \hat{Z} \in T|_{\hat{x}} \hat{M}, \quad \hat{x} \in \hat{M}. \end{aligned}$$

On the other hand, if $A \in Q$, $X, Y, Z \in T|_x M$, we would then have

$$\begin{aligned} \hat{R}(AX, AY)(AZ) &= k(\hat{g}(AY, AZ)AX - \hat{g}(AX, AZ)(AY)) \\ &= A(k(g(Y, Z)X - g(X, Z)Y) = A(R(X, Y)Z). \end{aligned}$$

This implies that $\text{Rol}(X, Y)(A) = 0$ since Z was arbitrary. □

In the situation of the previous corollary, the control system $(\Sigma)_R$ is as far away from being controllable as possible: all the orbits $\mathcal{O}_{\mathcal{D}_R}(q)$, $q \in Q$, are integral manifolds of \mathcal{D}_R .

The next consequence of Proposition 5.17 can be seen as a (partial) generalization of the previous corollary and a special case of the Ambrose's theorem 5.42. The corollary gives a necessary and sufficient condition describing the case in which at least one \mathcal{D}_R -orbit is an integral manifold of \mathcal{D}_R . It will be used in the proof of Theorem 5.32 below.

Corollary 5.29 Suppose that (M, g) and (\hat{M}, \hat{g}) are complete. The following cases are equivalent:

(i) There exists a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is an integral manifold of \mathcal{D}_R .

(ii) There exists a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that

$$\text{Rol}(X, Y)(A) = 0, \quad \forall (x, \hat{x}; A) \in \mathcal{O}_{\mathcal{D}_R}(q_0), \quad X, Y \in T|_x M.$$

(iii) There is a complete Riemannian manifold (N, h) and Riemannian covering maps $F : N \rightarrow M, G : N \rightarrow \hat{M}$.

Proof. (i) \Rightarrow (ii): Notice that the restrictions of vector fields $\mathcal{L}_R(X), X \in \text{VF}(M)$, to the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ are smooth vector fields of that orbit. Thus $[\mathcal{L}_R(X), \mathcal{L}_R(Y)]$ is also tangent to this orbit for any $X, Y \in \text{VF}(M)$ and hence Proposition 5.17 implies the claim.

(ii) \Rightarrow (i): It follows, from Proposition 5.17, that $\mathcal{D}_R|_{\mathcal{O}_{\mathcal{D}_R}(q_0)}$, the restriction of \mathcal{D}_R to the manifold $\mathcal{O}_{\mathcal{D}_R}(q_0)$, is involutive. Since maximal connected integral manifolds of an involutive distribution are exactly its orbits, it follows that $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is an integral manifold of \mathcal{D}_R .

(i) \Rightarrow (iii): Let $N := \mathcal{O}_{\mathcal{D}_R}(q_0)$ and $h := (\pi_{Q, M}|_N)^*(g)$ i.e., for $q = (x, \hat{x}; A) \in N$ and $X, Y \in T|_x M$, define

$$h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q) = g(X, Y).$$

If $F := \pi_{Q, M}|_N$ and $G := \pi_{Q, \hat{M}}|_N$, we immediately see that F is a local isometry (note that $\dim(N) = n$) and the fact that G is a local isometry follows from the following computation: for $q = (x, \hat{x}; A) \in N, X, Y \in T|_x M$, one has

$$\hat{g}(G_*(\mathcal{L}_R(X)|_q), G_*(\mathcal{L}_R(Y)|_q)) = \hat{g}(AX, AY) = g(X, Y) = h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q).$$

The completeness of (N, h) can be easily deduced from the completeness of M and \hat{M} together with Proposition 5.6. Proposition II.1.1 in [25] proves that the maps F, G are in fact (surjective and) Riemannian coverings.

(iii) \Rightarrow (ii): Let $x_0 \in M$ and choose $z_0 \in N$ such that $F(z_0) = x_0$. Define $\hat{x}_0 = G(z_0) \in \hat{M}$ and $A_0 := G_*|_{z_0} \circ (F_*|_{z_0})^{-1}$ which is an element of $Q|_{(x_0, \hat{x}_0)}$ since F, G were local isometries. Write $q_0 = (x_0, \hat{x}_0; A_0) \in Q$.

Let $\gamma : [0, 1] \rightarrow M$ be an a.c. curve with $\gamma(0) = x_0$. Since F is a smooth covering map, there is a unique a.c. curve $\Gamma : [0, 1] \rightarrow N$ with $\gamma = F \circ \Gamma$ and $\Gamma(0) = z_0$. Define $\hat{\gamma} = G \circ \Gamma$ and $A(t) = G_*|_{\Gamma(t)} \circ (F_*|_{\Gamma(t)})^{-1} \in Q, t \in [0, 1]$. It follows that, for a.e. $t \in [0, 1]$,

$$\dot{\hat{\gamma}}(t) = G_*|_{\Gamma(t)} \dot{\Gamma}(t) = A(t) \dot{\gamma}(t).$$

Since F, G are local isometries, $\overline{\nabla}_{(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))} A(\cdot) = 0$ for a.e. $t \in [0, 1]$. Thus $t \mapsto (\gamma(t), \hat{\gamma}(t); A(t))$ is the unique rolling curve along γ starting at q_0 and defined on $[0, 1]$ and therefore curves of Q formed in this fashion fill up the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$. Moreover, since F, G are local isometries, it follows that for every $z \in N$ and $X, Y \in T|_{F(z)} M$, $\text{Rol}(X, Y)(G_*|_z \circ (F_*|_z)^{-1}) = 0$. These facts prove that the condition in (ii) holds and the proof is therefore finished. \square

We conclude this subsection with a necessary condition for complete controllability, which is an immediate consequence of Theorem 4.18.

Proposition 5.30 Assume that the manifolds M and \hat{M} are complete non-symmetric, simply connected, irreducible and $n \neq 8$. If the control system $(\Sigma)_R$ is completely controllable, then one of the holonomy groups of M or \hat{M} is equal to $\text{SO}(n)$ (w.r.t some orthonormal frames).

5.4.2 A "Rolling Along Loops" Characterization of Isometry

In this paragraph, we provide a general non-controllability result that will be used later on. It will be the converse of the following simple proposition.

Proposition 5.31 Suppose (M, g) and (\hat{M}, \hat{g}) have a common Riemannian covering space (N, h) with projections (local isometries) $F : N \rightarrow M$ and $G : N \rightarrow \hat{M}$. Then if there exist $x_0 \in M$, $\hat{x}_0 \in \hat{M}$ such that

$$F^{-1}(x_0) \subset G^{-1}(\hat{x}_0),$$

then for $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ with $A_0 = G_* \circ (F_*|_{q_0})^{-1}$ one has that for every loop $\gamma \in \Omega_{x_0}(M)$ based at x_0 the corresponding curve $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)$ on \hat{M} determined by the rolling curve starting from q_0 (exists and) is a loop based \hat{x}_0 i.e.,

$$\gamma \in \Omega_{x_0}(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) \in \Omega_{\hat{x}_0}(\hat{M}).$$

Proof. If $\gamma \in \Omega_{x_0}(M)$, let Γ be the unique lift of γ to N such that $\Gamma(0) = q_0$ and define $\hat{\gamma} = F \circ \Gamma$, $A(t) = G_* \circ (F_*|_{\Gamma(t)})^{-1}$. Then $q(t) = (\gamma(t), \hat{\gamma}(t); A(t))$ is an element of $Q|_{(\gamma(t), \hat{\gamma}(t))}$, since F, G are local isometries and moreover, $q_0 = q(0)$,

$$\begin{aligned} \dot{\hat{\gamma}}(t) &= \frac{d}{dt}(G \circ \Gamma)(t) = G_* \dot{\Gamma}(t) = (G_* \circ (F_*|_{\Gamma(t)})^{-1})(F_* \dot{\Gamma}(t)) = A(t) \dot{\gamma}(t) \\ \hat{\nabla}_{\dot{\hat{\gamma}}(t)}(A(t)P_0^t(\gamma)X) &= \hat{\nabla}_{G_* \dot{\Gamma}(t)}(G_* (F_*|_{\Gamma(t)})^{-1} P_0^t(\gamma)X) \\ &= G_* \nabla_{(F_*|_{\Gamma(t)})^{-1} \dot{\gamma}(t)}((F_*|_{\Gamma(t)})^{-1} P_0^t(\gamma)X) \\ &= (G_* \circ (F_*|_{\Gamma(t)})^{-1}) \nabla_{\dot{\gamma}(t)}(P_0^t(\gamma)X) = 0, \end{aligned}$$

for every $t \in [0, 1]$ and every $X \in T|_{x_0}M$. This proves that $q(t) = q_{\mathcal{D}_R}(\gamma, q_0)(t)$ and since γ is a loop based at x_0 , $F(\Gamma(1)) = \gamma(1) = x_0$, which means that $\Gamma(1) \in F^{-1}(x_0) \subset G^{-1}(\hat{x}_0)$ and thus $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(1) = \hat{\gamma}(1) = G(\Gamma(1)) = \hat{x}_0$. By definition, $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(0) = \hat{x}_0$ and hence $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) \in \Omega_{\hat{x}_0}(\hat{M})$. This completes the proof. \square

Conversely, we have the following theorem which is the main result of this subsection.

Theorem 5.32 Let (M, g) , (\hat{M}, \hat{g}) be complete Riemannian manifolds and suppose that there is a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that for every loop $\gamma \in \Omega_{x_0}(M)$ based at x_0 the corresponding curve $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)$ on \hat{M} determined by the rolling curve starting from q_0 is a loop based \hat{x}_0 i.e.,

$$\gamma \in \Omega_{x_0}(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) \in \Omega_{\hat{x}_0}(\hat{M}). \quad (50)$$

Then (M, g) and (\hat{M}, \hat{g}) have a common Riemannian covering space (N, h) such that if $F : N \rightarrow M$, $G : N \rightarrow \hat{M}$ are the corresponding covering maps, then

$$F^{-1}(x_0) \subset G^{-1}(\hat{x}_0).$$

Proof. For $u, v \in T|_{x_0}M$, a Jacobi field along the geodesic $t \mapsto \exp_{x_0}(tu) =: \gamma_u(t)$ is given by

$$Y_{u,v}(t) = \frac{\partial}{\partial s} \Big|_{s=0} \exp_{x_0}(t(u + sv)) = t(\exp_{x_0})_*|_{tu}(v),$$

together with the initial conditions: $Y_{u,v}(0) = 0$, $\nabla_{\dot{\gamma}_u(0)} Y_{u,v} = v$. Define a function $\hat{\omega}_{u,v} : [0, 1] \times [-1, 1] \rightarrow \hat{M}$ by

$$\begin{aligned} \hat{\omega}_{u,v}(t, s) &:= \\ \hat{\gamma}_{\mathcal{D}_R} \left(\tau \mapsto \exp_{x_0}((1 - \tau)(u + sv)), q_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(u + \sigma v), q_{\mathcal{D}_R}(\gamma_u, q_0)(1))(s) \right) (t). \end{aligned}$$

It is clear from Proposition 5.6 that for every $s \in [-1, 1]$ the map $t \mapsto \hat{\omega}_{u,v}(t, s)$ is a geodesic and moreover it is clear that $\hat{\omega}(t, 0) = \hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q_0)(1 - t)$. This implies that

$$\hat{Y}_{u,v}(t) := \frac{\partial}{\partial s} \Big|_{s=0} \hat{\omega}_{u,v}(1 - t, s), \quad t \in [0, 1],$$

defines a Jacobi field of (\hat{M}, \hat{g}) along the geodesic $\hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q_0)$. We now derive some properties of this Jacobi field.

We first observe that

$$\begin{aligned} \hat{Y}_{u,v}(1) &= \frac{\partial}{\partial s} \Big|_{s=0} \hat{\gamma}_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(u + \sigma v), q_{\mathcal{D}_R}(\gamma_u, q_0)(1))(s) \\ &= A_{\mathcal{D}_R}(\gamma_u, q_0)(1) \frac{\partial}{\partial s} \Big|_0 \exp_{x_0}(u + sv) \\ &= A_{\mathcal{D}_R}(\gamma_u, q_0)(1) Y_{u,v}(1). \end{aligned} \tag{51}$$

We now claim that $\hat{\omega}_{u,v}(1, s) = \hat{x}_0$ for all s . Indeed, we may write $\hat{\omega}_{u,v}(1, s)$ as

$$\hat{\omega}_{u,v}(1, s) = \hat{\gamma}_{\mathcal{D}_R} \left(\underbrace{(\tau \mapsto \exp_{x_0}((1 - \tau)(u + sv)) \cdot (\sigma \mapsto \exp_{x_0}(u + \sigma sv)) \cdot \gamma_u, q_0)(1)}_{=:(\star) \in \Omega_{x_0}(M)} \right) = \hat{x}_0$$

and since the expression (\star) is a loop on M based at x_0 , it follows from the assumption that the path defined on right of the first equality sign is a loop on \hat{M} based at \hat{x}_0 , hence its value at $t = 1$ is \hat{x}_0 . From this follows the second property of $\hat{Y}_{u,v}$, namely

$$\hat{Y}_{u,v}(0) = 0, \tag{52}$$

since $\hat{Y}_{u,v}(0) = \frac{\partial}{\partial s} \Big|_{s=0} \hat{\omega}_{u,v}(1, s) = \frac{\partial}{\partial s} \Big|_0 (s \mapsto x_0) = 0$.

This is a key property since it implies that $\hat{Y}_{u,v}$ has the form

$$\hat{Y}_{u,v}(t) = \frac{\partial}{\partial s} \Big|_0 \widehat{\exp}_{\hat{x}_0}(t(A_0 u + s\hat{v}(u, v))) = t(\widehat{\exp}_{\hat{x}_0})_*|_{tA_0 u}(\hat{v}(u, v)), \tag{53}$$

where $\hat{v}(u, v) := \hat{\nabla}_{\frac{d}{dt} \widehat{\exp}_{\hat{x}_0}(tA_0 u)} \hat{Y}_{u,v} \Big|_{t=0}$. It is clear that $(u, v) \mapsto \hat{v}(u, v)$ is a smooth map $(T|_{x_0}M)^2 \rightarrow T|_{\hat{x}_0}\hat{M}$. We also observed that $\hat{\gamma}_u(t) := \hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q_0) = \widehat{\exp}_{\hat{x}_0}(tA_0 u)$.

We next show the following relation.

Lemma 5.33 With the above notations,

$$\hat{\nabla}_{\hat{\gamma}_u(t)} \hat{Y}_{u,v}|_{t=1} = A_{\mathcal{D}_R}(\gamma_u, q_0)(1) \nabla_{\gamma_u(t)} Y_{u,v}|_{t=1}. \quad (54)$$

Proof. Writing $\partial_t := \frac{\partial}{\partial t} \exp_{x_0}(t(u+sv))$, $\partial_s := \frac{\partial}{\partial s} \exp_{x_0}(t(u+sv))$, $\hat{\partial}_t := \frac{\partial}{\partial t} \hat{\omega}(1-t, s)$, $\hat{\partial}_s := \frac{\partial}{\partial s} \hat{\omega}(1-t, s)$ and $\bar{\partial}_s = (\partial_s, \hat{\partial}_s)$, we have

$$\begin{aligned} \hat{\nabla}_{\hat{\gamma}_u(t)} \hat{Y}_{u,v}|_{t=1} &= \hat{\nabla}_{\hat{\partial}_t} \frac{\partial}{\partial s} \Big|_0 \hat{\omega}(1-t, s) \Big|_{t=1} = \hat{\nabla}_{\hat{\partial}_s} \frac{\partial}{\partial t} \Big|_1 \hat{\omega}(1-t, s) \Big|_{s=0} \\ &= \hat{\nabla}_{\hat{\partial}_s} \frac{\partial}{\partial t} \Big|_1 \hat{\gamma}_{\mathcal{D}_R} \left(\tau \mapsto \exp_{x_0}((1-\tau)(u+sv)), \right. \\ &\quad \left. q_{\mathcal{D}_R}((\sigma \mapsto \exp_{x_0}(u+\sigma sv)) \cdot \gamma_u, q_0)(1) \right) (1-t) \Big|_{s=0} \\ &= \hat{\nabla}_{\hat{\partial}_s} \left(A_{\mathcal{D}_R}((\sigma \mapsto \exp_{x_0}(u+\sigma sv)) \cdot \gamma_u, q_0)(1) \frac{\partial}{\partial t} \Big|_1 \exp_{x_0}(t(u+sv)) \right) \Big|_{s=0} \\ &= \hat{\nabla}_{\hat{\partial}_s} \left(A_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(u+\sigma v), q_{\mathcal{D}_R}(\gamma_u, q_0)(1))(s) \frac{\partial}{\partial t} \Big|_1 \exp_{x_0}(t(u+sv)) \right) \Big|_{s=0} \\ &= \underbrace{\left(\bar{\nabla}_{\bar{\partial}_s} A_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(u+\sigma v), q_{\mathcal{D}_R}(\gamma_u, q_0)(1))(s) \right) \Big|_{s=0}}_{=0} \frac{\partial}{\partial t} \Big|_1 \exp_{x_0}(t(u+sv)) \\ &\quad + A_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(u+\sigma v), q_{\mathcal{D}_R}(\gamma_u, q_0)(1))(0) \left(\nabla_{\partial_s} \frac{\partial}{\partial t} \Big|_1 \exp_{x_0}(t(u+sv)) \right) \Big|_{s=0} \\ &= A_{\mathcal{D}_R}(\gamma_u, q_0)(1) \nabla_{\partial_t} \underbrace{\frac{\partial}{\partial s} \Big|_0 \exp_{x_0}(t(u+sv))}_{=Y_{u,v}(t)} \Big|_{t=1} \\ &= A_{\mathcal{D}_R}(\gamma_u, q_0)(1) \nabla_{\hat{\gamma}_u(t)} Y_{u,v}|_{t=1}, \end{aligned}$$

which gives (54). □

The next technical result goes as follows.

Lemma 5.34 Consider $\hat{v}(u, v)$ defined by (53). Then,

$$\hat{v}(u, v) = A_0 v, \quad \forall u, v \in T|_{x_0} M. \quad (55)$$

Proof. Notice first that for any $\tau \in \mathbb{R}$,

$$Y_{\tau u, v}(t) = \frac{\partial}{\partial s} \Big|_0 \exp_{x_0}(t(\tau u + sv)) = \frac{1}{\tau} \frac{\partial}{\partial \sigma} \Big|_0 \exp_{x_0}(t\tau(u + \sigma v)) = \frac{1}{\tau} Y_{u, v}(t\tau),$$

where, in the first equality, we substituted $\sigma := \frac{s}{\tau}$. Therefore (51) implies that

$$\hat{Y}_{\tau u, v}(1) = A_{\mathcal{D}_R}(\gamma_{\tau u}, q_0)(1) Y_{\tau u, v}(1) = \frac{1}{\tau} A_{\mathcal{D}_R}(\gamma_u, q_0)(\tau) Y_{u, v}(\tau),$$

i.e.,

$$A_{\mathcal{D}_R}(\gamma_u, q_0)(\tau) Y_{u, v}(\tau) = \tau \hat{Y}_{\tau u, v}(1). \quad (56)$$

On one hand, from (54), (56) and (53) one has (recall that $\hat{\gamma}_u(t) = \gamma_{\mathcal{D}_R}(\gamma_u, q_0)(t) = \widehat{\exp}_{\hat{x}_0}(tA_0u)$)

$$\begin{aligned}\hat{\nabla}_{\hat{\gamma}_u(t)}\hat{Y}_{u,v}|_{t=1} &= A_{\mathcal{D}_R}(\gamma_u, q_0)(1)\nabla_{\hat{\gamma}_u(t)}Y_{u,v}|_{t=1} = \hat{\nabla}_{\hat{\gamma}_u(t)}(A_{\mathcal{D}_R}(\gamma_u, q_0)(t)Y_{u,v}(t))|_{t=1} \\ &= \hat{\nabla}_{\hat{\gamma}_u(t)}(t\hat{Y}_{tu,v}(1))|_{t=1} = \hat{Y}_{u,v}(1) + \hat{\nabla}_{\hat{\gamma}_u(t)}\hat{Y}_{tu,v}(1)|_{t=1} \\ &= \hat{Y}_{u,v}(1) + \hat{\nabla}_{\hat{\gamma}_u(t)}\frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(tu, v))\Big|_{t=1} \\ &= \hat{Y}_{u,v}(1) + \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(tu, v)))|_{t=1}.\end{aligned}$$

On the other hand, using only (53) one has

$$\begin{aligned}\hat{\nabla}_{\hat{\gamma}_u(t)}\hat{Y}_{u,v}|_{t=1} &= \hat{\nabla}_{\hat{\gamma}_u(t)}\frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(t(A_0u + s\hat{v}(u, v)))\Big|_{t=1} \\ &= \hat{\nabla}_{\hat{\gamma}_u(t)}(t(\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1} \\ &= (\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\hat{v}(u, v)) + \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1} \\ &= \frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(A_0u + s\hat{v}(u, v)) + \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1} \\ &= \hat{Y}_{u,v}(1) + \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1}.\end{aligned}$$

Combining these two formulas, whose left hand sides are equal, and canceling the common terms $\hat{Y}_{u,v}(1)$ we end up with

$$\hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(tu, v)))|_{t=1} = \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1}.$$

Here we can simplify the left hand side by the following computation: With the notation

$$D_s := \frac{\partial}{\partial s}\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(tu, v)), \quad D_t := \frac{\partial}{\partial t}\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(tu, v)),$$

(notice also that $D_t|_{s=0} = \dot{\hat{\gamma}}_u(t)$) we get

$$\begin{aligned}\hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(tu, v)))|_{t=1} &= \hat{\nabla}_{D_t}\frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(tu, v))\Big|_{t=1} \\ &= \hat{\nabla}_{D_s}\frac{\partial}{\partial t}\Big|_1\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(tu, v))\Big|_{s=0} \\ &= \hat{\nabla}_{D_s}(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u+s\hat{v}(u,v)}(A_0u + s\partial_1\hat{v}(u, v)(u))\Big|_{s=0} \\ &= \hat{\nabla}_{D_s}(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u+s\hat{v}(u,v)}(A_0u)\Big|_{s=0} + \hat{\nabla}_{D_s}(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u+s\hat{v}(u,v)}(s\partial_1\hat{v}(u, v)(u))\Big|_{s=0} \\ &= \hat{\nabla}_{D_s}\frac{\partial}{\partial t}\Big|_1\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(u, v))\Big|_{s=0} \\ &\quad - \hat{\nabla}_{D_s}\frac{\partial}{\partial t}\Big|_1\widehat{\exp}_{\hat{x}_0}(A_0u + s\hat{v}(u, v) + (1-t)s\partial_1\hat{v}(u, v)(u))\Big|_{s=0} \\ &= \hat{\nabla}_{D_t}\frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(tA_0u + s\hat{v}(u, v))\Big|_{t=1} \\ &\quad - \hat{\nabla}_{D_t}\frac{\partial}{\partial s}\Big|_0\widehat{\exp}_{\hat{x}_0}(A_0u + s\hat{v}(u, v) + (1-t)s\partial_1\hat{v}(u, v)(u))\Big|_{t=1} \\ &= \hat{\nabla}_{\hat{\gamma}_u(t)}((\widehat{\exp}_{\hat{x}_0})_*|_{tA_0u}(\hat{v}(u, v)))|_{t=1} \\ &\quad - \hat{\nabla}_{D_t}(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\hat{v}(u, v) + (1-t)\partial_1\hat{v}(u, v)(u))\Big|_{t=1},\end{aligned}$$

where $\partial_1 \hat{v}(u, v)(w)$ for $u, v, w \in T|_{x_0} M$ denotes the directional derivative of \hat{v} at (u, v) in the direction w . The last term on the right of the previous formula simplifies to

$$\begin{aligned} & \hat{\nabla}_{D_t}(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\hat{v}(u, v) + (1-t)\partial_1 \hat{v}(u, v)(u))\Big|_{t=1} \\ &= \hat{\nabla}_{D_t}\left((\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\hat{v}(u, v)) + (1-t)(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\partial_1 \hat{v}(u, v)(u))\right)\Big|_{t=1} \\ &= -(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\partial_1 \hat{v}(u, v)(u)). \end{aligned}$$

Combining the last three formulas, one obtains

$$(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(\partial_1 \hat{v}(u, v)(u)) = 0.$$

Thus for all u such that A_0u is not in the tangent conjugate locus $\hat{Q}_{\hat{x}_0}$ of $\widehat{\exp}_{\hat{x}_0}$, one has $\partial_1 \hat{v}(u, v)(u) = 0$. Moreover, since the complement of $\hat{Q}_{\hat{x}_0}$ is a dense subset of $T|_{\hat{x}_0} \hat{M}$, the continuity of $(u, v) \mapsto \partial_1 \hat{v}(u, v)(u)$ implies that

$$\partial_1 \hat{v}(u, v)(u) = 0, \quad \forall u, v \in T|_{\hat{x}_0} M.$$

But this implies that

$$\hat{v}(u, v) - \hat{v}(0, v) = \int_0^1 \frac{d}{dt} \hat{v}(tu, v) dt = \int_0^1 \frac{1}{t} \underbrace{\partial_1 \hat{v}(tu, v)(tu)}_{=0} dt = 0,$$

and hence we need only to know the values of $\hat{v}(0, v)$ to know all values of $\hat{v}(u, v)$. By the definition of $\hat{\omega}_{u,v}(t)$ one sees that

$$\begin{aligned} \hat{\omega}_{0,v}(t, s) &= \hat{\gamma}_{\mathcal{D}_R}\left(\tau \mapsto \exp_{x_0}((1-\tau)sv), q_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(\sigma v), \underbrace{q_{\mathcal{D}_R}(\gamma_0, q_0)(1)}_{=q_0})(s)\right)(t) \\ &= \hat{\gamma}_{\mathcal{D}_R}\left(\tau \mapsto \exp_{x_0}((1-\tau)sv), q_{\mathcal{D}_R}(\sigma \mapsto \exp_{x_0}(\sigma sv), q_0)(1)\right)(t) \\ &= \hat{\gamma}_{\mathcal{D}_R}(\tau \mapsto \exp_{x_0}(\tau sv), q_0)(1-t) = \widehat{\exp}_{\hat{x}_0}((1-t)sA_0v), \end{aligned}$$

which implies that

$$\hat{Y}_{0,v}(t) = \frac{\partial}{\partial s}\Big|_0 \hat{\omega}_{0,v}(1-t, s) = \frac{\partial}{\partial s}\Big|_0 \widehat{\exp}_{\hat{x}_0}(t(0 + sA_0v)),$$

and therefore, comparing to (53), one obtains

$$\hat{v}(0, v) = A_0v.$$

This finally proves (55) since by the above considerations, $\hat{v}(u, v) = \hat{v}(0, v) = A_0v$. \square

Equations (51), (53) and (55) show that (take $t = 1$)

$$(\widehat{\exp}_{\hat{x}_0})_*|_{A_0u}(A_0v) = A_{\mathcal{D}_R}(\gamma_u, q_0)(1)((\exp_{x_0})_*|_u(v)), \quad \forall u, v \in T|_{x_0} M. \quad (57)$$

We now show that (50) holds with q_0 replaced by any element of the fiber $\mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(x_0)$ of the orbit above x_0 .

Lemma 5.35 Write $F_{x_0} := \mathcal{O}_{\mathcal{D}_R}(q_0) \cap \pi_{Q,M}^{-1}(x_0)$. Then

$$q \in F_{x_0}, \gamma \in \Omega_{x_0}(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, q) \in \Omega_{\hat{x}_0}(\hat{M}). \quad (58)$$

Remark: $\pi_Q(F_{x_0}) = (x_0, \hat{x}_0)$.

Proof. Let $q \in F_{x_0}$. Then there is a $\omega \in \Omega_{x_0}(M)$ such that $q = q_{\mathcal{D}_R}(\omega, q_0)(1)$. Then if $\gamma \in \Omega_{x_0}(M)$,

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma, q)(1) = \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_{\mathcal{D}_R}(\omega, q_0)(1))(1) = \hat{\gamma}_{\mathcal{D}_R}(\gamma \cdot \omega, q_0)(1) = \hat{x}_0,$$

where the last equality follows from (50) since $\gamma \cdot \omega \in \Omega_{x_0}(M)$. Since $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q)(0) = \pi_{Q,\hat{M}}(q) = \hat{x}_0$ as remarked just before the proof, we have $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q) \in \Omega_{\hat{x}_0}(\hat{M})$. \square

Define U to be the subset of $T|_{x_0}M$ of points before the cut time i.e., if for $X \in T|_{x_0}M$, $\|X\|_g = 1$ we let $\tau(X) \in]0, \infty]$ denote the time such that the geodesic γ_X is optimal on $[0, \tau(X)]$ but not after, then

$$U = \{sX \mid X \in T|_{x_0}M, \|X\|_g = 1, 0 \leq s < \tau(X)\}.$$

Since (M, g) is complete, $\tilde{U} := \exp_{x_0}(U)$ is dense in M and $\exp_{x_0} : U \rightarrow \tilde{U}$ is a diffeomorphism.

We now have the following result.

Lemma 5.36 For each $q \in F_{x_0}$ let

$$\phi_q : \tilde{U} \rightarrow \hat{M}; \quad \phi_q = \widehat{\exp}_{\hat{x}_0} \circ A \circ (\exp_{x_0}|_U)^{-1}$$

where $q = (x_0, \hat{x}_0; A)$. Then each mapping ϕ_q is a local isometry $(\tilde{U}, g|_{\tilde{U}}) \rightarrow (\hat{M}, \hat{g})$ and $(\phi_q)_*|_{T|_{x_0}M} = A$.

Proof. Since $q = (x_0, \hat{x}_0; A) \in F_{x_0}$, the previous lemma implies that (57) holds with $q_0 = (x_0, \hat{x}_0; A_0)$ replaced by q . Therefore, if $x \in \tilde{U}$ and $X \in T|_xM$ write $u = (\exp_{x_0}|_U)^{-1}(x) \in T|_{x_0}M$ and $v = ((\exp_{x_0}|_U)^{-1})_*(X) \in T|_u(T|_{x_0}M) = T|_{x_0}M$ and (57) with q_0 replaced by q implies

$$\begin{aligned} \|(\phi_q)_*(X)\|_{\hat{g}} &= \|((\widehat{\exp}_{\hat{x}_0})_* \circ A \circ ((\exp_{x_0}|_U)^{-1})_*)(X)\|_{\hat{g}} \\ &= \|(\widehat{\exp}_{\hat{x}_0})_*|_{Au}(Av)\|_{\hat{g}} = \|A_{\mathcal{D}_R}(\gamma_u, q)(1)(\exp_{x_0})_*|_u(v)\|_{\hat{g}} \\ &= \|(\exp_{x_0})_*|_u(v)\|_g = \|(\exp_{x_0})_*|_u(((\exp_{x_0}|_U)^{-1})_*(X))\|_g \\ &= \|X\|_g, \end{aligned}$$

where the 4. equality follows from the fact that $A \in T|_{x_0}M \rightarrow T|_{\hat{x}_0}\hat{M}$ is an isometry. The claim $(\phi_q)_*|_{T|_{x_0}M} = A$ is obviously true. \square

We will now start proving that $\text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma, q_0)(t)) = 0$ for every piecewise C^1 -path (not necessarily a loop) γ on M such that $\gamma(0) = x_0$ and for all t . First we prove a special case of this (but with q_0 replaced by any $q \in F_{x_0}$).

Lemma 5.37 Let $q \in F_{x_0}$, $u \in T|_{x_0}M$ be a unit vector and let γ_u be the geodesic $t \mapsto \exp_{x_0}(tu)$, Then

$$\text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma_u, q)(t)) = 0, \quad \forall t \in [0, \tau(u)].$$

Proof. Write $q = (x_0, \hat{x}_0; A)$ and notice that by definition of U we have $tu \in U$ for all $t \in [0, \tau(u)[$.

Since ϕ_q of the previous lemma is a local isometry, it follows that

$$P_0^t(\phi_q(\gamma_u)) \circ A = (\phi_q)_* \circ P_0^t(\gamma_u), \quad \forall 0 \leq t < \tau(u).$$

Also, $\frac{d}{dt}\phi_q(\gamma_u)(t) = (\phi_q)_*\dot{\gamma}_u(t)$ for all t so we may conclude that

$$q_{\mathcal{D}_R}(\gamma_u, q)(t) = (\gamma_u(t), (\phi_q \circ \gamma_u)(t); (\phi_q)_*|_{\gamma_u(t)}), \quad \forall 0 \leq t < \tau(u).$$

Again, since ϕ_q is a local isometry, for all $X, Y, Z \in T|_xM$, $x \in \tilde{U}$ we have $(\phi_q)_*(R(X, Y)Z) = \hat{R}((\phi_q)_*X, (\phi_q)_*Y)((\phi_q)_*Z)$ i.e., $\text{Rol}(\cdot, \cdot)((\phi_q)_*|_x) = 0$ for all $x \in \tilde{U}$. But then

$$\text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma_u, q)(t)) = \text{Rol}(\cdot, \cdot)((\phi_q)_*|_{\gamma_u(t)}) = 0, \quad 0 \leq t < \tau(u).$$

Continuity of Rol and $q_{\mathcal{D}_R}(\gamma_u, q)$ now allows us to conclude that the above equation holds for all $0 \leq t \leq \tau(u)$. \square

Now we may prove the claim that was asserted before the previous lemma.

Lemma 5.38 Let $\gamma : [0, 1] \rightarrow M$ a piecewise C^1 -path on M such that $\gamma(0) = x_0$. Then

$$\text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma, q_0)(t)) = 0, \quad \forall t \in [0, 1].$$

Proof. It is clearly enough to prove the claim in the case $t = 1$. Choose any vector $u \in T|_{x_0}M$ such that $\gamma_u : [0, 1] \rightarrow M$ is the minimal geodesic from x_0 to $\gamma(1)$. Define $q := q_{\mathcal{D}_R}(\gamma_u^{-1} \cdot \gamma, q_0)(1)$ and notice that since $\gamma_u^{-1} \cdot \gamma \in \Omega_{x_0}(M)$, we have $q \in F_{x_0}$. Thus by the previous lemma,

$$\text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma_u, q)(1)) = \text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma_{\frac{u}{\|u\|_g}}, q)(\|u\|_g)) = 0,$$

since $\tau(\frac{u}{\|u\|_g}) = \|u\|_g$. But

$$q_{\mathcal{D}_R}(\gamma_u, q)(1) = q_{\mathcal{D}_R}(\gamma_u, q_{\mathcal{D}_R}(\gamma_u^{-1} \cdot \gamma, q_0)(1))(1) = q_{\mathcal{D}_R}(\gamma_u \cdot \gamma_u^{-1} \cdot \gamma, q_0)(1) = q_{\mathcal{D}_R}(\gamma, q_0)(1)$$

and hence

$$0 = \text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma_u, q)(1)) = \text{Rol}(\cdot, \cdot)(A_{\mathcal{D}_R}(\gamma, q_0)(1))$$

which concludes the proof. \square

Finally we may proceed to the proof of the theorem itself. Indeed, since $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is the set of all $q_{\mathcal{D}_R}(\gamma, q_0)(1)$ with all the possible piecewise C^1 -curves $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = x_0$, the previous lemma implies that the condition (ii) of Corollary 5.29 is satisfied. Thus there is a Riemannian manifold (N, h) and Riemannian

covering maps $F : N \rightarrow M$, $G : N \rightarrow \hat{M}$ i.e., (M, g) and (\hat{M}, \hat{g}) have a common Riemannian covering space.

Actually, by Corollary 5.29, we may take $N = \mathcal{O}_{\mathcal{D}_R}(q_0)$, $F = \pi_{Q,M}|_N$, $G = \pi_{Q,\hat{M}}|_N$ and hence if $q \in F^{-1}(x_0)$, then there exists a $\gamma \in \Omega_{x_0}(M)$ such that $q = q_{\mathcal{D}_R}(\gamma, q_0)(1)$ and hence $G(q) = \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(1) = \hat{x}_0$ since $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) \in \Omega_{\hat{x}_0}(\hat{M})$ by the assumption. This shows that $F^{-1}(x_0) \subset G^{-1}(\hat{x}_0)$ and concludes the proof. \square

Remark 5.39 The difficulty in the proof of the previous theorem is due to the fact that the contact points x_0, \hat{x}_0 are fixed i.e., we only assume that loops that are based at x_0 generate, by rolling, loops that are based at \hat{x}_0 .

If we were allowed to have an open neighbourhood of points on M with the property that loops based at these points generate loops on \hat{M} , one could prove that (M, g) and (\hat{M}, \hat{g}) have the same universal Riemannian covering by an easier argument than above.

More precisely, suppose there is a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and a neighbourhood U of x_0 which consists of points x such that whenever one rolls along a geodesic from x_0 to x followed by any loop at x , then the corresponding curve on \hat{M} , generated by rolling, is a geodesic followed by a loop based at the end point of this geodesic.

This means that there is a (possibly smaller) normal neighbourhood U of x_0 such that defining a local $\pi_{Q,M}$ -section \tilde{q} on U by

$$\tilde{q}(x) = (x, \hat{f}(x); \tilde{A}|_x) := q_{\mathcal{D}_R}((t \mapsto \exp_{x_0}(t \exp_{x_0}^{-1}(x))), q_0)(1),$$

then it holds that

$$\forall x \in U, \gamma \in \Omega_x(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, \tilde{q}(x)) \in \Omega_{\hat{f}(x)}(\hat{M}).$$

Notice that $q_0 = \tilde{q}(x_0)$. (In the case of the previous theorem, we had $U = \{x_0\}$, which is not open.)

We will now sketch an easy argument to reach the conclusion of the theorem under this stronger assumption.

Write $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)} = \pi_{Q,M}|_{\mathcal{O}_{\mathcal{D}_R}(q_0)}$ as usual. We show that the vertical bundle $V(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)})$ is actually trivial in the sense that all its fibers consist of one point only (the origin). From this one concludes that $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is an integral manifold of \mathcal{D}_R and hence $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)}$ is (complete and) a Riemannian covering map once the manifold $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is equipped with the Riemannian metric pulled back from that of M (or \hat{M}).

Take $x \in U$ and $v \in V|_{\tilde{q}(x)}(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)})$. This means that there is a smooth curve $s \mapsto \Gamma(s)$, $s \in [0, 1]$, in $\mathcal{O}_{\mathcal{D}_R}(q_0)$ such that $\pi_{Q,M}(\Gamma(s)) = x$ for all s and $\dot{\Gamma}(0) = v$.

One may then choose for each s a smooth path γ_s in M starting at x and defined on $[0, 1]$ such that $q_{\mathcal{D}_R}(\gamma_s, \tilde{q}(x))(1) = \Gamma(s)$. We have $\gamma_s \in \Omega_x(M)$ since

$$\gamma_s(1) = \pi_{Q,M}(q_{\mathcal{D}_R}(\gamma_s, \tilde{q}(x))(1)) = \pi_{Q,M}(\Gamma(s)) = x.$$

Thus by assumption,

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma_s, \tilde{q}(x)) \in \Omega_{\hat{f}(x)}(\hat{M})$$

from which

$$(\pi_{Q,\hat{M}})_* v = \frac{d}{ds} \Big|_0 \pi_{Q,\hat{M}}(\Gamma(s)) = \frac{d}{ds} \Big|_0 \hat{\gamma}_{\mathcal{D}_R}(\gamma_s, \tilde{q}(x))(1) = \frac{d}{ds} \Big|_0 (s \mapsto \tilde{f}(x)) = 0.$$

This proves that every element of $V|_{\tilde{q}(x)}(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)})$, $x \in U$, is of the form $\nu(B)|_{\tilde{q}(x)}$ where $B \in T^*|_x M \otimes T|_{\tilde{f}(x)} \hat{M}$ and $\hat{g}(BX, \tilde{A}|_x Y) + \hat{g}(\tilde{A}|_x X, BY) = 0$, $\forall X, Y \in T|_x M$.

Take a vector field of the form $q \mapsto \nu(B|_q)|_q$ on $\mathcal{O}_{\mathcal{D}_R}(q_0)$ defined along the image of \tilde{q} . Arguing as in Proposition 5.25 and using Eq. (47), we conclude that for every $X \in T|_{x_0} M$ we have

$$-\mathcal{L}_{\text{NS}}(B|_{q_0} X)|_{q_0} + \nu(\overline{\nabla}_{(X, A_0 X)} B|_{\tilde{q}(\cdot)}) \in T|_{q_0} \mathcal{O}_{\mathcal{D}_R}(q_0)$$

and hence, by what we just proved above, the image of this vector under $(\pi_{Q, \hat{M}})_*$ must be zero i.e., $B|_{q_0} X = 0$. Since this holds for all $X \in T|_{x_0} M$, it means that $B|_{q_0} = 0$ and hence we have that $V|_{q_0}(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)}) = \{0\}$. Thus the vertical bundle $V(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0)})$ has rank = 0 since its fiber is = $\{0\}$ at one point.

Remark 5.40 The assumption given by Formula (50) is a special case of a more general one: There is $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ and points $x_1 \in M$, $\hat{x}_1 \in \hat{M}$ such that

$$\gamma \in \Omega_{x_0, x_1}(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0) \in \Omega_{\hat{x}_0, \hat{x}_1}(\hat{M}), \quad (59)$$

where $\Omega_{x_0, x_1}(M)$ is used to denote the set of piecewise C^1 -curves from x_0 to x_1 in M with $\Omega_{\hat{x}_0, \hat{x}_1}(\hat{M})$ defined similarly for \hat{M} .

We actually reduce this setting to the one given in Theorem 5.32 as follows. Fix once and for all a curve $\omega : [0, 1] \rightarrow M$ s.t. $\omega(0) = x_0$, $\omega(1) = x_1$ and write $q_1 = q_{\mathcal{D}_R}(\omega, q_0)(1)$. Then $q_1 = (x_1, \hat{x}_1; A_1)$ by assumption given by Eq. (59), with $A_1 : T|_{x_1} M \rightarrow T|_{\hat{x}_1} \hat{M}$. Then if $\gamma \in \Omega_{x_1}(M)$ is any loop in M based at x_1 , one gets that $\gamma \cdot \omega \in \Omega_{x_0, x_1}(M)$ is a path from x_0 and x_1 . By assumption in Eq. (59) again, one has

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_1)(1) = \hat{\gamma}_{\mathcal{D}_R}(\gamma \cdot \omega, q_0)(1) = \hat{x}_1,$$

and since $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_1)(0) = \hat{x}_1$, we have obtained

$$\gamma \in \Omega_{x_1}(M) \implies \hat{\gamma}_{\mathcal{D}_R}(\gamma, q_1) \in \Omega_{\hat{x}_1}(\hat{M}).$$

Therefore any result obtained under Assumption (50) will also hold true under the assumption given by Formula (59).

5.4.3 The Ambrose's Theorem Revisited

The results developed so far allow us to somewhat simplify the proof of the Ambrose's theorem (see [25] Theorem III.5.1). In fact, the elaborate construction of the covering space X (of the manifold M) is no longer needed since we build this space by simple integrating the distribution \mathcal{D}_R . Actually, as in [25], we will first prove (a version of) the Cartan's theorem ([25] Theorem II.3.2) by using the rolling framework and then use that result and some "patching" to obtain the Ambrose's theorem. The considerations are in parallel to those found in [4], [23].

Definition 5.41 A continuous curve $\gamma : [0, a] \rightarrow M$ on a Riemannian manifold (M, g) is called *once broken geodesic*, broken at t_0 , if there is a $t_0 \in [0, a]$ such that $\gamma|_{[0, t_0]}$, $\gamma|_{[t_0, a]}$ are geodesics of (M, g) .

Notice that if $q = (x, \hat{x}; A) \in Q$ and γ is a once broken geodesic on M starting at x broken at t_0 , then $\hat{\gamma}_{\mathcal{D}_R}(\gamma, q)$ is a once broken geodesic on \hat{M} broken at t_0 .

Ambrose's theorem can now be stated as follows.

Theorem 5.42 (Ambrose) Let (M, g) , (\hat{M}, \hat{g}) be *complete* n -dimensional Riemannian manifolds and let $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. Suppose that M is simply connected and that, for any once broken geodesic $\gamma : [0, a] \rightarrow M$ starting from x_0 , we have

$$A_{\mathcal{D}_R}(\gamma, q_0)(t)(R(X, Y)Z) = \hat{R}(A_{\mathcal{D}_R}(\gamma, q_0)(t)X, A_{\mathcal{D}_R}(\gamma, q_0)(t)Y)(A_{\mathcal{D}_R}(\gamma, q_0)(t)Z), \quad (60)$$

for all $X, Y, Z \in T|_{\gamma(t)}M$ and $t \in [0, a]$. Then, if for any minimal geodesic $\gamma : [0, a] \rightarrow M$ starting from x_0 , one defines $\Phi(\gamma(t)) = \widehat{\exp}_{\hat{x}_0}(tA_0\dot{\gamma}(0))$, $t \in [0, a]$, it follows that the map $\Phi : M \rightarrow \hat{M}$ is a well-defined Riemannian covering.

Remark 5.43 The assumption of Ambrose's theorem is equivalent to the following: For any once broken geodesic $\gamma : [0, a] \rightarrow M$ starting from x_0 and for all $X, Y \in \text{VF}(M)$, $t \in [0, a]$,

$$\text{Rol}(X, Y)(A_{\mathcal{D}_R}(\gamma, q_0)(t)) = 0,$$

which by Proposition 5.17 is equivalent to

$$[\mathcal{L}_R(X), \mathcal{L}_R(Y)]|_{q_{\mathcal{D}_R}(\gamma, q_0)(t)} = \mathcal{L}_R([X, Y])|_{q_{\mathcal{D}_R}(\gamma, q_0)(t)}.$$

i.e., that the distribution \mathcal{D}_R is involutive at each point of Q of the form $q_{\mathcal{D}_R}(\gamma, q_0)(t)$. This should suggest that it is worthwhile to study the integrability of \mathcal{D}_R near the point $q_0 \in Q$, although we are not allowed to use Frobenius theorem.

On a Riemannian manifold (N, h) , we use d_h to denote the distance function (metric) on N induced by h and, for $y \in N$, $X \in T|_yN$, $r > 0$, we use $B_{d_h}(y, r) \subset N$ (resp. $B_h(X, r) \subset T|_yN$) to denote the open ball of radius r on N (resp. $T|_yN$) centered at y (resp. X) w.r.t d_h (resp. h).

The next result provides a local integral manifold of \mathcal{D}_R under milder assumptions than those given in the statement of Ambrose's theorem.

Theorem 5.44 (Cartan) Let (M, g) and (\hat{M}, \hat{g}) be (not necessarily complete) Riemannian manifolds. Consider $q = (x, \hat{x}; A) \in Q$ and $\epsilon > 0$ such that the exponential maps $\exp_x : B_g(0_x, \epsilon) \subset T|_xM \rightarrow B_{d_g}(x, \epsilon)$ and $\widehat{\exp}_{\hat{x}} : B(0_{\hat{x}}, \epsilon) \subset T|_{\hat{x}}\hat{M} \rightarrow B_{d_{\hat{g}}}(\hat{x}, \epsilon)$ are (defined and) diffeomorphisms. Then the following are equivalent:

- (i) For every (non-broken) *geodesic* $\gamma : [0, 1] \rightarrow B_{d_g}(x, \epsilon)$ starting from x , we have

$$\begin{aligned} & A_{\mathcal{D}_R}(\gamma, q)(t)(R(X, Y)Z) \\ &= \hat{R}(A_{\mathcal{D}_R}(\gamma, q)(t)X, A_{\mathcal{D}_R}(\gamma, q)(t)Y)(A_{\mathcal{D}_R}(\gamma, q)(t)Z) \end{aligned} \quad (61)$$

i.e., $\text{Rol}(X, Y)(A_{\mathcal{D}_R}(\gamma, q)(t))Z = 0$ for every $X, Y, Z \in T|_{\gamma(t)}M$ and $t \in [0, 1]$.

- (ii) For every (non-broken) *geodesic* $\gamma : [0, 1] \rightarrow B_{d_g}(x, \epsilon)$ starting from x , we have

$$\begin{aligned} & A_{\mathcal{D}_R}(\gamma, q)(t)(R(X, \dot{\gamma}(t))\dot{\gamma}(t)) \\ &= \hat{R}(A_{\mathcal{D}_R}(\gamma, q)(t)X, \dot{\gamma}_{\mathcal{D}_R}(\gamma, q)(t))\dot{\gamma}_{\mathcal{D}_R}(\gamma, q)(t) \end{aligned} \quad (62)$$

i.e., $\text{Rol}(X, \dot{\gamma}(t))(A_{\mathcal{D}_R}(\gamma, q)(t))\dot{\gamma}(t) = 0$ for every $X \in T|_{\gamma(t)}M$ and $t \in [0, 1]$ (except the break point of γ).

- (iii) There is a connected integral manifold N of \mathcal{D}_R passing through q such that $\pi_{Q,M}|_N \rightarrow B_{d_g}(x, \epsilon)$ (or $\pi_{Q,\hat{M}}|_N \rightarrow B_{d_{\hat{g}}}(\hat{x}, \epsilon)$) is a bijection.
- (iv) The map $\Phi := \widehat{\text{exp}}_{\hat{x}} \circ A \circ \text{exp}_x^{-1}|_{B_{d_g}(x, \epsilon)}$ is an isometric diffeomorphism (onto $B_{d_{\hat{g}}}(\hat{x}, \epsilon)$).

Moreover, if any of the above cases holds, then, for every $X \in B_g(0_x, \epsilon)$, it holds that

$$\begin{aligned} \Phi_*|_{\text{exp}_x(X)} &= P_0^1(s \mapsto \widehat{\text{exp}}_{\hat{x}}(sAX)) \circ A \circ P_1^0(s \mapsto \text{exp}_x(sX)) \\ &= P_0^1\left(s \mapsto \overline{\text{exp}}_{(x, \hat{x})}(s(X, AX))\right). \end{aligned} \quad (63)$$

Proof. (i) \Rightarrow (iii): By taking $Y = Z = \dot{\gamma}$, one has

$$A_{\mathcal{D}_R}(\gamma, q)(t)Y = A_{\mathcal{D}_R}(\gamma, q)(t)Z = \dot{\gamma}_{\mathcal{D}_R}(\gamma, q)(t),$$

for all $t \in [0, 1]$.

(ii) \Rightarrow (iv): Let $u, v \in T|_x M$, $\|u\|_g < \epsilon$ and define for $t \in [0, 1]$

$$Y_{u,v}(t) := \frac{d}{ds}\Big|_0 \text{exp}_x(t(u + sv)) = t(\text{exp}_x)_*|_{tu}v.$$

It is the Jacobi field on M along the geodesic $\gamma_u(t) := \text{exp}_x(tu)$, $t \in [0, 1]$, with $Y_{u,v}(0) = 0$, $\nabla_u Y_{u,v} = v$.

Proposition 5.6 implies that the rolling curve $q_{\mathcal{D}_R}(\gamma_u, q)$ along γ_u is given as

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q)(t) = \widehat{\text{exp}}_{\hat{x}}(tAu), \quad A_{\mathcal{D}_R}(\gamma_u, q)(t) = P_0^t(\hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q)) \circ A \circ P_t^0(\gamma_u).$$

On the other hand, the assumption implies that

$$\text{Rol}(Y_{u,v}(t), \dot{\gamma}_u(t))(A_{\mathcal{D}_R}(\gamma_u, q)(t))\dot{\gamma}_u(t) = 0, \quad t \in [0, 1]$$

and Proposition 5.20 imply that $\hat{Y}_{u,v} := A_{\mathcal{D}_R}(\gamma_u, q)Y_{u,v}$ is a Jacobi field on \hat{M} along the geodesic $\hat{\gamma}_{\mathcal{D}_R}(\gamma_u, q)$.

Clearly, $\hat{Y}_{u,v}(0) = 0$ and $\hat{\nabla}_{Au}\hat{Y}_{u,v} = A\nabla_u Y_{u,v} = Av$, from which it follows (by the uniqueness of solutions of second order ODEs) that $\hat{Y}_{u,v}$ must be the Jacobi field given by

$$\hat{Y}_{u,v}(t) = \frac{d}{ds}\Big|_0 \widehat{\text{exp}}_{\hat{x}}(tA(u + sv)) = t(\widehat{\text{exp}}_{\hat{x}})_*|_{tAu}(Av).$$

Setting $t = 1$, we see that

$$A_{\mathcal{D}_R}(\gamma_u, q)(1)(\text{exp}_x)_*|_u v = (\widehat{\text{exp}}_{\hat{x}})_*|_{Au}(Av),$$

for all $u, v \in T|_x M$ with $\|u\|_g < \epsilon$. In other words, this means that

$$\Phi_*|_y = (\widehat{\text{exp}}_{\hat{x}})_*|_{Au} \circ A \circ (\text{exp}_x^{-1})_*|_y = A_{\mathcal{D}_R}(\gamma_{\text{exp}_x^{-1}(y)}, q)(1),$$

for every $y \in B_{d_g}(x, \epsilon)$, where $B_{d_g}(x, \epsilon)$ is also equal to $\{\text{exp}_x(u) \in T|_x M \mid \|u\|_g < \epsilon\}$. Since $A_{\mathcal{D}_R}(\gamma_{\text{exp}_x^{-1}(y)}, q)(1) \in Q$, this means that $\Phi_*|_y$ is an isometry $T|_y M \rightarrow T|_{\Phi(y)} \hat{M}$ i.e., Φ is an isometry.

(iv) \Rightarrow (iii): This follows from Lemma 5.45 below.

(iii) \Rightarrow (i): Proposition 5.17 implies that $\text{Rol}(X, Y)(A') = 0$ for all $(x', \hat{x}'; A') \in N$ and $X, Y \in T|_{x'}M$.

On the other hand, the assumption implies that $f := (\pi_{Q, M})|_N^{-1}$ is a smooth local section of $\pi_{Q, M}$ defined on $B_{d_g}(x, \epsilon)$ whose image is the integral manifold N of \mathcal{D}_R .

Let $\gamma(t) = \exp_x(tu)$, $t \in [0, 1]$, be a geodesic of M with $\|u\|_g < \epsilon$. Then, since $f \circ \gamma$ is an integral curve of \mathcal{D}_R and $f(\gamma(0)) = q$, the rolling curve $q_{\mathcal{D}_R}(\gamma, q)$ is defined on $[0, 1]$ and is given by $q_{\mathcal{D}_R}(\gamma, q)(t) = f(\gamma(t))$ for all $t \in [0, 1]$. Hence $q_{\mathcal{D}_R}(\gamma, q)(t) \in N$ for all $t \in [0, 1]$, which implies that $\text{Rol}(X, Y)(A_{\mathcal{D}_R}(\gamma, q)(t))Z = 0$ for all $X, Y, Z \in T|_{\gamma(t)}M$. This completes the proof. \square

Lemma 5.45 Let $x_0 \in M$ and $\hat{x}_0 \in \hat{M}$ with corresponding open neighborhoods U and \hat{U} . Then there is a isometry onto $\Phi : U \rightarrow \hat{U}$ if and only if there is a smooth local $\pi_{Q, M}$ -section $f : U \rightarrow Q$, whose image is an integral manifold of \mathcal{D}_R and projects bijectively by $\pi_{Q, \hat{M}}$ onto \hat{U} .

Moreover, the correspondence $\Phi \leftrightarrow f$ is given by

$$\begin{aligned} f_\Phi(x) &= (x, \Phi(x); \Phi_*|_x), \\ \Phi_f(x) &= \pi_{Q, \hat{M}} \circ f. \end{aligned}$$

Proof. Let Φ be an isometry onto $U \rightarrow \hat{U}$ and define f_Φ as above. For every $x \in U$ and $u \in T|_xM$, let $\gamma_u(t) := \exp_x(tX)$. Since Φ is an isometry, $\Phi \circ \gamma_u$ is a \hat{g} -geodesic starting at $\Phi(x)$. Moreover, defining $A(t) = \Phi_*|_{\gamma_u(t)} \in Q$ and taking any $X_0 \in T|_xM$, $X(t) = P_0^t(\gamma_u)X_0$, we have $A(t)X(t) = P_0^t(\Phi \circ \gamma_u)(\Phi_*(X_0))$ and hence

$$(\bar{\nabla}_{(\dot{\gamma}_u(t), \frac{d}{dt}(\Phi \circ \gamma_u)(t))} A(t))X(t) = \hat{\nabla}_{\Phi_*\dot{\gamma}_u(t)}(A(t)X(t)) - A(t)\nabla_{\dot{\gamma}_u(t)}X(t) = 0,$$

which proves that $t \mapsto (\gamma_u(t), (\Phi \circ \gamma_u)(t); A(t)) =: q(t)$ is an integral curve of \mathcal{D}_R through $q(0) = (x, \Phi(x); \Phi_*|_x)$. On the other hand, $q(t) = f_\Phi(\gamma_u(t))$ and thus it follows that

$$(f_\Phi)_*(u) = \dot{q}(0) \in \mathcal{D}_R|_{q(0)}.$$

Hence the image of f_Φ is an integral manifold of \mathcal{D}_R and it clearly projects bijectively onto \hat{U} by $\pi_{Q, \hat{M}}$.

Conversely, suppose that $f : U \rightarrow Q$ is a local π_Q -section whose image is an integral manifold of \mathcal{D}_R and which projects onto \hat{U} . Define Φ_f as above. Then $\Phi_f : U \rightarrow \hat{U}$ and, for every $x \in U$ and $X \in T|_xM$, we have $f_*(X) = \mathcal{L}_R(X)|_{f(x)}$ and thus

$$\|(\Phi_f)_*X\|_{\hat{g}} = \left\| (\pi_{Q, \hat{M}})_*(f_*(X)) \right\|_{\hat{g}} = \left\| (\pi_{Q, \hat{M}})_*(\mathcal{L}_R(X)|_{f(x)}) \right\|_{\hat{g}} = \|f(x)X\|_{\hat{g}} = \|X\|_g,$$

where the final equality follows from the fact that $f(x) \in Q$. The fact that Φ is a bijection $U \rightarrow \hat{U}$ is clear. Hence the conclusion. \square

We can now provide an argument for Theorem 5.42. According to the assumptions done in the statement, Theorem 5.44 implies that there is an integral manifold of \mathcal{D}_R passing through $q_0 = (x_0, \hat{x}_0; A_0)$. Hence, we may choose the maximal connected integral manifold N of \mathcal{D}_R passing through q_0 (where N is the union of all connected integral manifold of \mathcal{D}_R passing through q_0 , see e.g. Lemma 3.19 in [14]).

Endow N with a Riemannian metric h given by: $h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q) = g(X, Y)$ for $q = (x, \hat{x}; A) \in N$ and $X, Y \in T|_x M$. It is then clear that

$$F := \text{pr}_1 \circ \pi_Q|_N : N \rightarrow M, \text{ and } G := \text{pr}_2 \circ \pi_Q|_N : N \rightarrow \hat{M},$$

are local isometries onto open subsets of M and \hat{M} (see also the proof of Corollary 5.29).

We next intend to prove that (N, h) is a complete Riemannian manifold. Here, we have to be more careful than in the proof of "(i) \Rightarrow (iii)" in Corollary 5.29 since we cannot assume that N is the whole orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$.

First of all, the facts that N is an integral manifold of \mathcal{D}_R and F is a local isometry imply that, for any $q = (x, \hat{x}; A) \in N$ and any g -geodesic $t \mapsto \gamma(t) = \exp_x(tX)$ on M starting at x , the rolling curve $t \mapsto q_{\mathcal{D}_R}(\gamma, q)(t)$ stays in N is a h -geodesic on N for t in a small interval containing 0.

Let us assume that N is not complete. Then there exists a h -geodesic $\Gamma : [0, T[\rightarrow N$ starting from q_0 where $[0, T[$ is the maximal non-negative interval of definition and $T < \infty$. Since F is a local isometry, $F \circ \Gamma$ is a g -geodesic on M and since Γ is an integral curve of \mathcal{D}_R , it follows that there is a unique $X \in T|_{x_0} M$ such that, for $t \in [0, T[$, one has

$$\begin{aligned} \Gamma(t) &= q_{\mathcal{D}_R}((s \mapsto \exp_{x_0}(sX)), q_0)(t) \\ &= (\exp_{x_0}(tX), \widehat{\exp}_{x_0}(tA_0X); P_0^t(s \mapsto \widehat{\exp}_{x_0}(sA_0X)) \circ A_0 \circ P_t^0(s \mapsto \exp_{x_0}(sX))). \end{aligned}$$

We write $(\gamma(t), \hat{\gamma}(t); A(t)) := \Gamma(t)$. Since M and \hat{M} are complete, the right hand side of the above equation makes sense for all $t \geq 0$ and we define Γ on $[T, \infty[$ by this formula. We emphasize that we assume Γ to be a geodesic on N only for $[0, T[$.

Write $q_T = (x_T, \hat{x}_T; A_T) := \Gamma(T)$. Choose $\epsilon > 0$ such that \exp_{x_T} and $\widehat{\exp}_{\hat{x}_T}$ are diffeomorphisms $B(0, \epsilon) \rightarrow B_{d_g}(x_T, \epsilon)$, $B(0, \epsilon) \rightarrow B_{d_g}(\hat{x}_T, \epsilon)$ respectively.

If ω is any geodesic $[0, 1] \rightarrow B_{d_g}(x_T, \epsilon)$ starting from x_T , then the concatenation $\omega \sqcup \gamma$ of ω and γ is a once broken geodesic starting from x_0 and therefore, Eq. (60) implies that the assumptions of Theorem 5.44, Case (i), are satisfied (with (ω, q_T) in place of (γ, q)). Indeed, for every $X, Y \in T|_{\omega(t)} M$ and t ,

$$\begin{aligned} \text{Rol}(X, Y)(A_{\mathcal{D}_R}(\omega, q_T)(t)) &= \text{Rol}(X, Y)(A_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q_0)(T))(t)) \\ &= \text{Rol}(X, Y)(A_{\mathcal{D}_R}(\omega \sqcup \gamma, q_0)(t + T)) = 0. \end{aligned}$$

Therefore, Case (iii) there implies the existence of a connected integral manifold \tilde{N} of \mathcal{D}_R passing through $q_T = \Gamma(T)$.

Since \tilde{N} is an integral manifold of \mathcal{D}_R and Γ is an integral curve of \mathcal{D}_R and since $\Gamma(T) \in \tilde{N}$, it follows that $\Gamma(t) \in \tilde{N}$ for all t in an open interval $]T - \eta, T + \eta[$ containing T . Since $\Gamma(t) \in N$ for $t \in [0, T[$, it follows that, for some $t_0 \in]T - \eta, T[$, we have $\Gamma(t_0) \in N \cap \tilde{N}$.

Thus $N \cap \tilde{N} \neq \emptyset$ and hence $N \cup \tilde{N}$ is a connected integral manifold of \mathcal{D}_R passing through q_0 which, because of the maximality of N , implies that $\tilde{N} \subset N$. This implies that Γ is a geodesic of N (since $F \circ \Gamma = \gamma$ is a geodesic of M and F is a local isometry) on the interval $[0, T + \eta[$, contradicting the choice of the finite time T . Thus (N, h) is complete.

Since $F = \text{pr}_1 \circ \pi_Q|_N$ and $G = \text{pr}_2 \circ \pi_Q|_N$ are local Riemannian isometries, it follows from Proposition II.1.1 in [25] that they are covering maps. Taking finally into

account that M is simply connected, one gets that F is an isometric diffeomorphism $N \rightarrow M$ and hence $G \circ F^{-1} : M \rightarrow \hat{M}$ is a Riemannian covering map.

Finally notice that if $\gamma : [0, a] \rightarrow M$ is a minimal geodesic starting from x_0 , then $(G \circ F^{-1})(\gamma(t)) = \widehat{\exp}_{\hat{x}_0}(tA_0\dot{\gamma}(0))$ and hence $\Phi = G \circ F^{-1}$.

6 Rolling Against a Space Form

This section is devoted to the special case of the rolling problem (R) with one of the Riemannian manifolds, usually (\hat{M}, \hat{g}) , being equal to a space form i.e., a simply connected complete Riemannian manifold of constant curvature. The possible cases are: (i) Euclidean space with Euclidean metric (zero curvature), (ii) Sphere (positive curvature) and (iii) Hyperbolic space (negative curvature), cf. e.g. [25].

As mentioned in the introduction, the rolling problem against a space form actually presents a fundamental feature: on the bundle $\pi_{Q,M} : Q \rightarrow M$ one can define a principal bundle structure that preserves the rolling distribution \mathcal{D}_R , and this renders the study of controllability of $(\Sigma)_R$ easier to handle.

We will first provide a detailed study for the rolling against an Euclidean space and then proceed to the case of space forms with non-zero curvature.

6.1 Rolling Against an Euclidean Space

In this section, we give a necessary and sufficient condition for the controllability of $(\Sigma)_R$ in the case that $\hat{M} = \mathbb{R}^n$ equipped with the Euclidean metric $\hat{g} = s_n$.

Recall that if V is a finite dimensional inner product space with h the inner product, the special Euclidean group of (V, h) also denoted $\text{SE}(V)$ is equal to $V \times \text{SO}(V)$, and is equipped with the group operation \star given by

$$(v, L) \star (u, K) := (Lu + v, L \circ K).$$

Here $\text{SO}(V)$ is defined with respect to the inner product h of V . In particular, we write $\text{SE}(n)$ for $\text{SE}(\mathbb{R}^n)$ with \mathbb{R}^n equipped with the standard inner product.

Now fix a point q_0 of $Q = Q(M, \mathbb{R}^n)$ of the form $q_0 = (x_0, 0; A_0)$ i.e., the initial contact point on M is equal to x_0 and, on \mathbb{R}^n , it is the origin. Since (\mathbb{R}^n, s_n) is flat, for any a.c. curve $t \mapsto \hat{x}(t)$ in \mathbb{R}^n and $\hat{X} \in \mathbb{R}^n$ we have $P_0^t(\hat{x}(t))\hat{X} = \hat{X}$, where we understand the canonical isomorphisms $T|_{\hat{x}(0)}\mathbb{R}^n \cong \mathbb{R}^n \cong T|_{\hat{x}(t)}\mathbb{R}^n$. It follows that we parameterize the rolling curves explicitly in the form:

$$q_{\mathcal{D}_R}(\gamma, (x_0, \hat{x}; A))(t) = \left(\gamma(t), \hat{x} + A \int_0^t P_s^0(\gamma)\dot{\gamma}(s)ds; AP_t^0(\gamma) \right), \quad (64)$$

where $\gamma \in \Omega_{x_0}(M)$.

From this it follows that for any $(x_0, 0; A_0), (x_0, \hat{x}; A) \in Q$ and $\gamma \in \Omega_{x_0}(M)$, the point $q_{\mathcal{D}_R}(\gamma, (x_0, \hat{x}; A))(1)$ is equal to

$$(x_0, \hat{x} + AA_0^{-1}\hat{\gamma}_{\mathcal{D}_R}(\gamma, (x_0, 0; A_0))(1); AA_0^{-1}A_{\mathcal{D}_R}(\gamma, (x_0, 0; A_0))(1)).$$

Let $\gamma \in \Omega_{x_0}(M)$ be a piecewise C^1 -loop of M based at x_0 . We define a map

$$\begin{aligned} \rho &= \rho_{q_0} : \Omega_{x_0}(M) \rightarrow \text{SE}(n); \\ \rho(\gamma) &= (\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(1), A_{\mathcal{D}_R}(\gamma, q_0)(1)A_0^{-1}), \end{aligned}$$

where $q_0 = (x_0, 0; A_0) \in Q$. Hence by Remark 5.4 and the above formulas we have

$$\begin{aligned}
\rho(\omega.\gamma) &= (\hat{\gamma}_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q_0)(1))(1), A_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q_0)(1))(1)A_0^{-1}) \\
&= (\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(1) + A_{\mathcal{D}_R}(\gamma, q_0)(1)A_0^{-1}\hat{\gamma}_{\mathcal{D}_R}(\omega, q_0)(1), \\
&\quad A_{\mathcal{D}_R}(\gamma, q_0)(1)A_0^{-1}A_{\mathcal{D}_R}(\omega, q_0)(1)A_0^{-1}) \\
&= (\hat{\gamma}_{\mathcal{D}_R}(\gamma, q_0)(1), A_{\mathcal{D}_R}(\gamma, q_0)(1)A_0^{-1}) \star (\hat{\gamma}_{\mathcal{D}_R}(\omega, q_0)(1), A_{\mathcal{D}_R}(\omega, q_0)(1)A_0^{-1}) \\
&= \rho(\gamma) \star \rho(\omega).
\end{aligned}$$

Thus ρ is a group anti-homomorphism $(\Omega_{x_0}(M), \cdot) \rightarrow (\text{SE}(n), \star)$. This proves that the elements of the form $\rho(\omega)$, $\omega \in \Omega_{x_0}(M)$, form a subgroup of $\text{SE}(n)$. We also see that

$$(\hat{\gamma}_{\mathcal{D}_R}(\gamma, q)(1), A_{\mathcal{D}_R}(\gamma, q)(1)) = (\hat{x}, A) \star (0, A_0)^{-1} \star \rho_{q_0}(\gamma) \star (0, A_0),$$

where $q = (x_0, \hat{x}; A)$, $q_0 = (x_0, 0; A_0) \in Q$ and $\gamma \in \Omega_{x_0}(M)$.

We also make the simple observation from Eq. (64) that the image of $\text{pr}_2 \circ \rho : \Omega_{x_0}(M) \rightarrow \text{SO}(n)$ is exactly $A_0 H|_{x_0} A_0^{-1}$, where $H|_{x_0}$ is the holonomy group of (M, g) at x_0 . Here $A_0 H|_{x_0} A_0^{-1} = H|_F$ with respect to the orthonormal frame $F = (A_0^{-1}e_1, \dots, A_0^{-1}e_n)$ where e_1, \dots, e_n is the standard basis of \mathbb{R}^n .

From these remarks the next proposition follows easily.

Proposition 6.1 Let $Q = Q(M, \mathbb{R}^n)$ and $q_0 = (x_0, \hat{x}_0; A_0) \in Q$. Then the map

$$\begin{aligned}
K_{q_0} : \pi_{Q, M}^{-1}(x_0) &\rightarrow \text{SE}(n); \\
(x_0, \hat{x}; A) &\mapsto (\hat{x} - \hat{x}_0, AA_0^{-1})
\end{aligned}$$

is a diffeomorphism which carries the fiber $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}^{-1}(x_0)$ of the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$ to a submanifold of $\text{SE}(n)$. In particular, if $\hat{x}_0 = 0$ we have that

$$K_{q_0}(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}^{-1}(x_0)) = \rho_{q_0}(\Omega_{x_0}(M))$$

which is a Lie subgroup of $\text{SE}(n)$.

We will make some standard observations of subgroups G of an Euclidean group $\text{SE}(V)$, where (V, h) is a finite dimensional inner product space. Call an element of G of the form (v, id_V) a *pure translation* of G and write $T = T(G)$ for the set that they form. Clearly T is a subgroup of G . As before, pr_1, pr_2 denote the projections $\text{SE}(V) \rightarrow V$ and $\text{SE}(V) \rightarrow \text{SO}(V)$. The natural action, also written by \star , of $\text{SO}(V)$ on V is defined as

$$(u, K) \star v := Kv + u, \quad (u, K) \in \text{SO}(V), v \in V.$$

Proposition 6.2 Let G be a Lie subgroup of $\text{SE}(V)$ with $\text{pr}_2(G) = \text{SO}(V)$. Then either of the following cases hold:

- (i) $G = \text{SE}(V)$ or
- (ii) there exists $v^* \in V$ which is a fixed point of G .

Proof. Suppose first that $T = T(G)$ is non-trivial i.e., there exists a pure translation $(v, \text{id}_V) \in T$, $v \neq 0$. Then for any $(w, A) \in G$ it holds that

$$\begin{aligned} G \ni (w, A)^{-1} \star (v, \text{id}_V) \star (w, A) &= (-A^{-1}w, A^{-1}) \star (v + w, A) \\ &= (A^{-1}(v + w) - A^{-1}w, \text{id}_V) = (A^{-1}v, \text{id}_V) \end{aligned}$$

which implies that

$$\begin{aligned} T \supset \{(A^{-1}v, \text{id}_V) \mid (w, A) \in G\} &= \{(A^{-1}v, \text{id}_V) \mid A \in \text{pr}_2(G) = \text{SO}(V)\} \\ &= S^{n-1}(0, \|v\|) \times \{\text{id}_V\} \end{aligned}$$

where $S^{n-1}(w, r)$, $w \in \mathbb{R}^n$, $r > 0$ is the sphere of radius r centered at $w \in V$ and $\|\cdot\| = h(\cdot, \cdot)^{1/2}$. If $w \in V$ such that $\|w\| \leq \|v\|$ then it is clear that there are $u, u' \in S^{n-1}(0, \|v\|)$ such that $u + u' = w$ (choose $u \in S^{n-1}(0, \|v\|) \cap S^{n-1}(w, \|v\|)$ and $u' = w - u$). Therefore

$$(w, \text{id}_V) = (u, \text{id}_V) \star (u', \text{id}_V) \in T$$

i.e., $\overline{B}(0, \|v\|) \subset T$ where $\overline{B}(w, r)$ is the closed ball of radius r centered at w . Thus for all $k \in \mathbb{N}$,

$$\begin{aligned} &\underbrace{\{\overline{B}(0, \|v\|) + \dots + \overline{B}(0, \|v\|)\}}_{k \text{ times}} \times \{\text{id}_V\} \\ &= \underbrace{\{\overline{B}(0, \|v\|) \times \{\text{id}_V\} \star \dots \star \overline{B}(0, \|v\|) \times \{\text{id}_V\}\}}_{k \text{ times}} \subset T. \end{aligned}$$

From this we conclude that $V \times \{\text{id}_V\} = T$.

Therefore we get the case (i) since

$$\begin{aligned} G = T \star G &= \{(u, \text{id}_V) \star (w, A) \mid u \in V, (w, A) \in G\} \\ &= \{(u + w, A) \mid u \in V, (w, A) \in G\} \\ &= \{(u, A) \mid u \in V, A \in \text{pr}_2(G) = \text{SO}(V)\} \\ &= V \times \text{SO}(V) = \text{SE}(V). \end{aligned}$$

The case that is left to investigate is the one where T is trivial i.e., $T = \{(0, \text{id}_V)\}$. In this case the smooth surjective Lie group homomorphism $\text{pr}_2|_G : G \rightarrow \text{SO}(V)$ is also injective. In fact, if $A = \text{pr}_2(v, A) = \text{pr}_2(w, A)$ for $(v, A), (w, A) \in G$ and $v \neq w$, then

$$G \ni (w, A) \star (v, A)^{-1} = (w, A) \star (-A^{-1}v, A^{-1}) = (w - v, \text{id}_V) \in T$$

and since $(w - v, \text{id}_V) \neq (0, \text{id}_V)$, this contradicts the triviality of T . It follows that $\text{pr}_2|_G$ is a Lie group isomorphism onto $\text{SO}(V)$ and hence a diffeomorphism. In particular, G is compact since $\text{SO}(V)$ is compact.

We next show that there exists $v^* \in V$ which is a fixed point of G . Indeed, taking arbitrary $v \in V$ and writing μ_H for the (right- and) left-invariant normalized (to 1) Haar measure of the compact group G , then we may define

$$v^* := \int_G (B \star v) d\mu_H(B).$$

Thus for $(w, A) \in G$,

$$\begin{aligned} (w, A) \star v^* &= w + Av^* = \int_G (w + A(B \star v)) d\mu_{\mathbb{H}}(B) = \int_G (((w, A) \star B) \star v) d\mu_{\mathbb{H}}(B) \\ &= \int_G (B \star v) d\mu_{\mathbb{H}}(B) = v^*, \end{aligned}$$

where, in the second equality, we have used the linearity of the integral and normality of the Haar measure and in the last phase the left invariance of the Haar measure. This proves that v^* is a fixed point of G and completes the proof. \square

Remark 6.3 With a slight modification, the previous proof actually gives the following generalisation of the last proposition: If G is a connected subgroup of $\text{SE}(V)$ such that the subgroup $\text{pr}_2(G)$ of $\text{SO}(V)$ acts transitively on the unit sphere of V then either (i) $G = V \times \text{pr}_2(G)$ or (ii) there is a fixed point v^* of G .

The previous proposition allows us prove the main theorem of this section.

Theorem 6.4 Suppose (M, g) is a complete Riemannian n -manifold and $(\hat{M}, \hat{g}) = (\mathbb{R}^n, s_n)$ is the Euclidean n -space. Then the rolling problem $(\Sigma)_R$ is completely controllable if and only if the holonomy group of (M, g) is $\text{SO}(n)$ (w.r.t. an orthonormal frame).

Proof. Suppose first that $(\Sigma)_R$ is completely controllable. Then for any given $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ we have that $\pi_{Q, M}^{-1}(x_0) = \pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}^{-1}(x_0)$. In particular, taking any $q_0 \in Q$ of the form $q_0 = (x_0, 0; A_0)$ (i.e., $\hat{x}_0 = 0$), we have by Proposition 6.1 that

$$\text{SE}(n) = K_{q_0}(\pi_{Q, M}^{-1}(x_0)) = K_{q_0}(\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}^{-1}(x_0)) = \rho_{q_0}(\Omega_{x_0}(M)).$$

Hence the image of $\text{pr}_2 \circ \rho_{q_0}$ is $\text{SO}(n)$ and, on the other hand, this image is also $A_0 H|_{x_0} A_0^{-1}$ as noted previously. This proves the necessity of the condition.

Assume now that the holonomy group of M is $\text{SO}(n)$ or, more precisely, that for any $x \in M$ we have $H|_x = \text{SO}(T|_x M)$. Let $q = (x, 0; A) \in Q$ and let $G_q := K_q(\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M}^{-1}(x))$ (see Proposition 6.1) which is a subgroup of $\text{SE}(n)$. Since

$$\text{SO}(n) = AH|_x A^{-1} = (\text{pr}_2 \circ \rho)(\Omega_x(M)) = \text{pr}_2(K_q(\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M}^{-1}(x))) = \text{pr}_2(G_q),$$

by Proposition 6.2, either (i) $G_q = \text{SE}(n)$ or (ii) there exists a fixed point $w_q^* \in \mathbb{R}^n$ of G_q .

If (i) is the case for some $q_0 = (x_0, 0; A_0) \in Q$, then, since K_{q_0} maps $\pi_{Q, M}^{-1}(q_0) \cap \mathcal{O}_{\mathcal{D}_R}(q_0)$ diffeomorphically onto $G_{q_0} = \text{SE}(n)$, it follows that $\pi_{Q, M}^{-1}(q_0) \cap \mathcal{O}_{\mathcal{D}_R}(q_0) = \pi_{Q, M}^{-1}(q_0)$ and hence $\mathcal{O}_{\mathcal{D}_R}(q_0) = Q$ (since $\pi_{\mathcal{O}_{\mathcal{D}_R}(q_0), M}$ is a subbundle of $\pi_{Q, M}$) i.e., $(\Sigma)_R$ is completely controllable.

Therefore suppose that (ii) holds i.e., for every $q \in Q$ of the form $q = (x, 0; A)$ there is a fixed point $w_q^* \in \mathbb{R}^n$ of G_q . We will prove that this implies that (M, g) is flat which is a contradiction since (M, g) does not have a trivial holonomy group.

Thus for any point of Q of the form $q = (x, 0; A)$ we have for all loops $\gamma \in \Omega_x(M)$ that

$$AP_1^0(\gamma)A^{-1}w_q^* + A \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds = w_q^*,$$

since ρ_q is a bijection onto G_q and w_q^* is a fixed point of G_q . In other words we have

$$(P_1^0(\gamma) - \text{id})A^{-1}w_q^* + \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds = 0.$$

Thus if $q = (x, 0; A)$ and $q' = (x, 0; A')$ are on the same π_Q fiber over $(x, 0)$, then

$$(P_1^0(\gamma) - \text{id})(A^{-1}w_q^* - A'^{-1}w_{q'}^*) = 0$$

for every $\gamma \in \Omega_x(M)$. On the other hand, since M has full holonomy i.e., $H|_x = \text{SO}(T|_x M)$, and $H|_x = \{P_1^0(\gamma) \mid \gamma \in \Omega_x(M)\}$, it follows from the above equation that

$$A^{-1}w_q^* = A'^{-1}w_{q'}^*.$$

This means that for every $x \in M$ there is a unique vector $V|_x \in T|_x M$ such that

$$V|_x = A^{-1}w_q^*, \quad \forall q \in \pi_Q^{-1}(x, 0).$$

Moreover, the map $V : M \rightarrow TM; x \mapsto V|_x$ is a vector field on M satisfying

$$P_1^0(\gamma)V|_x - V|_x = - \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds, \quad \forall \gamma \in \Omega_x(M). \quad (65)$$

It follows from this that, for any piecewise C^1 path $\gamma \in C_{\text{pw}}^1([0, 1], M)$, we have

$$V|_{\gamma(1)} = P_0^1(\gamma) \left(V|_{\gamma(0)} - \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds \right). \quad (66)$$

Indeed, if $\omega \in \Omega_{\gamma(1)}(M)$, then $\gamma^{-1}.\omega.\gamma \in \Omega_{\gamma(0)}(M)$ and therefore

$$\begin{aligned} & P_1^0(\gamma)P_1^0(\omega)P_0^1(\gamma)V|_{\gamma(0)} - V|_{\gamma(0)} = P_1^0(\gamma^{-1}.\omega.\gamma)V|_{\gamma(0)} - V|_{\gamma(0)} \\ &= - \int_0^1 P_s^0(\gamma^{-1}.\omega.\gamma) \frac{d}{ds}(\gamma^{-1}.\omega.\gamma)(s)ds \\ &= - \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds - P_1^0(\gamma) \int_0^1 P_s^0(\omega)\dot{\omega}(s)ds - P_1^0(\gamma)P_1^0(\omega) \int_0^1 P_s^0(\gamma^{-1}) \frac{d}{ds}\gamma^{-1}(s)ds \\ &= - \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds + P_1^0(\gamma)(P_1^0(\omega)V|_{\gamma(1)} - V|_{\gamma(1)}) \\ & \quad + P_1^0(\gamma)P_1^0(\omega)P_0^1(\gamma) \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds, \end{aligned}$$

that is

$$(P_1^0(\omega) - \text{id})P_0^1(\gamma) \left(V|_{\gamma(0)} - \int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds \right) = (P_1^0(\omega) - \text{id})V|_{\gamma(1)}.$$

Equation (66) then follows from this since $\{P_1^0(\omega) \mid \omega \in \Omega_{\gamma(1)}(M)\} = H|_{\gamma(1)} = \text{SO}(T|_{\gamma(1)}M)$.

Since (M, g) is complete, the geodesic $\gamma_X(t) = \exp_x(tX)$ is defined for all $t \in [0, 1]$. Inserting this to Eq. (66) and noticing that $P_s^0(\gamma_X)\dot{\gamma}_X(s) = X$ in this case for all $s \in [0, 1]$, we get

$$V|_{\gamma_X(1)} = P_0^1(\gamma_X)(V|_x - X),$$

Therefore, if $X = V|_x$ and $z := \gamma_X(1) = \exp_x(V|_x)$, we get

$$V|_z = 0. \quad (67)$$

Inserting this fact into Eq. (65), one gets

$$\int_0^1 P_s^0(\gamma)\dot{\gamma}(s)ds = 0, \quad \forall \gamma \in \Omega_z(M).$$

Fix $q^* = (z, 0; A_0) \in Q$ (for any isometry $A_0 : T|_z M \rightarrow T|_0 \mathbb{R}^n$). Eq. (64) implies that

$$\hat{\gamma}_{\mathcal{D}_R}(\gamma, (x_0, 0; A_0))(1) = 0, \quad \forall \gamma \in \Omega_z(M).$$

We now apply Theorem 5.32 to conclude that (M, g) has (\mathbb{R}^n, s_n) as a Riemannian covering (i.e., (M, g) is flat) and hence reach the desired contradiction mentioned above. Even though this allows to conclude the proof, we will also give below a direct argument showing this.

Equation (66) is trivially equivalent to

$$P_t^0(\gamma)V|_{\gamma(t)} = V|_{\gamma(0)} - \int_0^t P_s^0(\gamma)\dot{\gamma}(s)ds$$

where $\gamma \in C_{\text{pw}}^1([a, b], M)$, $a < b$, is arbitrary. Taking γ to be smooth and differentiating the above equation w.r.t to t (notice that both sides of the equation are in $T|_{\gamma(0)}M$ for all t), we get

$$P_t^0(\gamma)\nabla_{\dot{\gamma}(t)}V|_{\gamma(\cdot)} = -P_t^0(\gamma)\dot{\gamma}(t),$$

that is

$$\nabla_{\dot{\gamma}(t)}V|_{\gamma(\cdot)} = -\dot{\gamma}(t).$$

Since γ was an arbitrary smooth curve, this implies that V is a smooth vector field on M and

$$\nabla_X V = -X, \quad \forall X \in \text{VF}(M). \quad (68)$$

For any $X \in \text{VF}(M)$, the special curvature $R(X, V)V$ can be seen to vanish everywhere since

$$\begin{aligned} R(X, V)V &= \nabla_X \nabla_V V - \nabla_V \nabla_X V - \nabla_{[X, V]}V = -\nabla_X V + \nabla_V X + [X, V] \\ &= [V, X] + [X, V] = 0, \end{aligned}$$

where, in the second equality, we used (68).

For any $X \in T|_z M$, we write $\gamma_X(t) = \exp_z(tX)$ for the geodesic through z in the direction of X . It follows that

$$\begin{aligned} V|_{\gamma_X(t)} &= P_0^t(\gamma_X)(V|_z - \int_0^t P_s^0(\gamma_X)\dot{\gamma}_X(s)ds) \\ &= P_0^t(\gamma_X)(-\int_0^t X ds) = P_0^t(\gamma_X)(-tX) = -t\dot{\gamma}_X(t). \end{aligned} \quad (69)$$

Now for given $X, v \in T|_z M$ let $Y(t) = \frac{\partial}{\partial s}|_0 \exp_z(t(X + sv))$ be the Jacobi field along γ_X such that $Y(0) = 0, \nabla_{\dot{\gamma}_X(t)} Y|_{t=0} = v$. Then one has

$$\nabla_{\dot{\gamma}_X(t)} \nabla_{\dot{\gamma}_X} Y = R(\dot{\gamma}_X(t), Y(t)) \dot{\gamma}_X(t) = \frac{1}{t^2} R(V|_{\gamma_X(t)}, Y(t)) V|_{\gamma_X(t)} = 0,$$

for $t \neq 0$ which means that $t \mapsto \nabla_{\dot{\gamma}_X(t)} Y$ is parallel along γ_X i.e.,

$$\nabla_{\dot{\gamma}_X(t)} Y = P_0^t(\gamma_X) \nabla_{\dot{\gamma}_X(0)} Y = P_0^t(\gamma_X) v.$$

This allows us to compute

$$\begin{aligned} \frac{d^2}{dt^2} \|Y(t)\|_g^2 &= 2 \frac{d}{dt} g(\nabla_{\dot{\gamma}_X(t)} Y, Y(t)) \\ &= 2g(\underbrace{\nabla_{\dot{\gamma}_X(t)} \nabla_{\dot{\gamma}_X} Y}_{=0}, Y(t)) + 2g(\nabla_{\dot{\gamma}_X(t)} Y, \nabla_{\dot{\gamma}_X(t)} Y) \\ &= 2g(P_0^t(\gamma_X) v, P_0^t(\gamma_X) v) = 2 \|v\|_g^2 \end{aligned}$$

and hence for any t

$$\frac{d}{dt} \|Y(t)\|_g^2 = 2 \|v\|_g^2 t + \frac{d}{dt}|_0 \|Y(t)\|_g^2 = 2 \|v\|_g^2 t,$$

because $\frac{d}{dt}|_0 \|Y(t)\|_g^2 = 2g(\nabla_{\dot{\gamma}_X(0)} Y, Y(0)) = 0$ since $Y(0) = 0$. Again, since $Y(0) = 0$,

$$\|Y(t)\|_g^2 = \|v\|_g^2 t^2 + \|Y(0)\|_g^2 = \|v\|_g^2 t^2$$

which, when spelled out, means that $\|t(\exp_z)_*|_{tX}(v)\|_g = \|tv\|_g$ and hence, when $t = 1$,

$$\|(\exp_z)_*|_X(v)\|_g = \|v\|_g, \quad \forall X, v \in T|_z M. \quad (70)$$

This proves that \exp_z is a local isometry $(T|_z M, g|_z) \rightarrow (M, g)$ and hence a Riemannian covering. Thus (M, g) is flat and the proof is finished. \square

Remark 6.5 For results and proofs in similar lines to those of the above Proposition and Theorem, see Theorem IV.7.1, p. 193 and Theorem IV.7.2, p. 194 in [13].

6.2 Rolling Against a Non-Flat Space Form

In this subsection, we study the controllability problem of $(\Sigma)_R$ in the case where \hat{M} is a simply connected n -dimensional manifold with non zero constant curvature equal to $\frac{1}{k}$, with $k \neq 0$.

6.2.1 Standard Results on Space Forms

Following section V.3 of [13], we define the space form \hat{M}_k of curvature $\frac{1}{k}$ as a subset of \mathbb{R}^{n+1} , $n \in \mathbb{N}$, given by

$$\hat{M}_k := \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid x_1^2 + \dots + x_n^2 + kx_{n+1}^2 = k, x_{n+1} + \frac{k}{|k|} \geq 0\}.$$

Equip \hat{M}_k with a Riemannian metric \hat{g}_k defined as the restriction to \hat{M}_k of the non-degenerate symmetric $(0, 2)$ -tensor

$$s_{n,k} := (dx_1)^2 + \cdots + (dx_n)^2 + k(dx_{n+1})^2.$$

The condition $x_{n+1} + \frac{k}{|k|} \geq 0$ in the definition \hat{M}_k guarantees that \hat{M}_k is connected also when $k < 0$. If the dimension n is not clear from context, we write $(\hat{M}_{n,k}, \hat{g}_{n,k})$ for the above Riemannian manifolds.

Remark 6.6 (i) If $k = 1$ then $\hat{M}_1 = S^n$ (the usual Euclidean unit sphere in \mathbb{R}^{n+1}) and $s_{n,1}$ is the usual Euclidean metric s_{n+1} on \mathbb{R}^{n+1} . For a fixed $n \in \mathbb{N}$, the spaces \hat{M}_k for $k > 0$ are all diffeomorphic: the map $\phi_k : (x_1, \dots, x_n, x_{n+1}) \mapsto (\frac{x_1}{\sqrt{k}}, \dots, \frac{x_n}{\sqrt{k}}, x_{n+1})$ gives a diffeomorphism from \hat{M}_k onto \hat{M}_1 . Moreover, ϕ_k is a homothety since $\phi_k^* \hat{g}_1 = \frac{1}{k} \hat{g}_k$.

(ii) If $k = -1$ then $s_{n,-1}$ is the usual Minkowski "metric" on \mathbb{R}^{n+1} . For a fixed $n \in \mathbb{N}$, the spaces \hat{M}_k for $k < 0$ are all diffeomorphic: the map $\phi_k : (x_1, \dots, x_n, x_{n+1}) \mapsto (\frac{x_1}{\sqrt{-k}}, \dots, \frac{x_n}{\sqrt{-k}}, x_{n+1})$ gives a diffeomorphism from \hat{M}_k onto \hat{M}_{-1} . Moreover, ϕ_k is a homothety since $\phi_k^* \hat{g}_1 = -\frac{1}{k} \hat{g}_k$.

Let $G(n, k)$ be the Lie group of linear maps $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ that leave invariant the bilinear form

$$\langle x, y \rangle_{n,k} := \sum_{i=1}^n x_i y_i + k x_{n+1} y_{n+1},$$

for $x = (x_1, \dots, x_{n+1})$, $y = (y_1, \dots, y_{n+1})$ and having determinant $+1$. In other words, a linear map $B : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ belongs to $G(n, k)$ if and only if $\det(B) = +1$ and

$$\langle Bx, By \rangle_{n,k} = \langle x, y \rangle_{n,k}, \quad \forall x, y \in \mathbb{R}^{n+1},$$

or, equivalently,

$$B^T I_{n,k} B = I_{n,k}, \quad \det(B) = +1,$$

where $I_{n,k} = \text{diag}(1, 1, \dots, 1, k)$. In particular, $G(n, 1) = \text{SO}(n+1)$ and $G(n, -1) = \text{SO}(n, 1)$.

The Lie algebra of the Lie group $G(n, k)$ will be denoted by $\mathfrak{g}(n, k)$. Notice that an $(n+1) \times (n+1)$ real matrix B belongs to $\mathfrak{g}(n, k)$ if and only if

$$B^T I_{n,k} + I_{n,k} B = 0,$$

where $I_{n,k}$ was introduced above.

Sometimes we identify the form $s_{n,k}$ on \mathbb{R}^{n+1} with $\langle \cdot, \cdot \rangle_{n,k}$ using the canonical identification of the tangent spaces $T|_v \mathbb{R}^{n+1}$ with \mathbb{R}^{n+1} . Notice that if $\hat{x} \in \hat{M}_k$ and $V \in T|_{\hat{x}} \mathbb{R}^{n+1}$, then

$$V \in T|_{\hat{x}} \hat{M}_k \iff s_{n,k}(V, \hat{x}) = 0.$$

In fact, if we identify V as a vector (V_1, \dots, V_{n+1}) in \mathbb{R}^{n+1} , then the condition for V to be tangent to the hypersurface \hat{M}_k is

$$\begin{aligned} 0 &= \langle V, \text{grad}(x_1^2 + \dots + x_n^2 + kx_{n+1}) \rangle_{n+1} \\ &= \langle V, (x_1, \dots, x_n, kx_{n+1}) \rangle_{n+1} \\ &= \sum_{i=1}^n x_i V_i + kx_{n+1} V_{n+1} \\ &= s_{n,k}(V, \hat{x}), \end{aligned}$$

with $\langle \cdot, \cdot \rangle_{n+1}$ the standard Euclidean inner product of \mathbb{R}^{n+1} .

Remark 6.7 By using the bilinear form $\langle \cdot, \cdot \rangle_{n,k}$ one may restate the definition of \hat{M}_k by

$$\hat{M}_k = \left\{ \hat{x} \in \mathbb{R}^{n+1} \mid \langle \hat{x}, \hat{x} \rangle_{n,k} = k, x_{n+1} + \frac{k}{|k|} \geq 0 \right\}.$$

Remark 6.8 For convenience we recall a standard result ([13], Theorem V.3.1): The Riemannian manifold (\hat{M}_k, \hat{g}_k) has constant sectional curvature $\frac{1}{k}$ and the isometry group $\text{Iso}(\hat{M}_k, \hat{g}_k)$ is equal to $G(n, k)$.

We understand without mention that when considering the action of $G(n, k)$ on \hat{M}_k we consider the restriction of the maps of $G(n, k)$ onto the set \hat{M}_k .

6.2.2 Orbit Structure

Proposition 6.9 The bundle $\pi_{Q,M} : Q \rightarrow M$ is a principal $G(n, k)$ -bundle with a left action $\mu : G(n, k) \times Q \rightarrow Q$ defined by

$$\mu(B, q) = (x, B\hat{x}; B \circ A),$$

where $q = (x, \hat{x}; A)$ and in $B \circ A$ we understand the range $T|_{\hat{x}} \hat{M}_k$ of A to be identified with a linear subspace of \mathbb{R}^{n+1} in the canonical way.

Moreover, the action μ preserves the distribution \mathcal{D}_R i.e., for any $q \in Q$ and $B \in G(n, k)$,

$$(\mu_B)_* \mathcal{D}_R|_q = \mathcal{D}_R|_{\mu(B,q)}$$

where $\mu_B : Q \rightarrow Q; q \mapsto \mu(B, q)$.

Proof. Let us first check that for $(x, \hat{x}; A) \in Q$, $B \circ A : T|_x M \rightarrow \mathbb{R}^{n+1}$ can be viewed as an orientation preserving map $T|_x M \rightarrow T|_{B\hat{x}} \hat{M}_k$ and that really $(x, B\hat{x}; B \circ A)$ is an element of Q . First of all, $B\hat{x} \in \hat{M}_k$ when $\hat{x} \in \hat{M}_k$ as remarked above. Moreover, for $X \in T|_x M$,

$$s_{n,k}((B \circ A)(X), B\hat{x}) = s_{n,k}(AX, \hat{x}) = 0,$$

since $AX \in T|_{\hat{x}} \hat{M}_k$. Hence $B \circ A : T|_x M \rightarrow T|_{B\hat{x}} \hat{M}_k$. Similarly, for $X, Y \in T|_x M$,

$$\begin{aligned} \hat{g}_k((B \circ A)(X), (B \circ A)(Y)) &= s_{n,k}((B \circ A)(X), (B \circ A)(Y)) \\ &= s_{n,k}(AX, AY) = \hat{g}_k(AX, AY) = g(X, Y), \end{aligned}$$

and clearly $B \circ A$ preserves orientation (since $G(n, k)$ is connected). Thus $(x, B\hat{x}; B \circ A) \in Q$.

It is clear that μ is a well defined left $G(n, k)$ -action on Q , that it is free, maps each $\pi_{Q, M}$ -fiber to itself ($\pi_{Q, M} \circ \mu(B, q) = \pi_{Q, M}(q)$) and that it is transitive fiberwise (for each $q, q' \in \pi_{Q, M}^{-1}(x)$, $\mu(B, q) = q'$ for some $B \in G(n, k)$). It remains to check the claim that this action preserves \mathcal{D}_R in the sense stated above.

Let $B \in G(n, k)$ and $q_0 = (x_0, \hat{x}_0; A_0)$. The fact that $\text{Iso}(\hat{M}_k, \hat{g}) = G(n, k)$ means that defining $F : \hat{M}_k \rightarrow \hat{M}_k$; $F = B|_{\hat{M}_k}$ then $F \in \text{Iso}(\hat{M}_k, \hat{g})$. Clearly $F(\hat{x}_0) = B\hat{x}_0$ and $\hat{F}_*|_{\hat{x}_0} = B|_{T|_{\hat{x}_0}\hat{M}_k}$ and hence by Proposition 5.13

$$\mu(B, q_{\mathcal{D}_R}(\gamma, q_0)(t)) = \hat{F} \cdot q_{\mathcal{D}_R}(\gamma, q_0)(t) = q_{\mathcal{D}_R}(\gamma, \hat{F} \cdot q_0)(t) = q_{\mathcal{D}_R}(\gamma, \mu(B, q_0))(t),$$

for any smooth curve $\gamma : [0, 1] \rightarrow M$, $\gamma(0) = x_0$ and $t \in [0, 1]$. Taking derivative with respect to t at $t = 0$ and using the fact that, by definition, $q_{\mathcal{D}_R}(\gamma, q_0)$ is tangent to \mathcal{D}_R , we find that

$$\begin{aligned} (\mu_B)_* \mathcal{L}_R(\dot{\gamma}(0))|_{q_0} &= (\mu_B)_* \frac{d}{dt} \Big|_0 q_{\mathcal{D}_R}(\gamma, q_0)(t) = \frac{d}{dt} \Big|_0 \mu(B, q_{\mathcal{D}_R}(\gamma, q_0)(t)) \\ &= \frac{d}{dt} \Big|_0 q_{\mathcal{D}_R}(\gamma, \mu(B, q_0))(t) = \mathcal{L}_R(\dot{\gamma}(0))|_{\mu(B, q_0)}. \end{aligned}$$

This allows us to conclude. \square

We will denote the left action of $B \in G(n, k)$ on $q \in Q$ usually by $B \cdot q = \mu(B, q)$.

Proposition 6.10 For any given $q = (x, \hat{x}; A) \in Q$ there is a unique subgroup G_q of $G(n, k)$, called the holonomy group of \mathcal{D}_R , such that

$$G_q \cdot q = \mathcal{O}_{\mathcal{D}_R}(q) \cap \pi_{Q, M}^{-1}(x).$$

Also, if $q' = (x, \hat{x}'; A') \in Q$ is in the same $\pi_{Q, M}$ -fiber as q , then G_q and $G_{q'}$ are conjugate in $G(n, k)$ and all conjugacy classes of G_q in $G(n, k)$ are of the form $G_{q'}$. This conjugacy class will be denoted by G .

Moreover, $\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M} : \mathcal{O}_{\mathcal{D}_R}(q) \rightarrow M$ is a principal G -bundle over M .

Proof. These results follow from the general theory of principal bundle connections (cf. [11], [13]) but the argument is reproduced here for convenience.

Let $q' \in \mathcal{O}_{\mathcal{D}_R}(q) \cap \pi_{Q, M}^{-1}(x)$ and choose a $\gamma \in \Omega_x(M)$ such that $q' = q_{\mathcal{D}_R}(\gamma, q)(1)$. Since the $G(n, k)$ action is free and transitive on $\pi_{Q, M}^{-1}(x)$, it follows that there is a unique $B_q(\gamma) \in G(n, k)$ such that $B_q(\gamma) \cdot q = q'$. We define $G_q = \{B_q(\gamma) \mid \gamma \in \Omega_x(M)\}$ and note that for $\gamma, \omega \in \Omega_x(M)$ one has

$$\begin{aligned} (B_q(\gamma)B_q(\omega)) \cdot q &= B_q(\gamma) \cdot (B_q(\omega) \cdot q) = B_q(\gamma) \cdot q_{\mathcal{D}_R}(\omega, q)(1) = q_{\mathcal{D}_R}(\omega, B_q(\gamma) \cdot q)(1) \\ &= q_{\mathcal{D}_R}(\omega, q_{\mathcal{D}_R}(\gamma, q)(1))(1) = q_{\mathcal{D}_R}(\omega \cdot \gamma, q)(1) \in \mathcal{O}_{\mathcal{D}_R}(q) \cap \pi_{Q, M}^{-1}(x), \end{aligned}$$

which proves that $B_q(\gamma)B_q(\omega) = B_q(\omega \cdot \gamma) \in G_q$. Next if $\gamma^{-1} : [0, 1] \rightarrow M$ denotes the inverse path of γ i.e., $\gamma^{-1}(t) = \gamma(1 - t)$ for $t \in [0, 1]$, it follow that

$$(B_q(\gamma)B_q(\gamma^{-1})) \cdot q = q_{\mathcal{D}_R}(\gamma^{-1} \cdot \gamma, q)(1) = q,$$

i.e., $B_q(\gamma)^{-1} = B_q(\gamma^{-1}) \in G_q$. This shows that G_q is indeed a subgroup of $G(n, k)$. Moreover, it is clear that

$$G_q \cdot q = \mathcal{O}_{\mathcal{D}_R}(q) \cap \pi_{Q, M}^{-1}(x),$$

where the left hand side is $\{B \cdot q \mid B \in G_q\}$.

Let us prove the statement about the conjugacy class of G_q . Take $q' = (x, \hat{x}'; A') \in Q$. Because $G(n, k)$ acts transitively on the fibers, there exists a $B \in G(n, k)$ such that $q' = B \cdot q$. Therefore for any $\gamma \in \Omega_x(M)$,

$$\begin{aligned} (B^{-1}B_{q'}(\gamma)B) \cdot q &= (B^{-1}B_{q'}(\gamma)) \cdot q' = B^{-1} \cdot q_{\mathcal{D}_R}(\gamma, q') \\ &= q_{\mathcal{D}_R}(\gamma, B^{-1} \cdot q') = q_{\mathcal{D}_R}(\gamma, q) = B_q(\gamma) \cdot q, \end{aligned}$$

i.e., $B^{-1}B_{q'}(\gamma)B = B_q(\gamma)$ since the $G(n, k)$ action is free. This proves that $B^{-1}G_{q'}B = G_q$. Moreover, if there is a $B \in G(n, k)$ and a subgroup G' of $G(n, k)$ such that $B^{-1}G'B = G_q$, then defining $q' := B \cdot q$ one gets that $G' = G_{q'}$.

By Proposition 5.11, $\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M}$ is a smooth bundle and, by what has been said already, it is clear that G_q preserves the fibers $\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M}^{-1}(x) = \pi_{Q, M}^{-1}(x) \cap \mathcal{O}_{\mathcal{D}_R}(q)$ and the action is free. Recall that, if a map from some manifold to the ambient manifold is smooth and its image is contained in the orbit (as a set), then this map is also smooth as a map into the orbit (as a manifold) (cf. [14], Theorem 3.22 and Lemma 2.17). As a consequence, the action of G_q is also smooth. From this, one concludes that $\pi_{\mathcal{O}_{\mathcal{D}_R}(q), M}$ is a G_q -bundle and hence a G -bundle since the Lie groups in the conjugacy class are all isomorphic. \square

6.2.3 The Rolling Connection

Let $\pi_{TM \oplus \mathbb{R}} : TM \oplus \mathbb{R} \rightarrow M$ be the vector bundle over M where $\pi_{TM \oplus \mathbb{R}}(X, r) = \pi_{TM}(X)$. In this section we will prove the following result.

Theorem 6.11 There exists a vector bundle connection ∇^{Rol} of the vector bundle $\pi_{TM \oplus \mathbb{R}}$ that we call the *rolling connection*, and which we define as follows: for every $x \in M$, $Y \in T|_x M$, $X \in \text{VF}(M)$, $r \in C^\infty(M)$,

$$\nabla_Y^{\text{Rol}}(X, r) = \left(\nabla_Y X + r(x)Y, Y(r) - \frac{1}{k}g(X|_x, Y) \right), \quad (71)$$

such that in the case of M rolling against the space form \hat{M}_k , $k \neq 0$, the holonomy group G of \mathcal{D}_R is isomorphic to the holonomy group $H^{\nabla^{\text{Rol}}}$ of ∇^{Rol} .

Moreover, if one defines a fiber inner product h_k on $TM \oplus \mathbb{R}$ by

$$h_k((X, r), (Y, s)) = g(X, Y) + krs,$$

where $X, Y \in T|_x M$, $r, s \in \mathbb{R}$, then ∇^{Rol} is a metric connection in the sense that for every $X, Y, Z \in \text{VF}(M)$, $r, s \in C^\infty(M)$,

$$Z(h_k((X, r), (Y, s))) = h_k(\nabla_Z^{\text{Rol}}(X, r), (Y, s)) + h_k((X, r), \nabla_Z^{\text{Rol}}(Y, s)).$$

Before providing the proof of the theorem, we present the equations of parallel transport w.r.t ∇^{Rol} along a general curve and along a geodesic of M and also the curvature of ∇^{Rol} . Let $\gamma : [0, 1] \rightarrow M$ be an a.c. curve on M , $\gamma(0) = x$ and let

$(X_0, r_0) \in T|_x M \oplus \mathbb{R}$. Then the parallel transport $(X(t), r(t)) = (P^{\nabla^{\text{Rol}}})_0^t(\gamma)(X_0, r_0)$ of (X_0, r_0) is determined from the equations

$$\begin{cases} \nabla_{\dot{\gamma}(t)} X + r(t)\dot{\gamma}(t) = 0, \\ \dot{r}(t) - \frac{1}{k}g(\dot{\gamma}(t), X(t)) = 0, \end{cases} \quad (72)$$

for a.e. $t \in [0, 1]$. In particular, if γ is a geodesic on (M, g) , one may derive the following uncoupled second order differential equations for X and r ,

$$\begin{cases} \nabla_{\dot{\gamma}(t)} \nabla_{\dot{\gamma}(t)} X + \frac{1}{k}g(X(t), \dot{\gamma}(t))\dot{\gamma}(t) = 0, \\ \ddot{r}(t) + \frac{\|\dot{\gamma}(t)\|_g^2}{k}r(t) = 0, \end{cases} \quad (73)$$

for all t .

One easily checks by direct computation that the connection ∇^{Rol} on $\pi_{TM \oplus \mathbb{R}}$ has the curvature,

$$R^{\nabla^{\text{Rol}}}(X, Y)(Z, r) = (R(X, Y)Z - \frac{1}{k}(g(Y, Z)X - g(X, Z)Y), 0), \quad (74)$$

where $X, Y, Z \in \text{VF}(M)$, $r \in C^\infty(M)$.

We will devote the rest of the subsection to prove Theorem 6.11.

Proof. The rolling distribution \mathcal{D}_R is a principal bundle connection for the principal $G(n, k)$ -bundle $\pi_{Q, M} : Q \rightarrow M$ and hence there is a vector bundle $\xi : E \rightarrow M$ with fibers isomorphic to \mathbb{R}^{n+1} and a unique linear vector bundle connection $\nabla^{\text{Rol}} : \Gamma(\xi) \times \text{VF}(M) \rightarrow \Gamma(\xi)$ which induces the distribution \mathcal{D}_R on Q . This clearly implies that the holonomy group G of \mathcal{D}_R and $H^{\nabla^{\text{Rol}}}$ of ∇^{Rol} are isomorphic. We will eventually show that ξ is further isomorphic to $\pi_{TM \oplus \mathbb{R}}$ and give the explicit expression (71) for the connection of $\pi_{TM \oplus \mathbb{R}}$ induced by this isomorphisms from ∇^{Rol} on ξ .

There is a canonical non-degenerate metric $h_k : E \odot E \rightarrow M$ on the vector bundle ξ (positive definite when $k > 0$ and indefinite of Minkowskian type if $k < 0$) and the connection ∇^{Rol} is a metric connection w.r.t. to h_k i.e., for any $Y \in \text{VF}(M)$ and $s, \sigma \in \Gamma(\nu)$,

$$Y(h_k(s, \sigma)) = h_k(\nabla_Y^{\text{Rol}} s, \sigma) + h_k(s, \nabla_Y^{\text{Rol}} \sigma). \quad (75)$$

The construction of ξ goes as follows (see [11], section 2.1.3). Define a left $G(n, k)$ -group action β on $Q \times \mathbb{R}^{n+1}$ by

$$\beta(B, (q, v)) = (B \cdot q, Bv),$$

where $q \in Q$, $v \in \mathbb{R}^{n+1}$, $B \in G(n, k)$. The action β is clearly smooth, free and proper. Hence $E := (Q \times \mathbb{R}^{n+1})/\beta$ is a smooth manifold of dimension $n + (n+1) = 2n+1$. The β -equivalence classe (i.e., β -orbit) of $(q, v) \in Q \times \mathbb{R}^{n+1}$ is denoted by $[(q, v)]$. Then one defines $\xi([(q, v)]) = \pi_{Q, M}(q)$ which is well defined since the β -action preserves the fibers of $Q \times \mathbb{R}^{n+1} \rightarrow M$; $(q, v) \mapsto \pi_{Q, M}(q)$. We prove now that ξ is isomorphic, as a vector bundle over M , to

$$\begin{aligned} \pi_{TM \oplus \mathbb{R}} : TM \oplus \mathbb{R} &\rightarrow M, \\ (X, t) &\mapsto \pi_{TM}(X). \end{aligned}$$

Indeed, let $f \in \Gamma(\xi)$ and notice that for any $q \in Q$ there exists a unique $\bar{f}(q) \in \mathbb{R}^{n+1}$ such that $[(q, \bar{f}(q))] = f(\pi_{Q,M}(q))$ by the definition of the action β . Then $\bar{f} : Q \rightarrow \mathbb{R}^{n+1}$ is well defined and, for each $q = (x, \hat{x}; A)$, there are unique $X|_q \in T|_x M$, $r(q) \in \mathbb{R}$ such that

$$\bar{f}(q) = AX|_q + r(q)\hat{x}.$$

The maps $q \mapsto X|_q$ and $q \mapsto r(q)$ are smooth. We show that the vector $X|_q$ and the real number $r(q)$ depend only on x and hence define a vector field and a function on M . One has $[(x, \hat{x}; A), v] = [(x, \hat{y}; B), w]$ if and only if there is $C \in G(n, k)$ such that $C\hat{x} = \hat{y}$, $CA = B$ and $Cv = w$. This means that $C|_{\text{im}A} = BA^{-1}|_{\text{im}A} : T|_{\hat{x}}\hat{M}_k \rightarrow T|_{\hat{y}}\hat{M}_k$ (with $\text{im}A$ denoting the image of A) and this defines C uniquely as an element of $G(n, k)$ and also, by the definition of \bar{f} , $C\bar{f}(x, \hat{x}, A) = \bar{f}(x, \hat{y}, B)$. Therefore,

$$BX|_{(x, \hat{y}; B)} + r(x, \hat{y}; B)\hat{y} = C(AX|_{(x, \hat{x}; A)} + r(x, \hat{x}; A)\hat{x}) = BX|_{(x, \hat{x}; A)} + r(x, \hat{x}; A)\hat{y},$$

which shows that $X|_{(x, \hat{y}; B)} = X|_{(x, \hat{x}; A)}$, $r(x, \hat{y}; B) = r(x, \hat{x}; A)$ and proves the claim.

Hence for each $f \in \Gamma(\xi)$ there are unique $X_f \in \text{VF}(M)$ and $r_f \in C^\infty(M)$ such that

$$f(x) = [((x, \hat{x}; A), AX_f|_x + r_f(x)\hat{x})],$$

(here the right hand side does not depend on the choice of $(x, \hat{x}; A) \in \pi_{Q,M}^{-1}(x)$).

Conversely, given $X \in \text{VF}(M)$, $r \in C^\infty(M)$ we may define $f_{(X,r)} \in \Gamma(\xi)$ by

$$f_{(X,r)}(x) = [((x, \hat{x}; A), AX|_x + r(x)\hat{x})],$$

where the right hand side does not depend on the choice of $(x, \hat{x}; A) \in \pi_{Q,M}^{-1}(x)$.

Clearly, for $f \in \Gamma(\xi)$, one has $f_{(X_f, r_f)} = f$ and, for $(X, r) \in \text{VF}(M) \times C^\infty(M)$, one has $(X_{f_{(X,r)}}, r_{f_{(X,r)}}) = (X, r)$. This proves that the map defined by

$$\begin{aligned} \Gamma(\xi) &\rightarrow \text{VF}(M) \times C^\infty(M) \\ f &\mapsto (X_f, r_f) \end{aligned}$$

is a bijection. It is easy to see that it is actually a $C^\infty(M)$ -module homomorphism. Since $C^\infty(M)$ -modules $\Gamma(\xi)$ and $\text{VF}(M) \times C^\infty(M)$ are isomorphic and since $\text{VF}(M) \times C^\infty(M)$ is obviously isomorphic, as a $C^\infty(M)$ -module, to $\Gamma(\pi_{TM \oplus \mathbb{R}})$, it follows that ξ and $\pi_{TM \oplus \mathbb{R}}$ are isomorphic vector bundles over M .

We now describe the connection ∇^{Rol} and the inner product structure h_k on ξ and we determine to which objects they correspond to in the isomorphic bundle $\pi_{TM \oplus \mathbb{R}}$.

By Section 2.1.3 in [11] and the above notation, one defines for $f \in \Gamma(\xi)$, $Y \in T|_x M$, $x \in M$

$$\nabla_Y^{\text{Rol}} f|_x := [((x, \hat{x}; A), \mathcal{L}_R(Y)|_{(x, \hat{x}; A)} \bar{f})],$$

where $\bar{f} : Q \rightarrow \mathbb{R}^{n+1}$ is defined above and $\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} \bar{f}$ is defined componentwise (i.e., we let $\mathcal{L}_R(Y)|_{(x, \hat{x}; A)}$ to operate separately to each of the $n + 1$ component functions of \bar{f}). The definition does not depend on $(x, \hat{x}; A) \in \pi_{Q,M}^{-1}(x)$ as should be

evident from the above discussions. The inner product on ξ , on the other hand, is defined by

$$h_k([((x, \hat{x}; A), v)], [((x, \hat{y}; B), w)]) = g(X, Y) + krt,$$

where $v = AX + r\hat{x}$, $w = BY + t\hat{y}$. It is clear that h_k is well defined.

We slightly work out the expression for ∇^{Rol} . Let $f \in \Gamma(\xi)$, $Y \in T|_x M$, $x \in M$. Then $\bar{f}(y, \hat{y}, B) = BX_f|_y + r_f(y)\hat{y}$ where $X_f \in \text{VF}(M)$, $r_f \in C^\infty(M)$,

$$\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} \bar{f} = \mathcal{L}_R(Y)|_{(x, \hat{x}; A)} ((y, \hat{y}; B) \mapsto BX_f|_y) + Y(r_f)\hat{x} + r_f(x)AY$$

and choosing some path γ on M such that $\dot{\gamma}(0) = Y$, then $\dot{q}_{\mathcal{D}_R}(\gamma, q)(0) = \mathcal{L}_R(Y)|_q$, where $q = (x, \hat{x}; A)$ and therefore

$$\begin{aligned} s_{n,k}(\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} ((y, \hat{y}; B) \mapsto BX_f|_y), \hat{x}) &= s_{n,k}\left(\frac{d}{dt}\Big|_0 (A_{\mathcal{D}_R}(\gamma, q)(t)X_f|_{\gamma(t)}), \hat{x}\right) \\ &= \frac{d}{dt}\Big|_0 s_{n,k}(A_{\mathcal{D}_R}(\gamma, q)(t)X_f|_{\gamma(t)}, \hat{\gamma}_{\mathcal{D}_R}(\gamma, q)(t)) - s_{n,k}(AX_f|_x, AY) \\ &= -\hat{g}_k(AX_f|_x, AY) = -g(X_f|_x, Y) = s_{n,k}\left(-\frac{1}{k}g(X_f|_x, Y)\hat{x}, \hat{x}\right). \end{aligned}$$

Therefore,

$$\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} ((y, \hat{y}; B) \mapsto BX_f|_y) + \frac{1}{k}g(X_f|_x, Y)\hat{x} \in T|_{\hat{x}} \hat{M}_k,$$

and we write

$$\begin{aligned} \mathcal{L}_R(Y)|_{(x, \hat{x}; A)} \bar{f} &= \left(\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} ((y, \hat{y}; B) \mapsto BX_f|_y) + \frac{1}{k}g(X_f|_x, Y)\hat{x} + r_f(x)AY \right) \\ &\quad + (Y(r_f) - \frac{1}{k}g(X_f|_x, Y))\hat{x}. \end{aligned}$$

Correspondingly, using the isomorphism of ξ and $\pi_{TM \oplus \mathbb{R}}$, to the connection ∇^{Rol} and the non-degenerate metric h_k on ξ , there is a connection ∇^{Rol} and an indefinite metric h_k (with the same names as the ones on ξ) on $\pi_{TM \oplus \mathbb{R}}$ such that for $X \in \text{VF}(M)$, $r \in C^\infty(M)$ and $Y \in T|_x M$,

$$\begin{aligned} \nabla_Y^{\text{Rol}}(X, r) &= \left(A^{-1}(\mathcal{L}_R(Y)|_{(x, \hat{x}; A)} ((y, \hat{y}; B) \mapsto BX|_y) + \frac{1}{k}g(X|_x, Y)\hat{x}) + r(x)Y, \right. \\ &\quad \left. Y(r) - \frac{1}{k}g(X|_x, Y) \right), \end{aligned} \tag{76}$$

where $(x, \hat{x}; A) \in Q$ is arbitrary point of Q over x and

$$h_k((X, r), (Y, s)) = g(X, Y) + krs,$$

for $X, Y \in T|_x M$, $r, s \in \mathbb{R}$.

We will now prove the metric property (75) of the connection ∇^{Rol} . This will be done in the case of the bundle $\pi_{TM \oplus \mathbb{R}}$ but it gives the equivalent result on ξ .

If $(X, r), (Y, s) \in \Gamma(\pi_{TM \oplus \mathbb{R}})$ and $Z \in T|_x M$ then

$$\begin{aligned} Z(h_k((X, r), (Y, s))) &= Z(g(X, Y) + krs) \\ &= g(\nabla_Z X, Y|_x) + g(X|_x, \nabla_Z Y) + kZ(r)s(x) + kr(x)Z(s). \end{aligned}$$

On the other hand,

$$h_k(\nabla_Z^{\text{Rol}}(X, r), (Y, s)) = s_{n,k} \left(\mathcal{L}_R(Z)|_{(x,\hat{x};A)}((y, \hat{y}; B) \mapsto BX|_y) \right. \\ \left. + \frac{1}{k}g(X|_x, Z)\hat{x} + r(x)AZ, AY \right) + k(Z(r) - \frac{1}{k}g(X|_x, Z))s(x),$$

for any $q = (x, \hat{x}; A) \in \pi_{Q,M}^{-1}(x)$ and choosing a path γ s.t. $\dot{\gamma}(0) = Z$ we get

$$h(\nabla_Z^{\text{Rol}}(X, r), (Y, s)) = s_{n,k} \left(\frac{d}{dt} \Big|_0 (A_{\mathcal{D}_R}(\gamma, q)(t)X|_{\gamma(t)}, AY) \right. \\ \left. + r(x)g(Z, Y|_x) + (kZ(r) - g(X|_x, Z))s(x), \right)$$

from which we finally get

$$h_k(\nabla_Z^{\text{Rol}}(X, r), (Y, s)) + h_k((X, r), \nabla_Z^{\text{Rol}}(Y, s)) \\ = s_{n+1} \left(\frac{d}{dt} \Big|_0 (A_{\mathcal{D}_R}(\gamma, q)(t)X|_{\gamma(t)}, AY) + s_{n+1}(AX, \frac{d}{dt} \Big|_0 (A_{\mathcal{D}_R}(\gamma, q)(t)Y|_{\gamma(t)})) \right. \\ \left. + r(x)g(Z, Y|_x) + (kZ(r) - g(X|_x, Z))s(x) + s(x)g(Z, X|_x) \right. \\ \left. + (kZ(s) - g(Y|_x, Z))r(x) \right) \\ = \frac{d}{dt} \Big|_0 s_{n+1}(A_{\mathcal{D}_R}(\gamma, q)(t)X|_{\gamma(t)}, A_{\mathcal{D}_R}(\gamma, q)(t)Y|_{\gamma(t)}) + kZ(r)s(x) + kr(x)Z(s) \\ = \frac{d}{dt} \Big|_0 g(X|_{\gamma(t)}, Y|_{\gamma(t)}) + kZ(r)s(x) + kr(x)Z(s) \\ = g(\nabla_Z X, Y|_x) + g(X|_x, \nabla_Z Y) + kZ(r)s(x) + kr(x)Z(s),$$

which is exactly $Z(h_k((X, r), (Y, s)))$.

Motivated by Eq. (76), we make the following definition. If $Y \in T|_x M$ and $X \in \text{VF}(M)$ then define

$$\tilde{\nabla}_Y^{\text{Rol}} X := A^{-1}(\mathcal{L}_R(Y)|_{(x,\hat{x};A)}((y, \hat{y}; B) \mapsto BX|_y) + \frac{1}{k}g(X|_x, Y)\hat{x}),$$

where $(x, \hat{x}; A)$ is an arbitrary point on the fiber $\pi_{Q,M}^{-1}(x)$ over x . It is easily seen that it is \mathbb{R} -linear in X and Y and, for $f \in C^\infty(M)$,

$$\tilde{\nabla}_Y^{\text{Rol}}(fX) = Y(f)X|_x + f(x)\tilde{\nabla}_Y^{\text{Rol}} X,$$

so $\tilde{\nabla}^{\text{Rol}}$ is a connection on M . Moreover, from the above computations, we see that $\tilde{\nabla}^{\text{Rol}}$ is a metric connection with respect to g i.e., for $X, Y \in \text{VF}(M)$ and $Z \in T|_x M$,

$$Z(g(X, Y)) = g(\tilde{\nabla}_Z^{\text{Rol}} X, Y) + g(X, \tilde{\nabla}_Z^{\text{Rol}} Y).$$

We will prove that $\tilde{\nabla}^{\text{Rol}} = \nabla$ i.e., that $\tilde{\nabla}^{\text{Rol}}$ is the Levi-Civita connection of g . To do this, we show that the connection $\tilde{\nabla}^{\text{Rol}}$ is torsion-free.

Let $X, Y \in \text{VF}(M)$, $x \in M$. Then taking any $q = (x, \hat{x}; A) \in \pi_{Q,M}^{-1}(x)$ and any local smooth π_Q -section \tilde{A} such that $\tilde{A}|_x = A$ and $\overline{\nabla} A|_x = 0$, we compute

$$(\tilde{\nabla}_X^{\text{Rol}} Y - \tilde{\nabla}_Y^{\text{Rol}} X)|_x = A^{-1}(\mathcal{L}_R(X)|_{(x,\hat{x};A)}((y, \hat{y}; B) \mapsto BY|_y) + g(X|_x, Y|_x)\hat{x}) \\ - A^{-1}(\mathcal{L}_R(Y)|_{(x,\hat{x};A)}((y, \hat{y}; B) \mapsto BX|_y) + g(X|_x, Y|_x)\hat{x}) \\ = A^{-1}(\overline{\nabla}_{(Y, AY)}(\tilde{A}X) - \overline{\nabla}_{(Y, AY)}(\tilde{A}X)) \\ = (\nabla_X Y - \nabla_Y X)|_x = [X, Y]|_x.$$

Since $\tilde{\nabla}^{\text{Rol}}$ is a torsion-free metric connection w.r.t. g on M , it follows by uniqueness of Levi-Civita connection that

$$\tilde{\nabla}^{\text{Rol}} = \nabla.$$

Thus if $X \in \text{VF}(M)$, $r \in C^\infty(M)$ and $Y \in T|_x M$,

$$\nabla_Y^{\text{Rol}}(X, r) = \left(\nabla_Y X + r(x)Y, Y(r) - \frac{1}{k}g(X|_x, Y) \right).$$

This concludes the proof of Theorem 6.11. \square

Remark 6.12 Define a number δ_{ij}^k for $i, j = 1, \dots, n+1$ as follows,

$$\delta_{ij}^k := \begin{cases} 0, & i \neq j, \\ 1, & i = j = 1, \dots, n, \\ k, & i = j = n+1. \end{cases}$$

We say that a frame $(X_i, t_i)_{i=1}^{n+1}$ of $T|_x M \oplus \mathbb{R}$ is h_k -orthonormal if $h_k((X_i, t_i), (X_j, t_j)) = \delta_{ij}^k$. We may build the manifold $F_{\text{OON}}^{h_k}(\pi_{TM \oplus \mathbb{R}})$ of h_k -orthonormal frames in the standard way.

Now we will prove that the bundle $F_{\text{OON}}^{h_k}(\pi_{TM \oplus \mathbb{R}})$ of h_k -orthonormal frames of $\pi_{TM \oplus \mathbb{R}}$ is isomorphic to $\pi_{Q, M}$ as a bundle over M . The isomorphism $\Phi_k : \pi_{Q, M} \rightarrow F_{\text{OON}}^{h_k}(\pi_{TM \oplus \mathbb{R}})$ can be described as follows. Let $(x, \hat{x}; A) \in Q$. Then there are unique $(X_i, t_i) \in T|_x M \oplus \mathbb{R}$, $i = 1, \dots, n+1$ such that $e_i = AX_i + t_i \hat{x}$ where e_i , $i = 1, \dots, n+1$, is the standard basis of \mathbb{R}^{n+1} . One easily computes that

$$\begin{aligned} h_k((X_i, t_i), (X_j, t_j)) &= g(X_i, X_j) + kt_i t_j = s_{n,k}(AX_i, AX_j) + s_{n,k}(t_i \hat{x}, t_j \hat{x}) \\ &= s_{n,k}(e_i, e_j) = \delta_{ij}^k, \end{aligned}$$

since $s_{n,k}(AX_i, t_j \hat{x}) = 0$, $s_{n,k}(t_i \hat{x}, AX_j) = 0$. Thus define $\Phi(x, \hat{x}; A) := (X_i, t_i)_{i=1}^{n+1}$.

We will give a description the inverse map Φ^{-1} . Let $(X_i, t_i)_{i=1}^{n+1} \in F_{\text{OON}}^{h_k}(\pi_{TM \oplus \mathbb{R}})$. Then there are unique $a_i \in \mathbb{R}$ such that $\sum_{i=1}^{n+1} a_i (X_i, t_i) = (0, 1)$. We notice that $a_i = kt_i$ for all $i = 1, \dots, n$ and $a_{n+1} = t_{n+1}$, since

$$0 = g\left(\sum_{i=1}^{n+1} a_i X_i, X_j\right) = \sum_{i=1}^{n+1} a_i (\delta_{ij}^k - kt_i t_j),$$

and because $\sum_{i=1}^{n+1} a_i t_i = 1$. Hence $k \sum_{i=1}^n t_i^2 + t_{n+1}^2 = 1$. Define $\hat{x} := \sum_{i=1}^n (kt_i) e_i + t_{n+1} e_{n+1}$ for which $s_{n,k}(\hat{x}, \hat{x}) = k(k \sum_{i=1}^n t_i^2 + t_{n+1}^2) = k$ i.e., $\hat{x} \in \hat{M}_k$. Moreover, it is easy to see that each $e_i - t_i \hat{x}$ is $s_{n,k}$ -orthogonal to \hat{x} and hence we may define $A : T|_x M \rightarrow T|_{\hat{x}} \hat{M}_k$ by requiring that $AX_i = e_i - t_i \hat{x}$, $i = 1, \dots, n+1$. It can be shown that A is well defined by this formula and an orthogonal linear map i.e., $(x, \hat{x}; A) \in Q$. Also, evidently $\Phi(x, \hat{x}; A) = (X_i, t_i)_{i=1}^{n+1}$.

6.3 Controllability Results for Rolling Against a Non-Flat Space Form

It is now clear, thanks to Theorem 6.11, that the controllability of the rolling problem of a manifold M against a space form \hat{M}_k amounts to checking whether the

connection ∇^{Rol} of $\pi_{TM \oplus \mathbb{R}}$ has full holonomy or not i.e., whether $H^{\nabla^{\text{Rol}}} = G(n, k)$ or not.

For the rest of the section, we assume that k only takes the values 1 and -1 , and for notational purposes, we use the letter "c" instead of "k" and thus $c \in \{+1, -1\}$.

In Riemannian geometry, the reducibility of the Riemannian holonomy group is characterized (in the complete simply connected case) by the de Rham Theorem (see [25]). We aim at giving an analog of this result w.r.t. ∇^{Rol} in Theorem 6.14 below. Before doing so, we first prove a simpler result showing that the conclusion of Theorem 6.14 below is not trivial.

Proposition 6.13 Suppose that (M, g) is a space form of constant curvature equal to $c \in \{+1, -1\}$. Then the rolling connection ∇^{Rol} defined by the rolling problem (R) of (M, g) against $(\hat{M}_c, s_{n,c})$ (i.e., we roll (M, g) against itself) is reducible and, for each $x \in M$, the irreducible subspaces of the action of the holonomy group $H^{\nabla^{\text{Rol}}}|_x$ on $T|_x M \oplus \mathbb{R}$ are all 1-dimensional.

Proof. Let (p^1, \dots, p^{n+1}) be the canonical chart of \mathbb{R}^{n+1} where p^j is the projection onto the j -th factor and write $h = h_c$ for the inner product in $TM \oplus \mathbb{R}$. We will assume that the space form M is the subset \hat{M}_c of \mathbb{R}^{n+1} as defined previously. Define a vector field $Z := \sum_{i=1}^{n+1} p^i \frac{\partial}{\partial p^i}$ i.e., Z is equal to the half of the gradient in $(\mathbb{R}^{n+1}, s_{n,c})$ of the function $(p^1)^2 + \dots + (p^n)^2 + c(p^{n+1})^2$. Notice that Z is $s_{n,c}$ -orthogonal to the submanifold $M = \hat{M}_c$ of \mathbb{R}^{n+1} and hence $T|_x M$ is the $s_{n,c}$ -orthogonal complement of $Z|_x$ for $x \in M$. Moreover, $s_{n,c}(Z, Z) = c$.

Next we define, for $j = 1, \dots, n+1$, the vector fields

$$Y_j := \frac{\partial}{\partial p^j} - cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z \right) Z$$

and functions

$$r^j(x) = cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right).$$

The restrictions of Z, Y_j, r^j onto M will be denoted by the same letters. Notice that (Y_j, r^j) , $j = 1, \dots, n+1$ are h -orthogonal at each point of M and hence they form a global orthogonal frame of $\pi_{TM \oplus \mathbb{R}}$.

Denote, as usual, by ∇ the Levi-Civita connection of (M, g) . Take any vector fields $X = \sum_{i=1}^{n+1} X^i \frac{\partial}{\partial p^i} \in \text{VF}(M)$ and $U = \sum_{i=1}^{n+1} U^i \frac{\partial}{\partial p^i} \in \text{VF}(M)$ and let \tilde{U} be some extension of U onto a neighbourhood of M in \mathbb{R}^{n+1} with corresponding components \tilde{U}^i . Then we have for $x \in M$,

$$\nabla_X U|_x = \tilde{U}_*(X|_x) - cs_{n,c}(\tilde{U}_*(X|_x), Z|_x)Z|_x.$$

where we understand \tilde{U}_* as a map $T\mathbb{R}^{n+1} \rightarrow T\mathbb{R}^{n+1}$ using the obvious isomorphisms $T|_X(T\mathbb{R}^{n+1}) \rightarrow T|_x\mathbb{R}^{n+1}$ for each $X \in T|_x\mathbb{R}^{n+1}$.

Then we compute for any $x \in M$ and $X = \sum_{i=1}^{n+1} X^i \frac{\partial}{\partial p^i} \in T|_x M$

$$Z_*(X) = \sum_{i=1}^{n+1} X^i Z_* \left(\frac{\partial}{\partial p^i} \right) = \sum_{i=1}^{n+1} X^i \frac{\partial}{\partial p^i} = X,$$

and (notice that $(\frac{\partial}{\partial p^j})_* = 0$)

$$\begin{aligned} (Y_j)_*(X) &= \sum_{i=1}^{n+1} X^i (Y_j)_* \left(\frac{\partial}{\partial p^i} \right) = -cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z_*(X) \right) Z|_x - cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) Z_*(X) \\ &= -cs_{n,c} \left(\frac{\partial}{\partial p^j}, X \right) Z|_x - cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) X, \end{aligned}$$

from which we get

$$\begin{aligned} \nabla_X Y_j &= (Y_j)_*(X|_x) - cs_{n,c} \left((Y_j)_*(X|_x), Z|_x \right) Z|_x \\ &= -cs_{n,c} \left(\frac{\partial}{\partial p^j}, X \right) Z|_x - cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) X + cs_{n,c} \left(\frac{\partial}{\partial p^j}, X \right) Z|_x \\ &+ s_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) \underbrace{s_{n,c}(X, Z|_x)}_{=0} Z|_x = -cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) X. \end{aligned}$$

Moreover,

$$\begin{aligned} X(r^j) &= cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z_*(X) \right) = cs_{n,c} \left(\frac{\partial}{\partial p^j}, X \right) \\ s_{n,c}(X, Y_j) &= s_{n,c} \left(X, \frac{\partial}{\partial p^j} \right) - \underbrace{cs_{n,c}(X, Z|_x)}_{=0} s_{n,c} \left(Z, \frac{\partial}{\partial p^j} \right) = s_{n,c} \left(X, \frac{\partial}{\partial p^j} \right), \end{aligned}$$

and thus

$$\begin{aligned} \nabla_X^{\text{Rol}}(Y_j, r^j) &= (\nabla_X Y_j + r^j(x)X, X(r^j) - \frac{1}{c} s_{n,c}(X, Y_j|_x)) \\ &= \left(-cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) X + cs_{n,c} \left(\frac{\partial}{\partial p^j}, Z|_x \right) X, cs_{n,c} \left(\frac{\partial}{\partial p^j}, X \right) - cs_{n,c} \left(X, \frac{\partial}{\partial p^j} \right) \right) \\ &= (0, 0). \end{aligned}$$

This means that all the $\pi_{TM \oplus \mathbb{R}}$ -sections (Y_j, r^j) , $j = 1, \dots, n+1$ are ∇^{Rol} -parallel globally. In particular, for any $x \in M$ and loop $\gamma \in \Omega_x(M)$,

$$\frac{d}{dt} (P^{\nabla^{\text{Rol}}})_t^0(\gamma) \left((Y_j, p^j)|_{\gamma(t)} \right) = (P^{\nabla^{\text{Rol}}})_t^0(\gamma) \nabla_{\dot{\gamma}(t)} (Y_j, p^j) = 0,$$

which means that $(x = \gamma(0))$

$$(Y_j, p^j)|_{\gamma(t)} = (P^{\nabla^{\text{Rol}}})_0^t(\gamma) (Y_j, p^j)|_x, \forall t,$$

and hence

$$(P^{\nabla^{\text{Rol}}})_0^t(\gamma) (Y_j, p^j)|_x = (Y_j, p^j)|_{\gamma(1)} = (Y_j, p^j)|_x,$$

i.e., that the 1-dimensional subbundles spanned by each (Y_j, r^j) are invariant under the holonomy group of ∇^{Rol} . Thus we have proved what we claimed. \square

Below we will only consider the case of positive curvature $c = +1$ i.e., rolling against the unit sphere.

Theorem 6.14 Let (M, g) be a complete Riemannian manifold and (\hat{M}_1, s_{n+1}) be the unit sphere with the metric induced from the Euclidean metric of \mathbb{R}^{n+1} . If the rolling connection ∇^{Rol} (see (71)) corresponding to rolling of (M, g) against (\hat{M}_1, s_{n+1}) is reducible, then (\hat{M}_1, s_{n+1}) is a Riemannian covering of (M, g) .

Recall that the reducibility of the connection ∇^{Rol} means that its holonomy group, which is a subgroup of $G(n, c)$, is reducible i.e., there exists two nontrivial invariant subspaces $V_1, V_2 \notin \{\{0\}, \mathbb{R}^{n+1}\}$ of \mathbb{R}^{n+1} which are invariant by the action of this group.

Proof. In this case we have $c = +1$ (corresponding to the sphere space form) and we will write $h = h_1$ for the inner product on $TM \oplus \mathbb{R}$.

Fix once and for all a point $x_0 \in M$. The assumption that ∇^{Rol} is reducible means that there are two subspaces $V_1, V_2 \subset T|_{x_0}M \oplus \mathbb{R}$ which are nontrivial (i.e., $V_1, V_2 \notin \{\{0\}, T|_{x_0}M \oplus \mathbb{R}\}$) and invariant by the action of the holonomy group of ∇^{Rol} at x_0 . Since the holonomy group of ∇^{Rol} acts h -orthogonally on $T|_{x_0}M$, it follows that $V_1 \perp V_2$.

Define subbundles $\pi_{\mathcal{D}_j} : \mathcal{D}_j \rightarrow M$, $j = 1, 2$ of $\pi_{TM \oplus \mathbb{R}}$ such that for any $x \in M$ one chooses a piecewise C^1 curve $\gamma : [0, 1] \rightarrow M$ from x_0 to x and defines

$$\mathcal{D}_j|_x = (P^{\nabla^{\text{Rol}}})_0^1(\gamma)V_j, \quad j = 1, 2.$$

These definitions are independent of the chosen path γ since if ω is another such curve, then $\omega^{-1} \cdot \gamma \in \Omega_{x_0}(M)$ is a loop based at x_0 and hence by the invariance of V_j , $j = 1, 2$ under the holonomy of ∇^{Rol} ,

$$(P^{\nabla^{\text{Rol}}})_0^1(\gamma)V_j = (P^{\nabla^{\text{Rol}}})_0^1(\omega) \underbrace{(P^{\nabla^{\text{Rol}}})_0^1(\omega^{-1} \cdot \gamma)V_j}_{=V_j} = (P^{\nabla^{\text{Rol}}})_0^1(\omega)V_j.$$

Moreover, since parallel transport $(P^{\nabla^{\text{Rol}}})_0^1(\gamma)$ is an h -orthogonal map, it follows that $\mathcal{D}_1 \perp \mathcal{D}_2$ w.r.t the vector bundle metric h .

It is a standard fact that \mathcal{D}_j , $j = 1, 2$, are smooth embedded submanifolds of $TM \oplus \mathbb{R}$ and that the restriction of $\pi_{TM \oplus \mathbb{R}}$ to \mathcal{D}_j defines a smooth subbundle $\pi_{\mathcal{D}_j}$ as claimed. Moreover, it is clear that

$$\pi_{\mathcal{D}_1} \oplus \pi_{\mathcal{D}_2} = \pi_{TM \oplus \mathbb{R}},$$

and this sum is h -orthogonal.

We will now assume that both \mathcal{D}_j , $j = 1, 2$, have dimension at least 2. The case where one of them has dimension = 1 can be treated in a similar fashion and will be omitted. So we let $m+1 = \dim \mathcal{D}_1$ where $m \geq 1$ and then $n-m = (n+1) - (m+1) = \dim \mathcal{D}_2 \geq 2$ i.e., $1 \leq m \leq n-2$. Define for $j = 1, 2$

$$\mathcal{D}_j^M = \text{pr}_1(\mathcal{D}_j) = \{X \mid (X, r) \in \mathcal{D}_j\} \subset TM,$$

and

$$N_j = \{x \in M \mid (0, 1) \in \mathcal{D}_j|_x\} \subset M.$$

Trivially, $N_1 \cap N_2 = \emptyset$. Also, N_j , $j = 1, 2$, are closed subsets of M since they can be written as $N_j = \{x \in M \mid p_j^\perp(T|_x) = T|_x\}$ where $p_j^\perp : TM \oplus \mathbb{R} \rightarrow \mathcal{D}_j$ is the

h -orthogonal projection onto \mathcal{D}_j and T is the (smooth) constant section $x \mapsto (0, 1)$ of $\pi_{TM \oplus \mathbb{R}}$.

We next briefly sketch the rest of proof. We will show that N_j are nonempty totally geodesic submanifolds of M and, for any given $x_j \in N_j$, $j = 1, 2$, that (M, g) is locally isometric to the sphere

$$S = \{(X_1, X_2) \in T|_{x_1}^\perp N_1 \oplus T|_{x_2}^\perp N_2 \mid \|X_1\|_g^2 + \|X_2\|_g^2 = 1\},$$

with the metric $G := (g|_{T|_{x_1}^\perp N_1} \oplus g|_{T|_{x_2}^\perp N_2})|_S$. Here \perp denotes the orthogonal complement inside $T|_x M$ w.r.t. g . Since (S, G) is isometric to the Euclidean sphere $(\hat{M}_1, s_{n,1})$ this would finish the argument. The latter is rather long and we decompose it in a sequence of ten lemmas and we start with the first one.

Lemma 6.15 The sets N_j , $j = 1, 2$, are non-empty.

Proof. Note first that $N_1 \cup N_2 \neq M$ since otherwise $N_1 = M \setminus N_2$ would be open and closed and similarly for N_2 . But then if, say, $N_1 \neq \emptyset$ we have $N_1 = M$ by connectedness of M i.e., the point $(0, 1) \in \mathcal{D}_1|_x$ for all $x \in M$. Then for all $x \in M$, $X \in \text{VF}(M)$ one has, by the invariance of \mathcal{D}_1 by the holonomy of ∇^{Rol} and by (71),

$$\mathcal{D}_1|_x \ni \nabla_{X|_x}^{\text{Rol}}(0, 1) = (X|_x, 0),$$

which implies that $\mathcal{D}_1 = TM \oplus \mathbb{R}$, a contradiction.

Let $x' \in M \setminus (N_1 \cup N_2)$ be arbitrary. Choose a basis $(X_0, r_0), \dots, (X_m, r_m)$ of $\mathcal{D}_1|_{x'}$. Then at least one of the numbers r_0, \dots, r_m is non-zero, since otherwise one would have $(X_i, r_i) = (X_i, 0) \perp (0, 1)$ for all i and thus $\mathcal{D}_1|_{x'} \perp (0, 1)$ i.e., $(0, 1) \in \mathcal{D}_2|_{x'}$ i.e., $x' \in N_2$ which is absurd. We assume that it is r_0 which is non-zero. By taking appropriate linear combinations of (X_i, r_i) , $i = 0, \dots, m$ (and by Gram-Schmidt's process), one may change the basis (X_i, r_i) , $i = 0, \dots, m$, of $\mathcal{D}_1|_x$ so that $r_1, \dots, r_m = 0$, $r_0 \neq 0$ and that $(X_0, r_0), (X_1, 0), \dots, (X_m, 0)$ are h -orthonormal. Also, X_0, \dots, X_m are non-zero: for X_1, \dots, X_m this is evident, and for X_0 it follows from the fact that if $X_0 = 0$, then $r_0 = 1$ and hence $x' \in N_1$, which contradicts our choice of x' .

Now let $\gamma : \mathbb{R} \rightarrow M$ be the unit speed geodesic with $\gamma(0) = x'$, $\dot{\gamma}(0) = \frac{X_0}{\|X_0\|_g}$. Parallel translate (X_i, r_i) along γ by ∇^{Rol} to get $\pi_{\mathcal{D}_1}$ -sections $(X_i(t), r_i(t))$ along γ . In particular, from (73) one gets

$$\ddot{r}_i(t) + r_i(t) = 0,$$

with $r_0(0) \neq 0$, $r_1(0) = \dots = r_m(0) = 0$. From the second equation in (72) one obtains $\dot{r}_i(0) = g(\dot{\gamma}(0), X_i(0)) = \|X_0\|_g^{-1} g(X_0, X_i)$ and thus $\dot{r}_i(0) = 0$ for $i = 1, \dots, m$ since $(X_i, 0)$ is h -orthogonal to (X_0, r_0) . Moreover, $\dot{r}_0(0) = \|X_0\|_g$. Hence $r_i(t) = 0$ for all t and $i = 1, \dots, m$ and $r_0(t) = \|X_0\|_g \sin(t) + r_0 \cos(t)$. In particular, at $t = t_0 := \arctan(-\frac{r_0}{\|X_0\|_g})$ one has $r_i(t_0) = 0$ for all $i = 0, \dots, m$ which implies that $\mathcal{D}_1|_{\gamma(t_0)} \perp (0, 1)$ i.e., $\gamma(t_0) \in N_2$. This proves that N_2 is non-empty. The same argument with \mathcal{D}_1 and \mathcal{D}_2 interchanged shows that N_1 is non-empty. \square

Lemma 6.16 For any $x \in M$ and any unit vector $u \in T|_x M$,

$$(P^{\nabla^{\text{Rol}}})_0^t(\dot{\gamma}_u)(0, 1) = (-\sin(t)\dot{\gamma}_u(t), \cos(t)). \quad (77)$$

Proof. Here and in what follows, $\gamma_u(t) := \exp_x(tu)$. Write

$$(X_0(t), r_0(t)) := (P^{\nabla^{\text{Rol}}})_0^t(\gamma_u)(0, 1).$$

The second equation in (72) implies that $\dot{r}_0(0) = g(\dot{\gamma}_u(0), X_0(0)) = g(u, 0) = 0$ and, since $r_0(0) = 1$, the second equation in (73) gives

$$r_0(t) = \cos(t).$$

Notice that, for all $t \in \mathbb{R}$,

$$\begin{aligned} & \nabla_{\dot{\gamma}_u(t)}(-\sin(t)\dot{\gamma}_u(t)) + r_0(t)\dot{\gamma}_u(t) \\ &= \nabla_{\dot{\gamma}_u(t)}(-\sin(t))\dot{\gamma}_u(t) - \sin(t)\nabla_{\dot{\gamma}_u(t)}\dot{\gamma}_u(t) + \cos(t)\dot{\gamma}_u(t) \\ &= -\cos(t)\dot{\gamma}_u(t) - 0 + \cos(t)\dot{\gamma}_u(t) = 0, \end{aligned}$$

i.e., $-\sin(t)\dot{\gamma}_u(t)$ solves the same first order ODE as $X_0(t)$, namely $\nabla_{\dot{\gamma}_u(t)}X_0 + r_0(t)\dot{\gamma}_u(t) = 0$ for all t by the first equation in (72). Moreover, since

$$(-\sin(t)\dot{\gamma}_u(t))|_{t=0} = 0 = X_0(0),$$

it follows that $X_0(t) = -\sin(t)\dot{\gamma}_u(t)$, which, combined with the fact that $r_0(t) = \cos(t)$ proven above, gives (77). \square

Lemma 6.17 The sets N_j , $j = 1, 2$, are complete, totally geodesic submanifolds of (M, g) and $\mathcal{D}_j^M|_x = T|_x N_j$, $\forall x \in N_j$, $j = 1, 2$.

Proof. We show this for N_1 . The same argument then proves the claim for N_2 . Let $x \in N_1$ and $u \in \mathcal{D}_1^M|_x$ a unit vector. Since $(0, 1) \in \mathcal{D}_1|_x$, Eq. (77) implies that

$$\mathcal{D}_1|_{\gamma_u(t)} \ni (P^{\nabla^{\text{Rol}}})_0^t(\gamma_u)(0, 1) = (-\sin(t)\dot{\gamma}_u(t), \cos(t))$$

Next notice that

$$\begin{aligned} \nabla_{\dot{\gamma}_u(t)}^{\text{Rol}}(\cos(t)\dot{\gamma}_u(t), \sin(t)) &= (-\sin(t)\dot{\gamma}_u(t) + \sin(t)\dot{\gamma}_u(t), \cos(t) - g(\dot{\gamma}_u(t), \cos(t)\dot{\gamma}_u(t))) \\ &= (0, 0), \end{aligned}$$

and hence, since $(\cos(t)\dot{\gamma}_u(t), \sin(t))|_{t=0} = (u, 0) \in \mathcal{D}_1|_x$ (this is so because $u \in \mathcal{D}_1^M|_x$, hence there is some $r \in \mathbb{R}$ such that $(u, r) \in \mathcal{D}_1|_x$ and since $(0, 1) \in \mathcal{D}_1|_x$ because $x \in N_1$, then $\mathcal{D}_1|_x \ni (u, r) - r(0, 1) = (u, 0)$), we have, for all $t \in \mathbb{R}$,

$$(\cos(t)\dot{\gamma}_u(t), \sin(t)) = (P^{\nabla^{\text{Rol}}})_0^t(u, 0) \in \mathcal{D}_1|_{\gamma_u(t)}.$$

Hence for all $t \in \mathbb{R}$,

$$\mathcal{D}_1|_{\gamma_u(t)} \ni \sin(t)(\cos(t)\dot{\gamma}_u(t), \sin(t)) + \cos(t)(-\sin(t)\dot{\gamma}_u(t), \cos(t)) = (0, 1).$$

This proves that any geodesic starting from a point of N_1 with the initial direction from \mathcal{D}_1^M stays in N_1 forever. Hence, once it has been shown that N_1 is a submanifold of M with tangent space $T|_x N_1 = \mathcal{D}_1^M|_x$ for all $x \in N_1$, then automatically N_1 is totally geodesic and complete.

Let $x \in N_1$. If one takes an open neighbourhood U of x and local $\pi_{\mathcal{D}_2}$ -sections $(X_{m+1}, r_{m+1}), \dots, (X_n, r_n)$ which form a basis of \mathcal{D}_2 over U , then it is clear that $N_1 \cap U = \{x \in U \mid r_{m+1}(x) = \dots = r_n(x) = 0\}$

Thus let $(X_{m+1}, r_{m+1}), \dots, (X_n, r_n) \in \mathcal{D}_2|_x$ be a basis of $\mathcal{D}_2|_x$. Choose $\epsilon > 0$ such that \exp_x is a diffeomorphism from $B_g(0, \epsilon)$ onto its image U_ϵ and define for $y \in U_\epsilon$, $j = m + 1, \dots, n$,

$$(X_j, r_j)|_y = (P^{\nabla^{\text{Rol}}})_0^1(\tau \mapsto \exp_x(\tau \exp_x^{-1}(y)))(X_j, r_j).$$

Then (X_j, r_j) are local $\pi_{\mathcal{D}_2}$ -sections and it is clear that

$$N_1 \cap U_\epsilon = \{y \in U_\epsilon \mid r_{m+1}(y) = \dots = r_n(y) = 0\}.$$

Moreover, from (72),

$$\nabla r_j|_x = X_j|_x, \quad j = m + 1, \dots, n,$$

which are linearly independent. Hence, by taking $\epsilon > 0$ possibly smaller, we may assume that the local vector fields ∇r_j , $j = m + 1, \dots, n$, are linearly independent on U_ϵ . But this means that $N_1 \cap U_\epsilon = \{y \in U_\epsilon \mid r_{m+1}(y) = \dots = r_n(y) = 0\}$ is a smooth embedded submanifold of U_ϵ with tangent space

$$\begin{aligned} T|_x N_1 &= \{X \in T|_x M \mid g(\nabla r_j, X) = 0, \quad j = m + 1, \dots, n\} \\ &= \{X \in T|_x M \mid g(X_j, X) = 0, \quad j = m + 1, \dots, n\} \\ &= \mathcal{D}_1^M|_x. \end{aligned}$$

Since $x \in N_1$ was arbitrary, this proves that N_1 is indeed an embedded submanifold of M and $T|_x N_1 = \mathcal{D}_1^M|_x$ for all $x \in N_1$. □

Lemma 6.18 Let $d_i(x) := d_g(N_i, x)$, $x \in M$. Then in the set where d_i is smooth,

$$(\nabla \cos(d_i(\cdot)), \cos(d_i(\cdot))) \in \mathcal{D}_i^M, \quad (78)$$

where ∇ is the gradient w.r.t g .

Proof. Let $x \in M \setminus N_1$. Choose $y \in N_1$, $u \in (T|_y N_1)^\perp$ such that $\gamma_u : [0, d_i(x)] \rightarrow M$ is the minimal normal unit speed geodesic from N_1 to x . Since $(0, 1) \in \mathcal{D}_1|_y$ (because $y \in N_1$), it follows that the parallel translate of $(0, 1)$ along γ_u stays in \mathcal{D}_1 which, in view of (77), gives

$$\begin{aligned} \mathcal{D}_1|_x \ni (P^{\nabla^{\text{Rol}}})_0^{d_1(x)}(\gamma_u)(0, 1) &= (-\sin(d_1(x))\dot{\gamma}_u(d_1(x)), \cos(d_1(x))) \\ &= (-\sin(d_1(x))\nabla(d_1(\cdot))|_x, \cos(d_1(x))) \\ &= (\nabla \cos(d_1(\cdot))|_x, \cos(d_1(x))), \end{aligned}$$

where the last two equalities hold true if x is not in the cut nor the conjugate locus of N_1 (nor is x in N_1 , by assumption). Working in the complement of these points, which is a dense subset of M and using a continuity argument, we may assure that the result holds true everywhere where d_i is smooth. The same argument proves the formula (78) for d_2 as well. □

Lemma 6.19 For every $Y \in \text{VF}(M)$, one has

$$g(R(Y, \nabla d_1(\cdot)) \nabla d_1(\cdot), Y) = g(Y, Y) - (\nabla_Y(d_1(\cdot)))^2, \quad (79)$$

wherever $d_1(\cdot)$ is smooth.

Proof. It is known (see [24]) that for any $Y, Z \in \text{VF}(M)$, $d_1(\cdot)$ satisfies a PDE

$$\begin{aligned} -g(R(Y|_y, \nabla d_1(y)) \nabla d_1(y), Z|_y) &= \text{Hess}^2(d_1(\cdot))(Y|_y, Z|_y) \\ &\quad + (\nabla_{\nabla d_1(y)} \text{Hess}(d_1(\cdot)))(Y|_y, Z|_y), \end{aligned}$$

for every $y \in M$ such that d_1 is smooth at y (and this is true in a dense subset of M). In particular, $y \notin N_1$. Also, since the set of points $y \in M$ where $\cos(d_1(y)) = 0$ or $\sin(d_1(y)) = 0$ is clearly Lebesgue zero-measurable, we may assume that $\cos(d_1(y)) \neq 0$ and $\sin(d_1(y)) \neq 0$.

Notice that $(X_0, r_0) := (\nabla \cos(d_1(\cdot)), \cos(d_1(\cdot)))$ belongs to \mathcal{D}_1 and has h -norm equal to 1. We may choose in a neighbourhood U of y vector fields $X_1, \dots, X_m \in \text{VF}(U)$ such that $(X_0, r_0), (X_1, 0), \dots, (X_m, 0)$ is an h -orthonormal basis of \mathcal{D}_1 over U . Assume also that (X_0, r_0) is smooth on U . This implies that there are smooth one-forms ω_j^i , $i, j = 0, \dots, m$ defined by (set here $r_1 = \dots = r_m = 0$)

$$\nabla_Y^{\text{Rol}}(X_i, r_i) = \sum_{i=0}^m \omega_i^j(Y)(X_j, r_j), \quad Y \in \text{VF}(M),$$

or, more explicitly,

$$\begin{cases} \nabla_Y X_j + r_j Y = \sum_{i=0}^m \omega_j^i(Y) X_i \\ Y(r_j) - g(Y, X_j) = \sum_{i=0}^m \omega_j^i(Y) r_i, \end{cases}$$

Since $(X_0, r_0), \dots, (X_m, r_m)$ are h -orthonormal, it follows that $\omega_j^i = -\omega_i^j$. The fact that $r_1 = \dots = r_m = 0$ implies that

$$-g(Y, X_j) = \omega_j^0(Y) r_0, \quad j = 1, \dots, m$$

i.e.,

$$\omega_0^j(Y) = \frac{g(Y, X_j)}{\cos(d_1(\cdot))}.$$

But then one has that (notice that $\omega_0^0 = 0$)

$$\nabla_Y X_0 + r_0 Y = \sum_{j=1}^m \omega_0^j(Y) X_j,$$

which simplifies to

$$\begin{aligned} & -\sin(d_1(\cdot)) \nabla_Y \nabla d_1(\cdot) - \cos(d_1(\cdot)) \nabla_Y(d_1(\cdot)) \nabla d_1(\cdot) \\ &= -\cos(d_1(\cdot)) Y + \frac{1}{\cos(d_1(\cdot))} \sum_{j=1}^m g(X_j, Y) X_j, \end{aligned}$$

or

$$\begin{aligned}\nabla_Y \nabla d_1(\cdot) &= -\cot(d_1(\cdot)) \nabla_Y(d_1(\cdot)) \nabla d_1(\cdot) + \cot(d_1(\cdot)) Y \\ &\quad - \frac{1}{\sin(d_1(\cdot)) \cos(d_1(\cdot))} \sum_{j=1}^m g(X_j, Y) X_j.\end{aligned}$$

Writing $S(Y) := \nabla_Y \nabla d_1(\cdot) = \text{Hess}(d_1(\cdot))(Y, \cdot)$, one obtains

$$\begin{aligned}(\nabla_{\nabla d_1(\cdot)} S)(Y) &= \nabla_{\nabla d_1(\cdot)}(S(Y)) - S(\nabla_{\nabla d_1(\cdot)} Y) \\ &= \frac{1}{\sin^2(d_1(\cdot))} \nabla_Y(d_1(\cdot)) \nabla d_1(\cdot) - \cot(d_1(\cdot)) g(\nabla_{\nabla d_1(\cdot)} Y, \nabla d_1(\cdot)) \nabla d_1(\cdot) \\ &\quad - \frac{1}{\sin^2(d_1(\cdot))} Y - \left(\frac{1}{\cos^2(d_1(\cdot))} - \frac{1}{\sin^2(d_1(\cdot))} \right) \sum_{j=1}^m g(Y, X_j) X_j \\ &\quad - \frac{1}{\sin(d_1(\cdot)) \cos(d_1(\cdot))} \sum_{j=1}^m (g(Y, \nabla_{\nabla d_1(\cdot)} X_j) X_j + g(Y, X_j) \nabla_{\nabla d_1(\cdot)} X_j) \\ &\quad + \cot(d_1(\cdot)) \underbrace{\nabla_{\nabla_{\nabla d_1(\cdot)} Y} d_1(\cdot)}_{=g(\nabla d_1(\cdot), \nabla_{\nabla d_1(\cdot)} Y)} \nabla d_1(\cdot),\end{aligned}$$

where we used that $\nabla_{\nabla d_1(\cdot)}(d_1(\cdot)) = g(\nabla d_1(\cdot), \nabla d_1(\cdot)) = 1$. On the other hand,

$$\begin{aligned}\text{Hess}^2(d_1(\cdot))(Y, \cdot) &= S^2(Y) = S(S(Y)) \\ &= S\left(-\cot(d_1(\cdot)) \nabla_Y(d_1(\cdot)) \nabla d_1(\cdot) + \cot(d_1(\cdot)) Y\right. \\ &\quad \left. - \frac{1}{\sin(d_1(\cdot)) \cos(d_1(\cdot))} \sum_{j=1}^m g(X_j, Y) X_j\right) \\ &= -\cot^2(d_1(\cdot)) \nabla_Y(d_1(\cdot)) \nabla d_1(\cdot) + \cot^2(d_1(\cdot)) Y - \frac{2}{\sin^2(d_1(\cdot))} \sum_{j=1}^m g(X_j, Y) X_j \\ &\quad + \frac{1}{\sin^2(d_1(\cdot)) \cos^2(d_1(\cdot))} \sum_{j=1}^m g(X_j, Y) X_j,\end{aligned}$$

where we used that $\nabla d_1(\cdot), X_1, \dots, X_m$ are g -orthonormal (recall that $X_0 = -\sin(d_1(\cdot)) \nabla d_1(\cdot)$).

Thus, for any $Y, Z \in \text{VF}(M)$, one has on U that

$$\begin{aligned}&-g(R(Y, \nabla d_1(\cdot)) \nabla d_1(\cdot), Z) \\ &= -g(Y, Z) + \left(\frac{1}{\sin^2(d_1(\cdot))} - \cot^2(d_1(\cdot)) \right) \nabla_Y(d_1(\cdot)) \nabla_Z(d_1(\cdot)) \\ &\quad - \frac{1}{\sin(d_1(\cdot)) \cos(d_1(\cdot))} \sum_{j=1}^m (g(Y, \nabla_{\nabla d_1(\cdot)} X_j) g(X_j, Z) + g(Y, X_j) g(\nabla_{\nabla d_1(\cdot)} X_j, Z)).\end{aligned}$$

We also set $Z = Y$ and hence get

$$\begin{aligned}-g(R(Y, \nabla d_1(\cdot)) \nabla d_1(\cdot), Y) &= -g(Y, Y) + \nabla_Y(d_1(\cdot)) \nabla_Y(d_1(\cdot)) \\ &\quad - \frac{2}{\sin(d_1(\cdot)) \cos(d_1(\cdot))} \sum_{j=1}^m g(Y, \nabla_{\nabla d_1(\cdot)} X_j) g(X_j, Y).\end{aligned}$$

Here

$$\begin{aligned}
& \sum_{j=1}^m g(Y, \nabla_{\nabla d_1(\cdot)} X_j) g(X_j, Y) = -\frac{1}{\sin(d_1(\cdot))} \sum_{j=1}^m g(Y, \nabla_{X_0} X_j) g(X_j, Y) \\
&= -\frac{1}{\sin(d_1(\cdot))} \sum_{j=1}^m g(Y, \sum_{i=1}^m \omega_j^i(X_0) X_i) g(X_j, Y) \\
&= -\frac{1}{\sin(d_1(\cdot))} \sum_{i,j=1}^m \underbrace{\omega_j^i(X_0)}_{(\star)_1} \underbrace{g(Y, X_i) g(X_j, Y)}_{(\star)_2} = 0,
\end{aligned}$$

where expression $(\star)_1$ is skew-symmetric in (i, j) while $(\star)_2$ is symmetric on (i, j) . Hence the sum is zero. We finally obtain

$$g(R(Y, \nabla d_1(\cdot)) \nabla d_1(\cdot), Y) = g(Y, Y) - (\nabla_Y(d_1(\cdot)))^2,$$

as claimed. It is clear that this formula now holds at every point of M where $d_1(\cdot)$ is smooth and for any $Y \in \text{VF}(M)$. In particular, if Y is a unit vector g -perpendicular to $\nabla d_1(\cdot)$ at a point y of M , then $\nabla_Y d_1(\cdot)|_y = g(\nabla d_1(\cdot)|_y, Y|_y) = 0$ and hence

$$\sec(Y, d_1(\cdot))|_y = +1.$$

□

Lemma 6.20 For every $x \in N_1$, a unit vector $u \in (T|_x N_1)^\perp$ and $v \in T|_x M$ with $v \perp u$,

$$\|(\exp_x)_*|_{tu}(v)\|_g = \left| \frac{\sin(t)}{t} \right| \|v\|_g, \quad t \in \mathbb{R}. \quad (80)$$

In particular, for all unit vectors $u_1, u_2 \in (T|_x N_1)^\perp$ one has

$$\exp_x(\pi u_1) = \exp_x(\pi u_2).$$

Proof. Let $Y_{u,v}(t) = \frac{\partial}{\partial s}|_0 \exp_x(t(u + sv))$ be the Jacobi field along $\gamma_u(t) = \exp_x(tu)$ such that $Y_{u,v}(0) = 0$, $\nabla_{\dot{\gamma}_u(0)} Y_{u,v} = v$. Since $v \perp u$, it follows from the Gauss lemma (see [25]) that $Y_{u,v}(t) \perp \dot{\gamma}_u(t)$ for all t . Moreover, the assumption $u \in (T|_x N_1)^\perp$ implies that $\nabla d_1(\cdot)|_{\gamma_u(t)} = \dot{\gamma}_u(t)$ and thus $\nabla_{Y_{u,v}(t)}(d_1(\cdot)) = g(\dot{\gamma}_u(t), Y_{u,v}(t)) = 0$.

By polarization, one may write (79) into the form

$$R(Z(t), \dot{\gamma}_u(t)) \dot{\gamma}_u(t) = Z(t) - g(Z(t), \dot{\gamma}_u(t)) \dot{\gamma}_u(t),$$

for any vector field Z along γ_u . In particular,

$$\nabla_{\dot{\gamma}_u} \nabla_{\dot{\gamma}_u} Y_{u,v} = -R(Y_{u,v}, \dot{\gamma}_u) \dot{\gamma}_u = -Y_{u,v},$$

since $g(Y_{u,v}(t), \dot{\gamma}_u(t)) = 0$ for all t . On the other hand, the vector field $Z(t) = \sin(t) P_0^t(\gamma_u) v$ satisfies along γ_u

$$\begin{aligned}
& \nabla_{\dot{\gamma}_u(t)} \nabla_{\dot{\gamma}_u} Z = -Z(t), \quad \forall t \\
& Z(0) = 0, \quad \nabla_{\dot{\gamma}_u} Z|_{t=0} = v,
\end{aligned}$$

i.e., the same initial value problem as $Y_{u,v}$. This implies that

$$Y_{u,v}(t) = \sin(t)P_0^t(\gamma_u)v, \quad (81)$$

from which we obtain (80) because $Y_{u,v}(t) = t(\exp_x)_*|_{tu}(v)$.

The last claim follows from the fact that the map $\exp_x|_S : S \rightarrow M$ where $S = \{u \in (T|_x N_1)^\perp \mid \|u\| = \pi\}$ is a constant map. Indeed, if $u \in S$, $v \in T|_u S$ and we identify v as an element of $T|_x M$ as usual, then by what we have just proved (note that $u = \pi \frac{u}{\|u\|_g}$),

$$\|(\exp_x)_*|_u(v)\|_g = \frac{\sin(\pi)}{\pi} \|v\|_g = 0.$$

Hence $\exp_x|_S$ has zero differential on all over S which is connected, since its dimension is $n - m - 1 \geq 1$ by assumption. Hence $\exp_x|_S$ is a constant map. \square

Lemma 6.21 For every $x \in N_1$ and unit normal vector $u \in (T|_x N_1)^\perp$, the geodesic $t \mapsto \gamma_u(t)$ meets N_2 exactly at $t \in (\mathbb{Z} + \frac{1}{2})\pi$. The same holds with the roles of N_1 and N_2 interchanged.

Proof. Let $x \in N_1$ and $u \in (T|_x N_1)^\perp$ be a unit vector normal vector to N_1 . For $(X, r) \in \mathcal{D}_1|_x$ define $(X(t), r(t)) = (P^{\nabla^{\text{Rol}}})_0^t(\gamma_u)(X, r)$. Then by (72), (73) we have (notice that $g(u, X) = 0$ since $u \in (T|_x N_1)^\perp = (\mathcal{D}_1^M|_x)^\perp$ and $X \in \mathcal{D}_1^M|_x$)

$$r(t) = r(0) \cos(t).$$

Hence, $(X(t), r(t))$ is h -orthogonal to $(0, 1)$ if and only if $r(t) = 0$ i.e., $r(0) \cos(t) = 0$. This proves that $(0, 1) \perp \mathcal{D}_1|_{\gamma_u(t)}$ i.e., $(0, 1) \in \mathcal{D}_2|_{\gamma_u(t)}$ i.e., $\gamma_u(t) \in N_2$ if and only if $t \in (\frac{1}{2} + \mathbb{Z})\pi$ (obviously, there is a vector $(X, r) \in \mathcal{D}_1|_x$ with $r \neq 0$). \square

Lemma 6.22 The submanifolds N_1, N_2 are isometrically covered by Euclidean spheres of dimensions m and $n - m$, respectively, and the fundamental groups of N_1 and N_2 are finite and have the same number of elements. More precisely, for any $x \in N_1$ define

$$S_x = \{u \in (T|_x N_1)^\perp \mid \|u\|_g = 1\},$$

equipped with the restriction of the metric $g|_x$ of $T|_x M$. Then

$$S_x \rightarrow N_2; \quad u \mapsto \exp_x\left(\frac{\pi}{2}u\right),$$

is a Riemannian covering. The same claim holds with N_1 and N_2 interchanged.

Proof. Denote by C_1 the component of N_1 containing x . We will show first that $C_1 = N_1$ i.e., N_1 is connected.

Let $y_1 \in N_1$. Since C_1 is a closed subset of M , there is a minimal geodesic γ_v in M from C_1 to y_1 with $\dot{\gamma}_v(0) = v$ a unit vector, $x_1 := \gamma_v(0) \in C_1$ and $\gamma_v(d) = y_1$, with $d := d_g(y_1, C_1)$. By minimality, $v \in (T|_{x_1} C_1)^\perp = (T|_{x_1} N_1)^\perp$. Hence by Lemma 6.21 the point $x_2 := \exp_{x_1}(\frac{\pi}{2}v) = \gamma_v(\frac{\pi}{2})$ belongs to N_2 . Since the set

$$S_{x_2} = \{u \in (T|_{x_2} N_2)^\perp \mid \|u\|_g = 1\}.$$

is connected (its dimension is $m \geq 1$, by assumption that we made before), Lemma 6.21 implies that $\exp_{x_2}(\frac{\pi}{2}S_{x_2})$ is contained in a single component C'_1 of N_1 . Writing $u := \dot{\gamma}_v(\frac{\pi}{2})$, we have $\pm u \in S_{x_2}$ so

$$C'_1 \ni \exp_{x_2}(-\frac{\pi}{2}u) = \exp_{x_2}\left(-\frac{\pi}{2}\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_1}(tv)\right) = \exp_{x_1}((\frac{\pi}{2}-t)v)|_{t=\frac{\pi}{2}} = x_1,$$

and since also $x_1 \in C_1$, it follow that $C'_1 = C_1$. But this implies that

$$\gamma_v(\pi) = \exp_{x_1}(\pi v) = \exp_{x_2}\left(\frac{\pi}{2}\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_1}(tv)\right) = \exp_{x_2}(\frac{\pi}{2}u) \in C_1.$$

It also follows from $u \in (T|_{x_2}N_2)^\perp$ that $\dot{\gamma}_v(\pi) = \frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_2}(tu) \in (T|_{\gamma_v(\pi)}N_1)^\perp$. Since $\exp_{x_2}((d-\frac{\pi}{2})u) = y_1 \in N_1$, Lemma 6.21 implies that $d-\frac{\pi}{2} \in (\frac{1}{2} + \mathbb{Z})\pi$, from which, since $d \geq 0$, we get $d \in \mathbb{N}_0\pi$, where $\mathbb{N}_0 = \{0, 1, 2, \dots\}$.

By taking $x'_2 = \gamma_v(\frac{3}{2}\pi) \in N_2$ we may show similarly that $\gamma_v(2\pi) \in C_1$ and by induction we get $\gamma_v(k\pi) \in C_1$ for every $k \in \mathbb{N}_0$. In particular, since $d \in \mathbb{N}_0\pi$, we get $y_1 = \gamma_v(d) \in C_1$. Since $y_1 \in N_1$ was arbitrary, we get $N_1 \subset C_1$ which proves the claim.

By repeating the argument with N_1 and N_2 interchanged, we see that N_2 is connected.

Eq. (80) shows that, taking $u \in S_x$ and $v \in T|_u S_x$, i.e., $v \perp u$, $v \perp T|_x N_1$,

$$\left\|\frac{d}{dt}\Big|_0 \exp_x\left(\frac{\pi}{2}(u+tv)\right)\right\|_g = \left\|(\exp_x)_* \Big|_{\frac{\pi}{2}u} \left(\frac{\pi}{2}v\right)\right\|_g = \|v\|_g.$$

This shows that $u \mapsto \exp_x(\frac{\pi}{2}u)$ is a local isometry $S_x \rightarrow N_2$. In particular, the image is open and closed in N_2 , which is connected, hence $u \mapsto \exp_x(\frac{\pi}{2}u)$ is onto N_2 . According to Proposition II.1.1 in [25], $u \mapsto \exp_x(\frac{\pi}{2}u)$ is a covering $S_x \rightarrow N_2$.

Similarly, for any $y \in N_2$ the map $S_y \rightarrow N_1$; $u \mapsto \exp_y(\frac{\pi}{2}u)$ is a Riemannian covering.

Finally, let us prove the statement about fundamental groups. Fix a point $x_i \in N_i$ and write $\phi_i(u) = \exp_{x_i}(\frac{\pi}{2}u)$, $i = 1, 2$, for maps $\phi_1 : S_{x_1} \rightarrow N_2$, $\phi_2 : S_{x_2} \rightarrow N_1$. The fundamental groups $\pi_1(N_1)$, $\pi_1(N_2)$ of N_1 , N_2 are finite since their universal coverings are the (normal) spheres S_{x_2} , S_{x_1} which are compact. Also, $\phi_1^{-1}(x_2)$ and $\phi_2^{-1}(x_1)$ are in one-to-one correspondence with $\pi_1(N_2)$ and $\pi_1(N_1)$ respectively.

Define $\Phi_1 : \phi_1^{-1}(x_2) \rightarrow S_{x_2}$; $\Phi_1(u) = -\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_1}(tu) \in S_{x_2}$ and similarly $\Phi_2 : \phi_2^{-1}(x_1) \rightarrow S_{x_1}$; $\Phi_2(u) = -\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_2}(tu) \in S_{x_1}$. Clearly, for $u \in \phi_1^{-1}(x_2)$,

$$\phi_2(\Phi_1(u)) = \exp_{x_2}\left(-\frac{\pi}{2}\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_1}(tu)\right) = \exp_{x_1}((\frac{\pi}{2}-t)u)|_{t=\frac{\pi}{2}} = x_1,$$

i.e., Φ_1 maps $\phi_1^{-1}(x_2) \rightarrow \phi_2^{-1}(x_1)$. Similarly Φ_2 maps $\phi_2^{-1}(x_1) \rightarrow \phi_1^{-1}(x_2)$. Finally, Φ_1 and Φ_2 are inverse maps to each other since for $u \in \phi_1^{-1}(x_2)$,

$$\Phi_2(\Phi_1(u)) = -\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_2}\left(-t\frac{d}{ds}\Big|_{\frac{\pi}{2}}\exp_{x_1}(su)\right) = -\frac{d}{dt}\Big|_{\frac{\pi}{2}}\exp_{x_1}((\frac{\pi}{2}-t)u) = u,$$

and similarly $\Phi_1(\Phi_2(u)) = u$ for $u \in \phi_2^{-1}(x_1)$. □

For the sake of simplicity, we will finish the proof of Theorem 6.14 under the assumption that N_2 is simply connected and indicate in Remark 6.25 following the proof how to handle the general case.

The fact that N_2 is simply connected is clearly equivalent to saying that

$$S_x \rightarrow N_2; \quad u \mapsto \exp_x\left(\frac{\pi}{2}u\right),$$

defined in Lemma 6.22 is an isometry for some (and hence every) $x \in N_1$. It then follows from Lemma 6.22 that N_1 is (simply connected and) isometric to a sphere as well.

We next get the following.

Lemma 6.23 Fix $x_i \in N_j$, $j = 1, 2$ and let

$$S_{x_1} = \{u \in (T|_{x_1}N_1)^\perp \mid \|u\|_g = 1\}, \quad S_{x_2} = \{u \in (T|_{x_2}N_2)^\perp \mid \|u\|_g = 1\},$$

the unit normal spheres to N_1, N_2 at x_1, x_2 respectively. Consider first the maps

$$\begin{aligned} f_1 : S_{x_1} &\rightarrow N_2 & f_2 : S_{x_2} &\rightarrow N_1 \\ f_1(u) &= \exp_{x_1}\left(\frac{\pi}{2}u\right) & f_2(v) &= \exp_{x_2}\left(\frac{\pi}{2}v\right), \end{aligned} \tag{82}$$

and the map w which associates to $(u, v) \in S_{x_1} \times S_{x_2}$ the unique element of $S_{f_2(v)}$ such that $\exp_{f_2(v)}\left(\frac{\pi}{2}w(u, v)\right) = f_1(u)$. Finally let

$$\begin{aligned} \Psi :]0, \frac{\pi}{2}[\times S_{x_1} \times S_{x_2} &\rightarrow M \\ \Psi(t, u, v) &= \exp_{f_2(v)}(tw(u, v)). \end{aligned} \tag{83}$$

Suppose that $\tilde{S} :=]0, \frac{\pi}{2}[\times S_{x_1} \times S_{x_2}$ is endowed with the metric \tilde{g} such that

$$\tilde{g}|_{(t,u,v)} = dt^2 + \sin^2(t)g|_{T|_uS_{x_1}} + \cos^2(t)g|_{T|_vS_{x_2}}.$$

Then Ψ is a local isometry.

Proof. We use G to denote the *geodesic vector field* on TM i.e., for $u \in TM$ we have

$$G|_u := \ddot{\gamma}_u(0) = \frac{d^2}{dt^2}\Big|_0 \exp_{\pi_{TM}(u)}(tu).$$

Then the projections on M by π_{TM} of its integral curves are geodesics. Indeed, first we notice that

$$G|_{\dot{\gamma}_u(t)} = \frac{d^2}{ds^2}\Big|_0 \exp_{\gamma_u(t)}(s\dot{\gamma}_u(t)) = \frac{d^2}{ds^2}\Big|_0 \gamma_u(t+s) = \ddot{\gamma}_u(t),$$

and hence, if Γ be a curve on TM defined by $\Gamma(t) = \dot{\gamma}_u(t)$, then

$$\dot{\Gamma}(t) = \ddot{\gamma}_u(t) = G|_{\dot{\gamma}_u(t)} = G|_{\Gamma(t)},$$

and $\Gamma(0) = u$. Hence Γ satisfies the same initial value problem as $t \mapsto \Phi_G(t, u)$, which implies that

$$\Phi_G(t, u) = \dot{\gamma}_u(t), \quad \forall t \in \mathbb{R}, \quad u \in TM,$$

and in particular,

$$(\pi_{TM} \circ \Phi_G)(t, u) = \gamma_u(t), \quad \forall t \in \mathbb{R}, u \in TM.$$

For every $u \in TM$ there is a direct sum decomposition $H_u \oplus V_u$ of $T|_u TM$ where $V_u = V|_u(\pi_{TM})$ is the π_{TM} -vertical fiber over u and H_u is defined as

$$H_u = \left\{ \frac{d}{dt} \Big|_0 P_0^t(\gamma_X)u \mid X \in T|_{\pi_{TM}(u)} M \right\}.$$

We write the elements of $T|_u TM$ w.r.t. this direct sum decomposition as (A, B) where $A \in H_u$, $B \in V_u$. It can now be shown that (see [25] Lemma 4.3, Chapter II)

$$((\Phi_G)_t)_*|_u(A, B) = (Z_{(A,B)}(t), \nabla_{\dot{\gamma}_u(t)} Z_{(A,B)}), \quad (A, B) \in T|_u TM, \quad u \in TM,$$

where $Z_{(A,B)}$ is the unique Jacobi field along geodesic γ_u such that $Z_{(A,B)}(0) = A$, $\nabla_{\dot{\gamma}_u(0)} Z_{(A,B)} = B$.

We are now ready to prove the claim. First observe that

$$\Psi(t, u, v) = (\pi_{TM} \circ \Phi_G)(t, w(u, v))$$

and hence, for $(\frac{\partial}{\partial t}, X_1, X_2) \in T\tilde{S}$,

$$\begin{aligned} \Psi_*\left(\frac{\partial}{\partial t}, X_1, X_2\right) &= (\pi_{TM})_*\left(\frac{\partial}{\partial t} \Phi_G(t, w(u, v)) + ((\Phi_G)_t)_*|_{w(u,v)} w_*(X_1, X_2)\right) \\ &= (\pi_{TM})_*\left(G|_{\Phi_G(t, w(u, v))} + (Z_{w_*(X_1, X_2)}(t), \nabla_{\frac{\partial}{\partial t}(\pi_{TM} \circ \Phi_G)(t, w(u, v))} Z_{w_*(X_1, X_2)})\right) \\ &= \dot{\gamma}_{w(u,v)}(t) + Z_{w_*(X_1, X_2)}(t). \end{aligned}$$

On the other hand,

$$(\pi_{TM} \circ \Phi_G)\left(\frac{\pi}{2}, w(u, v)\right) = f_1(u),$$

from where

$$(f_1)_*|_u(X_1) = Z_{w_*(X_1, X_2)}\left(\frac{\pi}{2}\right).$$

Similarly, since

$$(\pi_{TM} \circ \Phi_G)(0, w(u, v)) = \pi_{TM}(w(u, v)) = f_2(v),$$

we get

$$(f_2)_*|_v(X_2) = Z_{w_*(X_1, X_2)}(0).$$

As in the proof of Lemma 6.20, we see that the Jacobi equation that $Z_{w_*(X_1, X_2)}$ satisfies is $\nabla_{\dot{\gamma}_{w(u,v)}(t)} \nabla_{\dot{\gamma}_{w(u,v)}} Z_{w_*(X_1, X_2)} = -Z_{w_*(X_1, X_2)}(t)$. It is clear that this implies that $Z_{w_*(X_1, X_2)}$ has the form

$$Z_{w_*(X_1, X_2)}(t) = \sin(t)P_0^t(\gamma_{w(u,v)})V_1 + \cos(t)P_0^t(\gamma_{w(u,v)})V_2,$$

for some $V_1, V_2 \in T|_{f_2(u)}M$. Now, taking into account the boundary values of $Z_{w_*(X_1, X_2)}(t)$ at $t = 0$ and $t = \frac{\pi}{2}$ as derived above, we get

$$\begin{aligned} V_1 &= P_{\frac{\pi}{2}}^0(\gamma_{w(u,v)})((f_1)_*|_u(X_1)), \\ V_2 &= (f_2)_*|_v(X_2). \end{aligned}$$

Define

$$\begin{aligned} Y_1(t) &= \sin(t)P_0^t(\gamma_{w(u,v)})V_1 = \sin(t)P_{\frac{\pi}{2}}^t(\gamma_{w(u,v)})((f_1)_*|_u(X_1)), \\ Y_2(t) &= \cos(t)P_0^t(\gamma_{w(u,v)})V_2 = \cos(t)P_0^t(\gamma_{w(u,v)})((f_2)_*|_v(X_2)), \end{aligned}$$

which means that

$$Z = Y_1 + Y_2.$$

Notice that Y_1 and Y_2 are Jacobi fields along $\gamma_{w(u,v)}$.

Since $w(u, v) \in (T|_{f_2(v)}N_1)^\perp$ and $\dot{\gamma}_{w(u,v)}(\frac{\pi}{2}) \in (T|_{f_1(u)}N_2)^\perp$ and

$$Y_1(\frac{\pi}{2}) = (f_1)_*|_u(X_1) \in T|_{f_1(u)}N_2, \quad Y_2(0) = (f_2)_*|_v(X_2) \in T|_{f_2(v)}N_1,$$

it follows that

$$Y_1, Y_2 \perp \gamma_{w(u,v)}.$$

We claim that moreover

$$Y_1 \perp Y_2.$$

Indeed, since $(f_2)_*|_v(X_2) \in T|_{f_2(v)}N_1$ and $(0, 1) \in \mathcal{D}_1|_{f_2(v)}$ (by definition of N_1), we have $((f_2)_*|_v(X_2), 0) \in \mathcal{D}_1|_{f_2(v)}$ and hence, for all t ,

$$(Z_1(t), r_1(t)) := (P^{\nabla^{\text{Rol}}})_0^t(\gamma_{w(u,v)})((f_2)_*|_v(X_2), 0) \in \mathcal{D}_1.$$

On the other hand, r_1 satisfies $\ddot{r}_1 + r_1 = 0$ with initial conditions $r_1(0) = 0$ and $\dot{r}_1(0) = g(\dot{\gamma}_{w(u,v)}(0), Z_1(0)) = g(w(u, v), (f_2)_*|_v(X_2)) = 0$ so $r_1(t) = 0$ for all t . Thus $Z_1(t)$ satisfies $\nabla_{\dot{\gamma}_{w(u,v)}(t)}Z_1 = 0$ i.e., $Z_1(t) = P_0^t(\gamma_{w(u,v)})((f_2)_*|_v(X_2))$. Similarly, if $w'(u, v) := -\frac{d}{dt}\Big|_{\frac{\pi}{2}} \exp_{f_2(v)}(tw(u, v)) = -\dot{\gamma}_{w(u,v)}(\frac{\pi}{2})$,

$$(Z_2(\frac{\pi}{2} - t), r_2(\frac{\pi}{2} - t)) := (P^{\nabla^{\text{Rol}}})_0^t(\gamma_{w'(u,v)})((f_1)_*|_u(X_1), 0) \in \mathcal{D}_2,$$

and we have $r_2(\frac{\pi}{2} - t) = 0$ and $Z_2(\frac{\pi}{2} - t) = P_0^t(\gamma_{w'(u,v)})((f_1)_*|_u(X_1))$ i.e., $Z_2(t) = P_{\frac{\pi}{2}}^t(\gamma_{w(u,v)})((f_1)_*|_u(X_1))$. But since $\mathcal{D}_1 \perp \mathcal{D}_2$ w.r.t. h , we have that $(Z_1, r_1) \perp (Z_2, r_2)$ w.r.t. h i.e., $g(Z_1(t), Z_2(t)) = 0$ for all t (since $r_1(t) = r_2(t) = 0$). Thus,

$$\begin{aligned} g(Y_1(t), Y_2(t)) &= \sin(t) \cos(t) g(P_{\frac{\pi}{2}}^t(\gamma_{w(u,v)})((f_1)_*|_u(X_1)), P_0^t(\gamma_{w(u,v)})((f_2)_*|_v(X_2))) \\ &= \sin(t) \cos(t) g(Z_2(t), Z_1(t)) = 0 \end{aligned}$$

This proves the claim, i.e., $Y_1 \perp Y_2$.

Since $\|w(u, v)\|_g = 1$, one has

$$\begin{aligned} \left\| \Psi_* \left(\frac{\partial}{\partial t}, X_1, X_2 \right) \right\|_g^2 &= \left\| \dot{\gamma}_{w(u, v)}(t) + Y_1(t) + Y_2(t) \right\|_g^2 \\ &= \left\| \dot{\gamma}_{w(u, v)}(t) \right\|_g^2 + \|Y_1(t)\|_g^2 + \|Y_2(t)\|_g^2 \\ &= 1 + \sin^2(t) \|(f_1)_*|_u(X_1)\|_g^2 + \cos^2(t) \|(f_2)_*|_v(X_2)\|_g^2. \end{aligned}$$

Finally, since

$$(f_1)_*|_u(X_1) = (\exp_{x_1})_*|_{\frac{\pi}{2}u} \left(\frac{\pi}{2} X_1 \right) \text{ and } (f_2)_*|_v(X_2) = (\exp_{x_2})_*|_{\frac{\pi}{2}v} \left(\frac{\pi}{2} X_2 \right),$$

Eq. (80) implies that

$$\begin{aligned} \|(f_1)_*|_u(X_1)\|_g &= \left| \sin\left(\frac{\pi}{2}\right) \right| \|X_1\|_g = \|X_1\|_g, \\ \|(f_2)_*|_v(X_2)\|_g &= \left| \sin\left(\frac{\pi}{2}\right) \right| \|X_2\|_g = \|X_2\|_g, \end{aligned}$$

and therefore

$$\begin{aligned} \left\| \Psi_* \left(\frac{\partial}{\partial t}, X_1, X_2 \right) \right\|_g^2 &= 1 + \sin^2(t) \|X_2\|_g^2 + \cos^2(t) \|X_1\|_g^2 \\ &= \tilde{g}|_{(t, u, v)} \left(\frac{\partial}{\partial t}, X_1, X_2 \right), \end{aligned}$$

i.e., Ψ is a local isometry $\tilde{S} \rightarrow M$. □

We next need one extra lemma.

Lemma 6.24 The manifold M has constant constant curvature equal to 1.

Proof. By Lemma 6.23, we know that $\Psi : \tilde{S} \rightarrow M$ is a local isometry. Now (\tilde{S}, \tilde{g}) has constant curvature = 1 since it is isometric to an open subset of the unit sphere (cf. [24] Chapter 1, Section 4.2). The image $\Psi(\tilde{S})$ of Ψ is clearly a dense subset of M (indeed, $\Psi(\tilde{S}) = M \setminus (N_1 \cup N_2)$), which implies that M has constant curvature = 1. □

This completes the proof the theorem in the case $1 \leq m \leq n - 2$, since a complete Riemannian manifold (M, g) with constant curvature = 1 is covered, in a Riemannian sense, by the unit sphere i.e., \hat{M}_1 . The cases $m = 0$ and $m = n - 1$ i.e., $\dim \mathcal{D}_1 = 1$ and $\dim \mathcal{D}_2 = 1$, respectively, are treated exactly in the same way as above, but in this case N_1 is a discrete set which might not be connected. □

Remark 6.25 The argument can easily be modified to deal with the case where N_2 (nor N_1) is not simply-connected. The simplifying assumption of simply connectedness of N_1 and N_2 made previously just serves to render the map $w(\cdot, \cdot)$ globally defined on $S_{x_1} \times S_{x_2}$. Otherwise we must define w only locally and, in its definition, make a choice corresponding to different sheets (of which there is a finite number).

Remark 6.26 As mentioned in the introduction, the following issue to address is that of an irreducible holonomy group of the rolling connection ∇^{Rol} i.e., for a given $x \in M$, the only non-trivial subspace of $T|_x M \oplus \mathbb{R}$ left invariant by parallel transport w.r.t ∇^{Rol} along loops based at $x \in M$ is $T|_x M \oplus \mathbb{R}$.

7 Rolling of Spaces of Different Dimensions

7.1 Definitions of the State Space and the Rolling Distributions

Definition 7.1 Let (M, g) , (\hat{M}, \hat{g}) be Riemannian manifolds of dimensions $n = \dim(M) \geq 2$ and $\hat{n} = \dim(\hat{M}) \geq 2$, not necessarily equal. Then one defines:

(i) if $n \leq \hat{n}$,

$$Q(M, \hat{M}) := \{A \in T^*M \otimes T\hat{M} \mid \hat{g}(AX, AY) = g(X, Y), X, Y \in T|_x M, x \in M\},$$

the set of isometric infinitesimal immersions. This defines a smooth manifold of $T^*M \otimes T\hat{M}$ of dimension

$$\dim(Q) := n + \hat{n} + n(\hat{n} - n) + \frac{n(n-1)}{2} = n + \hat{n} + n\hat{n} - \frac{n(n+1)}{2}.$$

(ii) If $n \geq \hat{n}$,

$$Q(M, \hat{M}) = \{A \in T^*M \otimes T\hat{M} \mid \hat{g}(AX, AY) = g(X, Y), X, Y \in (\ker A)^\perp, x \in M, A \text{ is onto a tangent space of } \hat{M}\},$$

where L^\perp is the orthogonal subspace of $L \subset T|_x M$ w.r.t. g . This defines a smooth submanifold of $T^*M \otimes T\hat{M}$ of dimension

$$\dim(Q) = n + \hat{n} + \hat{n}(n - \hat{n}) + \frac{\hat{n}(\hat{n} - 1)}{2} = n + \hat{n} + n\hat{n} - \frac{\hat{n}(\hat{n} + 1)}{2}.$$

If $n = \hat{n}$ and M, \hat{M} are oriented we also demand in (i) and (ii) that the elements of Q to preserve the orientations. Hence we recover the definition used before.

One defines distributions $\mathcal{D}_{\text{NS}}, \mathcal{D}_{\text{R}}$ on Q and the lifts $\mathcal{L}_{\text{NS}}, \mathcal{L}_{\text{R}}$ as before. In both cases the dimension of \mathcal{D}_{NS} is $n + \hat{n}$ and that of \mathcal{D}_{R} is n . Notice that by the above definition the dimension of $Q(M, \hat{M})$ is the same as that of $Q(\hat{M}, M)$. These manifolds are actually diffeomorphic as the next proposition shows.

Before proceeding, we introduce some notations. Given (M, g) and (\hat{M}, \hat{g}) as before, we write $Q = Q(M, \hat{M})$ and $\hat{Q} = Q(\hat{M}, M)$. We write $\mathcal{D}_{\text{NS}}, \mathcal{D}_{\text{R}}, \mathcal{L}_{\text{NS}}$ and \mathcal{L}_{R} on Q as before but on \hat{Q} we write the corresponding objects as $\widehat{\mathcal{D}}_{\text{NS}}, \widehat{\mathcal{D}}_{\text{R}}, \widehat{\mathcal{L}}_{\text{NS}}$ and $\widehat{\mathcal{L}}_{\text{R}}$. Thus $\dim \mathcal{D}_{\text{R}} = n$ but $\dim \widehat{\mathcal{D}}_{\text{R}} = \hat{n}$. As before, for $q = (x, \hat{x}; A) \in Q$ we write $A^{\overline{T}} : T|_{\hat{x}} \hat{M} \rightarrow T|_x M$ the (g, \hat{g}) -transpose of A i.e., $g(X, A^{\overline{T}} \hat{Y}) = \hat{g}(AX, \hat{Y})$ for all $X \in T|_x M, \hat{Y} \in T|_{\hat{x}} \hat{M}$.

Proposition 7.2 For every $(x, \hat{x}; A) \in Q$, one has $(\hat{x}, x; A^{\overline{T}}) \in \hat{Q}$ and the application

$$\overline{T} : Q \rightarrow \hat{Q}; \quad \overline{T}(x, \hat{x}; A) = (\hat{x}, x; A^{\overline{T}}),$$

is a diffeomorphism. Moreover, this diffeomorphism \overline{T} is an isometry of fiber bundles $\pi_Q \rightarrow \pi_{\hat{Q}}$ that preserves the no-spinning distributions on these manifolds i.e.,

$$\overline{T}_* \mathcal{D}_{\text{NS}} = \widehat{\mathcal{D}}_{\text{NS}}.$$

Proof. Suppose w.l.o.g. that $n \leq \hat{n}$. It is clear that $A^{\overline{T}}A = \text{id}_{T|_x M}$ for every $(x, \hat{x}; A) \in Q$ and $\ker(A^{\overline{T}}) = \text{im}(A)^\perp$ and thus if $\hat{X}, \hat{Y} \in \ker(A^{\overline{T}})^\perp = \text{im}(A)$, one gets

$$g(A^{\overline{T}}\hat{X}, A^{\overline{T}}\hat{Y}) = g(A^{\overline{T}}AX, A^{\overline{T}}AY) = g(X, Y) = \hat{g}(AX, AY) = \hat{g}(\hat{X}, \hat{Y}),$$

where $X, Y \in T|_x M$ were such that $AX = \hat{X}$, $AY = \hat{Y}$. This proves that $\overline{T}(x, \hat{x}; A)$ is actually an element of \hat{Q} .

Let then $\hat{q} = (\hat{x}, x; B) \in \hat{Q}$ and define

$$\overline{S}(\hat{q}) = (x, \hat{x}; B^{\overline{T}}) \in T^*M \otimes TM.$$

Since $\text{im}(B^{\overline{T}}) = \ker(B)^\perp$, we have for $X, Y \in T|_x M$,

$$g(B^{\overline{T}}X, B^{\overline{T}}Y) = \hat{g}(BB^{\overline{T}}X, BB^{\overline{T}}Y) = \hat{g}(X, Y),$$

directly from the definition of \hat{Q} and since $BB^{\overline{T}} = \text{id}_{T|_x M}$ (since $n \leq \hat{n}$). This shows that $\overline{S} : \hat{Q} \rightarrow Q$.

Moreover, one clearly has that \overline{T} and \overline{S} are maps inverse to each other. They are obviously smooth, hence Q and \hat{Q} are diffeomorphic. Also, \overline{T} is actually a bundle isomorphism $\pi_Q \rightarrow \pi_{\hat{Q}}$ whose inverse as a bundle isomorphism is \overline{S} .

Finally, observe that if $\gamma, \hat{\gamma}$ are smooth paths in M, \hat{M} starting at x_0, \hat{x}_0 , respectively, at $t = 0$, and if $q_0 = (x_0, \hat{x}_0; A_0) \in Q(M, \hat{M})$ then

$$(P_0^t(\hat{\gamma}) \circ A_0 \circ P_0^t(\gamma))^{\overline{T}} = P_0^t(\gamma) \circ A_0^{\overline{T}} \circ P_0^t(\hat{\gamma}),$$

so

$$\overline{T}(\gamma(t), \hat{\gamma}(t); P_0^t(\hat{\gamma}) \circ A_0 \circ P_0^t(\gamma)) = (\hat{\gamma}(t), \gamma(t); P_0^t(\gamma) \circ \overline{T}(x_0, \hat{x}_0; A_0) \circ P_0^t(\hat{\gamma})),$$

which immediately shows, by differentiating $\frac{d}{dt}\Big|_0$ and using the definition of \mathcal{L}_{NS} , that

$$\overline{T}_*|_{q_0} \mathcal{L}_{\text{NS}}(X, \hat{X})\Big|_{q_0} = \widehat{\mathcal{L}}_{\text{NS}}(\hat{X}, X)\Big|_{\overline{T}(q_0)}$$

where $X = \dot{\gamma}(0)$, $\hat{X} = \dot{\hat{\gamma}}(0)$. This proves in particular that \overline{T}_* maps \mathcal{D}_{NS} isomorphically onto $\widehat{\mathcal{D}}_{\text{NS}}$. This completes the proof. \square

Corollary 7.3 In the case $n \leq \hat{n}$, one has for $q_0 = (x_0, \hat{x}_0; A_0) \in Q(M, \hat{M})$ and $X \in T|_x M$,

$$\overline{T}_*|_{q_0} \mathcal{L}_R(X)|_{q_0} = \widehat{\mathcal{L}}_R(A_0 X)|_{\Phi(q_0)}.$$

In particular, $\overline{T}_* \mathcal{D}_R \subset \widehat{\mathcal{D}}_R$.

Proof. Indeed, for $X \in T|_{x_0} M$ and $q_0 = (x_0, \hat{x}_0; A_0)$ one has

$$\overline{T}_*|_{q_0} \mathcal{L}_R(X)|_{q_0} = \overline{T}_*|_{q_0} \mathcal{L}_{NS}(X, A_0 X)|_{q_0} = \widehat{\mathcal{L}}_{NS}(A_0 X, X)|_{\overline{T}(q_0)} = \widehat{\mathcal{L}}_R(A_0 X)|_{\overline{T}(q_0)},$$

since $X = (A_0^{\overline{T}})(A_0 X) = \overline{T}(q_0)(A_0 X)$. Hence \overline{T} maps \mathcal{D}_R of $Q(M, \hat{M})$ into $\widehat{\mathcal{D}}_R$ of $Q(\hat{M}, M)$. □

Remark 7.4 Recall that the distribution \mathcal{D}_R on $Q(M, \hat{M})$ has dimension n and $\widehat{\mathcal{D}}_R$ on $Q(\hat{M}, M)$ has dimension \hat{n} . Hence the inclusion $\overline{T}_* \mathcal{D}_R \subset \widehat{\mathcal{D}}_R$ is strict whenever $n < \hat{n}$. This shows that the model of rolling of manifolds of different dimensions against each other is not symmetric with respect to the order of the manifolds M and \hat{M} .

We can now provide a description of the vertical fiber $V|_q(\pi_Q)$ for a point $q = (x, \hat{x}; A) \in Q$.

Proposition 7.5 If $q = (x, \hat{x}; A) \in Q$, then the vertical fiber $V|_q(\pi_Q)$ is given by

$$V|_q(\pi_Q) = \begin{cases} \nu(\{B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M} \mid A^{\overline{T}} B \in \mathfrak{so}(T|_x M)\})|_q, & \text{if } n \leq \hat{n}, \\ \nu(\{B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M} \mid B A^{\overline{T}} \in \mathfrak{so}(T|_{\hat{x}} \hat{M})\})|_q, & \text{if } n \geq \hat{n}. \end{cases}$$

Proof. Let $q = (x, \hat{x}; A) \in Q$ and $B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M}$. Proving the proposition amounts to show that $\nu(B)|_q$ (which is *a priori* only an element of $V|_q(\pi_{T^*M \otimes T\hat{M}})$) belongs to $V|_q(\pi_Q)$ if and only if

- (i) $A^{\overline{T}} B \in \mathfrak{so}(n)$, if $n \leq \hat{n}$,
- (ii) $B A^{\overline{T}} \in \mathfrak{so}(\hat{n})$, if $n \geq \hat{n}$.

Choose first a π_Q -vertical curve $q(t) = (x, \hat{x}; A(t))$ inside Q such that $A(0) = A$ i.e., $q(t) \in \pi_Q^{-1}(x, \hat{x}) \subset T^*|_x M \otimes T|_{\hat{x}} \hat{M}$.

In the case (i), we have $\hat{g}(A(t)X, A(t)Y) = g(X, Y)$ for all $X, Y \in T|_x M$ so by differentiating, at $t = 0$, $g(AX, BY) + g(BX, AY) = 0$ for all $X, Y \in T|_x M$, where $B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M}$ is such that $A'(0) = \nu(B)|_q$. This condition can be written as $g(B^{\overline{T}} A X, Y) + g(A^{\overline{T}} B X, Y) = 0$ for all X, Y and hence $B^{\overline{T}} A + A^{\overline{T}} B = 0$. The result follows, since for a given $(x, \hat{x}; A) \in Q$, the set of $B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M}$ s.t. $A^{\overline{T}} B \in \mathfrak{so}(T|_x M)$ has dimension equal to $\dim \pi_Q^{-1}(x, \hat{x})$.

In the case (ii), we have $g(A(t)X, A(t)Y) = g(X, Y)$ for all $X, Y \in (\ker A(t))^\perp = \text{im}(A(t)^{\overline{T}})$. Choose $\hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}$. Then $g(A(t)^{\overline{T}} \hat{X}, A(t)^{\overline{T}} \hat{Y}) = \hat{g}(\hat{X}, \hat{Y})$, since $A(t)A(t)^{\overline{T}} = \text{id}_{T|_{\hat{x}} \hat{M}}$, and so by differentiating at $t = 0$, we get $g(A^{\overline{T}} \hat{X}, B^{\overline{T}} \hat{Y}) + g(B^{\overline{T}} \hat{X}, A^{\overline{T}} \hat{Y}) = 0$, where $B \in T^*|_x M \otimes T|_{\hat{x}} \hat{M}$ is such that $A'(0) = \nu(B)|_q$. This clearly means that $B A^{\overline{T}} + A B^{\overline{T}} = \text{id}_{T|_{\hat{x}} \hat{M}}$ and the result follows. □

Remark 7.6 The case (ii) considered above could be handled by using the diffeomorphism $\bar{T} : Q \rightarrow \hat{Q}$ introduced in Proposition 7.2. Indeed, if $n \geq \hat{n}$, we may apply (i) on \hat{Q} to obtain that for $q' = (\hat{x}, x; A') \in \hat{Q}$, we have that $V|_{q'}(\pi_{\hat{Q}})$ consists of $B' \in T^*|_{\hat{x}}\hat{M} \otimes T|_x M$ such that $A'^T B' \in \mathfrak{so}(T|_{\hat{x}}\hat{M})$. But taking $q = (x, \hat{x}; A)$, $q' = \bar{T}(q)$, $B \in T^*|_x M \otimes T|_{\hat{x}}\hat{M}$ and $B' = B^T$, this means $AB^T \in \mathfrak{so}(T|_{\hat{x}}\hat{M})$ i.e., $BA^T \in \mathfrak{so}(T|_{\hat{x}}\hat{M})$.

7.2 Controllability Results

7.2.1 Rolling Problem (NS)

Since Theorem 4.9 and Corollary 4.11 evidently hold as such in the case of non-equal dimensions (i.e., $n \neq \hat{n}$), we will be more interested to see how Theorem 4.16 could be formulated. We first need a definition.

Definition 7.7 For $n, \hat{n} \geq 2$, we define

$$\mathrm{SO}(n; \hat{n}) := \begin{cases} \{A \in (\mathbb{R}^n)^* \otimes \mathbb{R}^{\hat{n}} \mid A^T A = \mathrm{id}_{\mathbb{R}^n}\}, & \text{if } n < \hat{n}, \\ \{A \in (\mathbb{R}^n)^* \otimes \mathbb{R}^{\hat{n}} \mid A A^T = \mathrm{id}_{\mathbb{R}^{\hat{n}}}\}, & \text{if } n > \hat{n}, \\ \mathrm{SO}(n), & \text{if } n = \hat{n}, \end{cases}$$

where $(\mathbb{R}^n)^* \otimes \mathbb{R}^{\hat{n}}$ is the set of $n \times \hat{n}$ real matrices and A^T denotes the usual transpose of matrices.

Theorem 7.8 Fix some orthonormal frames F, \hat{F} of M, \hat{M} at $x \in M, \hat{x} \in \hat{M}$ and let $\mathfrak{h} = \mathfrak{h}|_F \subset \mathfrak{so}(n)$, $\hat{\mathfrak{h}} = \hat{\mathfrak{h}}|_{\hat{F}} \subset \mathfrak{so}(\hat{n})$ be the holonomy Lie-algebras of M, \hat{M} w.r.t to these frames. Then the control system $(\Sigma)_{NS}$ is completely controllable if and only if for every $A \in \mathrm{SO}(n; \hat{n})$,

$$A\mathfrak{h} - \hat{\mathfrak{h}}A = \begin{cases} \{B \in (\mathbb{R}^n)^* \otimes \mathbb{R}^{\hat{n}} \mid A^T B \in \mathfrak{so}(n)\}, & \text{if } n \leq \hat{n}, \\ \{B \in (\mathbb{R}^n)^* \otimes \mathbb{R}^{\hat{n}} \mid B A^T \in \mathfrak{so}(\hat{n})\}, & \text{if } n \geq \hat{n}. \end{cases}$$

7.2.2 Rolling Problem (R)

Notice that Proposition 5.17 still holds when $n = \dim M$ is not equal to $\hat{n} = \dim \hat{M}$. The rolling curvature of \mathcal{D}_R on Q is denoted as before but that of $\widehat{\mathcal{D}}_R$ on \hat{Q} is written as $\widehat{\mathrm{Rol}}$ i.e.,

$$\widehat{\mathrm{Rol}}(\hat{X}, \hat{Y})(B) = B\hat{R}(\hat{X}, \hat{Y}) - R(B\hat{X}, B\hat{Y})B,$$

for $(\hat{x}, x; B) \in \hat{Q}$ and $\hat{X}, \hat{Y} \in T|_{\hat{x}}\hat{M}$.

As a consequence, we have a generalization of Corollary 5.28.

Corollary 7.9 Use the notations introduced previously and assume that $n \leq \hat{n}$. Then the following two cases are equivalent:

- (i) \mathcal{D}_R is involutive,
- (ii) (M, g) and (\hat{M}, \hat{g}) have constant and equal curvature.

Also, if $n < \hat{n}$, then there is an equivalence between the two cases below:

- (1) $\widehat{\mathcal{D}}_R$ is involutive,
- (2) (M, g) and (\hat{M}, \hat{g}) are both flat.

Proof. For some of the notations, see the proof of Corollary 5.28.

(i) \implies (ii): Assume that \mathcal{D}_R is involutive. This is equivalent to the vanishing of Rol i.e.,

$$A(R(X, Y)Z) = \hat{R}(AX, AY)(AZ), \quad \forall (x, \hat{x}; A) \in Q, \quad X, Y, Z \in T|_x M,$$

which implies

$$\begin{aligned} \sigma_{(X, Y)} &= g(R(X, Y)Y, X) = \hat{g}(A(R(X, Y)Y), AX) \\ &= \hat{g}(\hat{R}(AX, AY)(AY), AX) = \hat{\sigma}_{(AX, AY)}, \end{aligned}$$

for every X, Y orthonormal in $T|_x M$ and $(x, \hat{x}; A) \in Q$.

Let $x \in M$, $\hat{x} \in \hat{M}$ be arbitrary points and $X, \hat{X} \in T|_x M$ and $\hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}$ be arbitrary pairs of orthonormal vectors.

Choose any vectors $X_3, \dots, X_n \in T|_x M$ and $\hat{X}_3, \dots, \hat{X}_{\hat{n}} \in T|_{\hat{x}} \hat{M}$ such that X, Y, X_3, \dots, X_n and $\hat{X}, \hat{Y}, \hat{X}_3, \dots, \hat{X}_{\hat{n}}$ are positively oriented orthonormal frames.

Since $n \leq \hat{n}$, we may define $q = (x, \hat{x}; A) \in Q$ by

$$AX = \hat{X}, \quad AY = \hat{Y}, \quad AX_i = \hat{X}_i, \quad i = 3, \dots, n,$$

to obtain that $\sigma_{(X, Y)} = \hat{\sigma}_{(\hat{X}, \hat{Y})}$. Thus (M, g) and (\hat{M}, \hat{g}) have equal and constant curvature, since the orthonormal pairs X, Y and \hat{X}, \hat{Y} were arbitrary and chosen independently from one another.

(ii) \implies (i): Since (M, g) , (\hat{M}, \hat{g}) both have equal constant curvature, say $k \in \mathbb{R}$, we have

$$\begin{aligned} R(X, Y)Z &= k(g(Y, Z)X - g(X, Z)Y), \quad X, Y, Z \in T|_x M, \quad x \in M, \\ \hat{R}(\hat{X}, \hat{Y})\hat{Z} &= k(\hat{g}(\hat{Y}, \hat{Z})\hat{X} - \hat{g}(\hat{X}, \hat{Z})\hat{Y}), \quad \hat{X}, \hat{Y}, \hat{Z} \in T|_{\hat{x}} \hat{M}, \quad \hat{x} \in \hat{M}. \end{aligned}$$

On the other hand, if $(x, \hat{x}; A) \in Q$, $X, Y, Z \in T|_x M$ we get

$$\begin{aligned} \hat{R}(AX, AY)(AZ) &= k(\hat{g}(AY, AZ)(AX) - \hat{g}(AX, AZ)(AY)) \\ &= A(k(g(Y, Z)X - g(X, Z)Y) = A(R(X, Y)Z). \end{aligned}$$

This implies that $\text{Rol}(X, Y)(A) = 0$ since Z was arbitrary. Hence \mathcal{D}_R is involutive.

(2) \implies (1): In this case $R = 0$ and $\hat{R} = 0$ so that clearly $\widehat{\text{Rol}}(\hat{X}, \hat{Y})(B)\hat{Z} = B(\hat{R}(\hat{X}, \hat{Y})\hat{Z}) - R(B\hat{X}, B\hat{Y})(B\hat{Z}) = 0$ for all $\hat{X}, \hat{Y}, \hat{Z} \in T|_{\hat{x}} \hat{M}$ and $(\hat{x}, x; B) \in \hat{Q}$. This proves that $\widehat{\mathcal{D}}_R$ is involutive.

(1) \implies (2): Assume that $\widehat{\mathcal{D}}_R$ is involutive i.e.,

$$B(\hat{R}(\hat{X}, \hat{Y})\hat{Z}) = R(B\hat{X}, B\hat{Y})(B\hat{Z}), \quad \forall (\hat{x}, x; B) \in \hat{Q}, \quad \hat{X}, \hat{Y}, \hat{Z} \in T|_{\hat{x}} \hat{M}.$$

Then

$$\sigma_{(B\hat{X}, B\hat{Y})} = g(B(\hat{R}(\hat{X}, \hat{Y})\hat{Y}), B\hat{X}),$$

or

$$\sigma_{(X,Y)} = g(B(\hat{R}(B^T X, B^T Y)(B^T Y), X)) = \hat{\sigma}_{(B^T X, B^T Y)}.$$

Given any $x \in M$, $\hat{x} \in \hat{M}$, $X, Y \in T|_x M$ and $\hat{X}, \hat{Y} \in T|_{\hat{x}} \hat{M}$, choose some $X_3, \dots, X_n \in T|_x M$, $\hat{X}_3, \dots, \hat{X}_{\hat{n}} \in T|_{\hat{x}} \hat{M}$ such that

$$X, Y, X_3, \dots, X_n \text{ and } \hat{X}, \hat{Y}, \hat{X}_3, \dots, \hat{X}_{\hat{n}},$$

are positively oriented orthonormal frames. We define

$$\begin{aligned} B\hat{X} &= X, & B\hat{Y} &= Y, & B\hat{X}_i &= X_i, & i &= 3, \dots, n, \\ B\hat{X}_i &= 0, & i &= n+1, \dots, \hat{n} \end{aligned}$$

so that $q = (\hat{x}, x; B) \in \hat{Q}$, $B^T X = \hat{X}$, $B^T Y = \hat{Y}$ and hence $\sigma_{(X,Y)} = \hat{\sigma}_{(\hat{X}, \hat{Y})}$. Thus (M, g) , (\hat{M}, \hat{g}) have constant and equal curvature.

Suppose that the common constant curvature of (M, g) , (\hat{M}, \hat{g}) is $k \in \mathbb{R}$. We need to show that $k = 0$. Choose any $(\hat{x}, x; B) \in \hat{Q}$. Since $n < \hat{n}$, we may choose non-zero vectors $\hat{X} \in \ker B$ and $\hat{Y} \in (\ker B)^\perp$. Then

$$\begin{aligned} 0 &= \widehat{\text{Rol}}(\hat{X}, \hat{Y})(B)\hat{X} = k(\hat{g}(\hat{Y}, \hat{X})B\hat{X} - \hat{g}(\hat{X}, \hat{X})B\hat{Y}) - R(B\hat{X}, B\hat{Y})(B\hat{X}) \\ &= k(0 - \|\hat{X}\|_{\hat{g}}^2 B\hat{Y}) - 0 = -k \|\hat{X}\|_{\hat{g}}^2 B\hat{Y} \end{aligned}$$

and since $\|\hat{X}\|_{\hat{g}} \neq 0$ and $B\hat{Y} \neq 0$ (since $0 \neq \hat{Y} \in (\ker B)^\perp$), it follows that $k = 0$. This completes the proof. \square

We may also easily generalize Corollary 5.29. The use will be made of Gauss-formula, which relates the curvature of a submanifold to that of the ambient Riemannian manifold and O'Neill-formulas, which relate the various curvatures related to Riemannian submersions (see [25], Propositions 3.8, 6.1, 6.2 and Corollary 6.3, Chapter II). Since the proof is slightly less trivial, we state this as a proposition.

Proposition 7.10 Suppose that (M, g) and (\hat{M}, \hat{g}) are complete with $\dim M = n$, $\dim \hat{M} = \hat{n}$. The following cases are equivalent:

- (i) There exists a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that $\mathcal{O}_{\mathcal{D}_R}(q_0)$ is an integral manifold of \mathcal{D}_R .
- (ii) There exists a $q_0 = (x_0, \hat{x}_0; A_0) \in Q$ such that

$$\text{Rol}(X, Y)(A) = 0, \quad \forall (x, \hat{x}; A) \in \mathcal{O}_{\mathcal{D}_R}(q_0), \quad X, Y \in T|_x M.$$

- (iii) There is a complete Riemannian manifold (N, h) , a Riemannian covering map $F : N \rightarrow M$ and a smooth map $G : N \rightarrow \hat{M}$ such that:

- (1) If $n \leq \hat{n}$, G is a Riemannian immersion that maps h -geodesics to \hat{g} -geodesics.
- (2) If $n \geq \hat{n}$, G is a Riemannian submersion such that the co-kernel distribution $(\ker G_*)^\perp \subset TN$ is involutive and the fibers $G^{-1}(\hat{x})$, $\hat{x} \in \hat{M}$, are totally geodesic submanifolds of (N, h) .

Moreover, in the case (iii)-(2), we may choose N to be simply connected and then (N, h) is a Riemannian product of (N_1, h_1) , (N_2, h_2) , where $\dim N_1 = \hat{n}$, $\dim N_2 = n - \hat{n}$, the space (N_1, g_1) is the universal Riemannian covering of (\hat{M}, \hat{g}) and G is given by

$$G : N = N_1 \times N_2 \rightarrow \hat{M}; \quad G(y_1, y_2) = \hat{\pi}(y_1)$$

where $\hat{\pi} : N_1 \rightarrow \hat{M}$ is a Riemannian covering map.

Proof. (i) \iff (ii): This is proved with exactly the same argument that was used in the proof of Corollary 5.29.

(i) \Rightarrow (iii): Let $N := \mathcal{O}_{\mathcal{D}_R}(q_0)$ and $h := (\pi_{Q,M}|_N)^*(g)$ i.e., for $q = (x, \hat{x}; A) \in N$ and $X, Y \in T|_x M$, define

$$h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q) = g(X, Y).$$

If $F := \pi_{Q,M}|_N$ and $G := \pi_{Q,\hat{M}}|_N$, we immediately see that F is a local isometry (note that $\dim(N) = n$). The completeness of (N, h) follows from the completeness of M and \hat{M} using Proposition 5.6 which holds in verbatim also in the case where $n \neq \hat{n}$. Hence by Proposition II.1.1 in [25], F is a (surjective) Riemannian covering.

Suppose then that $n \leq \hat{n}$. Then for $q = (x, \hat{x}; A) \in N$, $X, Y \in T|_x M$, one has

$$\hat{g}(G_*(\mathcal{L}_R(X)|_q), G_*(\mathcal{L}_R(Y)|_q)) = \hat{g}(AX, AY) = g(X, Y) = h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q),$$

i.e., G is a Riemannian immersion. Moreover, if $\bar{\Gamma} : [0, 1] \rightarrow N$ is an h -geodesic, it is tangent to \mathcal{D}_R and since it projects by F to a g -geodesic γ , it follows that $\bar{\Gamma} = q_{\mathcal{D}_R}(\gamma, \bar{\Gamma}(0))$ and Proposition 5.6 shows that $G \circ \bar{\Gamma} = \hat{\gamma}_{\mathcal{D}_R}(\gamma, \bar{\Gamma}(0))$ is a \hat{g} -geodesic. This proves (iii)-(1).

On the other hand, if $n \geq \hat{n}$, then for $q = (x, \hat{x}; A) \in N$, any $X \in T|_x M$ s.t. $\mathcal{L}_R(X)|_q \in (\ker G_*|_q)^\perp$ and $Z \in \ker A$, we have $G_*(\mathcal{L}_R(Z)|_q) = AZ = 0$ i.e., $\mathcal{L}_R(Z)|_q \in \ker(G_*|_q)$ from which $g(X, Z) = h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Z)|_q) = 0$. This shows that $X \in (\ker A)^\perp$ and therefore, for all $X, Y \in T|_x M$ such that $\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q \in (\ker G_*|_q)^\perp$, we get $\hat{g}(G_*(\mathcal{L}_R(X)|_q), G_*(\mathcal{L}_R(Y)|_q)) = h(\mathcal{L}_R(X)|_q, \mathcal{L}_R(Y)|_q)$ as above. This proves that $G : N \rightarrow \hat{M}$ is a Riemannian submersion.

For any $\bar{X}, \bar{Y} \in \text{VF}(N)$ orthonormal and tangent to $(\ker G_*)^\perp$ around a point $q \in N$, we have $\sigma_{(\bar{X}, \bar{Y})}^h = \hat{\sigma}_{(G_*\bar{X}, G_*\bar{Y})}$ (σ^h is the sectional curvature on N) in that neighbourhood because F is a Riemannian covering map and because

$$\begin{aligned} \hat{\sigma}_{(G_*\bar{X}, G_*\bar{Y})} &= \hat{g}(\hat{R}(G_*\bar{X}, G_*\bar{Y})(G_*\bar{Y}), G_*\bar{X}) = \hat{g}(\hat{R}(AF_*\bar{X}, AF_*\bar{Y})(AF_*\bar{Y}), AF_*\bar{X})) \\ &= \hat{g}(AR(F_*\bar{X}, F_*\bar{Y})F_*\bar{Y}, AF_*\bar{X}) = g(R(F_*\bar{X}, F_*\bar{Y})F_*\bar{Y}, A^T AF_*\bar{X}) \\ &= g(R(F_*\bar{X}, F_*\bar{Y})F_*\bar{Y}, F_*\bar{X}) = \sigma_{(F_*\bar{X}, F_*\bar{Y})}, \end{aligned}$$

since $\text{Rol} = 0$, $F_*\bar{X} \in (\ker A)^\perp$ on N and where we wrote $A = G_* \circ (F_*|_q)^{-1}$ for $q = (x, \hat{x}; A)$ in the chosen neighbourhood. By Corollary 6.3, Chapter II in [25], it follows that for any $\bar{X}, \bar{Y} \in \text{VF}(N)$ tangent to $(\ker G_*)^\perp$ in an open set, $[\bar{X}, \bar{Y}]$ is tangent to $(\ker G_*)^\perp$ in that open set. Thus $(\ker G_*)^\perp$ is involutive.

We still need to prove that the G -fibers are totally geodesic. Let $q = (x, \hat{x}; A) \in Q$, $V \in \ker G_*|_q = T|_q(G^{-1}(\hat{x}))$. Then $V = \mathcal{L}_R(u)|_q$ for some $u \in T|_x M$ and if γ is the g -geodesic starting from x with the initial velocity u , then $\Gamma := q_{\mathcal{D}_R}(\gamma, q)$ is

the h -geodesic with initial velocity V (since F is a Riemannian covering) and also $\hat{\gamma} := \hat{\gamma}_{\mathcal{D}_R}(\gamma, q)$ is a \hat{g} -geodesic by Proposition 5.6 with initial velocity $\dot{\hat{\gamma}}(0) = A\dot{\gamma}(0) = Au = G_*V = 0$, by the choice of V . But this means that $\hat{\gamma}$ is a constant curve, $\hat{\gamma}(\cdot) \equiv \hat{x}$ for all t , which implies that $G(\Gamma(t)) = \hat{\gamma}(t) = \hat{x}$ for all t i.e., $\Gamma(t) \in G^{-1}(\hat{x})$. This proves that every fiber $G^{-1}(\hat{x})$, $\hat{x} \in \hat{M}$, is a totally geodesic submanifold of (N, h) and so we have finally proved (iii)-(2).

(iii) \Rightarrow (ii): Let $x_0 \in M$ and choose $z_0 \in N$ such that $F(z_0) = x_0$. Define $\hat{x}_0 = G(z_0) \in \hat{M}$ and $A_0 := G_*|_{z_0} \circ (F_*|_{z_0})^{-1} : T|_{x_0}M \rightarrow T|_{\hat{x}_0}\hat{M}$.

The fact that $q_0 = (x_0, \hat{x}_0; A_0)$ belongs to Q can be seen as follows: if (iii)-(1) holds, we have

$$\begin{aligned} \hat{g}(A_0X, A_0Y) &= \hat{g}(G_*|_{z_0}((F_*|_{z_0})^{-1}X), G_*|_{z_0}((F_*|_{z_0})^{-1}Y)) \\ &= h((F_*|_{z_0})^{-1}X, (F_*|_{z_0})^{-1}Y) = g(X, Y), \end{aligned}$$

where we used that G is a Riemannian immersion and that F is a Riemannian covering map. On the other hand, if (iii)-(2) holds and if $X, Y \in (\ker A_0)^\perp$, clearly $(F_*|_{z_0})^{-1}X, (F_*|_{z_0})^{-1}Y \in (\ker G_*|_{z_0})^\perp$ and hence also $\hat{g}(A_0X, A_0Y) = g(X, Y)$ since G is a Riemannian submersion.

Let $\gamma : [0, 1] \rightarrow M$ be a smooth curve with $\gamma(0) = x_0$. Since F is a smooth covering map, there is a unique smooth curve $\Gamma : [0, 1] \rightarrow N$ with $\gamma = F \circ \Gamma$. Define $\hat{\gamma} = G \circ \Gamma$ and $A(t) = G_*|_{\Gamma(t)} \circ (F_*|_{\Gamma(t)})^{-1}$, $t \in [0, 1]$. As before, it follows that $A(t) \in Q$ for all $t \in [0, 1]$ and

$$\dot{\hat{\gamma}}(t) = G_*|_{\Gamma(t)}\dot{\Gamma}(t) = A(t)\dot{\gamma}(t). \quad (84)$$

We claim that, for all $t \in [0, 1]$,

$$\overline{\nabla}_{(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))}A(\cdot) = 0, \quad (85)$$

and

$$\text{Rol}(\cdot, \cdot)(A(t)) = 0. \quad (86)$$

Indeed, suppose now that (iii)-(1) holds. This means that, for every $z \in N$, there is a neighbourhood U of z in N such that $G(U)$ is a totally geodesic submanifold of (\hat{M}, \hat{g}) and $G : U \rightarrow \hat{M}$ is an isometric embedding. Now if X is a vector field parallel along γ in M , then since F is a Riemannian covering, there is a unique vector field \overline{X} parallel along Γ in (N, h) such that $F_*\overline{X} = X$. For any $t_0 \in [0, 1]$, choose U as above around $\Gamma(t_0)$. Then near t_0 we have that $G_*\overline{X}$ is parallel to $\hat{\gamma}$ in (\hat{M}, \hat{g}) . This proves that

$$0 = \hat{\nabla}_{\dot{\hat{\gamma}}(t)}(G_*\overline{X}(t)) = \hat{\nabla}_{\dot{\hat{\gamma}}(t)}(A(\cdot)X(\cdot)) = (\overline{\nabla}_{(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))}A(\cdot))X(t),$$

and since $X(t)$ was an arbitrary field parallel along γ , we have $\overline{\nabla}_{(\dot{\gamma}(t), \dot{\hat{\gamma}}(t))}A(\cdot) = 0$ i.e., (85).

Since, locally, the shape operator of $G(N)$ w.r.t (\hat{M}, \hat{g}) vanishes and $G(N)$ is locally Riemannian embedded submanifold of (\hat{M}, \hat{g}) , we also have $G_*(R^h(\overline{X}, \overline{Y})\overline{Z}) = \hat{R}(G_*\overline{X}, G_*\overline{Y})(G_*\overline{Z})$ for all $\overline{X}, \overline{Y}, \overline{Z} \in T|_zN$, $z \in N$ (see [25], Proposition 3.8, Chapter II) and hence for all $X, Y, Z \in T|_{\gamma(t)}M$,

$$\begin{aligned} A(t)(R(X, Y)Z) &= G_*(F_*|_{\Gamma(t)})^{-1}(R(X, Y)Z) = G_*(R^h(\overline{X}, \overline{Y})\overline{Z}) \\ &= \hat{R}(G_*\overline{X}, G_*\overline{Y})(G_*\overline{Z}) = \hat{R}(A(t)X, A(t)Y)(A(t)Z), \end{aligned}$$

where $\overline{X} = (F_*|_{\Gamma(t)})^{-1}X$ etc. This proves (86).

On the other hand, suppose (iii)-(2) holds. First we see that Eq. (86) follows from [25], Proposition 6.2, Chapter II (the operators A and T there vanish, by assumptions on N and G) and the fact that F is a Riemannian covering.

To prove (85) we proceed as follows. Taking the simply connected covering of N , lifting the metric h and composing G and F with the projection from this covering to N , we see that the conditions (iii)-(2) still hold and thus we may assume that N was simply connected in the first place. Take any piecewise C^1 curve ω on N and let $V_0 \in \ker G_*|_{\omega(0)}$, $X_0 \in (\ker G_*)^\perp|_{\omega(0)}$. If $Z(t)$ is the parallel translate of $X_0 + V_0$ along ω , we get from [25], Proposition 6.1, Chapter II (again, the operators A and T there vanish by assumptions) that

$$0 = \nabla_{\dot{\omega}(t)}^h Z(t) = (\nabla_{\dot{\omega}(t)}^h Z(t)^\perp)^\perp + (\nabla_{\dot{\omega}(t)}^h Z(t)^T)^T,$$

where for $Y \in TN$ we wrote Y^T and Y^\perp for the components of Y in the distributions $(\ker G_*)^\perp$ and $\ker G_*$, respectively (this notation is in accordance with the notation in the referred result of [25] and is not completely compatible with ours). This proves that $Z(t)^T$ and $Z(t)^\perp$ are fields parallel to ω and since $Z(0)^T = X_0$, $Z(0)^\perp = V_0$, we have that $Z(t)^T$ and $Z(t)^\perp$ are the parallel translates of X_0 and V_0 , respectively. But this implies that

$$(P^{\nabla^h})_0^t(\omega)(\ker G_*|_{\omega(0)}) = \ker G_*|_{\omega(t)}, \quad (P^{\nabla^h})_0^t(\omega)((\ker G_*|_{\omega(0)})^\perp) = (\ker G_*|_{\omega(t)})^\perp,$$

i.e., $TN = \ker G_* \oplus (\ker G_*)^\perp$ is a splitting to TN into two subbundles that are invariant under ∇^h -parallel transport.

Since N is simply connected and complete, it follows from de Rham's Theorem (see [25], Theorem 6.11, Chapter II) that $(N, h) = (N_1, h_1) \times (N_2, h_2)$, a Riemannian product, where (N_1, h_1) and (N_2, h_2) are both complete and simply connected and $TN_1 = (\ker G_*)^\perp$, $TN_2 = \ker G_*$.

To see now that Eq. (85) holds, let X be a vector field parallel along γ in M , write $\Gamma = (\Gamma_1, \Gamma_2)$, take $\overline{X} = (\overline{X}_1, \overline{X}_2)$ (w.r.t $TN = TN_1 \oplus TN_2$) to be the unique lift of X onto a vector field along Γ in N and compute

$$0 = A(t)\nabla_{\dot{\gamma}(t)}X(\cdot) = G_*\nabla_{(\dot{\Gamma}_1(t), \dot{\Gamma}_2(t))}^h \overline{X}(\cdot) = G_*\nabla_{\dot{\Gamma}_1(t)}^h \overline{X}_1 + G_*\nabla_{\dot{\Gamma}_2(t)}^h \overline{X}_2 = G_*\nabla_{\dot{\Gamma}_1(t)}^h \overline{X}_1,$$

since $\nabla_{\dot{\Gamma}_2(t)}^h \overline{X}_2 \in TN_2 = \ker G_*$. On the other hand, $G^{y_2} : N_1 \rightarrow \hat{M}$; $y_1 \mapsto G(y_1, y_2)$ is a local isometry for any $y_2 \in N_2$ and hence

$$\begin{aligned} 0 &= G_*\nabla_{\dot{\Gamma}_1(t)}^h \overline{X}_1 = (G^{\Gamma_2(t)})_*\nabla_{\dot{\Gamma}_1(t)}^h \overline{X}_1 = \hat{\nabla}_{(G^{\Gamma_2(t)})_*\dot{\Gamma}_1(t)}((G^{\Gamma_2(t)})_*\overline{X}_1) \\ &= \hat{\nabla}_{\dot{\gamma}(t)}(G_*\overline{X}) = \hat{\nabla}_{\dot{\gamma}(t)}(A(\cdot)X(\cdot)), \end{aligned}$$

since $\dot{\gamma}(t) = G_*\dot{\Gamma}(t) = G_*\dot{\Gamma}_1(t) = (G^{\Gamma_2(t)})_*\dot{\Gamma}_1(t)$ and $G_*\overline{X} = G_*\overline{X}_1 = (G^{\Gamma_2(t)})_*\overline{X}_1$. Thus (85) holds and this finishes the proof of (85)-(86) in the case that (iii)-(2) holds.

Thus we have shown, because of (84) and (85), that $t \mapsto (\gamma(t), \hat{\gamma}(t); A(t))$ is the unique rolling curve along γ starting at $q_0 = (x_0, \hat{x}_0; A_0)$ and defined on $[0, 1]$ and therefore curves of Q formed in this fashion fill up the orbit $\mathcal{O}_{\mathcal{D}_R}(q_0)$. Therefore, Eq. (86) implies that Rol vanishes on $\mathcal{O}_{\mathcal{D}_R}(q_0)$ which means that we are in case (ii).

To prove the last claim in the statement of the proposition, we continue the deduction done above in the case that condition (iii)-(2) holds. Since for any $y_2 \in N_2$,

$G^{y_2} : N_1 \rightarrow \hat{M}$ is a local Riemannian isometry and N_1 is simply connected and complete, it follows that G^{y_2} is a universal Riemannian covering of \hat{M} . We show that the map G^{y_2} is independent of the choice of $y_2 \in N_2$ i.e., that $G(y_1, y_2) = G(y_1, y'_2)$ for all $y_1 \in N_1$ and $y_2, y'_2 \in N_2$. Indeed, take any smooth path Γ_2 in N_2 from $\Gamma_2(0) = y_2$ to $\Gamma_2(1) = y'_2$. Then, $\frac{d}{dt}(G_{y_1} \circ \Gamma_2(t)) = G_* \dot{\Gamma}_2(t) = 0$ for all t since $\dot{\Gamma}_2(t) \in TN_2 = \ker G_*$. This shows that $G_{y_1} \circ \Gamma_2$ is a constant curve in \hat{M} and thus $G(y_1, y_2) = G_{y_1}(\Gamma_2(0)) = G_{y_1}(\Gamma_2(1)) = G(y_1, y'_2)$.

We fix $y'_2 \in N_2$ and define $\hat{\pi} := G^{y'_2} : N_1 \rightarrow \hat{M}$ which is a universal Riemannian covering. By what we just proved, it holds that $G(y_1, y_2) = G(y_1, y'_2) = \hat{\pi}(y_1)$ which establishes the claim. \square

A Fiber and Local Coordinates Point of View

Let $F = (X_i)$, $\hat{F} = (\hat{X}_i)$ be (not necessarily orthonormal) local frames of M and \hat{M} defined on the open subsets U, \hat{U} , respectively. We have local frames of 1-forms $F^* = ((\theta^i), U)$, $\hat{F}^* = ((\hat{\theta}^i), \hat{U})$ dual to these frames i.e., defined by $\theta^j(X_i) = \delta_i^j$, $\hat{\theta}^j(\hat{X}_i) = \delta_i^j$. The Christoffel symbols $\Gamma_{ij}^k = (\Gamma_F)_{ij}^k$, $\hat{\Gamma}_{ij}^k = (\hat{\Gamma}_{\hat{F}})_{ij}^k$ of $\nabla, \hat{\nabla}$ w.r.t the frames F, \hat{F} are defined by (see [28], p. 266) $\nabla_{X_i} X_j = \sum_k \Gamma_{ij}^k X_k$, $\hat{\nabla}_{\hat{X}_i} \hat{X}_j = \sum_k \hat{\Gamma}_{ij}^k \hat{X}_k$. Any $(x, \hat{x}; A) \in T^*_x(M) \otimes T|_{\hat{x}}(\hat{M})$ with $(x, \hat{x}) \in U \times \hat{U}$ can be written in the form

$$A = \sum_{i,j} A_j^i \theta^j|_x \otimes \hat{X}_i|_{\hat{x}},$$

i.e., $(\text{pr}_2 \circ \tau_{F, \hat{F}})(A) = [A_j^i]$ (see section 3.2). Moreover, if F, \hat{F} are orthonormal frames, then $A \in Q$ if and only if $[A_j^i] \in \text{SO}(n)$.

Let $t \mapsto \gamma(t)$, $t \mapsto \hat{\gamma}(t)$, $t \in I$, be smooth curves in U, \hat{U} , respectively, such that $\gamma(0) = x_0$, $\hat{\gamma}(0) = \hat{x}_0$, where I is a compact interval containing 0. Moreover, take $q_0 = (x_0, \hat{x}_0; A_0) \in T^*M \otimes T\hat{M}$. The no-spinning condition (12) (i.e., the parallel translation equation) for the curve $t \mapsto q(t) = (\gamma(t), \hat{\gamma}(t); A(t))$ (i.e., $A(t) = P_0^t(\gamma, \hat{\gamma})A$) starting at q can be written as

$$\frac{dA_j^i}{dt}(t) - \sum_{k,m} \Gamma_{m,j}^k(\gamma(t)) A_k^i(t) v^m(t) + \sum_{k,m} \hat{\Gamma}_{m,k}^i(\hat{\gamma}(t)) A_j^k(t) \hat{v}^m(t) = 0, \quad (87)$$

where $t \in I$ and

$$\dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}, \quad \dot{\hat{\gamma}}(t) = \sum_i \hat{v}^i(t) \hat{X}_i|_{\hat{\gamma}(t)}.$$

This shows immediately that the equation for no-spinning is a linear ODE in A_j^i and thus the solution with the initial condition $A(0) = A_0$ exists for the whole interval I where $\gamma, \hat{\gamma}$ are defined. The control system $(\Sigma)_{NS}$ can now be written locally in the form

$$\begin{cases} \dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}, \\ \dot{\hat{\gamma}}(t) = \sum_i \hat{v}^i(t) \hat{X}_i|_{\hat{\gamma}(t)}, \\ \frac{dA_j^i}{dt}(t) = \sum_{k,m} \Gamma_{m,j}^k(\gamma(t)) A_k^i(t) v^m(t) - \sum_{k,m} \hat{\Gamma}_{m,k}^i(\hat{\gamma}(t)) A_j^k(t) \hat{v}^m(t) \end{cases}, \quad t \in I,$$

or

$$\begin{cases} \dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}, \\ \dot{\hat{\gamma}}(t) = \sum_i \hat{v}^i(t) \hat{X}_i|_{\hat{\gamma}(t)}, \\ \dot{A}(t) = \sum_i v^i \mathcal{L}_{NS}(X_i, 0)|_{A(t)} + \sum_i \hat{v}^i \mathcal{L}_{NS}(0, \hat{X}_i)|_{A(t)}, \end{cases} \quad t \in I,$$

where the controls $v = (v^1, \dots, v^n)$, $\hat{v} = (\hat{v}^1, \dots, \hat{v}^n)$ are elements of $L^1_{\text{loc}}(I, \mathbb{R}^n)$ (actually an open subset of it since the images of $\gamma, \hat{\gamma}$ should belong to U, \hat{U} , respectively). From this local form, we see that $(\Sigma)_{NS}$ is a driftless control affine system.

The curve $t \mapsto q(t) = (\gamma(t), \hat{\gamma}(t); A(t))$ is a rolling curve i.e., satisfies conditions (11) and (13) if and only if

$$\hat{v}^i(t) = \sum_k A_k^i(t) v^k(t), \quad (88)$$

$$\frac{dA_j^i}{dt}(t) - \sum_m \left(\sum_k \Gamma_{m,j}^k(\gamma(t)) A_k^i(t) - \sum_{k,l} \hat{\Gamma}_{l,k}^i(\hat{\gamma}(t)) A_j^k(t) A_m^l(t) \right) v^m(t) = 0, \quad (89)$$

where $\dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}$, $\dot{\hat{\gamma}}(t) = \sum_i \hat{v}^i(t) \hat{X}_i|_{\hat{\gamma}(t)}$. Supposing that \hat{U} is a domain of a coordinate chart $\hat{\phi} = (\hat{x}^1, \dots, \hat{x}^n)$ of \hat{M} and taking as the frame \hat{F} the coordinate fields $\hat{X}_i = \frac{\partial}{\partial \hat{x}^i}$, the previous equation can be written as

$$\frac{d\hat{\gamma}^i}{dt}(t) = \sum_k A_k^i(t) v^k(t), \quad (90)$$

$$\frac{dA_j^i}{dt}(t) = \sum_m \left(\sum_k \Gamma_{m,j}^k(\gamma(t)) A_k^i(t) - \sum_{k,l} \hat{\Gamma}_{l,k}^i(\hat{\gamma}(t)) A_j^k(t) A_m^l(t) \right) v^m(t), \quad (91)$$

where $\hat{\gamma}^i = \hat{x}^i \circ \hat{\gamma}$ and $t \in I$. This system is nonlinear in $\hat{\gamma}^i, A_j^i$ and thus the existence of solutions, for a given initial condition $\hat{\gamma}(0) = \hat{x}_0$, $A(0) = A_0$ cannot be guaranteed on a given compact interval $I \ni 0$ where γ is defined (even in a case where one is able to get $\hat{U} = \hat{M}$).

Moving back to a general frame \hat{F} (i.e., we are not assuming that it consists of coordinate vector fields), the local form of the control system $(\Sigma)_R$ can be written as

$$\begin{cases} \dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}, \\ \dot{\hat{\gamma}}(t) = \sum_i v^i(t) A(t) X_i|_{\gamma(t)}, \\ \frac{dA_j^i}{dt}(t) = \sum_m \left(\sum_k \Gamma_{m,j}^k(\gamma(t)) A_k^i(t) - \sum_{k,l} \hat{\Gamma}_{l,k}^i(\hat{\gamma}(t)) A_j^k(t) A_m^l(t) \right) v^m(t), \end{cases} \quad t \in I,$$

or

$$\begin{cases} \dot{\gamma}(t) = \sum_i v^i(t) X_i|_{\gamma(t)}, \\ \dot{\hat{\gamma}}(t) = \sum_i v^i(t) A(t) X_i|_{\gamma(t)}, \\ \dot{A}(t) = \sum_i v^i(t) \mathcal{L}_R(X_i)|_{A(t)}, \end{cases} \quad t \in I,$$

where the controls $v = (v^1, \dots, v^n)$ are elements of $L_{\text{loc}}^1(I, \mathbb{R}^n)$ as above. From this local form, we see that $(\Sigma)_R$ is a driftless control affine system.

B Sasaki-metric on $T^*M \otimes T\hat{M}$ and Q

The no-spinning distribution \mathcal{D}_{NS} can be given a natural sub-Riemannian structure (see e.g. [19]) with the sub-Riemannian metric $h_{\text{NSD}} := (\pi_Q^*(\bar{g}))|_{\mathcal{D}_{\text{NS}}}$ since $(\pi_Q)_*|_{\mathcal{D}_{\text{NS}}|_{(x, \hat{x}; A)}}$ is a linear isomorphism $T|_{(x, \hat{x}; A)}Q \rightarrow T|_{(x, \hat{x})}(M \times \hat{M})$ at each point $q = (x, \hat{x}; A) \in Q$ and $(\pi_Q)_*(\mathcal{L}_{\text{NS}}(X, \hat{X})|_q) = (X, \hat{X})$, for every $q = (x, \hat{x}; A) \in Q$ and $X \in T|_xM$, $\hat{X} \in T|_{\hat{x}}\hat{M}$.

Actually, we have more since there is a Sasaki-metric \bar{g}_1^1 on the whole tensor space $T^*(M) \otimes T(\hat{M})$ given in the following.

Definition B.1 The *Sasaki-metric* \bar{g}_1^1 on $T^*(M) \otimes T(\hat{M})$ is defined by

$$\begin{aligned} \bar{g}_1^1(\xi, \eta) = & \bar{g}(\mathcal{L}_{\text{NS}}|_q^{-1}(\text{pr}_1(\xi)), \mathcal{L}_{\text{NS}}|_q^{-1}(\text{pr}_1(\eta))) \\ & + ((g^* \otimes \hat{g}) \circ \tau)(\nu|_q^{-1}(\text{pr}_2(\xi)), \nu|_q^{-1}(\text{pr}_2(\eta))), \end{aligned} \quad (92)$$

where

$$q = (x, \hat{x}; A) \in T^*(M) \otimes T(\hat{M}), \quad \xi, \eta \in T|_q(T^*(M) \otimes T(\hat{M})),$$

pr_1, pr_2 are projections of the decomposition

$$T(T^*(M) \otimes T(\hat{M})) = \mathcal{D}_{\text{NS}} \oplus V(\pi_{T^*(M) \otimes T(\hat{M})}),$$

(see (20)) onto the first and second factors, $\mathcal{L}_{\text{NS}}|_q^{-1}, \nu|_q^{-1}$ are the inverse maps of

$$\begin{aligned} T|_{(x, \hat{x})}(M \times \hat{M}) & \rightarrow T|_q(T^*(M) \otimes T(\hat{M})) \\ \bar{X} & \mapsto \mathcal{L}_{\text{NS}}(\bar{X})|_q, \end{aligned} \quad (93)$$

and

$$\begin{aligned} (T^*(M) \otimes T(\hat{M}))|_{(x, \hat{x})} & \rightarrow V|_q(\pi_{T^*(M) \otimes T(\hat{M})}) \\ B & \mapsto \nu(B)|_q. \end{aligned} \quad (94)$$

Note that $g^* : T^*(M) \otimes T^*(M) \rightarrow \mathbb{R}$ is the dual metric induced by g and finally τ is the \mathbb{R} -linear isomorphism

$$(T^*M \otimes T\hat{M}) \otimes (T^*M \otimes T\hat{M}) \rightarrow (T^*M \otimes T^*M) \otimes (T\hat{M} \otimes T\hat{M})$$

uniquely determined by

$$\tau((\omega \otimes \hat{X}) \otimes (\theta \otimes \hat{Y})) = (\omega \otimes \theta) \otimes (\hat{X} \otimes \hat{Y}).$$

Denote by \bar{g}_Q the restriction (i.e., the pull-back) of the metric \bar{g}_1^1 onto Q .

Let us now use the local frames and notation as in Appendix A. Writing $\xi, \eta \in T|_q(T^*M \otimes T\hat{M})$, $q = (x, \hat{x}; A)$, as

$$\begin{aligned} \xi & = \sum_i (\xi^i X_i|_x + \hat{\xi}^i \hat{X}_i|_{\hat{x}}) + \sum_{i,j} \xi_{\nu_j}^i \theta^j|_x \otimes \hat{X}_i|_{\hat{x}}, \\ \eta & = \sum_i (\eta^i X_i|_x + \eta \xi^i \hat{X}_i|_{\hat{x}}) + \sum_{i,j} \eta_{\nu_j}^i \theta^j|_x \otimes \hat{X}_i|_{\hat{x}}, \end{aligned}$$

one gets

$$\begin{aligned} \text{pr}_2(\xi) &= \xi - \mathcal{L}_{\text{NS}} \sum_i (\xi^i X_i|_x + \hat{\xi}^i \hat{X}_i|\hat{x})|_q \\ &= \sum_{i,j} \left(\xi_{\nu_j}^i - \sum_{k,m} \mathbf{\Gamma}_{m,j}^k A_k^i \xi^m + \hat{\mathbf{\Gamma}}_{m,k}^i A_j^k \hat{\xi}^m \right) \theta^i|_x \otimes \hat{X}_j|\hat{x}, \end{aligned}$$

the similar formula holding for $\text{pr}_2(\eta)$ and hence

$$\begin{aligned} \bar{g}_1^1 &= \sum_{i,j} \xi^i \eta^j g(X_i|_x, X_j|_x) + \sum_{i,j} \hat{\xi}^i \hat{\eta}^j g(\hat{X}_i|\hat{x}, \hat{X}_j|\hat{x}) \\ &\quad + \sum_{i,j,\alpha,\beta} \left(\xi_{\nu_j}^i - \sum_{k,m} \mathbf{\Gamma}_{m,j}^k A_k^i \xi^m + \hat{\mathbf{\Gamma}}_{m,k}^i A_j^k \hat{\xi}^m \right) \\ &\quad \cdot \left(\eta_{\nu_\beta}^\alpha - \sum_{k,m} \mathbf{\Gamma}_{m,\beta}^k A_k^\alpha \xi^m + \hat{\mathbf{\Gamma}}_{m,k}^\alpha A_\beta^k \hat{\xi}^m \right) g^*(\theta^i, \theta^\alpha) \hat{g}(\hat{X}_j, \hat{X}_\beta). \end{aligned}$$

Moreover, with this choice of the Riemannian metric on $T^*M \otimes T\hat{M}$ and Q we have the following result.

Proposition B.2 Let $U, V \in C^\infty(\pi_{T^*M \otimes T\hat{M}}, \pi_{T^*M \otimes T\hat{M}})$, $\bar{X} \in T|_{(x_0, \hat{x}_0)}(M \times \hat{M})$ and $q_0 = (x_0, \hat{x}_0; A_0)$. Then the Sasaki-metric \bar{g}_1^1 has the following properties:

- (i) Letting $\text{tr} = \text{tr}_{T|_{x_0}^* M \otimes T|_{x_0} M}$ denote the trace of linear maps $T|_{x_0} M \rightarrow T|_{x_0} M$ and \bar{T} the (g, \hat{g}) -transpose of the linear maps $T|_{x_0} M \rightarrow T|_{\hat{x}_0} \hat{M}$, one has

$$\bar{g}_1^1|_{q_0}(\nu(U(A_0))|_{q_0}, \nu(V(A_0))|_{q_0}) = \text{tr}(U(A_0) \bar{T} V(A_0)) \quad (95)$$

- (ii) Choosing a smooth local $\pi_{T^*M \otimes T\hat{M}}$ -section \tilde{A} s.t. $\tilde{A}|_{(x_0, \hat{x}_0)} = A_0$ and $\bar{\nabla}_{\bar{Y}} \tilde{A} = 0$ for all $\bar{Y} \in T|_{(x_0, \hat{x}_0)}(M \times \hat{M})$,

$$\begin{aligned} &\mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(\bar{g}_1^1(\nu(U(\cdot)), \nu(V(\cdot)))) \\ &= \bar{g}_1^1(\nu(\bar{\nabla}_{\bar{X}} U(\tilde{A}))|_{q_0}, \nu(V(A_0))|_{q_0}) + \bar{g}_1^1(\nu(U(A_0))|_{q_0}, \nu(\bar{\nabla}_{\bar{X}} V(\tilde{A}))|_{q_0}). \end{aligned} \quad (96)$$

The same result holds if we throughout replace $T^*M \otimes T\hat{M}$ by Q and \bar{g}_1^1 by \bar{g}_Q with the exception that $U, V \in C^\infty(\pi_Q, \pi_{T^*M \otimes T\hat{M}})$ s.t. $U(A), V(A) \in \mathcal{A}\mathfrak{so}(T|_x M)$ for all $(x, \hat{x}; A) \in Q$.

Proof. Let $(X_i), (\hat{X}_i)$ be smooth g, \hat{g} -orthonormal frames of vector fields M, \hat{M} defined on the neighborhoods U, \hat{U} of x_0, \hat{x}_0 . Denote by $(\theta^i), (\hat{\theta}^i)$ the corresponding dual frames. Then there are unique functions $a_j^i, b_j^i \in C^\infty(\pi_{T^*M \otimes T\hat{M}}^{-1}(U \times \hat{U}))$ s.t.

$$U(A) = \sum_{i,j} a_j^i(x, \hat{x}; A) \theta^j|_x \otimes \hat{X}_i|\hat{x}, \quad V(A) = \sum_{i,j} b_j^i(x, \hat{x}; A) \theta^j|_x \otimes \hat{X}_i|\hat{x},$$

and thus (below we will denote $a_j^i(x_0, \hat{x}_0; A_0), b_j^i(x_0, \hat{x}_0; A_0)$ simply by a_j^i, b_j^i)

$$\begin{aligned}
& \bar{g}_1^1|_{q_0}(\nu(U(A_0))|_{q_0}, \nu(V(A_0))|_{q_0}) \\
&= ((g^* \otimes \hat{g}) \circ \tau) \left(\sum_{i,j,k,l} a_j^i b_l^k (\theta^j|_{x_0} \otimes \hat{X}_i|_{\hat{x}_0}) \otimes (\theta^l|_{x_0} \otimes \hat{X}_k|_{\hat{x}_0}) \right) \\
&= \sum_{i,j,k,l} a_j^i b_l^k g^*(\theta^j|_{x_0} \otimes \theta^l|_{x_0}) \otimes \hat{g}(\hat{X}_j|_{\hat{x}_0} \otimes \hat{X}_k|_{\hat{x}_0}) = \sum_{i,j,k,l} a_j^i b_l^k \delta^{jl} \delta_{jk} \\
&= \sum_{i,j} a_j^i b_j^i = \sum_i \hat{g}(U(A_0)X_i, V(A_0)X_i) = \text{tr}(U(A_0)\bar{T}V(A_0)).
\end{aligned}$$

This proves (i).

Next, by the definition of \mathcal{L}_{NS} and the choice of \tilde{A} , we have

$$\begin{aligned}
& \mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(\bar{g}_1^1(\nu(U(\cdot)), \nu(V(\cdot)))) = \bar{X}\bar{g}_1^1(\nu(U(\tilde{A}))|_{\tilde{A}}, \nu(V(\tilde{A}))|_{\tilde{A}}) \\
&= \bar{X} \left(\sum_{i,j} a_j^i(\tilde{A})b_j^i(\tilde{A}) \right) = \sum_{i,j} \bar{X}(a_j^i(\tilde{A}))b_j^i(q_0) + \sum_{i,j} a_j^i(q_0)\bar{X}(b_j^i(\tilde{A})).
\end{aligned}$$

Assuming for simplicity that $(X_i), (\hat{X}_i)$ were chosen so that $\nabla_Y X_i = 0, \hat{\nabla}_{\hat{Y}} \hat{X}_i = 0$ for all i and $Y \in T|_{x_0}M, \hat{Y} \in T|_{\hat{x}_0}\hat{M}$ (and hence $\nabla_Y \theta^i = 0$ for all i and $Y \in T|_{x_0}M$), we get

$$\bar{\nabla}_{\bar{X}}(U(\tilde{A})) = \mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(a_j^i \theta^j|_{x_0} \otimes \hat{X}_i|_{\hat{x}_0}), \quad \bar{\nabla}_{\bar{X}}(V(\tilde{A})) = \mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(b_j^i \theta^j|_{x_0} \otimes \hat{X}_i|_{\hat{x}_0}).$$

This shows that

$$\begin{aligned}
& \sum_{i,j} \mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(a_j^i)b_j^i(q_0) + \sum_{i,j} a_j^i(q_0)\mathcal{L}_{\text{NS}}(\bar{X})|_{q_0}(b_j^i) \\
&= \text{tr}((\bar{\nabla}_{\bar{X}}(U(\tilde{A})))\bar{T}V(A_0)) + \text{tr}(U(A_0)\bar{T}(\bar{\nabla}_{\bar{X}}(V(\tilde{A})))) \\
&= \bar{g}_1^1(\nu(\bar{\nabla}_{\bar{X}}U(\tilde{A}))|_{q_0}, \nu(V(A_0))|_{q_0}) + \bar{g}_1^1(\nu(U(A_0))|_{q_0}, \nu(\bar{\nabla}_{\bar{X}}V(\tilde{A}))|_{q_0}).
\end{aligned}$$

□

Proposition B.3 The maps $\pi_{T^*(M) \otimes T(\hat{M})}$ and π_Q are surjective Riemannian submersions onto $M \times \hat{M}$. Hence the restrictions of the Levi-Civita connection $\nabla^{\bar{g}_1^1}$ and the Riemannian curvature $R^{\bar{g}_1^1}$ on the \mathcal{D}_{NS} -horizontal fields are respectively given by

$$\nabla_{\mathcal{L}_{\text{NS}}(\bar{X})}^{\bar{g}_1^1} \mathcal{L}_{\text{NS}}(\bar{Y})|_q = \mathcal{L}_{\text{NS}}(\bar{\nabla}_{\bar{X}}\bar{Y})|_q + \frac{1}{2} \nu(AR(X, Y) - \hat{R}(\hat{X}, \hat{Y})A)|_q,$$

and

$$\begin{aligned}
& \bar{g}_1^1(R^{\bar{g}_1^1}(\mathcal{L}_{\text{NS}}(\bar{X}), \mathcal{L}_{\text{NS}}(\bar{Y}))\mathcal{L}_{\text{NS}}(\bar{X}), \mathcal{L}_{\text{NS}}(\bar{Y})) \\
&= \bar{g}(\bar{R}(\bar{X}, \bar{Y})\bar{X}, \bar{Y}) + \frac{3}{4} \left\| \nu(AR(X, Y) - \hat{R}(\hat{X}, \hat{Y})A)|_q \right\|_{\bar{g}_1^1},
\end{aligned}$$

with $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$, $\bar{X} = (X, \hat{X})$ where $X \in \text{VF}(M), \hat{X} \in \text{VF}(\hat{M})$ and similarly for \bar{Y} . The same formulas hold if one replaces \bar{g}_1^1 with \bar{g}_Q .

Proof. The first statement is obvious by construction. For the statement about the connection, we use Koszul's formula (cf. [15]), to notice that

$$\nabla_{\mathcal{L}_{\text{NS}}(\bar{X})}^{\bar{g}_1^1} \mathcal{L}_{\text{NS}}(\bar{Y}) = \mathcal{L}_{\text{NS}}(\bar{\nabla}_{\bar{X}} \bar{Y}) + \frac{1}{2} [\mathcal{L}_{\text{NS}}(\bar{X}), \mathcal{L}_{\text{NS}}(\bar{Y})]^v,$$

where for $Z \in T(T^*M \otimes T\hat{M})$ we denote $Z = Z^h + Z^v$ with $Z^h \in \mathcal{D}_{\text{NS}}$ and $Z^v \in V(\pi_{T^*M \otimes T\hat{M}})$. The fact about the Riemannian curvature is deduced similarly (see [15]). \square

Theorem B.4 Suppose $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ is a smooth curve on $T^*(M) \otimes T(\hat{M})$ that is \mathcal{D}_{NS} -horizontal i.e., $\dot{q}(t) \in \mathcal{D}_{\text{NS}}$ for all t . Then the following are equivalent:

- (i) $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ is a geodesic of $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$
- (ii) $t \mapsto x(t)$ and $t \mapsto \hat{x}(t)$ are geodesics of (M, g) and (\hat{M}, \hat{g}) respectively.

Moreover, in this case $t \mapsto A(t)$ is given by parallel transport as follows:

$$A(t) = P_0^t(\hat{x}) \circ A(0) \circ P_t^0(x) = P_0^t(x, \hat{x})A(0). \quad (97)$$

The same facts hold if $(T^*M \otimes T\hat{M}, \bar{g}_1^1)$ is replaced by (Q, \bar{g}_Q) .

Notice that the claim of the theorem can also be written more compactly as follows: For any $q = (x, \hat{x}; A) \in T^*M \otimes T\hat{M}$,

$$\pi_{T^*M \otimes T\hat{M}} \circ \exp_q^{\bar{g}_1^1} \circ \mathcal{L}_{\text{NS}}|_q = \overline{\text{exp}}_{(x, \hat{x})}, \quad (98)$$

with a similar formula holding when $T^*M \otimes T\hat{M}$ is replaced by Q .

Proof. This follows from the fact that $\pi_{T^*M \otimes T\hat{M}}$ (resp. π_Q) is a Riemannian submersion. Nevertheless, for the sake of convenience we outline the easy proof here.

The fact that $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ is \mathcal{D}_{NS} -horizontal implies that $\dot{q}(t) = \mathcal{L}_{\text{NS}}(\dot{x}(t), \dot{\hat{x}}(t))|_{q(t)}$ for all t . Thus, by Proposition B.3, we get

$$\nabla_{\dot{q}(t)}^{\bar{g}_1^1} \dot{q} = \mathcal{L}_{\text{NS}}(\nabla_{\dot{x}(t)} \dot{x}, \hat{\nabla}_{\dot{\hat{x}}(t)} \dot{\hat{x}})|_{q(t)},$$

since $R(\dot{x}(t), \dot{x}(t)) = 0$, $\hat{R}(\dot{\hat{x}}(t), \dot{\hat{x}}(t)) = 0$. The claim follows from this since $\mathcal{L}_{\text{NS}}(\cdot)|_{q(t)}$ is a linear isomorphism for each t . Also, Eq. (97) follows easily from the definition of \mathcal{D}_{NS} and Eq. (15). \square

Corollary B.5 The \mathcal{D}_{NS} -horizontal curve on Q is a geodesic of (Q, \bar{g}_Q) if and only if it is a geodesic of $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$.

Theorem B.6 The Riemannian manifolds (M, g) , (\hat{M}, \hat{g}) are complete Riemannian manifolds if and only if $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$ or (Q, \bar{g}_Q) is complete.

Proof. The completeness of $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$ (resp. (Q, \bar{g}_Q)) implies the completeness of (M, g) , (\hat{M}, \hat{g}) since $\pi_{T^*(M) \otimes T(\hat{M})}$ (resp. π_Q) is a Riemannian submersion onto $(M \times \hat{M}, \bar{g})$ and pr_1, pr_2 are Riemannian submersions from $M \times \hat{M}$ onto M and

\hat{M} , respectively (recall that Riemannian submersions map geodesics to geodesics). This proves the direction “ \Leftarrow ”.

Thus assume that (M, g) , (\hat{M}, \hat{g}) are complete, which is equivalent to the completeness of $(M \times \hat{M}, \bar{g})$. Let $(x_n, \hat{x}_n; A_n) \in T^*(M) \otimes T(\hat{M})$ be a Cauchy-sequence. Then (x_n, \hat{x}_n) is a Cauchy sequence in $M \times \hat{M}$ and hence converges to a point $(y, \hat{y}) \in M \times \hat{M}$ since $M \times \hat{M}$ is a complete (metric) space. Choose a local trivialization $\tau : \pi_{T^*(M) \times T(\hat{M})}^{-1}(U \times \hat{U}) \rightarrow (U \times \hat{U}) \times \mathfrak{gl}(n)$ of $T^*(M) \times T(\hat{M})$ induced by some coordinate charts $((x^i), U)$, $((\hat{x}^i), \hat{U})$ (see Appendix A) of M , \hat{M} around y , \hat{y} respectively. By Proposition II.1.1 in [25], the metric $d_{\bar{g}_1^1}$ induced by \bar{g}_1^1 on $T^*(M) \otimes T(\hat{M})$, and hence on $(U \times \hat{U}) \times \mathfrak{gl}(n)$ through τ , gives the original manifold topology. Choose an open neighbourhood $V \times \hat{V}$ of (y, \hat{y}) such that $\overline{V \times \hat{V}}$ is a compact subset of $U \times \hat{U}$. Then $(\overline{V \times \hat{V}}) \times \mathfrak{gl}(n)$ is a complete space that contains a Cauchy-sequence $((x_n, \hat{x}_n), a_n) = \tau(x_n, \hat{x}_n; A_n)$ for n large enough. Hence it converges to $(y, \hat{y}, a) \in (\overline{V \times \hat{V}}) \times \mathfrak{gl}(n)$ and thus $(x_n, \hat{x}_n; A_n)$ converges to $\tau^{-1}(y, \hat{y}, a) = (y, \hat{y}; A)$. This proves the completeness of $T^*(M) \otimes T(\hat{M})$. We thus get the completeness of Q since Q is a closed submanifold of $T^*(M) \otimes T(\hat{M})$. \square

Remark B.7 More generally, suppose $\pi : E \rightarrow N$ is a smooth bundle, (E, G) , (N, g) are Riemannian manifolds, π is a Riemannian submersion and the typical fiber of π is complete (i.e., all the fibers $\pi^{-1}(x)$ are complete subsets of E). Then the argument of the previous proof applies and shows that E is a complete Riemannian manifold if and only if N is a complete Riemannian manifold.

We record the following result.

Proposition B.8 Let N be an integral manifold of \mathcal{D}_R and equip it with the Riemannian metric $\bar{g}_N := \bar{g}_1^1|_N$. Then (N, \bar{g}_N) is a totally geodesic submanifold of $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$.

The same claim holds if one replaces $(T^*M \otimes T\hat{M}, \bar{g}_1^1)$ by (Q, \bar{g}_Q) and assumes that $N \subset Q$.

Proof. The assumptions immediately imply that the projection

$$\pi_N := \text{pr}_1 \circ \pi_{T^*(M) \otimes T(\hat{M})}|_N,$$

is a local isometric diffeomorphism from (N, \bar{g}_N) into (M, g) since $\text{pr}_1 \circ \pi_{T^*(M) \otimes T(\hat{M})}$ maps \mathcal{D}_R isometrically onto TM by the definition of \bar{g}_1^1 and \mathcal{D}_R .

Now if $t \mapsto (x(t), \hat{x}(t); A(t))$, $t \in]a, b[$, is a geodesic of N then (since it is tangent to \mathcal{D}_R) we have $\hat{\nabla}_{\hat{x}(t)} \dot{x}(t) = A(t)\dot{x}(t)$ and $t \mapsto x(t) = \pi_N(x(t), \hat{x}(t); A(t))$, $t \in]a, b[$, is a geodesic of (M, g) . We have

$$\hat{\nabla}_{\hat{x}(t)} \dot{\hat{x}}(t) = \hat{\nabla}_{\hat{x}(t)} (A(\cdot)\dot{x}) = (\bar{\nabla}_{(\hat{x}(t), \dot{\hat{x}}(t))} A)\dot{x}(t) + A(t)\nabla_{\hat{x}(t)} \dot{x},$$

and once we use the facts that $\nabla_{\dot{x}(t)} \dot{x} = 0$ (since x is a geodesic on M) and $\bar{\nabla}_{(\hat{x}(t), \dot{\hat{x}}(t))} A = 0$ (by the definition of \mathcal{D}_{NS}) to conclude that $\hat{\nabla}_{\hat{x}(t)} \dot{\hat{x}}(t) = 0$ i.e., $t \mapsto \hat{x}(t)$, $t \in]a, b[$, is a geodesic of \hat{M} . Thus Theorem B.4 implies that $t \mapsto (x(t), \hat{x}(t); A(t))$ is a (\mathcal{D}_R -horizontal) geodesic of $(T^*(M) \otimes T(\hat{M}), \bar{g}_1^1)$. The proof is complete. \square

C The Rolling Problem Embedded in \mathbb{R}^N

In this section, we compare the rolling model defined by the state space $Q = Q(M, \hat{M})$, whose dynamics is governed by the conditions (12)-(13) (or, equivalently, by \mathcal{D}_R), with the rolling model of two n -dimensional manifolds embedded in \mathbb{R}^N as given in [27] (Appendix B). See also [8], [10].

Let us first fix $N \in \mathbb{N}$ and introduce some notations. The special Euclidean group of \mathbb{R}^N is the set $\text{SE}(N) := \text{SO}(N) \times \mathbb{R}^N$ equipped with the group operation \star given by

$$(p, A) \star (q, B) = (Aq + p, AB), \quad (p, A), (q, B) \in \text{SE}(N).$$

We identify $\text{SO}(N)$ with the subgroup $\{0\} \times \text{SO}(N)$ of $\text{SE}(N)$, while \mathbb{R}^N is identified with the normal subgroup $\mathbb{R}^N \times \{\text{id}_{\mathbb{R}^N}\}$ of $\text{SE}(N)$. With these identifications, the action \star of the subgroup $\text{SO}(N)$ on the normal subgroup \mathbb{R}^N is given by

$$(p, A) \star q = Aq + p, \quad (p, A) \in \text{SE}(N), \quad p \in \mathbb{R}^N.$$

Let \mathcal{M} and $\hat{\mathcal{M}} \subset \mathbb{R}^N$ be two (embedded) submanifolds of dimension n . For every $z \in \mathcal{M}$, we identify $T|_z\mathcal{M}$ with a subspace of \mathbb{R}^N (the same holding in the case of $\hat{\mathcal{M}}$) i.e., elements of $T|_z\mathcal{M}$ are derivatives $\dot{\sigma}(0)$ of curves $\sigma : I \rightarrow \mathcal{M}$ with $\sigma(0) = z$ ($I \ni 0$ a nontrivial real interval).

The *rolling of \mathcal{M} against $\hat{\mathcal{M}}$ without slipping or twisting* in the sense of [27] is realized by a smooth curves $G : I \rightarrow \text{SE}(N)$; $G(t) = (p(t), U(t))$ (I a nontrivial real interval) called the *rolling map* and $\sigma : I \rightarrow \mathcal{M}$ called the *development curve* such that the following conditions (1)-(3) hold for every $t \in I$:

- (1) (a) $\hat{\sigma}(t) := G(t) \star \sigma(t) \in \hat{\mathcal{M}}$ and
 (b) $T|_{\hat{\sigma}(t)}(G(t) \star \mathcal{M}) = T|_{\hat{\sigma}(t)}\hat{\mathcal{M}}$.
- (2) No-slip: $\dot{G}(t) \star \sigma(t) = 0$.
- (3) No-twist: (a) $\dot{U}(t)U(t)^{-1}T|_{\hat{\sigma}(t)}\hat{\mathcal{M}} \subset (T|_{\hat{\sigma}(t)}\hat{\mathcal{M}})^\perp$ (tangential no-twist),
 (b) $\dot{U}(t)U(t)^{-1}(T|_{\hat{\sigma}(t)}\hat{\mathcal{M}})^\perp \subset T|_{\hat{\sigma}(t)}\hat{\mathcal{M}}$ (normal no-twist).

The orthogonal complements are taken w.r.t. the Euclidean inner product of \mathbb{R}^N . In condition (2) we define the action ' \star ' of $\dot{G}(t) = (\dot{U}(t), \dot{p}(t))$ on \mathbb{R}^N by the same formula as for the action ' \star ' of $\text{SE}(N)$ on \mathbb{R}^N .

The two manifolds M and \hat{M} are embedded inside \mathbb{R}^N by embeddings $\iota : M \rightarrow \mathbb{R}^N$ and $\hat{\iota} : \hat{M} \rightarrow \mathbb{R}^N$ and their metrics g and \hat{g} are induced from the Euclidean metric s_N of \mathbb{R}^N i.e., $g = \iota^*s_N$ and $\hat{g} = \hat{\iota}^*s_N$. In the above setting, we take now $\mathcal{M} = \iota(M)$, $\hat{\mathcal{M}} = \hat{\iota}(\hat{M})$.

For $z \in \mathcal{M}$ and $\hat{z} \in \hat{\mathcal{M}}$, consider the linear orthogonal projections

$$P^T : T|_z\mathbb{R}^N \rightarrow T|_z\mathcal{M} \text{ and } P^\perp : T|_z\mathbb{R}^N \rightarrow T|_z\mathcal{M}^\perp,$$

and

$$\hat{P}^T : T|_{\hat{z}}\mathbb{R}^N \rightarrow T|_{\hat{z}}\hat{\mathcal{M}} \text{ and } \hat{P}^\perp : T|_{\hat{z}}\mathbb{R}^N \rightarrow T|_{\hat{z}}\hat{\mathcal{M}}^\perp,$$

respectively.

For $X \in T|_z\mathbb{R}^N$ and $Y \in \Gamma(\pi_{T\mathbb{R}^N}|_{\mathcal{M}})$ (here $\pi_{T\mathbb{R}^N}|_{\mathcal{M}}$ is the pull-back bundle of $T\mathbb{R}^N$ over \mathcal{M}), we use $\nabla_X^\perp Y$ to denote $P^\perp(\nabla_X^{s_N} Y)$ and one proceeds similarly

$\hat{\nabla}_{\hat{X}}^\perp \hat{Y} = \hat{P}^\perp(\nabla_{\hat{X}}^{s_N} \hat{Y})$ for $\hat{X} \in T|_{\hat{z}} \mathbb{R}^N$ and $Y \in \Gamma(\pi_{T\mathbb{R}^N}|_{\hat{\mathcal{M}}})$. We notice that, for any $z \in \mathcal{M}$, $X \in T|_z \mathcal{M}$ and $Y \in \text{VF}(\mathcal{M})$, we have

$$\nabla_X^{s_N} Y = \iota_*(\nabla_{\iota_*^{-1}(X)} \iota_*^{-1}(Y)) + \nabla_X^\perp Y,$$

and similarly on $\hat{\mathcal{M}}$.

Notice that ∇^\perp and $\hat{\nabla}^\perp$ determine (by restriction) connections of vector bundles $\pi_{T\mathcal{M}^\perp} : T\mathcal{M}^\perp \rightarrow \mathcal{M}$ and $\pi_{T\hat{\mathcal{M}}^\perp} : T\hat{\mathcal{M}}^\perp \rightarrow \hat{\mathcal{M}}$. These connections can then be used in an obvious way to determine a connection $\bar{\nabla}^\perp$ on the vector bundle

$$\pi_{(T\mathcal{M}^\perp)^* \otimes T\mathcal{M}^\perp} : (T\mathcal{M}^\perp)^* \otimes T\mathcal{M}^\perp \rightarrow \mathcal{M} \times \hat{\mathcal{M}}.$$

Let us take any rolling map $G : I \rightarrow \text{SE}(N)$, $G(t) = (p(t), U(t))$ and development curve $\sigma : I \rightarrow \mathcal{M}$ and define $x = \iota^{-1} \circ \sigma$. We will go through the meaning of each of the above conditions (1)-(3).

- (1) (a) Since $\hat{\sigma}(t) \in \hat{\mathcal{M}}$, we may define a smooth curve $\hat{x} := \hat{\iota}^{-1} \circ \hat{\sigma}$.
(b) One easily sees that

$$U(t)T|_{\hat{\sigma}(t)} \mathcal{M} = T|_{\hat{\sigma}(t)}(G(t) \star \mathcal{M}) = T|_{\hat{\sigma}(t)} \hat{\mathcal{M}}.$$

Thus $A(t) := \hat{\iota}_*^{-1} \circ U(t) \circ \iota_*|_{T|_{x(t)} \mathcal{M}}$ defines a map $T|_{x(t)} \mathcal{M} \rightarrow T|_{\hat{x}(t)} \hat{\mathcal{M}}$, which is also orthogonal i.e., $A(t) \in Q|_{(x(t), \hat{x}(t))}$ for all t . Moreover, if $B(t) := U(t)|_{T|_{\sigma(t)} \mathcal{M}^\perp}$, then $B(t)$ is a map $T|_{\sigma(t)} \mathcal{M}^\perp \rightarrow T|_{\hat{\sigma}(t)} \hat{\mathcal{M}}^\perp$ and, by a slight abuse of notation, we can write $U(t) = A(t) \oplus B(t)$.

Thus Condition (1) just determines a smooth curve $t \mapsto (x(t), \hat{x}(t); A(t))$ inside the state space $Q = Q(M, \hat{M})$.

- (2) We compute

$$\begin{aligned} 0 &= \dot{G}(t) \star \sigma(t) = \dot{U}(t)\sigma(t) + \dot{p}(t) \\ &= \frac{d}{dt}(G(t) \star \sigma(t)) - U(t)\dot{\sigma}(t) = \dot{\hat{\sigma}}(t) - U(t) \circ \iota_* \circ \iota_*^{-1} \circ \dot{\sigma}(t), \end{aligned}$$

which, once composed with $\hat{\iota}_*^{-1}$ from the left, gives $0 = \dot{\hat{x}}(t) - A(t)\dot{x}(t)$. This is exactly the no-slip condition, Eq. (13).

- (3) Notice that, on $\mathbb{R}^N \times \mathbb{R}^N = \mathbb{R}^{2N}$, the sum metric $s_N \oplus s_N$ is just s_{2N} . Moreover, if $\gamma : I \rightarrow \mathbb{R}^N$ is a smooth curve, then smooth vector fields $X : I \rightarrow T(\mathbb{R}^N)$ along γ can be identified with smooth maps $X : I \rightarrow \mathbb{R}^N$ and with this observation one has: $\dot{X}(t) = \nabla_{\dot{\gamma}(t)}^{s_N} X$.

- (a) Since $U(t) = A(t) \oplus B(t)$, we get, for $t \mapsto \hat{X}(t) \in T|_{\hat{\sigma}(t)} \hat{\mathcal{M}}$, that

$$\begin{aligned} \dot{U}(t)U(t)^{-1}\hat{X}(t) &= \nabla_{(\hat{\sigma}, \hat{\sigma})(t)}^{s_{2N}} \hat{X}(\cdot) - U(t)\nabla_{(\hat{\sigma}, \hat{\sigma})(t)}^{s_{2N}} (U(\cdot)^{-1}\hat{X}(\cdot)) \\ &= P^T(\hat{\nabla}_{\hat{\sigma}(t)}^{s_N} \hat{X}(\cdot)) + \hat{\nabla}_{\hat{\sigma}(t)}^\perp \hat{X}(\cdot) \\ &\quad - U(t)(P^T(\nabla_{\hat{\sigma}(t)}^{s_N} (A(\cdot)^{-1}\hat{X}(\cdot))) + \nabla_{\hat{\sigma}(t)}^\perp (A(\cdot)^{-1}\hat{X}(\cdot))) \\ &= (\bar{\nabla}_{(\hat{x}, \hat{x})(t)} A(\cdot))A(t)^{-1}(\hat{\iota}_*^{-1}\hat{X}(t)) + (\hat{\nabla}_{\hat{\sigma}(t)}^\perp \hat{X}(\cdot) - B(t)\nabla_{\hat{\sigma}(t)}^\perp (A(\cdot)^{-1}\hat{X}(\cdot))), \end{aligned}$$

from which it is clear that the tangential no-twist condition corresponds to the condition that $\bar{\nabla}_{(\hat{x}(t), \dot{\hat{x}}(t))} A(\cdot) = 0$. This means exactly that $t \mapsto (x(t), \hat{x}(t); A(t))$ is tangent to \mathcal{D}_{NS} for all $t \in I$. Thus, the tangential no-twist condition (3)-(a) is equivalent to the no-spinning condition, Eq. (11).

(b) Choose $t \mapsto \hat{X}^\perp(t) \in T|_{\hat{\sigma}(t)} \hat{\mathcal{M}}^\perp$ and calculate as above

$$\begin{aligned} \dot{U}(t)U(t)^{-1}\hat{X}^\perp(t) &= P^T(\nabla_{\dot{\hat{\sigma}}(t)}^{s_N} \hat{X}^\perp(\cdot)) + \hat{\nabla}^\perp \dot{\hat{\sigma}}(t) \\ &\quad - U(t)(P^T(\nabla_{\dot{\sigma}(t)}^{s_N} (B(\cdot)^{-1}\hat{X}(\cdot))) + \nabla_{\dot{\sigma}(t)}^\perp (B(\cdot)^{-1}\hat{X}(\cdot))) \\ &= \left(P^T(\nabla_{\dot{\hat{\sigma}}(t)}^{s_N} \hat{X}^\perp(\cdot) - A(t)P^T(\nabla_{\dot{\sigma}(t)}^{s_N} (B(\cdot)^{-1}\hat{X}(\cdot))) \right) \\ &\quad + (\bar{\nabla}_{(\hat{\sigma}(t), \dot{\hat{\sigma}}(t))}^\perp B(\cdot))B(t)^{-1}\hat{X}(t), \end{aligned}$$

and hence we see that the normal no-twist condition (3)-(b) corresponds to the condition that

$$\bar{\nabla}_{(\sigma(t), \dot{\sigma}(t))}^\perp B(\cdot) = 0, \quad \forall t.$$

In a similar spirit to how Definition 3.13 was given, one easily sees that this condition just amounts to say that B maps parallel translated normal vectors to \mathcal{M} to parallel translated normal vectors to $\hat{\mathcal{M}}$. More precisely, if $X_0 \in T\mathcal{M}^\perp$ and $X(t) = (P^{\nabla^\perp})_0^t(\sigma)X_0$ is a parallel translate of X_0 along σ w.r.t. to the connection ∇^\perp (notice that $X(t) \in T|_{\sigma(t)}\mathcal{M}^\perp$ for all t), then the normal no-twist condition (3)-(b) requires that $t \mapsto B(t)X(t)$ (which is the same as $U(t)X(t)$) is parallel to $t \mapsto \hat{\sigma}(t)$ w.r.t the connection $\hat{\nabla}^\perp$ i.e., for all t ,

$$B(t)((P^{\nabla^\perp})_0^t(\sigma)X_0) = (P^{\hat{\nabla}^\perp})_0^t(\hat{\sigma})(B(0)X_0).$$

We formulate the preceding remarks to a proposition.

Proposition C.1 Let $\iota : M \rightarrow \mathbb{R}^N$ and $\hat{\iota} : \hat{M} \rightarrow \mathbb{R}^N$ be smooth embeddings and let $g = \iota^*(s_N)$ and $\hat{g} = \hat{\iota}^*(s_N)$. Fix points $x_0 \in M$, $\hat{x}_0 \in \hat{M}$ and an element $B_0 \in \text{SO}(T|_{\iota(x_0)}\mathcal{M}^\perp, T|_{\hat{\iota}(\hat{x}_0)}\hat{\mathcal{M}}^\perp)$. Then, there is a bijective correspondence between the smooth curves $t \mapsto (x(t), \hat{x}(t); A(t))$ of Q tangent to \mathcal{D}_{NS} (resp. \mathcal{D}_{R}), satisfying $(x(0), \hat{x}(0)) = (x_0, \hat{x}_0)$ and the pairs of smooth curves $t \mapsto G(t) = (p(t), U(t))$ of $\text{SE}(N)$ and $t \mapsto \sigma(t)$ of \mathcal{M} which satisfy the conditions (1), (3) (resp. (1),(2),(3) i.e., rolling maps) and $U(0)|_{T|_{\sigma(0)}\mathcal{M}^\perp} = B_0$.

Proof. Let $t \mapsto q(t) = (x(t), \hat{x}(t); A(t))$ to be a smooth curve in Q such that $(x(0), \hat{x}(0)) = (x_0, \hat{x}_0)$. Denote $\sigma = \iota \circ x$, $\hat{\sigma} = \hat{\iota} \circ \hat{x}$ and let $B(t) = (P^{\bar{\nabla}^\perp})_0^t((\sigma, \hat{\sigma}))B_0$ be the parallel translate of B_0 along $t \mapsto (\sigma(t), \hat{\sigma}(t))$ w.r.t the connection $\bar{\nabla}^\perp$. We define

$$U(t) := (\hat{\iota}_* \circ A(t) \circ \iota_*^{-1}) \oplus B(t) : T|_{\sigma(t)}\mathcal{M} \rightarrow T|_{\hat{\sigma}(t)}\hat{\mathcal{M}},$$

and $p(t) = \hat{\sigma}(t) - U(t)\sigma(t)$. Then, by the above remarks, the smooth curve $t \mapsto G(t) = (p(t), U(t))$ satisfies Conditions (1),(3) (resp. (1),(2),(3)) if $t \mapsto q(t)$ is tangent to \mathcal{D}_{NS} (resp. \mathcal{D}_{R}). This clearly gives the claimed bijective correspondence. \square

References

- [1] Alouges, F., Chitour Y., Long, R. *A motion planning algorithm for the rolling-body problem*, accepted for publication in IEEE Trans. on Robotics.
- [2] Agrachev A., Sachkov Y., *An Intrinsic Approach to the Control of Rolling Bodies*, Proceedings of the Conference on Decision and Control, Phoenix, 1999, pp. 431 - 435, vol.1.
- [3] Agrachev, A., Sachkov, Y., *Control Theory from the Geometric Viewpoint*, Encyclopaedia of Mathematical Sciences, 87. Control Theory and Optimization, II. Springer-Verlag, Berlin, 2004.
- [4] Blumenthal, R., Hebda, J., *The Generalized Cartan-Ambrose-Hicks Theorem*, Geom. Dedicata 29 (1989), no. 2, 163–175.
- [5] Bryant, R. and Hsu, L., *Rigidity of integral curves of rank 2 distributions*, Invent. Math. 114 (1993), no. 2, 435–461.
- [6] Bor, G., Montgomery, R., *G_2 and the “Rolling Distribution”*, arXiv:math/0612469, 2006.
- [7] Chelouah, A. and Chitour, Y., *On the controllability and trajectories generation of rolling surfaces*. Forum Math. 15 (2003) 727-758.
- [8] Molina, M., Grong, E., Markina, I., Leite, F., *An intrinsic formulation of the rolling manifolds problem*, arXiv:1008.1856, 2010.
- [9] Helgason, S., *Differential Geometry, Lie Groups, and Symmetric Spaces*, Pure and Applied Mathematics, 80. Academic Press, Inc., New York-London, 1978.
- [10] Hüper, K., Silva Leite, F. *On the Geometry of Rolling and Interpolation Curves on S^n , SO_n , and Grassmannian Manifolds*, Journal of Dynamical and Control Systems, Vol 13, No 4., 2007.
- [11] Joyce, D.D., *Riemannian Holonomy Groups and Calibrated Geometry*, Oxford University Press, 2007.
- [12] Jurdjevic, V. *Geometric control theory*, Cambridge Studies in Advanced Mathematics, 52. Cambridge University Press, Cambridge, 1997.
- [13] Kobayashi, S., Nomizu, K., *Foundations of Differential Geometry, Vol. I*, Wiley-Interscience, 1996.
- [14] Kolář, I., Michor, P., Slovák, J. *Natural operations in differential geometry*, Springer-Verlag, Berlin, 1993.
- [15] Lee, J., *Introduction to smooth manifolds*, Graduate Texts in Mathematics, 218. Springer-Verlag, New York, 2003.
- [16] Marigo, A. and Bicchi A., *Rolling bodies with regular surface: controllability theory and applications*, IEEE Trans. Automat. Control 45 (2000), no. 9, 1586–1599.

- [17] Marigo, A. and Bicchi A., *Planning motions of polyhedral parts by rolling*, Algorithmic foundations of robotics. *Algorithmica* 26 (2000), no. 3-4, 560–576.
- [18] Montgomery, D., Samelson, H., *Transformation Groups of Spheres*, The Annals of Mathematics, Second Series, Vol. 44, No. 3 (Jul., 1943), pp. 454-470.
- [19] Montgomery, R., *A Tour of Subriemannian Geometries, Their Geodesics and Applications*, American Mathematical Society, 2006.
- [20] Murray, R., Li, Z. and Sastry, S. *A mathematical introduction to robotic manipulation*, CRC Press, Boca Raton, FL, 1994.
- [21] Obata, M., *On Subgroups of the Orthogonal Group*, Transactions of the American Mathematical Society, Vol. 87, No. 2 (Mar., 1958), pp. 347-358.
- [22] Ozeki, H., *Infinitesimal holonomy groups of bundle connections*, Nagoya Math. J. 10 (1956), 105–123.
- [23] Pawel, K., Reckziegel, H., *Affine Submanifolds and the Theorem of Cartan-Ambrose-Hicks*, Kodai Math. J, 2002.
- [24] Petersen, P., *Riemannian Geometry*, Second edition. Graduate Texts in Mathematics, 171. Springer, New York, 2006.
- [25] Sakai, T., *Riemannian Geometry*, Translations of Mathematical Monographs, 149. American Mathematical Society, Providence, RI, 1996.
- [26] Sasaki, S., *On Differential Geometry of Tangent Bundles of Riemannian Manifolds*, Tôhoku Math. J. (2), 10, 1958, 338–354
- [27] Sharpe, R.W., *Differential Geometry: Cartan's Generalization of Klein's Erlangen Program*, Graduate Texts in Mathematics, 166. Springer-Verlag, New York, 1997.
- [28] Spivak, M., *A Comprehensive Introduction to Differential Geometry*, Vol. 2, Publish or Perish, 3rd edition, 1999.
- [29] Warner, F.W., *Foundations of Differentiable Manifolds*, Graduate Texts in Mathematics, 94. Springer-Verlag, New York-Berlin, 1983.